
Geology:

Extraterrestrial soils



Extraterrestrial soils may be defined as any of the solid granular crustal features of planets and moon other than those soils on planet Earth. Despite four decades of space exploration, which greatly expanded our understanding of the <u>Solar System</u>, there is considerable debate as to the loose covers of rocky planets and moons are <u>soils</u> in a pedological sense [11][1,2]. On Earth, <u>soil</u> form thanks to the combined action of at least five factors: parent rock, <u>climate</u>, <u>topography</u>, living <u>organisms</u> and time.

A few other factors can concur to drive pedogenesis[3]. However, the necessity of biota as unavoidable soil forming factor is debated. In fact, important parts of Earth, such as the hyperarid <u>Atacama Desert</u> of Chile and the Dry Valleys of <u>Antarctica</u>, host virtually life-free soils with advanced horizonation. Actually, although most people invokes the ability to support plant growth in its <u>natural environment</u> as condicio sine qua non for soil, a scientific definition considers soil to be any in situ weathered veneer of a planetary surface that retains information on its climatic and geochemical history. A current or past mineral weathering is hence the pivotal requisite for soil.

Table of Contents

1 Weathering processes 2 Within the Earth's solar system 3 Summary 4 See also 5 References 6 Further Readings 7 External links

Weathering processes

On Earth, weathering is promoted by liquid water, which is the solvent where most of reactions happen and a carrier of matter and <u>energy</u>. <u>Oxygen</u> or, in anaerobic environments, weaker electron acceptor such as nitrates, sulphates and ferric <u>iron</u> allow the altering action of proton donors. Several sources of energy are finally the motor of pedogenesis. Outside Earth most water probably occurs as ice, but weathering might be caused by thin layers of liquid water at the rock-ice interface, as documented to happen in frozen soils on Earth[4]. Furthermore, there are several polar solvents able to replace liquid water, such as sulphuric, hydrofluoric and hydrocyanic acids, <u>ammonia</u>, methanol and hydrazine[5]. Finally, a variety of energy sources can drive chemical reactions in space: thermal, osmotic and ionic gradients, solar wind, magnetosphere energy and radioactivity are the most important among those detected. What soil genesis cannot prescind from is a rocky parent material, which in the Solar System occurs just on the other three inner planets – Mercury, Venus and Mars –, a few planets' moons, and the largest asteroids. Any consideration on the occurrence of soils on exoplanets – those orbiting other stars – some of which are expected to be solid, is nonsense given

their remoteness that do not allow direct observation.

Within the Earth's solar system

Mercury, which is devoid of any atmosphere, is greatly affected by meteorite and solar bombardment and shows evidences of maturation[6]. Venus, on the contrary, has a dense atmosphere, mainly composed of carbon dioxide and sulphuric acid droplets, which protects the surface from erosion by cosmic particles but is effective at degrading rocks into secondary weathering products[7]. The recent direct investigations of three rovers on Mars, one of which in 2008 even opened the first soil profile outside Earth (Fig. 1), clearly demonstrated that chemical weathering combined with leaching occur in many geological settings of this planet[8]. The Earth's moon is the only celestial body samples of which were returned and studied on Earth. Lunar surface shows a thin rim of weathering on grains and even horizonation induced by meteorite impacts, chemical interactions of impactgenerated and sputtered ions, and irradiation by solar wind, which darkens and reddens powders in proportion to their Fe content[9]. Additionally, opposite to what believed until a little time ago, a minor hydrated phase occurs in the upper millimetres of lunar surface[10]. The largely prevailing icy nature - as in the cases of Phobos and Deimos, the two small moons of Mars, or Titan, the largest moon of Saturn – or volcanism that continuously reworks the crust, thus reducing the time available for any pedogenic processes – as in the case of Io, the only telluric moon of Jupiter – make the presence of soils on moons improbable, although not yet impossible. Asteroids are rocky bodies that mainly lie in a belt between Mars and Jupiter and that are believed to be left over from the beginning of the Solar System. They have round or irregular shapes up to several 100 km across, but often are much smaller. Such a small size deprives them of any internal heat and, as a consequence, volcanism and tectonics that could rejuvenate their surface. However, too little is known of the Asteroids' nature to make deductions about the occurrence of soils on them. After capturing samples from the asteroid 25143 Itokawa, the Japanese spacecraft Hayabusa is returning to the Earth and will soon provide the first direct information on an asteroid's skin and its stage of alteration.

Summary

To summarise, our nearest planetary neighbours, Venus, Mars and our moon, possess weathered mantles that should be considered to be soils in a pedological sense, while Mercury and some large asteroids have loose portions of surface that only a future better knowledge may eventually grant of the rank of soils.

See also

- Extremophile
- Moon shrinkage

References

- 1. Markevitz, D. (1997) Soil without life? Nature 389, 435.
- 2. Banin, A. (2005) The Enigma of the Martian Soil. Science 309, 888-890.
- 3. Certini, G., Scalenghe, R. (2006) Soil formation on Earth and beyond: the role of additional soil forming factors. In: G. Certini, R. Scalenghe (eds.), Soils. Basic Concepts and Future Challenges. Cambridge University Press, Cambridge, UK, pp. 193-210.
- 4. Ugolini, F.C., Anderson, D.M. (1973) Ionic migration and weathering in frozen Antarctic soils. Soil Science 115, 461-470.
- 5. Schulze-Makuch, D., Irwin, L.N. (2004). Life in the Universe: Expectations and Constraints. Springer, Berlin.
- 6. Robinson, M.S. et 12 al., (2008). Reflectance and color variations on Mercury: regolith

processes and compositional heterogeneity. Science 321, 66-69.

- Barsukov, V. L., Borunov, S. P., Volkov, V. P., Zolotov, M. Yu., Sidorov, Yu. I., Khodakovsky, I. L., 1986. Mineral composition of Venus soil at Venera 13, Venera 14 and Vega 2 landing sites: thermodynamic prediction. Lunar and Planetary Science 17, 28-29.
- 8. Amundson, R., Ewing, S., Dietrich, W., Setter, B., Owen, J., Chadwick, O., et al. (2008). On the in situ aqueous alteration of soils on Mars. Geochimica et Cosmochimica Acta 72, 3845-3864.
- Taylor, L.A., Pieters, C., Patchen, A., Taylor, D.H.S., Morris, R.V., Keller, L.P., Mckay, D.S. (2010) Mineralogical and chemical characterization of lunar highland soils: Insights into the space weathering of soils on airless bodies. Journal of Geophysical Research-Planets 115, E02002.
- 10. Pieters, C. M. et 28 al. (2009) Character and spatial distribution of OH/H2O on the surface of the Moon seen by M 3 on Chandrayaan-1. Science 326, 568-572.

Further Readings

- Certini, G., Scalenghe, R., Amundson, R. (2009) A view of extraterrestrial soils. European Journal of Soil Science 60: 1078-1092.
- Baker, V.R. (2008) Planetary landscape systems: A limitless frontier. Earth Surface Processes and Landforms 33, 1341-1353.
- Matson, D.L., Johnson, T.V., Veeder, G.J. (1977) Soil maturity and planetary regoliths: The Moon, Mercury, and the asteroids. pp. 1001-011. Proceedings of the 8th Lunar Science Conference, Houston, TX USA.
- Sanderson, K. (2007) Alien Earth. Nature 445, 10-11.

External links

• www.nasa.gov/mission_pages/exploration/news/presskits/living_on_the_moon.html

Fig. 1. Centimetric soil profiles at 68°N in the Martian arctic plain URL www.nasa.gov/mission_pages/phoenix/images/phx-17062.html [Courtesy of NASA/JPL-Caltech/University of Arizona/Texas A&M University/Max Planck Institute]

Citation

<u>Giacomo Certini, Riccardo Scalenghe</u> (Lead Author);<u>John Shroder</u>, <u>C Michael Hogan</u> (Topic Editor) "Extraterrestrial soils". In: Encyclopedia of Earth. Eds. Cutler J. Cleveland (Washington, D.C.: Environmental Information Coalition, National Council for Science and the Environment). [First published in the Encyclopedia of Earth April 15, 2010; Last revised Date December 16, 2011; Retrieved November 16, 2012 <<u>http://www.eoearth.org/article/Extraterrestrial_soils?topic=49478</u>>

The Authors



Dr. Giacomo Certini is a Tenured Researcher, chair of Pedology, at the College of Agriculture and Forestry, University of Florence. His main scientific interests are: forest pedology,

influence of plants on pedogenesis, mineral weathering in soil, carbon sequestration in forest soils, effects of forest fires on soil properties. He is author/co-author of about 30 papers published in International journals (ISI). He has co-edited the book Soils. Basic Concepts and Future Challenges, contributed by ... (Full Bio)



Riccardo Scalenghe was born in Torino, Italy on May 07, 1965. He received his PhD in Pisa, Italy in 1996. He teaches Land Evaluation and Soil Geography, and is the author or co-author of 68 scientific papers. Some publications: Scalenghe R., Zanini E. and Nielsen D.R. (2000) Modeling soil development in a post-incisive chronosequence. Soil. Sci., 165:455-462. Scalenghe R., Edwards A.C., Ajmone Marsan F., and Barberis E. (2002) The effect of reducing conditions on P solubility for a diverse rang ... (Full Bio)

0 Comments

Add Comment



You must be logged in to post a comment. Click here to login

<u>NCSE</u> Boston University <u>Trunity</u>

Unless otherwise noted, all text is available under the terms of the <u>Creative Commons Attribution-Share Alike license</u>. <u>Privacy Policy</u> | <u>Neutrality Policy</u> Supported by the <u>Environmental Information Coalition</u> and the <u>National Council for Science and the</u> Environment.