

Original Article

Euromediterranean Biomedical Journal
for young doctors
(formerly: Capsula Eburnea)

ANATOMICAL VARIATIONS OF THE INTERNAL JUGULAR VEIN: THE ROLE OF ULTRASONOGRAPHY

Adriana Lena, Francesco Raffaele, Giorgio Lo Iacono,
Girolamo Geraci, Massimo Cajozzo

SUMMARY

Purpose: In many places, especially in emergency department, central venous catheter is still inserted using anatomical landmark guidance with a success rate up to 97.6% and complications up to 15%. This study was aimed to determine by the support of ultrasonography (US) the anatomical variations of the internal jugular vein (IJV) in relation with other structures of the neck, such as the common carotid artery (CCA).

Material and Methods: 830 patients requiring central vein catheterization (CVC) were included in the analysis. The position of the IJV in relation to the other structures of the neck was demonstrated by portable ultrasonography.

Results: The mean diameter of IJV was 10.3 mm in right and 10.5 mm in left side of neck, in male ($p > 0.05$) and 9.1 mm in right and 10.5 mm in left side of neck, in female ($p > 0.05$). The mean distance between IJV and CCA was 1.9 mm in right and 1.7 mm in left side of neck in male, and 2.0 mm in right and 2.2 mm in left side of neck in female. The mean distance of IJV from the skin surface was 9.8 mm in right and 10.0 mm in left side of neck in male, and 12.1 mm in right and 12.5 mm in left side of neck in female. On 25.54 % we observed variations of internal jugular vein site. On 3.97 % we observed a small caliber of internal jugular vein that could complicate the catheterization of the vein. On 1.8 % was diagnosed a thrombus of internal jugular vein, that is considered as an absolute contraindication for a CVC.

Conclusion: Different patients had anatomical variations that are important and should be knowned, in order to reduce the possibility of severe complications.

Introduction

Central venous catheterization (CVC) is a common procedure used for long term central intravenous therapy, central venous pressure monitoring, dialysis and apheresis (1-3).

The most commonly used technique is percutaneous catheterisation of the internal jugular vein (IJV) which can be found by the identification of an anatomical landmark (Sedillot triangle, formed by medial and lateral heads of the sternocleido-mastoid muscle and base by medial end of the clavicle) (Figure 1), or by Ultrasonography scanning.

The aim of the present paper is to determine the rate of anatomical variations of IJV, in relation with the other structures of the neck, particularly the CCA (4-7), in a cohort of patients who underwent US-guided CVC, in order to underline the importance of anatomical knowledge to reduce the complications related to this procedure, such as difficult or

Address of the authors

Department of General Surgery, University of Palermo

Send correspondence to: Adriana Lena, adrylena86@gmail.com

Received: September 12th, 2014 — Revised: November 29th, 2014 — Accepted: December 21th 2014

failed cannulation, arterial puncture and hematoma, and hemo/pneumothorax (8).

Material and Methods

Between January 1998 and March 2013, in our Unit of General and Thoracic Surgery, we performed US-guided CVC on 830 patients affected by different diseases.

The patients were 456 male and 374 female, the age ranged between 20 and 75 years (mean: 55y.o.).

The IJV and its relationship with other anatomical structures of the neck was identified by a "Prisma-Diasonics" ultrasound instrument with high frequency linear transducer (10 Mhz).

Each patient was placed supine with slight Trendelenburg position, with a 30° neck rotation on contra-lateral side of the procedure. The physician was standing in the same side of the region where to place the catheter. We performed transversal and longitudinal scans on both sides of the neck for each patient, at the Sedillot triangle, at the level of the base of the neck, and we recorded the location of IJV in relation to the CCA (Figure 2) (it was identified as lateral, antero-lateral, anterior, medial, and posterior) and the diameter of IJV and its distance from the skin. Small sized IJV was defined as a diameter ≤ 5 mm (9). During the catheterization the needle has




Figure 1: yellow and orange arrows: respectively medial and lateral heads of the sternocleidomastoid muscle; red arrow: clavicle; star: Sedillot triangle




Figure 2: relation between IJV and CA. US guided images

been inserted with an angle of between 30 and 45 degrees. Data were analyzed using Student's T test.

Results

Analyzing the 830 US-guided CVCs we observed:

- the mean diameter of IJV was 10.3 mm in right and 10.5 mm in left side of neck, in male ($p>0.05$) and 9.1 mm in right and

10.5 mm in left side of neck, in female ($p>0.05$);

- the mean distance between IJV and CCA was 1.9 mm in right and 1.7 mm in left side of neck in male ($p>0.05$), and 2.0 mm in right and 2.2 mm in left side of neck in female ($p>0.05$) (figure 3);

- the mean distance of IJV from the skin surface was 9.8 mm in right and 10.0 mm




Figure 3: Data about distance between IJV and CCA. Data are expressed in mm




Figure 4: Data about depth of IJV from the skin surface. Data are expressed in mm

in left side of neck in male ($p>0.05$), and 12.1 mm in right and 12.5 mm in left side of neck in female ($p>0.05$) (figure 4);

On 212/830 patients (25.54 %) we observed variations of internal jugular vein site: medialization of the vein 20%, antero-medialization 80% (Figures 5A, 5B; Table 1).

On 33/830 cases (3.97 %) we observed a small caliber of internal jugular vein that could complicate the catheterization (table 2).

On 15/830 cases (1.8 %) was diagnosed a thrombus of internal jugular vein, that is considered as an absolute contraindication for a CVC.

On the remaining 570 (68.67 %) patients no variations of IJV were observed.

Discussion

Usually the IJV is the most superficial and lateral element of the neurovascular bundle (NVB) of the neck and collects the blood from the brain, the superficial part of the face and the neck. It runs down the side of the neck in a vertical direction, lying at first lateral to the internal carotid artery (ICA) and then lateral to the CCA (10). At the root of the neck, the IJV unites with the subclavian vein to form the brachiocephalic vein, direct to the right atrium. The IJV position is almost constant, and this is the reason why it is commonly used for CVC (11).

In previous paper published by our group (12-14) we recorded an average catheterization time of 4 minutes, with 98.7% of




Figure 5: (A) Left variations of the IJV in male and female; (B) Right variations of the IJV in male and female

success rate, 0% of major complications, 4.1% of minor complications such as vagal hypotension or catheter dislocation.

However, experienced operators can use different manoeuvres during the procedure to facilitate the access of the IJV, such as rotation of the head, extension of the neck, breath-holding (15), and identification of anatomic landmarks (16) especially in association with ultrasound guidance (17, 18)

In conclusion our study underline the potential problems of catheterization related to the anatomical variations of the IJV; for this reason the knowledge of the anatomy it is crucial for the correct and safe procedure (19-21). The use of ultrasonography it can be very helpful to improve the anatomical knowledge.

Therefore, according to The American Society of Anesthesiologists Task Force of 2012 (22), and based on our experience (23-27), we can confirm that US-guided procedure may avoid major complications (such as pneumothorax reported in literature) thanks to the precise localization of the anatomical structures, especially helpful in high-risk (27) patients.

References

1. Zimmermann JJ, Strauss RH. History and current application of intravenous therapy in children. *Pediatr Emerg Care* 1989; 5:120.
2. Scott WT, Bergamini MB. Long term venous access: indications and choice of site and catheter. *Seminars in vascular surgery* 1997; 10 (3): 130-134.
3. Newman KA, Reed WP, Schimpff SC, Bustamante CI, & Wade JC. Hickman catheters in association with intensive cancer chemotherapy. *Supportive care in Cancer* 1993; 1:92-97.
4. Denys BG, Uretsky BF, Reddy PS. Ultrasound assistant cannulation of the internal jugular vein. A prospective comparison to the external landmark guided technique. *Circulation*. May 1993; 87(5):1557-62.
5. Lim CL, Keshava SN, Lea M. Anatomical variations of the internal jugular veins and their relationship to the carotid arteries: a CT evaluation. *Australas Radiol*. 2006 Aug; 50 (4): 314-8.
6. Shoja MM, Ardalan MR, Tubbs RS, Loukas M, Vahedinia S, Jabbari R, Jalilvand M, Shakeri A. The relationship between the internal jugular vein and common carotid artery in the carotid sheath: the effects of age, gender and side. *An Anat*. 2008; 190 (4):339-43. Epub 2008 May 27.

	Male (n: 456)		Female (n: 374)	
	Left	Right	Left	Right
Lateral	401	395	325	327
Anterior	48	54	45	41
Posterior	3	5	2	4
Medial	4	2	2	2

Table 1: Data about position of IJV in relation to CCA

	Male			Female		
	Left	Right	Total	Left	Right	Total
Mean	10.5	10.3	10.4	9.5	9.1	9.3
± SD	1.22	1.26	1.24	1.24	1.18	1.21

Table 2: Data about diameter of IJV. Data are expressed in mm

7. Turba UC, Uflacker R, Hannegan C, Selby JB. Anatomic relationship of the internal jugular vein and the common carotid artery applied to percutaneous transjugular procedures. *Cardiovasc Intervent Radiol*. 2005 May-Jun; 28(3): 303-6.
8. Roth B, Marciniak B, Engelhardt T, Bissonnette B. Anatomic relationship between the internal jugular vein and the carotid artery in preschool children – an ultrasonographic study. *Paediatr Anaesth*. 2008 Aug; 18(8):752-6.
9. Mey U, Glasmacher A, Hahn C, et al. Evaluation of an ultrasound-guided technique for central venous access via the internal jugular vein in 493 patients. *Support Care Cancer* 2003; 11:148-55.
10. Caridi JG, Hawkins IF Jr, Wiechmann BN, et al. (1998) Sonography guidance, when using the right internal jugular vein for central vein access. *AJR Am J Roentgenol* 133: 467-476.
11. Denys BG, Uretsky BF, Reddy PS. Ultrasound assistant cannulation of the internal jugular vein. A prospective comparison to the external landmark guided technique. *Circulation*. May 1993; 87(5): 1557-62.
12. M. Cajozzo, G. Cocchiara, G. Grecom R. Vaglica, G. Pezzano, G. Quintini, G. Modica. Comparison of Central Venous Catheterization with and without Ultrasound Guide. *Transfusion and Apheresis Science*. 2004; vol 31-3: 199-202.
13. Verghese ST, Patel RI, Hannallah RS. Central venous cannulation in infants and children. A comparison of internal and external jugular vein approaches. *Paediatr Anaesth*, 1993; 3:95-9.
14. M. Cajozzo, G. Cocchiara, G. Greco, R. Vaglica, E. Bartolotta, L. Platia, G. Modica. Ultrasound (U.S.) central venous catheterization of internal jugular vein on over 65 years old patients versus blind technique. *Journal of Surgical Oncology*. 2004 vol. 88, 4: 267-268.
15. Armstrong PJ, Sutherland R, Scott DH (1994) The effect of position and different manoeuvres on internal jugular vein diameter size. *Acta Anaesthesiologica Scand* 38:229-231.
16. Heath KJ, Woulfe J, Lownie S, et al. (1998) A devastating complication of inadvertent carotid artery puncture. *Anesthesiology* 89:1273-1275.
17. Mitchell SE, Clark RA (1979) Complications of central venous catheterization. *AJR Am J Roentgenol* 133:467-476.
18. Docktor B, So CB, Saliken JC, et al. (1996) Ultrasound monitoring in cannulation of the internal jugular vein: Anatomic and technical considerations. *Can Assoc Radiol J* 47: 195-201.
19. Caridi JG, Hawkins IF Jr, Wiechmann BN, et al. (1998) Sonography guidance, when using the right internal jugular vein for central vein access. *AJR Am J Roentgenol* 133: 467-476.
20. Skolnick ML (1994) The role of sonography in the placement and management of jugular and subclavian central venous catheters. *AJR Am J Roentgenol* 163:291-295.
21. Conz PA, Dissegna D, Rodighiero MP, et al. (1997) Cannulation of the internal jugular vein: Comparison of the classic Seldinger Technique and an ultrasound guided method. *J Nephrol* 311-313.
22. Practice guidelines for central venous access – a report by the American society of Anesthesiologists task force on central venous access. *Anesthesiology* 2012; 116 (3): 539-73.
23. Parmar SB, Parikh SN, Mehta HK. Anatomical variations of the internal jugular vein in relation to carotid artery: An ultrasound study. *Int J Med Sci Public Health* 2013; 2:223-228.
24. Abraham E, Shapiro M, Podolsky S. Central venous catheterization in the emergency setting. *Crit. Care Med* 1983; 11:515-517.
25. Bolin GW, Anderson DJ, Anderson KC. Percutaneous central venous catheterization performed by medical house officers: a prospective study. *Cathet Cardiovasc Diagn* 1982; 8:23-29.
26. Sznajder JL, Zveibil FR, Bitterman H. Central vein catheterization. *Arch. Intern Med* 1986; 146:259-261.
- M. Napolitano, A. Malato, F. Raffaele, M. Palazzolo, G. Lo Iacono, R. Pinna, G. Geraci, G. Modica, G. Saccullo, S. Siragusa, M. Cajozzo. Ultrasonography-guided central venous catheterisation in haematological patients with severe thrombocytopenia. *Blood Transfus*. 2013 October; 11(4): 506-509.