

## [FRI0299] PREVALENCE OF VERTEBRAL FRACTURES IN OSTEOPOROTIC HIP FRACTURED PATIENTS: THE FOCUS STUDY

*G. Arioli<sup>1</sup>, P. Falaschi<sup>2</sup>, G. Letizia Mauro<sup>3</sup>, G. Iolascon<sup>4</sup>, M. Invernizzi<sup>5</sup>, M. Noale<sup>6</sup>, M. Manfredini<sup>1</sup>, G. Bernardi<sup>1</sup>, C. Cisari<sup>5</sup>. <sup>1</sup>Department of Neurosciences, Hospital "Carlo Poma", Mantova; <sup>2</sup>Geriatric Unit, "Sapienza" University of Rome, S. Andrea Hospital, Rome; <sup>3</sup>Department of Rehabilitation, University of Palermo, Palermo; <sup>4</sup>Department of Medical and Surgical Specialties and Dentistry, Second University of Naples, Naples; <sup>5</sup>Physical and Rehabilitative Medicine, Department of Health Sciences, University of Eastern Piedmont Novara; <sup>6</sup>Neuroscience Institute, Aging Branch, CNR, Padua, Italy*

**Background:** Osteoporosis is a multifactorial systemic skeletal disease, characterized by low bone mass and microarchitectural modifications of bone tissue, with a consequent increase in fragility fractures [1]. Vertebral fractures are the most prevalent osteoporotic fractures and osteoporotic hip fractures are the most serious complication of osteoporosis resulting in increased mortality and high socio-economic cost [2,3]. The coexistence of these two pathological conditions in elderly patients has been previously described, leading to even worse functional outcomes than each one alone [4].

**Objectives:** To determine the prevalence of vertebral fractures in osteoporotic hip fractured women and to evaluate the relationship between prevalence of vertebral fractures and pre-existing factors such as autonomy in daily life activity, quality of walking, numbers of falls, cognitive aspects and comorbidities.

**Methods:** 946 osteoporotic hip fractured women aged more than 60 years and with an X-ray evaluation of spine were consecutively enrolled in 25 Orthopaedic, Physical Medicine and geriatric centers in Italy. After spine X-ray morphometry patients were divided in two groups: previous vertebral fracture (F) and no previous fracture (NF). Moreover anamnestic, demographical and outcome related data (ADL, IADL, CIRS, SPMSQ, FAC and RANKIN scale) were collected.

**Results:** Prevalent vertebral fractures were present in 502 (54%) patients. 119 (13.7%) patients had at least one severe fracture. The F compared to NF group showed statistically significant worse scores regarding the pre hip fracture values of RANKIN, CIRS, SPMSQ, IADL and the overall number of falls ( $p < 0.001$ ). Moreover the F group showed statistically significant lower values of serum 25(OH)D than NF group ( $p < 0.001$ ).

**Conclusions:** Previous Vertebral fractures in hip fractured patients are a common issue and negatively influence several functional and cognitive outcome measures in these patients.

### References:

- NIH Consensus Development Panel on Osteoporosis Prevention, Diagnosis and Therapy. Osteoporosis prevention, diagnosis and therapy. *JAMA*. 2001;285:785–795
- Sattui SE, Saag KG. Fracture mortality: associations with epidemiology and osteoporosis treatment. *Nat Rev Endocrinol*. 2014 Oct;10(10):592–602.
- Svedbom A, Hernlund E, Ivergård M, Compston J, Cooper C, Stenmark J, McCloskey EV, Jönsson B, Kanis JA; EU Review Panel of IOF. Osteoporosis in the European Union: a compendium of country-specific reports. *Arch Osteoporos*. 2013;8(1-2):137.
- Gonnelli S, Caffarelli C, Maggi S et al. The assessment of vertebral fractures in elderly women with recent hip fractures: the BREAK Study. *Osteoporosis Int* 2013;24:1151–1159

**Disclosure of Interest:** None declared

**DOI:** 10.1136/annrheumdis-2015-eular.3834

**Citation:** *Ann Rheum Dis* 2015;74(Suppl2): 533