


REVIEW ARTICLE

Obstetrics

Increasing global temperatures threaten gains in maternal and newborn health in Africa: A review of impacts and an adaptation framework

Matthew F. Chersich¹  | Fiona Scorgie¹ | Veronique Filippi² | Stanley Luchters³ |
Climate Change and Heat-Health Study Group

¹Wits Reproductive Health and HIV Institute (Wits RHI), Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

²Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK

³Centre for Sexual Health and HIV/AIDS Research (CeSHHAR), Harare, Zimbabwe

Correspondence

Matthew F. Chersich, Wits RHI, Klein St & Esselen St, Hillbrow, Johannesburg, 2001, South Africa.

Email: mchersich@wrhi.ac.za

Abstract

Anatomical, physiologic, and socio-cultural changes during pregnancy and child-birth increase vulnerability of women and newborns to high ambient temperatures. Extreme heat can overwhelm thermoregulatory mechanisms in pregnant women, especially during labor, cause dehydration and endocrine dysfunction, and compromise placental function. Clinical sequelae include hypertensive disorders, gestational diabetes, preterm birth, and stillbirth. High ambient temperatures increase rates of infections, and affect health worker performance and healthcare seeking. Rising temperatures with climate change and limited resources heighten concerns. We propose an adaptation framework containing four prongs. First, behavioral changes such as reducing workloads during pregnancy and using low-cost water sprays. Second, health system interventions encompassing Early Warning Systems centered around existing community-based outreach; heat-health indicator tracking; water supplementation and monitoring for heat-related conditions during labor. Building modifications, passive and active cooling systems, and nature-based solutions can reduce temperatures in facilities. Lastly, structural interventions and climate financing are critical. The overall package of interventions, ideally selected following cost-effectiveness and thermal modeling trade-offs, needs to be co-designed and co-delivered with affected communities, and take advantage of existing maternal and child health platforms. Robust-applied research will set the stage for programs across Africa that target pregnant women. Adequate research and climate financing are now urgent.

KEYWORDS

Africa, climate change, heat, maternal and newborn health, pregnancy, reproductive health, temperature

A complete list of the CRIC Study Investigators can be found in the Appendix at the end of the article.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *International Journal of Gynecology & Obstetrics* published by John Wiley & Sons Ltd on behalf of International Federation of Gynecology and Obstetrics.

1 | BACKGROUND

“As a species, we are expert problem solvers. But we haven't yet applied ourselves to this problem [climate change] with the focus it requires.” Sir David Attenborough, 2019.

1.1 | Extreme heat in Africa

Africa is the continent hardest hit by climate change, has the fewest resources to respond,¹ is least responsible for the crisis, and to date has had limited success in securing adequate climate financing and reparations for losses and damages from climate change.² Heat-health burdens of disease are mounting as the average global temperature rises—it is already 1.2°C above pre-industrial levels, and a further degree or more warming appears inevitable.³ Yet, temperature increases are spread unevenly across the globe. Increases in many parts of Africa are well above the global average.² Heatwaves are becoming more frequent, intense, and longer lasting, and heat-wave conditions will occur on at least 50% of days during the warm season in many cities near the Sahel by 2100, assuming the current trajectory of emissions continues.⁴ Already many of the temperatures being recorded in Africa are close to the limits of “liveability”, and physical labor or “workability” is not possible for parts of the year.⁵ Studies in Burkina Faso,⁶ Ghana,⁷ Kenya,⁸ South Africa,⁹ and Tanzania¹⁰ have shown considerable spikes in overall population-level mortality on days with higher than average temperatures. Additionally, the number of emergency department visits increased at higher temperatures in Botswana,¹¹ as did hospital admissions in Uganda,¹² and diarrheal rates in children under 5 years of age in South Africa.^{7,9} Studies in Ethiopia, South Africa, and Uganda have documented heat impacts on pregnancy outcomes.¹³⁻¹⁵

Many women in Africa have little or no protection against exposure to extreme heat events during pregnancy. Rapid urbanization on the continent means that increasing numbers of pregnant women reside in informal housing in urban heat islands, where temperatures can be several degrees higher than in surrounding areas. Health facilities, poorly built brick dwellings, and informal housing are frequently 4°C–6°C warmer indoors than outdoors.^{16,17} Additionally, in many rural areas in Africa, pregnant women have little or no means of reducing heat exposure, and even drinking water may be scarce or non-potable.¹⁸ Levels of knowledge about the harmful effects of high ambient temperatures are low in Africa,¹⁹ and cultural, social, and religious practices such as outdoor church services, may increase exposure risks.

1.2 | Pathophysiologic processes and indirect impacts of high ambient temperatures on maternal and newborn health

Worldwide, around 200 studies have documented the negative impacts of exposure to high ambient temperatures on maternal and

newborn health.^{13,20,21} The threshold for harmful temperature exposure in pregnant women, however, varies by climate zone, health condition, and individual risk profiles, among other factors. Generally, relationships between temperature and adverse health outcomes are U- or J-shaped, with the base of the exposure-outcome curve commonly at relatively mild temperatures around 20°C, and the arms of the curves becoming exponentially steeper with each degree increase in temperature or additional day of a heatwave.¹³ Even at relatively mild temperatures, health consequences begin to mount and the harmful impacts of heat exposure are not restricted to heatwaves. The term “extreme heat” is used in this article to denote heat waves and “high ambient temperatures” to encapsulate the full range of temperature patterns that result in adverse health outcomes related to heat, either through direct or indirect pathways. These pathways and their pathophysiologic, infectious disease and health systems mechanisms are delineated below (Figure 1).

High ambient temperatures may overwhelm the capacity of maternal thermoregulatory mechanisms to dissipate heat in pregnancy and labor (Figure 1: Box 1). The physiologic and anatomical changes in pregnancy that challenge the maternal thermoregulation peak in advanced pregnancy and labor, including the considerable endogenous heat generated by fetal metabolism, strain from additional weight gain in pregnancy, fat deposits that retain heat, and the major exertion of labor and childbirth, which generates significant amounts of heat. This endogenous thermal load makes it difficult for pregnant or parturient women to maintain normothermia during periods of high ambient temperatures. We highlight four interlinked pathophysiologic processes that may explain the harmful impacts of heat when thermoregulatory capacity is exceeded during pregnancy or labor.

The first pathway involves elevated maternal body temperatures. In normal labor, maternal temperature increases by about 0.02°C per hour, but at a considerably faster pace among women who are primiparous,²² have prolonged labor,²³ or who are obese.²⁴ Additionally, maternal core temperature increase during fever from malaria or other infections, following epidural analgesia,²⁵ or if thermoregulatory mechanisms such as perspiration and hyperventilation cannot compensate for the additional burden of endogenous heat during periods with high ambient temperatures. Because fetal temperature is consistently 0.3°C–0.5°C higher than the maternal core temperature,²⁶ fetal temperatures can rise to dangerous levels if maternal temperatures increase. A systematic review summing 36 studies showed that maternal hyperthermia around the time of childbirth increased the odds of neonatal brain injuries, such as cerebral palsy, by 2.5-fold.²⁷ Neuronal injury in the fetus related to hyperthermia is ascribed to “energy failure”, where the amount of oxygen and substrates delivered to brain tissue in the fetus are insufficient to meet an increased demand during hyperthermia, leading to brain tissue acidosis, ischemic injury, and long-term neurologic sequelae in affected children.²⁸ In those studies, body temperatures above 39°C in the mother had particularly severe consequences for the fetus, but even mildly elevated maternal temperatures (37.1°C–37.5°C) carried substantial risks for adverse birth outcomes.²⁹




FIGURE 1 Pathophysiologic processes and indirect impacts of high ambient temperatures on maternal and newborn health.

Second, maternal dehydration during a heat event can lead to electrolyte imbalances and cause endothelial, oxidative, and inflammatory sequelae,³⁰ undermining already burdened cardiovascular and renal systems during pregnancy. Levels of mild dehydration, as measured by urine specific gravity, can reach 40% among women during pregnancy and childbirth, even during periods with normal temperatures.^{31,32} Within the fetus, dehydration can manifest as severe oligohydramnios during extreme heat events.³³ Third, high ambient temperatures can interrupt endocrine system function, increasing oxidative stress, inflammation, and the release of heat-shock proteins, which affect placental adaptation to hypoxia, cell death regulation, and cytoskeletal rearrangements.^{34,35} Extreme heat events in pregnancy have been linked with abnormal glucose tolerance and gestational diabetes mellitus, both of which can negatively affect maternal and newborn health.³⁶ At high ambient temperatures, adrenaline and stress hormones rise, adversely affecting mental health and well-being, of concern given the already heightened risk for mental health conditions during pregnancy and postpartum.^{37,38} Lastly, in a fourth mechanism, heat stress affects placental growth, development, and function.³⁹ Chronic thermal stress is associated with “placental shrinkage” or reduction in volume and weight of the placenta, as shown in human³⁹ and animal⁴⁰ studies. This limits placental transport efficiency, including for oxygen and nutrients. During more acute periods of thermal stress, shunting of blood to the body surface reduces uterine blood flow and causes placental hypoperfusion.⁴¹

The four interlinked pathophysiologic processes described above manifest clinically in a broad spectrum of conditions (Figure 1). Maternal health conditions linked with high ambient temperatures include hypertensive disorders of pregnancy (e.g., pre-eclampsia and eclampsia),^{42,43} including in a recent study in South Africa,¹⁴ preterm prelabor rupture of membranes,⁴⁴ placental abruption,⁴⁵ antenatal

and postpartum hemorrhage,^{42,43} renal disease,⁴³ and increased rates of emergency cesarean sections^{24,43} (Figure 1: Box 2). Data are not yet available on the impacts of high ambient temperatures on maternal deaths, though this is likely as heat exposure increases, the risk for all the principal causes of maternal deaths. Less frequent causes of these deaths in Africa also appear heat sensitive, for example, in one study in the USA the risk of a cardiovascular event during childbirth rose by 11% with each 1°C increase in temperature.⁴⁶

Heat exposure can lead to increased uterine contractions,^{47,48} fetal tachycardia and reduced fetal movements, as demonstrated in studies among pregnant women exposed to high temperatures in thermal chambers⁴⁹ or through intrauterine temperature monitors in women during labor.⁵⁰ Clinically, during periods of high ambient temperatures there are increased rates of fetal distress,⁴² preterm birth (increased in a meta-analysis by 1.16-fold during heatwaves, for example),¹³ low birth weight,¹³ stillbirths (increased in a meta-analysis by 1.24-fold at high versus low temperatures),¹³ meconium aspiration,⁴² neonatal jaundice,⁵¹ neonatal intensive care admissions,⁵² and perinatal mortality (increased 1.53-fold in one study when temperature exceeded the 95th centile).⁵³ Furthermore, several studies, including in Ethiopia⁵⁴ and Uganda,¹⁵ have suggested that exposure to high ambient temperatures in utero affects health throughout the life course.⁵⁵

Indirect impacts of high ambient temperatures include increased incidence of food-, water- and vector-borne infections (Figure 1: Box 3). High temperatures promote replication and survival of microbes and disease vectors. Infections of the genital tract, such as group B streptococcus,⁵⁶ which is the leading cause of bacterial pneumonia, sepsis, and meningitis in neonates, are especially concerning, in particular if labor is prolonged during heat extremes. Malaria infections and shifts in distribution to areas with non-immune populations have major implications for

maternal health as pregnant women are at high risk for complicated malaria.^{5,57}

High ambient temperatures place additional strains on health systems, including through reducing work performance of health providers (Figure 1: Box 4).⁵⁸ Further, levels of irritability and anxiety among health workers may rise as temperatures escalate,³⁸ potentially leading to patient disrespect. Extreme heat events threaten cold chains, and the safe storage of drugs and diagnostics. A study in Malawi showed that drug storage temperatures exceeded the recommended 25°C–30°C threshold for oxytocin in half of the 40 facilities assessed.⁵⁹ In another study in France, a large spike in cases of postpartum hemorrhage was ascribed to damage to an oxytocin batch during a heatwave, and rates of postpartum hemorrhage returned to baseline once a new batch of oxytocin was received.⁶⁰ High ambient temperatures may also undermine healthcare seeking behaviors during pregnancy.⁶¹ Many women wait in long queues to receive antenatal care, which is challenging during hot periods. Moreover, though no evidence is available, it is possible that during labor the burden of traveling by foot or in crowded, non-ventilated vehicles to facilities during periods of high ambient temperatures may make some women favor home over facility birth. Women giving birth under the care of a traditional birth attendant or a family member may lack protections during periods with high ambient temperatures: a study in Bangladesh, for example, found that newborns born at home during periods of extreme heat were 1.14-fold more likely to have neonatal illnesses (e.g., serious infections) than those born at home on cooler days.⁶² Additionally, periods of high ambient temperatures, accompanied by soil drying, can reduce crop and livestock yields, raising risk for malnutrition and reducing income for pregnant women. In turn, reduced income may limit a woman's ability to meet the out-of-pocket costs of accessing maternity care.

Taken together, it is clear that the escalations in temperatures owing to climate change will have major consequences for maternal and child health in Africa, and progressively so as temperatures escalate. Establishing large-scale programs to address these burdens is a major priority, but requires a robust evidence base to inform funding allocations. In the section that follows we present an approach to interventional research and program design that has been developed by the Climate Change and Health Group.

2 | AN APPROACH TO HEAT-HEALTH RESEARCH AND PROGRAMMING FOR SECURING MATERNAL AND NEWBORN HEALTH

In recent decades, it has become apparent that, with adequate intervention, health risks from extreme heat can be ameliorated. Within many high-income countries, heat-related mortality and morbidity have declined with successive heatwaves as adaptation measures have improved, approaching zero excess deaths in some areas.^{5,63} These declines may be attributable to interventions encompassing four domains,




FIGURE 2 Framework for reducing impacts of high ambient temperatures on maternal and newborn health.

namely (1) behavioral changes; (2) health system inputs, including the development of new health services; (3) changes to the built environment, in particular air conditioning^{64–66}; and (4) structural and policy interventions (Figure 2). We describe the salient features of these domains below, and how they relate to maternal and newborn health.⁶⁷

2.1 | Behavior change interventions

Information campaigns and behavior change interventions that target pregnant women, female family members (e.g., mothers-in-law), male partners, community leaders, and other stakeholders could support self-care interventions during pregnancy.⁶⁸ Such interventions include maintaining hydration during hot periods, accessing cool areas, or seeking shade; using low-cost water sprays; wearing appropriate clothing, ideally cotton; and self-monitoring for symptoms of heat illness, especially dizziness, heavy sweating, fatigue, and clammy skin with goosebumps.^{64–66} Many pregnant women continue to do physical work even late in gestation, including walking long distances to collect water and firewood. Interventions are needed to shift these workloads to other family members during pregnancy. Promotion of hydration and water supplementation during periods of high ambient temperatures, such as in “Water-Rest-Shade” initiatives, can have major benefits.⁶⁹

Coverage of antenatal care and community-based services are high in many parts of Africa, meaning that pregnant women have frequent contacts with the health system, providing multiple opportunities for reinforcement of behavior changes. Several mHealth initiatives have specifically targeted pregnant women (e.g., MomConnect SMS program in South Africa⁷⁰ and Wired Mothers in Zanzibar, Kenya⁷¹), offering an additional medium for heat-health messaging. Further, preparedness and complication readiness—key tenets of heat planning responses—have been at the heart of health promotion in maternal health, and could be adapted to encompass heat-related interventions. Birth preparedness, for example, includes helping women to plan in advance for their transport during labor,^{72,73} which may be particularly important during periods of high ambient temperatures.

2.2 | Health system and service interventions

The considerable investments in Africa in infrastructure, services, and human resources for maternal health provide a platform for interventions such as Early Warning Systems. Warning systems form the mainstay of heat-health services in high-income countries. Here, when weather forecasts predict that a pre-specified temperature threshold will be exceeded, warnings are issued and a series of tiered interventions are triggered. Then, when the heat event occurs, health services focus on protecting high-risk groups, including pregnant women and infants.^{64–66} This light include home “check-ups” of pregnant women by community health workers, and facilitating access to “cooling areas” in the community or in a maternity facility. Heightened monitoring during childbirth for potential complications of heat exposure is prudent, such as more frequent assessments of water intake and hydration status, labor duration, blood loss, and signs of infection, especially group B streptococcus. In hot climates, water supplementation, even in people with very mild dehydration, can have considerable positive impacts.⁷⁴ Other health system interventions could include integrating indicators within the district health systems to track the burden of heat-related conditions in pregnant women and newborns, and ensuring that the cold chain for drugs is maintained during extreme heat events, especially for oxytocin.

2.3 | Built environment, cooling, and nature-based solutions

Most women in Africa give birth in a health facility, meaning that providing an environment for childbirth that protects pregnant women and newborns against high ambient temperatures may have a high impact (Figure 3). Passive cooling can provide high levels of sustained protection. This may include minor modifications to labor wards to alter interactions between a building envelope and the natural elements. Simple facade fixes such as awnings, overhangs, louvres, and insulated walls and roofs are highly effective at reducing solar radiation. “Heat proofing” includes enhancing heat resistance of buildings through increasing reflectivity of surfaces (enhanced

albedo) and the provision of localized solutions such as mist sprays and evaporative materials. White reflective paint for rooftops can reflect almost all solar radiation.⁶⁶

Indigenous knowledge around building design and urban planning holds much promise, with, for example, traditional earthen housing construction with straw roof materials providing substantial thermal comfort advantages over poorly insulated concrete housing and sheet iron roofs. Natural ventilation strategies such as chimneys and air vents (e.g., “whirly birds”) in homes and maternity wards could facilitate cool air circulation and dissipate heat. The effectiveness of passive cooling, however, varies widely, depending on the location of the sun and wind direction, among other factors. Careful thermal modeling simulations and cost-effectiveness evaluations can assist in selecting the optimum building modifications.

Air conditioning is referred to as “tantamount to a potentially life-saving medical device” for groups with high heat vulnerability.⁶⁶ But, equally, the widespread use of air conditioning is “a clear example of maladaptation to climate change”. Hence, notwithstanding the multiple drawbacks of air conditioning (e.g., using as much as 8.5% of total electricity consumption and more than half of electricity on hot days, high energy costs, noise pollution, heat generation, power outages, and its signaling of widening global “thermal inequities”),⁵ if used appropriately, it may have an important role in protecting maternal and newborn health. Use of air conditioning in a few designated “cool spaces”, such as labor wards, can provide protection during extreme heat events and may be cost-effective. There are direct parallels between the concept of Maternity Waiting Homes, which serve to shelter and protect pregnant women, and the notion of having dedicated cooling “shelters” in maternity hospitals. Many lessons from waiting homes around acceptability, recipient selection, and implementation may also be applicable to “cooling centres”.⁷² Maintaining comfortable temperatures in facilities would also help to reduce thermal discomfort for maternity health workers, shoring up their performance, and potentially averting irritability or aggressive behaviors. Electric and hand-held fans with light water spraying offer low-cost, potentially scalable alternatives to air conditioning.⁵

Nature-based “green and blue” solutions around maternity facilities and at the homes of pregnant women can reduce temperatures




FIGURE 3 Potential built environment, cooling and nature-based solutions.

by several degrees, and are often low cost and considered an essential part of “bottom-up” community-led adaptations. Interventions may include street trees, green roofs, green walls, ponds and other water features, where feasible. Shading from tree canopies reduces radiant heat loads, a key determinant of outdoor thermal comfort around antenatal clinics, for example. Moreover, trees and other vegetation produce considerable cooling from evapotranspiration. Greening also has several environmental and health co-benefits, including noise reduction, improvements in social well-being, and amelioration of the mental health impacts of high ambient temperatures.

2.4 | Structural, policy, and climate financing interventions

Major investments are needed to improve the social and economic conditions that underlie much of the vulnerability to climate change in Africa. Most especially, many of the factors that place pregnant women in Africa at risk during periods of high ambient temperatures relate to resource constraints, and the deficiencies in housing, nutrition, and social services that mark the continent. Resilience to climate shocks is low in most parts of Africa. Intense, prolonged heat waves can have devastating impacts on crop and livestock resources, placing pregnant women and newborns at high risk for malnutrition. Rates of metabolism are raised markedly during periods of high temperatures, heightening nutritional needs. Clearly, pregnant women need to be a high priority population for food support. Cash transfers provided to pregnant women have multiple benefits for pregnant women and newborns, aside from improving food security.⁷⁵ Of major concern, few countries in Africa have developed policies around climate change and health. The CHANCE network established within the EU Horizons ENBEL project aims to address this key gap. A conducive policy environment is essential for developing programs and services around behavioral, health systems, and the built environment as described above.

Commitments to provide adequate climate financing were a central part of the Paris Accord and subsequent meetings of global leaders. Scant progress, however, has been made to meeting these commitments. For example, only a small fraction of the US\$ 100 billion per year promised by high-income countries has been provided.⁷⁶ Climate financing earmarked for enhancing resilience in pregnant women and for programming to provide heat-related services for pregnant women may go a long way towards addressing the growing burden of heat-related conditions among pregnant women in Africa.

3 | RESEARCH AND PROGRAMMATIC PRIORITIES IN AFRICA

Climate change threatens to undo the hard-fought gains made by maternal and child health programs in Africa over the past few decades.⁷⁷ These gains are fragile: maternal and newborn mortality in Africa remains the highest worldwide, with two out of every three maternal deaths in the world occurring in sub-Saharan

Africa for example,⁷⁸ and rates of preterm birth in Africa remain the highest globally.⁷⁹ Clearly, adaptation interventions that result in even small reductions, for example, risk for preterm births, will have major cost savings and lifelong benefits for affected individuals.

A cluster of factors and health system opportunities make pregnant women a highly suitable target group for heat-related interventions in Africa. Considerable investments in shoring up maternal and child health services—at both facility and community level—on the continent means that these services provide an ideal platform for heat-health interventions. Indeed, maternal and child health services form the cornerstone of primary health care in Africa and have proven historically to be adept at incorporating new programmatic areas, as shown by HIV mother-to-child transmission programs.⁸⁰

What is needed are intervention packages that draw on a range of behavioral, health systems, and built environment solutions, optimized in each setting. These packages need to be systematically co-designed with pregnant and postpartum women, health workers, male partners, and other stakeholders. Qualitative, ethnographic research, and participatory co-design processes are key for tailoring heat adaptation interventions to different socio-cultural and geographical contexts. Selection of the package of interventions requires careful consideration of the cost-effectiveness trade-offs between potential interventions. Integrating heat-related interventions within existing programs for maternal and newborn health may enhance sustainability in the long run. Importantly, many health workers do not yet have the knowledge and skills required to protect pregnant women and newborns against high ambient temperatures, and training is needed, both pre- and in-service.

Evidence from rigorous studies is required to ensure that climate financing aid to assist African countries to adapt to climate change is used optimally. The limited research to date on climate change adaptation in Africa, however, means that there are major gaps in evidence. Research among pregnant women is the ideal place to begin building an evidence base, which can then be extended to other vulnerable groups on the continent. To our knowledge, worldwide, only one study has so far set out to examine heat adaptation interventions in pregnant women (the CHAMNHA study, funded by the Belmont Forum).⁸¹ In fact, globally, across all population groups, most assessments to date of effectiveness of heat adaptation interventions have been based on post-hoc analyses of trends in health surveillance data after heat waves.^{52,63}

Clearly, formulating, testing, costing, and scaling up solutions to protect pregnant women against high ambient temperatures is a major public health priority. There is also a moral imperative to do so: women in Africa have made a negligible contribution to global carbon emissions, yet may be among the populations most affected by climate change and have the fewest resources to cope, especially while pregnant. Large-scale research projects among pregnant women in Africa and then heat-health programs at scale are long overdue.

AUTHOR CONTRIBUTIONS

MFC, FS, VL and SL contributed to the conceptualization of the paper, writing of drafts of the articles and preparation of the final manuscript. Climate Change and Heat-Health Study Group members contributed to the draft writing and editing of the manuscript.

ACKNOWLEDGMENTS

This research was supported by the Natural Environment Research Council (NERC; grant numbers NE/T013613/1, NE/T01363X/1); Research Council of Norway (RCN; grant number 312601); and The Swedish Research Council for Health, Working Life and Welfare in collaboration with the Swedish Research Council (Forte; grant number 2019-01570); coordinated through a Belmont Forum partnership in the CHAMNHA project.

CONFLICT OF INTEREST

MFC and FS hold investments in the fossil fuel industry through their pension funds as obligated by the Wits Health Consortium. The University of the Witwatersrand holds investments in the fossil fuel industry through their endowments and other financial reserves. No other conflicts of interest declared.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Matthew F. Chersich  <https://orcid.org/0000-0002-4320-9168>

REFERENCES

- Opoku SK, Leal Filho W, Hubert F, Adejumo O. Climate change and health preparedness in Africa: Analysing trends in six African countries. *Int J Environ Res Public Health*. 2021;18(9):4672.
- IPCC. *Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K. Mintenbeck, A. Alegria, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press. 2022.
- IPCC. In: Masson-Delmotte V, Zhai P, Pirani A, et al., eds. *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press. 2021.
- Varela R, Rodríguez-Díaz L, de Castro M. Persistent heat waves projected for Middle East and North Africa by the end of the 21st century. *PLoS One*. 2020;15(11):e0242477.
- Watts N, Amann M, Arnell N, et al. The 2020 report of the lancet countdown on health and climate change: responding to converging crises. *Lancet*. 2021;397(10269):129-170.
- Bunker A, Sewe MO, Sie A, Rocklov J, Sauerborn R. Excess burden of non-communicable disease years of life lost from heat in rural Burkina Faso: a time series analysis of the years 2000-2010. *BMJ Open*. 2017;7(11):e018068.
- Azongo DK, Awine T, Wak G, Binka FN, Oduro AR. A time series analysis of weather variability and all-cause mortality in the Kasena-Nankana districts of northern Ghana, 1995-2010. *Glob Health Action*. 2012;5:14-22.
- Egondi T, Kyobutungi C, Rocklov J. Temperature variation and heat wave and cold spell impacts on years of life lost among the urban poor population of Nairobi, Kenya. *Int J Environ Res Public Health*. 2015;12(3):2735-2748.
- Scovronick N, Sera F, Acquavota F, et al. The association between ambient temperature and mortality in South Africa: a time-series analysis. *Environ Res*. 2018;161:229-235.
- Mrema S, Shamte A, Selemani M, Masanja H. The influence of weather on mortality in rural Tanzania: a time-series analysis 1999-2010. *Glob Health Action*. 2012;5:33-43.
- van der Linden N, Longden T, Richards JR, et al. The use of an 'acclimatisation' heatwave measure to compare temperature-related demand for emergency services in Australia, Botswana, Netherlands, Pakistan, and USA. *PLoS One*. 2019;14(3):e0214242.
- Bishop-Williams KE, Berrang-Ford L, Sargeant JM, et al. Understanding weather and hospital admissions patterns to inform climate change adaptation strategies in the healthcare sector in Uganda. *Int J Environ Res Public Health*. 2018;15(11):2402.
- Chersich MF, Pham MD, Areal A, et al. Associations between high temperatures in pregnancy and risk of preterm birth, low birth weight, and stillbirths: systematic review and meta-analysis. *BMJ*. 2020;371:m3811.
- Part C, le Roux J, Chersich MF, Sawry S, Filippi V, Roos N, et al. Ambient temperature during pregnancy and risk of maternal hypertensive disorders: A time-to-event study in Johannesburg, South Africa. *Environ Res (under review)* 2022;212:113596.
- Duchoslav J. Prenatal temperature shocks reduce cooperation: evidence from public goods games in Uganda. *Front Behav Neurosci*. 2017;11:249.
- Naicker N, Teare J, Balakrishna Y, Wright CY, Mathee A. Indoor temperatures in low cost housing in Johannesburg, South Africa. *Int J Environ Res Public Health*. 2017;14(11):1410.
- Wright CY, Street RA, Cele N, et al. Indoor temperatures in patient waiting rooms in eight rural primary health care centers in Northern South Africa and the related potential risks to human health and wellbeing. *Int J Environ Res Public Health*. 2017;14(1):43.
- Kouanda S, CHAMNHA Consortium. *How Heatwaves Affect Maternal and Newborn Care Behaviours: Perspectives from Women and Health Providers. FIGO Conference October 25 2021 Session: Putting out the Fire! – Managing the Risks of Extreme Heat on Maternal and Newborn Health Arising from Climate Change*. 2021.
- Kutywayo A, Chersich MF, Naidoo N, et al. Knowledge, attitudes and impacts of climate change in secondary school learners in South Africa. *Jamba (Submitted)*. 2022;14(1):1162.
- Haghighi MM, Wright CY, Ayer J, et al. Impacts of high environmental temperatures on congenital anomalies: a systematic review. *Int J Environ Res Public Health*. 2021;18:4910.
- Chersich MF. *Impacts of Extreme Heat During Pregnancy: A Systematic Review. 33rd Annual Conference of the International Society for Environmental Epidemiology Session entitled: "Climate change, and maternal and newborn health outcomes: New methods to address heat impacts in vulnerable populations"* 2021.
- Marx GF, Loew DA. Tympanic temperature during labour and parturition. *Br J Anaesth*. 1975;47(5):600-602.
- Schouten FD, Wolf H, Smit BJ, Bekedam DJ, de Vos R, Wahlen I. Maternal temperature during labour. *Int J Obstet Gynaecol*. 2008;115(9):1131-1137.
- Frölich MA, Esame A, Zhang K, Wu J, Owen J. What factors affect intrapartum maternal temperature? A prospective cohort study: maternal intrapartum temperature. *Anesthesiology*. 2012;117(2):302-308.
- Fusi L, Steer PJ, Maresh MJ, Beard RW. Maternal pyrexia associated with the use of epidural analgesia in labour. *Lancet*. 1989;1(8649):1250-1252.
- Kasiteropoulou D, Topalidou A, Downe S. A computational fluid dynamics modelling of maternal-fetal heat exchange and blood flow in the umbilical cord. *PLoS One*. 2020;15(7):e0231997.

27. Morton S, Kua J, Mullington CJ. Epidural analgesia, intrapartum hyperthermia, and neonatal brain injury: a systematic review and meta-analysis. *Br J Anaesth*. 2021;126(2):500-515.
28. Kasdorf E, Perlman JM. Hyperthermia, inflammation, and perinatal brain injury. *Pediatr Neurol*. 2013;49(1):8-14.
29. Lieberman E, Cohen A, Lang J, Frigoletto F, Goetzl L. Maternal intrapartum temperature elevation as a risk factor for cesarean delivery and assisted vaginal delivery. *Am J Public Health*. 1999;89(4):506-510.
30. Halonen JI, Zanobetti A, Sparrow D, Vokonas PS, Schwartz J. Associations between outdoor temperature and markers of inflammation: a cohort study. *Environ Health*. 2010;9:42.
31. Aziz MM, Kulkarni A, Tunde-Agbede O, Benito CW, Oyelese Y. Are women with threatened preterm labor more dehydrated than women without it? *Journal of Obstetric, Gynecologic, and Neonatal Nursing: JOGNN*. 2018;47(5):602-607.
32. Mulyani EY, Hardinsyah, Briawan D, Santoso BI. Hydration status of pregnant women in West Jakarta. *Asia Pac J Clin Nutr*. 2017;26(Suppl 1):S26-s30.
33. Luton D, Alran S, Fourchette V, Sibony O, Oury JF. Paris heat wave and oligohydramnios. *Am J Obstet Gynecol*. 2004;191(6):2103-2105.
34. Kanninen TT, Sisti G, Witkin SS. Induction of the 70 kDa heat shock protein stress response inhibits autophagy: possible consequences for pregnancy outcome. *J Matern Fetal Neonatal Med*. 2016;29(1):159-162.
35. MacPhee DJ, Miskiewicz EI. The potential functions of small heat shock proteins in the uterine musculature during pregnancy. *Adv Anat Embryol Cell Biol*. 2017;222:95-116.
36. Molina-Vega M, Gutiérrez-Repiso C, Muñoz-Garach A, et al. Relationship between environmental temperature and the diagnosis and treatment of gestational diabetes mellitus: an observational retrospective study. *Sci Total Environ*. 2020;744:140994.
37. Chersich MF, Swift CP, Edelstein I, et al. Violence in hot weather: will climate change exacerbate rates of violence in South Africa? *S Afr Med J*. 2019;109(7):447-449.
38. Lin Y, Hu W, Xu J, et al. Association between temperature and maternal stress during pregnancy. *Environ Res*. 2017;158:421-430.
39. Wang J, Liu X, Dong M, et al. Associations of maternal ambient temperature exposures during pregnancy with the placental weight, volume and PFR: a birth cohort study in Guangzhou, China. *Environ Int*. 2020;139:105682.
40. Limesand SW, Camacho LE, Kelly A, Antolic AT. Impact of thermal stress on placental function and fetal physiology. *Anim Reprod*. 2018;15(1):886-898.
41. Pirhonen JP, Vähä-Eskeli KK, Seppänen A, Vuorinen J, Erkkola RU. Does thermal stress decrease uterine blood flow in hypertensive pregnancies? *Am J Perinatol*. 1994;11(5):313-316.
42. Cil G, Cameron TA. Potential climate change health risks from increases in heat waves: abnormal birth outcomes and adverse maternal health conditions. *Risk Anal*. 2017;37(11):2066-2079.
43. Qu Y, Zhang W, Ryan I, et al. Ambient extreme heat exposure in summer and transitional months and emergency department visits and hospital admissions due to pregnancy complications. *Sci Total Environ*. 2021;777:146134.
44. Gat R, Kachko E, Kloog I, et al. Differences in environmental factors contributing to preterm labor and PPRM - population based study. *Environ Res*. 2021;196:110894.
45. Rammah A, Whitworth KW, Han I, Chan W, Hess JW, Symanski E. Temperature, placental abruption and stillbirth. *Environ Int*. 2019;131:105067.
46. Agha SB, Tchouassi DP, Bastos ADS, Sang R. Assessment of risk of dengue and yellow fever virus transmission in three major Kenyan cities based on Stegomyia indices. *PLoS Negl Trop Dis*. 2017;11(8):20.
47. Khamis Y, Shaala S, Damarawy H, Romia A, Topozada M. Effect of heat on uterine contractions during normal labor. *Int J Gynaecol Obstet*. 1983;21(6):491-493.
48. Morishima HO, Glaser B, Niemann WH, James LS. Increased uterine activity and fetal deterioration during maternal hyperthermia. *Am J Obstet Gynecol*. 1975;121(4):531-538.
49. Vähä-Eskeli K, Erkkola R. The effect of short-term heat stress on uterine contractility, fetal heart rate and fetal movements at late pregnancy. *Eur J Obstet Gynecol Reprod Biol*. 1991;38(1):9-14.
50. Cronjé HS. Intra-uterine temperature measurements during fetal tachycardia. *S Afr Med J*. 1977;52(12):476-478.
51. Scrafford CG, Mullany LC, Katz J, et al. Incidence of and risk factors for neonatal jaundice among newborns in southern Nepal. *Tropical Med Int Health*. 2013;18(11):1317-1328.
52. Kakkad K, Barzaga ML, Wallenstein S, Azhar GS, Sheffield PE. Neonates in Ahmedabad, India, during the 2010 heat wave: a climate change adaptation study. *J Environ Public Health*. 2014;2014:946875.
53. Basagana X, Sartini C, Barrera-Gomez J, et al. Heat waves and cause-specific mortality at all ages. *Epidemiology*. 2011;22(6):765-772.
54. Randell H, Gray C, Grace K. Stunted from the start: Early life weather conditions and child undernutrition in Ethiopia. *Soc Sci Med*. 2020;261:113234.
55. Chersich M, Kovats S, Part C, Samuels L, Hajat S, Solarin I, et al. Systematic review of the effect of ambient heat on maternal health outcomes. *International Society for Environmental Epidemiology (ISEE) 2021 Conference 2021*; <https://ehp.niehs.nih.gov/doi/abs/10.1289/isee.2021.O-SY-092>.
56. Dadvand P, Basagana X, Figueras F, Sunyer J, Nieuwenhuijsen MJ. Climate and group B streptococci colonisation during pregnancy: present implications and future concerns. *BJOG*. 2011;118(11):1396-1400.
57. Rocklöv J, Dubrow R. Climate change: an enduring challenge for vector-borne disease prevention and control. *Nat Immunol*. 2020;21(5):479-483.
58. International Labour Organization. *Working on a Warmer Planet: The Effect of Heat Stress on Productivity and Decent Work*. 2019; https://www.ilo.org/global/publications/books/WCMS_711919/lang-en/index.htm.
59. Hagen N, Khuluza F, Heide L. Quality, availability and storage conditions of oxytocin and misoprostol in Malawi. *BMC Pregnancy Childbirth*. 2020;20(1):184.
60. Parpex G, Khediri Z, Michel P, Visbecq JN, Duviquet MJ, Poncelet C. Postpartum hemorrhage: could oxytocin be the cause? Results from a morbidity and mortality review to enhance quality, safety, and relevance of care. *Eur J Obstet Gynecol Reprod Biol*. 2021;258:299-303.
61. Lusambili A, Khaemba P, Chabeda S, Scorgie F, Nakstad B, Kadio K, et al. *Community Perspectives on Maternal and Newborn Experiences of Heat Stress: A qualitative Inquiry in Kilifi, Kenya*. 12th European Congress on Tropical Medicine and International Health 2021.
62. Mannan I, Choi Y, Coutinho AJ, et al. Vulnerability of newborns to environmental factors: findings from community based surveillance data in Bangladesh. *IJERPH*. 2011;8(8):3437-3452.
63. Chung Y, Yang D, Gasparrini A, et al. Changing susceptibility to non-optimum temperatures in Japan, 1972-2012: the role of climate, demographic, and socioeconomic factors. *Environ Health Perspect*. 2018;126(5):57002.
64. Public Health England, NHS England, Local government association, Met Office. *Heatwave Plan for England: Protecting Health and Reducing Harm From Severe Heat and Heatwaves*. Updated 2021.
65. WMO, WHO. *Heatwaves and health: guidance on warning-system development*. WMO-No. 1142. 2015.
66. WHO Regional Office for Europe. *Heat and health in the WHO European Region: updated evidence for effective prevention*. Licence: CC BY-NC-SA 3.0 IGO. 2021.
67. Chersich MF. *Maternal health and environmental heat*. 14th European Public Health Conference 10 November 2021 Session Entitled *Public Health Measures to Address the Impact of Climate Change on Population Health 2021*.

68. WHO. *WHO Consolidated Guideline on Self-Care Interventions for Health*. 2021.
69. Wegman DH, Apelqvist J, Bottai M, et al. Intervention to diminish dehydration and kidney damage among sugarcane workers. *Scand J Work Environ Health*. 2018;44(1):16-24.
70. Kabongo EM, Mukumbang FC, Delobelle P, Nicol E. Combining the theory of change and realist evaluation approaches to elicit an initial program theory of the MomConnect program in South Africa. *BMC Med Res Methodol*. 2020;20(1):282.
71. Lund S, Nielsen BB, Hemed M, et al. Mobile phones improve antenatal care attendance in Zanzibar: a cluster randomized controlled trial. *BMC Pregnancy Childbirth*. 2014;14:29.
72. WHO. WHO recommendations on health promotion interventions for maternal and newborn health. 2015. https://www.who.int/maternal_child_adolescent/documents/health-promotion-interventions/en/
73. Magoma M, Requejo J, Campbell O, Cousens S, Meriardi M, Filippi V. The effectiveness of birth plans in increasing use of skilled care at delivery and postnatal care in rural Tanzania: a cluster randomised trial. *Tropical Med Int Health*. 2013;18(4):435-443.
74. Fadda R, Rapinett G, Grathwohl D, et al. Effects of drinking supplementary water at school on cognitive performance in children. *Appetite*. 2012;59(3):730-737.
75. Chersich MF, Luchters S, Blaauw D, et al. Safeguarding maternal and child health in South Africa by starting the child support Grant before birth: design lessons from pregnancy support programmes in 27 countries. *S Afr Med J*. 2016;106(12):1192-1210.
76. Green Climate Fund. <https://www.greencclimatefund/about/resource-mobilisation>.
77. Garcia DM, Sheehan MC. Extreme weather-driven disasters and children's health. *Int J Health Serv*. 2016;46(1):79-105.
78. Graham W, Woodd S, Byass P, et al. Diversity and divergence: the dynamic burden of poor maternal health. *Lancet*. 2016;388(10056):2164-2175.
79. Chawanpaiboon S, Vogel JP, Moller AB, et al. Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *Lancet Glob Health*. 2019;7(1):e37-e46.
80. Chersich MF, Newbatt E, Ng'oma K, de Zoysa I. UNICEF's contribution to the adoption and implementation of option B+ for preventing mother-to-child transmission of HIV: a policy analysis. *Glob Health*. 2018;14(1):55.
81. CHAMNA Project. *Climate, Heat and Maternal and Neonatal Health In Africa: A Belmont Forum Project*. <https://www.wlshtmacuk/research/centres-projects-groups/chamna>.

How to cite this article: Chersich MF, Scorgie F, Filippi V, Luchters S, Climate Change and Heat-Health Study Group. Increasing global temperatures threaten gains in maternal and newborn health in Africa: A review of impacts and an adaptation framework. *Int J Gynecol Obstet*. 2022;00:1-9. doi:[10.1002/ijgo.14381](https://doi.org/10.1002/ijgo.14381)

APPENDIX

CLIMATE CHANGE AND HEAT-HEALTH STUDY GROUP

Andrea Huggett (Wits RHI), Euphemia Sibanda (CeSHHAR), Craig Parker (Wits RHI), Darshnika Lakhoo (Wits RHI), Gloria Maimela (Wits RHI), Helen Rees (Wits RHI), Ijeoma Solarin (Wits RHI), Lois Harden (Department of Physiology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa), Robyn Hetem (School of Animal, Plant and Environmental Sciences, Faculty of Sciences, University of the Witwatersrand, Johannesburg, South Africa), and Webster Mavhu (CeSHHAR).