

This is an author version of the contribution published in
Topology and its Applications

The definitive version is available at
<http://www.sciencedirect.com/science/article/pii/S0166864111002501>
doi:10.1016/j.topol.2011.02.013

Functorial topologies and finite-index subgroups of abelian groups

Dikran Dikranjan

dikran.dikranjan@uniud.it

Dipartimento di Matematica e Informatica,

Università di Udine,

Via delle Scienze, 206 - 33100 Udine, Italy

Anna Giordano Bruno

anna.giordanobruno@uniud.it

Dipartimento di Matematica e Informatica,

Università di Udine,

Via delle Scienze, 206 - 33100 Udine, Italy

Dedicated to the seventieth birthday of Eraldo Giuli

Abstract

In the general context of functorial topologies, we prove that in the lattice of all group topologies on an abelian group, the infimum between the Bohr topology and the natural topology is the profinite topology. The profinite topology and its connection to other functorial topologies is the main objective of the paper. We are particularly interested in the poset $\mathcal{C}(G)$ of all finite-index subgroups of an abelian group G , since it is a local base for the profinite topology of G . We describe various features of the poset $\mathcal{C}(G)$ (its cardinality, its cofinality, etc.) and we characterize the abelian groups G for which $\mathcal{C}(G) \setminus \{G\}$ is cofinal in the poset of all subgroups of G ordered by inclusion. Finally, for pairs of functorial topologies \mathcal{T}, \mathcal{S} we define the equalizer $\mathcal{E}(\mathcal{T}, \mathcal{S})$, which permits to describe relevant classes of abelian groups in terms of functorial topologies.

1 Introduction

The concept of functorial topology was introduced by Charles [2]. He proposed a method for constructing such topologies, which was generalized by Fuchs [20, Vol. I, p. 33]. Later on, functorial topologies were subject of study by many authors, among them Boyer and Mader [1], Mader [27] and Mader and Mines [28].

Let U be the forgetful functor $U : \mathbf{TopAb} \rightarrow \mathbf{Ab}$, where \mathbf{Ab} is the category of all abelian groups and their morphisms and \mathbf{TopAb} is the category of all topological abelian groups and their morphisms.

Definition 1.1. A functorial topology is a functor $\mathcal{T} : \mathbf{Ab} \rightarrow \mathbf{TopAb}$ such that $U\mathcal{T} = 1_{\mathbf{Ab}}$.

Equivalently, following [20], a functorial topology is a class $\mathcal{T} = \{\mathcal{T}_G : G \in \mathbf{Ab}\}$, where (G, \mathcal{T}_G) is a topological group for every $G \in \mathbf{Ab}$, and every homomorphism $G \rightarrow H$ in \mathbf{Ab} is continuous $(G, \mathcal{T}_G) \rightarrow (H, \mathcal{T}_H)$. So, a functorial topology is a functor $\mathcal{T} : \mathbf{Ab} \rightarrow \mathbf{TopAb}$ such that $\mathcal{T}(G) = (G, \mathcal{T}_G)$ for any $G \in \mathbf{Ab}$, where \mathcal{T}_G denotes the topology on G , and $\mathcal{T}(\phi) = \phi$ for any morphism ϕ in \mathbf{Ab} [1].

By [1, Theorem 2.2], any functorial topology defined on a full subcategory of \mathbf{Ab} extends to \mathbf{Ab} , so that there is no need to introduce functorial topologies for full subcategories of \mathbf{Ab} .

A functorial topology \mathcal{T} is *linear* if \mathcal{T}_G is linear for every $G \in \mathbf{Ab}$ (recall that a group topology is *linear* if it has a local base consisting of open subgroups); moreover, \mathcal{T} is *ideal* if \mathcal{T} maps surjective homomorphisms to open (continuous) homomorphisms [1], and it is *hereditary* if $\mathcal{T}_H = \mathcal{T}_G \upharpoonright_H$ for every $G \in \mathbf{Ab}$ and every subgroup H of G .

Functorial topologies exist in abundance. The discrete topology δ and the indiscrete topology ι are functorial topologies that are both hereditary and ideal. In this paper we mainly focus our attention on the following three functorial topologies: for an abelian group G ,

- the *profinite topology* γ_G has all finite-index subgroups as a base of the neighborhoods of 0;
- the *natural topology* (sometimes called also *\mathbb{Z} -adic topology*) ν_G has the countable family of subgroups $\{mG : m \in \mathbb{N}_+\}$ as a base of the neighborhoods of 0;
- the *Bohr topology* \mathcal{P}_G is the initial topology of all homomorphisms $G \rightarrow \mathbb{T} = \mathbb{R}/\mathbb{Z}$, namely, the characters of G , where \mathbb{T} is equipped with the compact quotient topology of \mathbb{R}/\mathbb{Z} .

We also consider the p -adic topology ν^p , which can be viewed as a natural local version of ν and analogous local versions γ^p of γ and \mathcal{P}^p of \mathcal{P} (see Example 4.1).

Unlike the Bohr topology, the profinite and the natural topology are linear topologies. On the other hand, the natural topology and the profinite topology are ideal but not hereditary (see Example 2.7), while the Bohr topology is both ideal and hereditary (see Lemma 2.9(b)).

A topological abelian group (G, τ) is *totally bounded* if for any non-empty open subset U of (G, τ) there exists a finite subset F of G such that $U + F = G$. If τ is totally bounded and Hausdorff, it is said to be *precompact*. The completion \tilde{G} of any precompact abelian group G is compact [33], so the precompact abelian groups are precisely the subgroups of the compact abelian groups. In this paper, we use the fact that the Bohr topology on an abelian group G is the maximal totally bounded group topology on G ; this is a deep fact, deducible from Peter-Weyl's theorem for compact abelian groups. See [7], [18], [25] and [26] for the remarkable properties of this topology.

Our choice to concentrate mainly on the profinite, the natural and the Bohr topology, with a special emphasis on the connections between the Bohr topology and the profinite topology, is motivated by the fact that the functorial aspect of the Bohr topology has not been sufficiently brought to light neither in topology nor in algebra.

In Section 2 we investigate the properties of the profinite topology, mainly its relationship with the natural and the Bohr topology. In particular, it is known that $\gamma_G \leq \inf\{\nu_G, \mathcal{P}_G\}$ for any abelian group G (for a proof see Lemma 2.10). The first of the main theorems of this paper, which is Theorem 2.13, shows that actually equality holds, that is, in the lattice of all group topologies of an abelian group, the profinite topology is the infimum of the natural topology and the Bohr topology. Since the natural topology is metrizable, one may be left with the misperception that the profinite topology and the Bohr topology are very close due to the equality $\gamma_G = \inf\{\nu_G, \mathcal{P}_G\}$. We see in Theorem 3.10 that the behavior of the subgroups shows a substantial difference between these topologies. Namely, while every subgroup is closed in the Bohr topology, the abelian groups G in which every subgroup is γ_G -closed form a quite small class (this class consists precisely of the Pontryagin duals of the so called “exotic tori” introduced in [15]).

In [11], the adjoint algebraic entropy for endomorphisms ϕ of abelian groups G was introduced making use of the family $\mathcal{C}(G)$ of all finite-index subgroups of G (see Section 3 for the precise definition). Indeed, the adjoint algebraic entropy of endomorphisms ϕ of abelian groups G measures to what extent ϕ moves the finite-index subgroups of G . So, in the context of the adjoint algebraic entropy, it is worth studying the poset of finite-index subgroups of abelian groups G , calculating various invariants of it (as size, cofinality, etc.). In this direction, [11, Theorem 3.3] (see Theorem 3.1 below) characterizes the abelian groups G with countable $\mathcal{C}(G)$; abelian groups with this property are called *narrow*. In particular, every endomorphisms ϕ of a narrow abelian group G has $\text{ent}^*(\phi) = 0$ [11, Proposition 3.7]. The surprising dichotomy discovered in [11] (namely, $\mathcal{C}(G)$ is either countable or has size at least \mathfrak{c}) is fully explained by Theorem 3.3 (see also Lemma 3.2), since $|\mathcal{C}(G)|$ coincides with the size of the torsion part of a compact abelian group (namely, the Pontryagin dual of G).

Indeed, in Section 3 we study the cardinality of $\mathcal{C}(G)$ of an abelian group G in the general setting. The family $\mathcal{C}(G)$ forms a semilattice with respect to intersections and with top element G . One can look at $\mathcal{C}(G)$ also as a filter-base that gives rise to the profinite topology γ_G of the abelian group G . More precisely, we show that when $\mathcal{C}(G)$ is infinite, its cardinality coincides with the weight and the local weight of (G, γ_G) . So, the purely algebraic object $\mathcal{C}(G)$ is strictly related to the topological invariants of the profinite topology of G (see below for the definitions of these topological invariants).

Making use of the results from Section 2, for any abelian group G we characterize the size and the cofinality of $\mathcal{C}(G)$, that is, we compute the weight and the local weight of (G, γ_G) in Theorem 3.3. Furthermore, Theorem 3.6 characterizes the density character of (G, γ_G) , using the fact that it coincides with the density character and the weight of (G, ν_G) . In another direction, in Theorem 3.10 we describe the abelian groups G for which the poset $\mathcal{C}(G) \setminus \{G\}$ is cofinal in the larger poset $\mathcal{S}(G)$ of all subgroups of G ordered by inclusion.

Inspired by the fact that the narrow abelian groups form precisely the class of abelian groups for which the profinite and the natural topology coincide (see Theorem 3.1), in Section 4 we define the equalizer $\mathcal{E}(\mathcal{T}, \mathcal{S})$ of a pair of functorial topologies \mathcal{T}, \mathcal{S} . Moreover, we describe its basic properties and arrive in this way to the standard correspondence between functorial topologies and classes of abelian groups stable under isomorphisms, finite products and subgroups. In this section we provide also more examples of functorial topologies to better illustrate the usefulness of the equalizer.

We dedicate this paper to the seventieth birthday of Eraldo Giuli, for his relevant contributions in the field

of categorical topology and in particular, the closure operators in the sense of [3, 12, 13, 17], of which the functorial topologies in the category of abelian groups are a relevant inspiring example.

Notation and terminology

We denote by \mathbb{Z} , \mathbb{N} , \mathbb{N}_+ , \mathbb{Q} and \mathbb{R} respectively the set of integers, the set of natural numbers, the set of positive integers, the set of rationals and the set of reals. For $m \in \mathbb{N}_+$, we use $\mathbb{Z}(m)$ for the finite cyclic group of order m . Consider on $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ the norm given by $\|r + \mathbb{Z}\| = \min\{d(r, m) : m \in \mathbb{Z}\}$, with d the usual metric of \mathbb{R} .

All groups in this paper are abelian. For an abelian group G and $m \in \mathbb{Z}$, we let $mG = \{mx : x \in G\}$ and $G^1 = \bigcap_{m \in \mathbb{N}_+} mG$, the *first Ulm subgroup* of G . Obviously, $(G/G^1)^1 = 0$. We say that an abelian group G is *divisible*, if $G^1 = G$ (i.e., $G = mG$ for every $m \in \mathbb{N}_+$). We denote by $D(G)$ the biggest divisible subgroup of G (namely, the one generated by all divisible subgroups of G). Obviously, $D(G) \subseteq G^1$. We call G *reduced* if $D(G) = 0$. We denote by $r_0(G)$ the torsion-free rank of G and, for a prime p , $r_p(G)$ denotes the p -rank of G , that is, $\dim_{\mathbb{F}_p} G[p]$, where $G[p] = \{x \in G : px = 0\}$ is the p -socle of G and \mathbb{F}_p is the field with p elements. More generally, for $n \in \mathbb{N}_+$, let $G[n] = \{x \in G : nx = 0\}$.

For a subset M of a topological space X , we denote by \overline{M} the closure of M . For a topological group (G, τ) and a subgroup H of G , let τ_q denote the quotient topology of τ on G/H . Moreover, the quotient group $(G/\overline{\{0\}}, \tau_q)$ is the largest Hausdorff quotient group of (G, τ) ; we call it the *Hausdorff reflection* of (G, τ) (as $(G, \tau) \mapsto (G/\overline{\{0\}}, \tau_q)$ defines a reflection of the category of all topological groups into the full subcategory of all Hausdorff topological groups).

We denote by δ_G and ι_G respectively the discrete and the indiscrete topology of an abelian group G . We denote by $(G, \tau)^*$ the *dual group* of a topological abelian group (G, τ) , that is, $(G, \tau)^*$ is the abelian group of all continuous characters $(G, \tau) \rightarrow \mathbb{T}$, endowed with the discrete topology. In particular, $G^* = \text{Hom}(G, \mathbb{T})$.

For a topological abelian group (G, τ) , the *weight* $w(G, \tau)$ is the minimum cardinality of a base of (G, τ) , and the *local weight* (or, *character*) $\chi(G, \tau)$ is the minimum cardinality of a local base of (G, τ) . If (G, τ) is a totally bounded abelian group, then $w(G, \tau) = \chi(G, \tau)$ [4]. Finally, the *density character* $d(G, \tau)$ is the minimum cardinality of a dense subset of (G, τ) .

2 The profinite, the natural and the Bohr topology

The following are basic properties of functorial topologies.

Lemma 2.1. *Let \mathcal{T} be a functorial topology. Then:*

- (a) $\mathcal{T}_{G_1 \times G_2} = \mathcal{T}_{G_1} \times \mathcal{T}_{G_2}$ for every $G_1, G_2 \in \mathbf{Ab}$;
- (b) $\mathcal{T}_G \geq \prod_{i \in I} \mathcal{T}_{G_i}$ for every arbitrary family $\{G_i : i \in I\}$ in \mathbf{Ab} with $G = \prod_{i \in I} G_i$;
- (c) $\mathcal{T}_H \geq \mathcal{T}_G \upharpoonright_H$ for every $G \in \mathbf{Ab}$ and every subgroup H of G ; equality holds for every G and H if and only if \mathcal{T} is hereditary;
- (d) $(\mathcal{T}_G)_q \geq \mathcal{T}_{G/H}$ for every $G \in \mathbf{Ab}$ and every subgroup H of G ; equality holds for every G and H if and only if \mathcal{T} is ideal.

Proof. (a) Consider the projections $p_j : (G, \mathcal{T}_{G_1 \times G_2}) \rightarrow (G_j, \mathcal{T}_{G_j})$ for $j = 1, 2$, which are continuous by the definition of functorial topology. Then, for every neighborhood $U_1 \times U_2$ of 0 in $(G_1 \times G_2, \mathcal{T}_1 \times \mathcal{T}_2)$ there exists a neighborhood W of 0 in $(G_1 \times G_2, \mathcal{T}_{G_1 \times G_2})$ such that $p_j(W) \subseteq U_j$ for $j = 1, 2$, that is, $W \subseteq U_1 \times U_2$. Hence, $\mathcal{T}_{G_1 \times G_2} \geq \mathcal{T}_{G_1} \times \mathcal{T}_{G_2}$. To prove the converse inequality consider the inclusions $i_j : (G_i, \mathcal{T}_{G_i}) \rightarrow (G_1 \times G_2, \mathcal{T}_{G_1 \times G_2})$, for $j = 1, 2$, which are continuous by the definition of functorial topology. Then $\mathcal{T}_{G_1 \times G_2} \leq \inf\{\mathcal{T}_{G_1} \times \delta_{G_2}, \delta_{G_1} \times \mathcal{T}_{G_2}\} = \mathcal{T}_{G_1} \times \mathcal{T}_{G_2}$.

To prove (b) proceed as in the first part of the proof of item (a). For (c) and (d) it suffices to note that by definition the inclusion $(H, \mathcal{T}_H) \hookrightarrow (G, \mathcal{T}_G)$ and the projection $(G, \mathcal{T}_G) \rightarrow (G/H, \mathcal{T}_{G/H})$ are continuous. \square

We introduce a partial order between functorial topologies by letting $\mathcal{T} \leq \mathcal{S}$ whenever $\mathcal{T}_G \leq \mathcal{S}_G$ for every abelian group G . This makes the class \mathfrak{ft} of all functorial topologies a large complete lattice with top element δ and bottom element ι .

2.1 The profinite topology vs the natural topology

There is an important connection (see [20]) between the first Ulm subgroup G^1 of an abelian group G and the family $\mathcal{C}(G)$, namely

$$G^1 = \bigcap_{N \in \mathcal{C}(G)} N. \quad (2.1)$$

Now we recall a notion closely related to these two concepts:

Definition 2.2. *An abelian group G is residually finite if G is isomorphic to a subgroup of a direct product of finite abelian groups.*

Obviously, every residually finite abelian group is reduced.

Clearly, the natural topology is metrizable whenever it is Hausdorff. As a consequence of (2.1), Lemma 2.3 shows that the profinite and the natural topology are simultaneously Hausdorff (respectively, indiscrete). Moreover, we see that this occurs precisely when the abelian group is residually finite (respectively, divisible) (which are merely algebraic properties).

Lemma 2.3. *Let G be an abelian group. Then $G^1 = \overline{\{0\}}^{\gamma_G} = \overline{\{0\}}^{\nu_G}$, so $(G/G^1, \gamma_{G/G^1})$ is the Hausdorff reflection of (G, γ_G) and $(G/G^1, \nu_{G/G^1})$ is the Hausdorff reflection of (G, ν_G) . Moreover, the following conditions are equivalent:*

- (a) γ_G is Hausdorff (respectively, indiscrete);
- (b) G is residually finite (respectively, divisible);
- (c) $G^1 = 0$ (respectively, $G^1 = G$);
- (d) ν_G is Hausdorff (respectively, indiscrete).

Proof. The first two assertions are obvious and imply the equivalences (a) \Leftrightarrow (c) \Leftrightarrow (d).

To prove that (b) and (c) are equivalent, first note that clearly $G^1 = G$ precisely when G is divisible. Now assume that G is residually finite. Then G is isomorphic to a subgroup of $F = \prod_{i \in I} F_i$, where each F_i is a finite abelian group. Since $F^1 = 0$, also $G^1 = 0$. To prove the converse implication suppose that $G^1 = 0$. It follows from (2.1) that G is isomorphic to a subgroup of $\prod_{N \in \mathcal{C}(G)} G/N$, where each G/N is obviously finite (for every $N \in \mathcal{C}(G)$ consider the canonical projection $G \rightarrow G/N$; then the diagonal homomorphism $G \rightarrow \prod_{N \in \mathcal{C}(G)} G/N$ is injective by the assumption $G^1 = 0$). \square

Remark 2.4. Let G be an abelian group. Every $N \in \mathcal{C}(G)$ contains G^1 , and the canonical projection $\pi : G \rightarrow G/G^1$ gives rise to a bijection between $\mathcal{C}(G/G^1)$ and $\mathcal{C}(G)$ by taking inverse images under π .

Now we characterize the abelian groups G with finite $\mathcal{C}(G)$.

Lemma 2.5. *Let G be an abelian group. Then the following conditions are equivalent:*

- (a) G/G^1 is finite;
- (b) $G/D(G)$ is finite;
- (c) the Hausdorff reflection of (G, γ_G) is finite;
- (d) $\mathcal{C}(G)$ is finite.

Proof. (a) \Rightarrow (b) As G^1 has finite index in G , G^1 admits no proper finite-index subgroup. Consequently, G^1 is divisible. Since $D(G) \subseteq G^1$, we conclude that $G^1 = D(G)$.

(b) \Rightarrow (a) Is clear, since $D(G) \subseteq G^1$.

(a) \Leftrightarrow (c) Is given by Lemma 2.3.

(a) \Rightarrow (d) Follows from Remark 2.4.

(d) \Rightarrow (a) Since $\mathcal{C}(G)$ is finite, G^1 has finite index in G . \square

If the equivalent conditions of Lemma 2.5 hold true for an abelian group G , then $G = D(G) \times F$, where F is a finite abelian group. This is why we call such a group *almost divisible*.

For reader's convenience, we collect in the next face some easy to prove properties of the profinite topology. Note that (a) and (c) are obvious and (b) is given by Lemma 2.3.

Fact 2.6. *Let G be an abelian group. Then:*

- (a) (G, γ_G) is totally bounded;
- (b) (G, γ_G) is precompact if and only if G is residually finite;
- (c) every surjective homomorphism $\phi : (G, \gamma_G) \rightarrow (H, \gamma_H)$ is continuous and open (i.e., the profinite topology is an ideal functorial topology); in particular, $w(G, \gamma_G) \geq w(H, \gamma_H)$.

Example 2.7. (a) Analogously to item (c) of Fact 2.6, the natural topology is an ideal functorial topology.

- (b) If D is a divisible abelian group, then ν_D is indiscrete, and so γ_D is indiscrete as well. Indeed, $mD = D$ for every $m \in \mathbb{N}_+$.
- (c) Consider $\mathbb{Z} \subseteq \mathbb{Q}$. By item (a) $\nu_{\mathbb{Q}}$ and $\gamma_{\mathbb{Q}}$ coincide with the indiscrete topology of \mathbb{Q} . Then $\nu_{\mathbb{Q}} \upharpoonright_{\mathbb{Z}}$ and $\gamma_{\mathbb{Q}} \upharpoonright_{\mathbb{Z}}$ coincide with the indiscrete topology of \mathbb{Z} , while $\nu_{\mathbb{Z}}$ and $\gamma_{\mathbb{Z}}$ are not indiscrete. This proves that the natural and the profinite topology are not hereditary.

We show now that the family of all γ_G -closed (respectively, γ_G -dense) subgroups of an abelian group G coincides with the family of all ν_G -closed (respectively, ν_G -dense) subgroups of G :

Lemma 2.8. *Let G be an abelian group and H a subgroup of G . Then:*

- (a) H is γ_G -closed if and only if H is ν_G -closed;
- (b) H is γ_G -dense if and only if H is ν_G -dense.

Proof. Since both γ_G and ν_G are ideal, their quotient topologies on G/H coincide with $\gamma_{G/H}$ and $\nu_{G/H}$ respectively.

(a) The subgroup H is γ_G -closed (respectively, ν_G -closed) if and only if $\gamma_{G/H}$ (respectively, $\nu_{G/H}$) is Hausdorff. By Lemma 2.3, $\gamma_{G/H}$ is Hausdorff precisely when $\nu_{G/H}$ is Hausdorff, and so H is γ_G -closed if and only if H is ν_G -closed.

(b) The subgroup H is γ_G -dense (respectively, ν_G -dense) if and only if $\gamma_{G/H}$ (respectively, $\nu_{G/H}$) is indiscrete. By Lemma 2.3, $\gamma_{G/H}$ is indiscrete precisely when $\nu_{G/H}$ is indiscrete, and so H is γ_G -dense if and only if H is ν_G -dense. \square

2.2 The profinite topology vs the Bohr topology

In the following lemma we give known useful properties of the Bohr topology. Let G be an abelian group. For every $\chi \in G^*$ and $0 < \varepsilon \leq 1$, let

$$U_G(\chi, \varepsilon) = \{x \in G : |\chi(x)| < \varepsilon\}.$$

Then $\{U_G(\chi, \varepsilon) : 0 < \varepsilon \leq 1, \chi \in G^*\}$ is a subbase of the neighborhoods of 0 in (G, \mathcal{P}_G) .

Lemma 2.9. *Let G be an abelian group. Then:*

- (a) \mathcal{P}_G is precompact;
- (b) \mathcal{P} is hereditary and ideal;
- (c) $w(G, \mathcal{P}_G) = 2^{|G|}$;
- (d) every subgroup of G is \mathcal{P}_G -closed;
- (e) $d(G, \mathcal{P}_G) = |G|$.

Proof. (a) As noted in the introduction, \mathcal{P}_G is totally bounded. Moreover, since the characters $\text{Hom}(G, \mathbb{T})$ separate the points of G , it follows that \mathcal{P}_G is Hausdorff.

(b) Let H be a subgroup of G and $\pi : G \rightarrow G/H$ the canonical projection. Let $0 < \varepsilon \leq 1$ and consider $U_H(\chi, \varepsilon)$ for a character χ of H . Since \mathbb{T} is divisible, there exists an extension $\tilde{\chi}$ of χ to G . Then $U_H(\chi, \varepsilon) = H \cap U_G(\tilde{\chi}, \varepsilon)$, and this proves that \mathcal{P} is hereditary.

To prove that \mathcal{P} is ideal, note that $\pi : (G, \mathcal{P}_G) \rightarrow (G/H, (\mathcal{P}_G)_q)$ is open by definition of quotient topology. Since $(\mathcal{P}_G)_q$ is precompact, being the quotient topology of the precompact topology \mathcal{P}_G (see (b)), and since $\mathcal{P}_{G/H}$ is the finest precompact topology on G/H (as noted in the introduction), we can conclude that $\mathcal{P}_{G/H} \geq (\mathcal{P}_G)_q$. In particular, $id_{G/H} : (G/H, (\mathcal{P}_G)_q) \rightarrow (G/H, \mathcal{P}_{G/H})$ is open, and hence $\pi : (G, \mathcal{P}_G) \rightarrow (G/H, \mathcal{P}_{G/H})$ is open being composition of two open endomorphisms. This shows that \mathcal{P} is ideal.

(c) In view of a theorem by Comfort and Ross [5], $w(G, \mathcal{P}_G) = |\text{Hom}(G, \mathbb{T})|$; now applying a result by Kakutani [24] we have $|\text{Hom}(G, \mathbb{T})| = 2^{|G|}$.

(d) For every subgroup H of G , since \mathcal{P} is ideal by (a), on the quotient G/H we have $(\mathcal{P}_G)_q = \mathcal{P}_{G/H}$, which is Hausdorff by (b). Hence, H is \mathcal{P}_G -closed.

(e) If D is a \mathcal{P}_G -dense subset of G , then $\langle D \rangle$ is a \mathcal{P}_G -dense subgroup of G . By item (b) $\langle D \rangle$ is also \mathcal{P}_G -closed and so $\langle D \rangle = G$. In particular, $|D| = |G|$. \square

We compare now the profinite topology with the natural topology and the Bohr topology, starting from the relatively easier relation given by Lemma 2.10.

Lemma 2.10. *In the lattice \mathfrak{Ft} of all functorial topologies, $\gamma \leq \inf\{\nu, \mathcal{P}\}$.*

Proof. Let G be an abelian group. We have to prove that $\gamma_G \leq \inf\{\nu_G, \mathcal{P}_G\}$. Since every finite-index subgroup of G contains a subgroup of the form mG , one has always $\gamma_G \leq \nu_G$. Let $H \in \mathcal{C}(G)$. Since H has finite index in G and H is \mathcal{P}_G -closed by Lemma 2.9(d), we have that H is \mathcal{P}_G -open. Hence, $\gamma_G \leq \mathcal{P}_G$. \square

The following proposition is a fundamental step for the proof of Theorem 2.13, which is the main result of this section.

Proposition 2.11. *Let G be an abelian group. The following conditions are equivalent:*

- (a) $\gamma_G = \mathcal{P}_G$;
- (b) G is bounded;
- (c) G^* is bounded;
- (d) G^* is torsion;
- (e) ν_G is discrete.

Proof. The implications (b) \Rightarrow (c) \Rightarrow (d) and the equivalence (b) \Leftrightarrow (e) are obvious.

(a) \Rightarrow (d) By the assumption, for every $\chi \in G^*$, the basic neighborhood $U_G(\chi, 1/4)$ contains some $N \in \mathcal{C}(G)$. Then $N \subseteq \ker \chi$, and so $\ker \chi \in \mathcal{C}(G)$. Therefore, $mG \subseteq \ker \chi$ for some $m \in \mathbb{N}_+$, i.e., χ is torsion. Hence, G^* is torsion.

(d) \Rightarrow (c) For every $n \in \mathbb{N}_+$ the subgroup $F_n = G^*[n]$ of G^* is closed and $G^* = \bigcup_{n \in \mathbb{N}_+} F_n$ by our hypothesis (d). Since G^* is a compact abelian group, it satisfies the Baire category theorem. Thus, there exists $n \in \mathbb{N}_+$ such that F_n has non-empty interior, hence F_n is open. Since G^* is compact, F_n must have finite index in G^* . Therefore, there exists $m \in \mathbb{N}_+$ such that $mG^* \subseteq F_n$, so $mnG^* = 0$, i.e., G^* is bounded.

(c) \Rightarrow (b) Assume that $nG^* = 0$ for some $n \in \mathbb{N}_+$. To show that $nG = 0$, pick an element $x \in G$. Then $\chi(nx) = 0$ for every $\chi \in G^*$. It follows that $nx = 0$, as it is a well-known fact that the characters of a discrete abelian group separate the points of the group. Hence $nG = 0$.

(d) \Rightarrow (a) Let $\chi \in G^*$ and $0 < \varepsilon \leq 1$. Then $U_G(\chi, \varepsilon)$ contains $\ker \chi$. Since G^* is torsion, there exists $m \in \mathbb{N}_+$ such that $m\chi = 0$, that is, $m\chi(G) = 0$. Therefore, $G/\ker \chi \cong \chi(G)$ is finite, so $\ker \chi$ has finite index in G . Hence, $\ker \chi$ is open in (G, γ_G) . Since $\chi \in G^*$ was chosen arbitrarily, this shows that $\mathcal{P}_G \leq \gamma_G$. Lemma 2.10 applies to conclude that $\gamma_G = \mathcal{P}_G$. \square

The following corollary of Proposition 2.11 shows that the weight of the profinite topology of a bounded abelian group G has the maximal possible value $2^{|G|}$.

Corollary 2.12. *If G is an infinite bounded abelian group, then $w(G, \gamma_G) = 2^{|G|}$. In particular, (G, γ_G) is non-metrizable. More precisely, it does not contain infinite compact sets (so in particular, no convergent non-trivial sequences).*

Proof. By Lemma 2.9(b), $w(G, \mathcal{P}_G) = 2^{|G|}$. Moreover, $\gamma_G = \mathcal{P}_G$ by Proposition 2.11. To conclude, recall that the Bohr topology admits no infinite compact sets (see [22]). \square

We have seen in Proposition 2.11 that for bounded abelian groups the profinite topology coincides with the Bohr topology. In particular, this means that the profinite topology is the infimum of the Bohr topology and the natural topology, since $\mathcal{P}_G = \gamma_G \leq \nu_G = \delta_G$ for any bounded abelian group G . The next theorem shows that the equality $\gamma_G = \inf\{\nu_G, \mathcal{P}_G\}$ holds for every abelian group G .

Theorem 2.13. *In the lattice \mathfrak{Ft} of all functorial topologies, $\gamma = \inf\{\nu, \mathcal{P}\}$.*

Proof. Let G be an abelian group. We have to prove that $\gamma_G = \inf\{\nu_G, \mathcal{P}_G\}$.

Lemma 2.10 gives the inequality $\gamma_G \leq \inf\{\nu_G, \mathcal{P}_G\}$.

Let now U be a neighborhood of 0 in \mathcal{P}_G and $n \in \mathbb{N}_+$. We prove that the typical neighborhood $U + nG$ of 0 in $\inf\{\mathcal{P}_G, \nu_G\}$ is a neighborhood of 0 in γ_G too. To this end consider the canonical projection $\pi : G \rightarrow G/nG$. Since G/nG is bounded, $\gamma_{G/nG} = \mathcal{P}_{G/nG}$ by Proposition 2.11. Moreover, $\pi : (G, \mathcal{P}_G) \rightarrow (G/nG, \mathcal{P}_{G/nG})$ is open, as the Bohr topology is ideal by Lemma 2.9(b). Therefore, $\pi(U) \in \mathcal{P}_{G/nG} = \gamma_{G/nG}$. Then $H_1 \subseteq \pi(U)$ for some $H_1 \in \mathcal{C}(G/nG)$. Consequently $H = \pi^{-1}(H_1) \in \mathcal{C}(G)$ and in particular H is a neighborhood of 0 in γ_G . Since $U + nG = \pi^{-1}(\pi(U))$, it follows that $H \subseteq U + nG$, which proves that $U + nG$ is a neighborhood of 0 in (G, γ_G) . This concludes the proof. \square

Remark 2.14. (a) If G and H are infinite bounded abelian groups, then for every continuous map $f : (G, \gamma_G) \rightarrow (H, \gamma_H)$ with $f(0) = 0$ there exist a homomorphism $\phi : G \rightarrow H$ and an infinite subset A of G containing 0, such that $f \upharpoonright_A = \phi \upharpoonright_A$. In particular, if G is of exponent p and H is of exponent q , where p and q are distinct primes, then there exists no homeomorphisms between (G, γ_G) and (H, γ_H) , considered as topological spaces [18, 25].

(b) Item (a) should be compared with the fact that two countable metrizable abelian groups are always homeomorphic considered as *topological spaces*; for example, the p -adic and the q -adic topologies on \mathbb{Z} are homeomorphic. Also the compact spaces \mathbb{J}_p and \mathbb{J}_q , provided with their natural topology, are homeomorphic (to the Cantor cube $\{0, 1\}^\omega$). Let us note that in spite of this homeomorphism, there exists no non-zero homomorphism $\mathbb{J}_p \rightarrow \mathbb{J}_q$ (this should be compared to (a)).

3 The poset of finite-index subgroups

This section is dedicated to the general problem of studying the poset of finite-index subgroups of an abelian group. This problem is motivated by the definition of the adjoint algebraic entropy of an endomorphism ϕ of an abelian group G , where the family $\mathcal{C}(G)$ of all finite-index subgroups of G plays a prominent role. Now we recall the precise definition of adjoint algebraic entropy. For $N \in \mathcal{C}(G)$ and $n \in \mathbb{N}_+$, the n -th ϕ -cotrajectory of N is

$$C_n(\phi, N) = \frac{G}{N \cap \phi^{-1}(N) \cap \dots \cap \phi^{-n+1}(N)}.$$

The *adjoint algebraic entropy* of ϕ with respect to N is

$$H^*(\phi, N) = \lim_{n \rightarrow \infty} \frac{\log |C_n(\phi, N)|}{n};$$

This limit exists and it is finite [11]. The *adjoint algebraic entropy* of $\phi : G \rightarrow G$ is

$$\text{ent}^*(\phi) = \sup\{H^*(\phi, N) : N \in \mathcal{C}(G)\}.$$

In [11, Theorem 7.4] a dichotomy for the values of the adjoint algebraic entropy is proved; indeed, it can take only the values 0 and ∞ . Moreover, [30] is dedicated to the characterization of abelian groups of zero adjoint algebraic entropy and [21] to the connection of the adjoint algebraic entropy with the topological entropy.

3.1 Size and cofinality of $\mathcal{C}(G)$

The poset $\mathcal{C}(G)$ of finite-index subgroups of an abelian group G is trivial if and only if G is divisible. Moreover, Lemma 2.5 describes the case when $\mathcal{C}(G)$ is finite; the abelian groups with this property are those that we have called almost divisible. The next result from [11] characterizes narrow abelian groups, that are the abelian groups G with countable $\mathcal{C}(G)$.

Theorem 3.1. [11, Theorem 3.3] *Let G be an abelian group. Then the following conditions are equivalent:*

- (a) G is narrow;
- (b) $|\mathcal{C}(G)| < \mathfrak{c}$;
- (c) G/pG is finite for every prime p ;
- (d) G/mG is finite for every $m \in \mathbb{N}_+$;
- (e) $\mathcal{C}(G)$ contains a countable decreasing cofinal chain;

(f) $\gamma_G = \nu_G$.

Note that (e) is equivalent to pseudometrizable of (G, γ_G) . The remarkable dichotomy hidden behind the equivalence between (d) and (f) (i.e., $\mathcal{C}(G)$ is either countable or has size at least \mathfrak{c}), discovered in [11, Theorem 3.3] becomes clear below. It is due to the fact that $|\mathcal{C}(G)|$ coincides with the cardinality of the torsion part of a compact abelian group (namely, $|\mathcal{C}(G)| = |t(\text{Hom}(G, \mathbb{T}))|$).

The next lemma plays a key role in the proofs of the results of this section.

Lemma 3.2. *Let G be an abelian group that is not almost divisible. Then:*

- (a) $(G, \nu_G)^* = (G, \gamma_G)^* = t(\text{Hom}(G, \mathbb{T})) = t(\text{Hom}(G/G^1, \mathbb{T}))$;
- (b) $|\mathcal{C}(G)| = w(G, \gamma_G) = \chi(G, \gamma_G) = |t(\text{Hom}(G, \mathbb{T}))|$.

Proof. (a) To prove the first two equalities we intend to show the following chain of inclusions

$$(G, \nu_G)^* \supseteq (G, \gamma_G)^* \supseteq t(\text{Hom}(G, \mathbb{T})) \supseteq (G, \nu_G)^*. \quad (3.1)$$

Since $\gamma_G \leq \nu_G$, the first inclusion $(G, \gamma_G)^* \subseteq (G, \nu_G)^*$ is obvious.

To prove the inclusion $t(\text{Hom}(G, \mathbb{T})) \subseteq (G, \gamma_G)^*$, note that for every $\chi \in t(\text{Hom}(G, \mathbb{T}))$ there exists $m \in \mathbb{N}_+$ such that $m\chi = 0$, so $\chi(mG) = m\chi(G) = 0$. In particular, $\chi(G)$ is finite and so $\ker \chi \in \mathcal{C}(G)$. Thus, $\chi : (G, \gamma_G) \rightarrow \mathbb{T}$ is continuous.

To prove the inclusion $(G, \nu_G)^* \subseteq t(\text{Hom}(G, \mathbb{T}))$, fix a neighborhood U of 0 in \mathbb{T} that contains no non-zero subgroups. For every continuous character $\chi : (G, \nu_G) \rightarrow \mathbb{T}$ there exists $m \in \mathbb{N}_+$ such that $\chi(mG) \subseteq U$. By the choice of U this yields $\chi(mG) = 0$, i.e., $m\chi = 0$. Then $\chi \in t(\text{Hom}(G, \mathbb{T}))$.

From the chain of inclusions (3.1), we obtain that $(G, \nu_G)^* = (G, \gamma_G)^* = t(\text{Hom}(G, \mathbb{T}))$.

To prove the last equality, we identify first $\text{Hom}(G/G^1, \mathbb{T})$ with a subgroup of $\text{Hom}(G, \mathbb{T})$ using the canonical homomorphism $\pi : G \rightarrow G/G^1$. In fact, the adjoint homomorphism $\pi^* : \text{Hom}(G/G^1, \mathbb{T}) \rightarrow \text{Hom}(G, \mathbb{T})$ is injective, as π is surjective. After this identification, we note that the inclusion $t(\text{Hom}(G, \mathbb{T})) \supseteq t(\text{Hom}(G/G^1, \mathbb{T}))$ is clear. So let $\chi \in t(\text{Hom}(G, \mathbb{T}))$. Then there exists $m \in \mathbb{N}_+$ such that $m\chi = 0$; in particular, $\ker \chi \supseteq G^1$ and so χ can be factorized as $\chi = \bar{\chi} \circ \pi$, where $\bar{\chi} : G/G^1 \rightarrow \mathbb{T}$ is the character induced by χ . Then χ can be considered as a torsion character of G/G^1 , and we have proved that $t(\text{Hom}(G, \mathbb{T})) \subseteq \text{Hom}(G/G^1, \mathbb{T})$. Then $t(\text{Hom}(G, \mathbb{T})) = t(\text{Hom}(G/G^1, \mathbb{T}))$.

(b) As recalled above, $w(L) = \chi(L)$ for any totally bounded abelian group L . So $w(G, \gamma_G) = \chi(G, \gamma_G)$ by Fact 2.6(a). As noted in Lemma 2.3, $(G/G^1, \gamma_{G/G^1})$ is the Hausdorff reflection of (G, γ_G) . Therefore, $w(G/G^1, \gamma_{G/G^1}) = w(G, \gamma_G)$ and these groups have the same dual group. Then we may assume without loss of generality that $G^1 = 0$ and so that (G, γ_G) is precompact by Fact 2.6(b). Consequently, $w(G, \gamma_G) = |(G, \gamma_G)^*|$, as $w(L) = |L^*|$ for any precompact abelian group L [5]. To conclude, $(G, \gamma_G)^* = t(\text{Hom}(G, \mathbb{T}))$ by (a).

It remains to prove that $|\mathcal{C}(G)| = |t(\text{Hom}(G, \mathbb{T}))|$. Let $\mathcal{C}_c(G)$ be the subfamily of those $N \in \mathcal{C}(G)$, such that G/N is (finite) cyclic. Note that every $N \in \mathcal{C}(G)$ is a finite intersection of subgroups from $\mathcal{C}_c(G)$. Moreover, $\mathcal{C}(G)$ is infinite by the hypothesis that G is not almost divisible, hence $|\mathcal{C}(G)| = |\mathcal{C}_c(G)|$. So it remains to verify that $|\mathcal{C}_c(G)| = |t(\text{Hom}(G, \mathbb{T}))|$.

Clearly, every $N \in \mathcal{C}_c(G)$ gives rise to a character $\chi_N : G \rightarrow \mathbb{T}$ by considering the finite cyclic group G/N as a subgroup of \mathbb{T} . This defines an injective map $\mathcal{C}_c(G) \rightarrow t(\text{Hom}(G, \mathbb{T}))$ as the character χ_N is obviously torsion. In the opposite direction we just assign to every $\chi \in t(\text{Hom}(G, \mathbb{T}))$ its kernel $N = \ker \chi \in \mathcal{C}_c(G)$. Let us see next that the map $t(\text{Hom}(G, \mathbb{T})) \rightarrow \mathcal{C}_c(G)$ defined by $\chi \mapsto \ker \chi$ is finitely-many-to-one. This will prove that $|\mathcal{C}_c(G)| = |t(\text{Hom}(G, \mathbb{T}))|$, as $\mathcal{C}_c(G)$ is infinite. Let $\chi, \eta \in t(\text{Hom}(G, \mathbb{T}))$ with $\ker \chi = \ker \eta = N$. Then there exist injective homomorphisms $\chi_1, \eta_1 : G/N \rightarrow \mathbb{T}$, such that $\chi = \chi_1 \circ \pi$ and $\eta = \eta_1 \circ \pi$, where $\pi : G \rightarrow G/N$ is the canonical homomorphism. Let $m = |G/N|$, then both $\chi(G)$ and $\eta(G)$ coincide with the unique cyclic subgroup C of \mathbb{T} of order m . Hence, there exists an automorphism ξ of C , such that $\eta_1 = \xi \circ \chi_1$. Since the automorphism group of C is finite, one has only finitely many distinct pairs χ, η with $\ker \chi = \ker \eta = N$. \square

Passing to the general case, the following theorem gives a precise formula for the cardinality and the cofinality of the family $\mathcal{C}(G)$ for an infinite abelian group G when $|\mathcal{C}(G)| \geq \omega$.

In the sequel, for an infinite cardinal κ we let $\log \kappa = \min\{\lambda : 2^\lambda \geq \kappa\}$ the logarithm of κ .

Theorem 3.3. *Let G be an abelian group that is not almost divisible. Then*

$$|\mathcal{C}(G)| = \chi(G, \gamma_G) = w(G, \gamma_G) = \omega \cdot \sup\{2^{|G/p^G|} : p \text{ prime}\}. \quad (3.2)$$

Proof. By Lemma 3.2(b), $|\mathcal{C}(G)| = \chi(G, \gamma_G) = w(G, \gamma_G)$, so it remains to prove only the last equality in (3.2).

Assume that G is narrow. According to Theorem 3.1 this is equivalent to finiteness of G/pG for every prime p . Then $\omega \cdot \sup\{2^{|G/pG|} : p \text{ prime}\} = \omega$. By Theorem 3.1 and Lemma 3.2, G narrow is equivalent to $w(G, \gamma_G) = \omega$ and also to $\chi(G, \gamma_G) = \omega$. This proves the desired equality.

Assume now that G is not narrow, and let $\kappa = w(G, \gamma_G)$. By Theorem 3.1, $\kappa \geq \mathfrak{c}$. Since every finite-index subgroup of G contains a subgroup of the form mG , and since Fact 2.6(c) can be applied for every $m \in \mathbb{N}_+$ for the projection $\pi_m : G \rightarrow G/mG$, we deduce that $\kappa \geq w(G, \gamma_{G/mG})$ for every $m \in \mathbb{N}_+$. Therefore, $\kappa \geq \sup\{w(G, \gamma_{G/mG}) : m \in \mathbb{N}_+\}$. On the other hand, if for each $m \in \mathbb{N}_+$ one fixes a base \mathcal{B}_m of neighborhoods of 0 in $(G/mG, \gamma_{G/mG})$ of minimum cardinality, then the family of sets $\bigcup_{m \in \mathbb{N}_+} \{\pi_m^{-1}(B) : B \in \mathcal{B}_m\}$ forms a base of neighborhoods of 0 in (G, γ_G) of size $\leq \sup\{w(G/mG, \gamma_{G/mG}) : m \in \mathbb{N}_+\}$. This proves that

$$w(G, \gamma_G) = \sup\{w(G/mG, \gamma_{G/mG}) : m \in \mathbb{N}_+\}. \quad (3.3)$$

Since G is not narrow, by Theorem 3.1 there exists $m \in \mathbb{N}_+$ such that G/mG is infinite. For every $m \in \mathbb{N}_+$ such that G/mG is infinite, $w(G/mG, \gamma_{G/mG}) = 2^{|G/mG|}$ by Corollary 2.12 and

$$\text{there exists a prime } p \text{ dividing } m \text{ such that } |G/mG| = |G/pG|. \quad (3.4)$$

Indeed, obviously $|G/mG| \geq |G/pG|$ for every such prime p . Next we note first that if G/pG is infinite, then $|G/p^nG| = |G/pG|$ for every $n \in \mathbb{N}_+$. Moreover, if $m = p_1^{k_1} \dots p_s^{k_s}$ with distinct primes p_i , then $mG = \bigcap_{i=1}^s p_i^{k_i}G$, hence $G/mG \hookrightarrow \bigoplus_{i=1}^s G/p_i^{k_i}G$. Therefore, $|G/mG| \leq \sup_{i=1}^s |G/p_iG|$.

Hence, (3.3) implies the desired equality. \square

Remark 3.4. When G is not narrow, there is another proof of this theorem requiring a better knowledge of Pontryagin duality. According to Lemma 3.2(b), $|\mathcal{C}(G)| = w(G, \gamma_G) = \chi(G, \gamma_G) = |t(\text{Hom}(G, \mathbb{T}))|$, so it suffices to prove that $|t(\text{Hom}(G, \mathbb{T}))| = \omega \cdot \sup\{2^{|G/pG|} : p \text{ prime}\}$. Since $G^* = \text{Hom}(G, \mathbb{T})$ is compact,

$$|t(G^*)| = r(t(G^*)) = \sup\{r_p(G^*) : p \text{ prime}\} = \sup\{2^{r_p(G/pG)} : p \text{ prime}\} = \sup\{2^{|G/pG|} : p \text{ prime}\}. \quad (3.5)$$

The next-to-the-last equality uses the fact that $r_p(G^*) = 2^{r_p(G/pG)}$, which can be obtained as follows (see also [16, Proposition 3.3.15]): $r_p(G^*) = r_p(G^*[p])$ and the compact group $G^*[p]$ of exponent p is the annihilator of pG , so isomorphic to the dual of the group G/pG . Since $G/pG \cong \bigoplus_{r_p(G/pG)} \mathbb{Z}(p)$, we conclude that $G^*[p] \cong \mathbb{Z}(p)^{r_p(G/pG)}$. Hence $r_p(G^*) = 2^{r_p(G/pG)}$ in case these cardinals are infinite, otherwise $r_p(G^*) = r_p(G/pG)$. Since at least one of the cardinals G/pG is infinite (by the assumption that G is narrow and the fact that if some G/mG is infinite, then also some G/pG is infinite as well, see (3.4) in the above proof).

Corollary 3.5. *Let G be a residually finite abelian group. If G is not narrow, then:*

- (a) $w(G, \gamma_G) \geq 2^\mathfrak{c}$ if and only if there exists a prime p such that $|G/pG| \geq \log 2^\mathfrak{c}$;
- (b) if $|G/pG| < \log 2^\kappa$ for all primes p and an infinite cardinal κ , then $w(G, \gamma_G) < 2^\kappa$;
- (c) [under MA] $w(G, \gamma_G) = \mathfrak{c}$, if $|G/pG| < \mathfrak{c}$ for all primes p .

Proof. (a) If $|G/pG| \geq \log 2^\mathfrak{c}$ for some prime p , then obviously $2^{|G/pG|} \geq 2^\mathfrak{c}$, so $w(G, \gamma_G) \geq 2^\mathfrak{c}$ by Theorem 3.3. Assume that $w(G, \gamma_G) \geq 2^\mathfrak{c}$ for all primes p . Then obviously $w(G, \gamma_G) = \sup\{2^{|G/pG|} : p \text{ prime}\}$ by Theorem 3.3. Assume that all $|G/pG| < \log 2^\mathfrak{c}$. Then

$$2^{|G/pG|} < 2^\mathfrak{c} \text{ for all primes } p. \quad (3.6)$$

By our assumption, $w(G, \gamma_G)$ cannot coincide with any $2^{|G/pG|}$, hence $\text{cf}(w(G, \gamma_G)) = \omega$. Thus the equality $w(G, \gamma_G) = 2^\mathfrak{c}$ is not possible. Since (3.6) rules out the strict inequality $w(G, \gamma_G) > 2^\mathfrak{c}$, we are left with $w(G, \gamma_G) < 2^\mathfrak{c}$, a contradiction.

(b) As all $|G/pG| < \log 2^\kappa$, $2^{|G/pG|} < 2^\kappa$ for all primes p . Then $w(G, \gamma_G) \leq 2^\kappa \leq \log 2^\kappa$. If $w(G, \gamma_G) = 2^\kappa$, then it cannot coincide with any $2^{|G/pG|}$, so $\text{cf}(w(G, \gamma_G)) = \omega$, while $\text{cf}(2^\kappa) > \omega$, a contradiction.

(c) Follows from Theorem 3.1(b) and from the fact that MA yields $\log 2^\mathfrak{c} = \mathfrak{c}$. \square

The equality (3.5) suggests to consider also the density character $d(G, \gamma_G)$. Let us recall that $d(G, \mathcal{P}_G) = |G|$ as every subgroup of G is \mathcal{P}_G -closed. In order to compute $d(G, \gamma_G)$ we observe first that $d(G, \gamma_G) = d(G, \nu_G)$ by Lemma 2.8.

Theorem 3.6. *Let G be an infinite residually finite abelian group. Then*

$$\log |G| \leq w(G, \nu_G) = d(G, \gamma_G) = d(G, \nu_G) = \omega \cdot \sup\{|G/pG| : p \text{ prime}\}. \quad (3.7)$$

Proof. The equality $d(G, \gamma_G) = d(G, \nu_G)$ follows from item (b) of Lemma 2.8. To prove the last equality in (3.7) fix an $m \in \mathbb{N}_+$. Let D be a ν_G -dense subset of G of size $d(G, \nu_G)$. Then its image in G/mG under the canonical homomorphism $\pi : G \rightarrow G/mG$ is a dense subset of the discrete group G/mG , so $\pi(D) = G/mG$. Thus $d(G, \nu_G) = |D| \geq |G/mG|$. Since (G, ν_G) is an infinite Hausdorff group, $d(G, \nu_G) \geq \omega$. This proves the inequality $d(G, \nu_G) \geq \omega \cdot \sup\{|G/pG| : p \text{ prime}\}$.

To prove the opposite inequality, for every $m \in \mathbb{N}_+$ fix a subset D_m of G such that

$$D_m + mG = G \quad \text{and} \quad |D_m| = |G/mG|. \quad (3.8)$$

Then the subgroup H_m of G generated by D_m has size $|H_m| = \omega \cdot |G/mG|$. So for the subgroup $H = \sum_{m \in \mathbb{N}_+} H_m$ of G one has $|H| \leq \omega \cdot \sup\{|G/mG| : m \in \mathbb{N}_+\}$. As noted in (3.4), if G/mG is infinite, there exists a prime p dividing m such that $|G/mG| = |G/pG|$. Therefore, $|H| \leq \omega \cdot \sup\{|G/pG| : p \text{ prime}\}$. Since (3.8) obviously implies $H + mG = G$ for every $m \in \mathbb{N}_+$, H is ν_G -dense in G . This proves the last equality in (3.7).

The equality $w(G, \nu_G) = d(G, \nu_G)$ follows from the equalities $w(G, \nu_G) = d(G, \nu_G)\chi(G, \nu_G)$ and $\chi(G, \nu_G) = \omega$. The first inequality in (3.7) follows from the inequality $|G| \leq 2^{w(G, \nu_G)}$. \square

Remark 3.7. One can relax the condition ‘‘infinite residually finite’’ to non-almost divisible, as in the previous theorem and lemma. This will be enough to ensure $d(G, \nu_G) \geq \omega$ (then $d(G, \nu_G)$ will have an infinite Hausdorff quotient group, so that $d(G, \nu_G) \geq \omega$ will hold anyway).

Theorem 3.6 gives a precise value of the density character $d(G, \gamma_G) = d(G, \nu_G)$, but in many cases the cardinal given in (3.7) coincides with $|G|$ (as in the case of the Bohr topology). In the next example we show that these cardinals need not coincide in general and the gap may be as big as possible (i.e., $|G| = 2^{d(G, \gamma_G)}$).

Example 3.8. Let p be a prime. Let $B = \bigoplus_{n \in \mathbb{N}_+} \mathbb{Z}(p^n)$, $\bar{B} = \prod_{n \in \mathbb{N}_+} \mathbb{Z}(p^n)$ and $G = t(\bar{B})$. Then one can easily see that $p^n G + B = G$. This means that B is ν_G -dense in G (as the natural topology of G coincides with its p -adic one). Since B is countable, this proves that $d(G, \nu_G) < |G|$ (as G has size \mathfrak{c}).

3.2 When $\mathcal{C}(G)$ is cofinal in the poset $\mathcal{S}(G)$

Let us recall the next notion following [15]:

Definition 3.9. *An abelian group G is strongly non-divisible if no proper quotient of G is divisible.*

Clearly, an abelian group is strongly non-divisible if and only if each of its quotient is reduced.

In the next theorem we describe a subclass \mathfrak{S} of the class of all residually finite abelian groups defined by the property that $\mathcal{C}(G)$ is cofinal in the poset $\mathcal{S}(G)$ of all subgroups of G ordered by inclusion. This remarkable class can be described also by many other equivalent properties:

Theorem 3.10. *Let G be an abelian group. Then the following conditions are equivalent:*

- (a) every quotient of G is residually finite;
- (b) every subgroup of G is γ_G -closed;
- (c) every subgroup of G is ν_G -closed;
- (d) G is strongly non-divisible;
- (e) $n = r_0(G)$ is finite and for every subgroup $F \cong \mathbb{Z}^n$ of G one has $G/F \cong \bigoplus_p B_p$, where each B_p is a bounded abelian p -group;
- (f) $n = r_0(G)$ is finite and for every subgroup F of G of rank n one has $G/F \cong \bigoplus_p B_p$, where each B_p is a bounded abelian p -group;
- (g) $\mathcal{C}(G) \setminus \{G\}$ is cofinal in the poset $\mathcal{S}(G)$ of all subgroups of G .

Proof. We prove first the equivalence of (a), (b), (c) and (d).

(a) \Rightarrow (b) Let H be a subgroup of G . Then G/H is residually finite by hypothesis. By Lemma 2.3, $(G/H, \gamma_{G/H})$ is Hausdorff, so $\{0_{G/H}\}$ is $\gamma_{G/H}$ -closed in G/H . Since H is an inverse image of $\{0_{G/H}\}$ under the continuous canonical homomorphism $G \rightarrow G/H$, we conclude that H is γ_G -closed.

(b) \Leftrightarrow (c) Follows from Lemma 2.8(a).

(c) \Rightarrow (d) Assume that H is a proper subgroup of G . To prove that G/H is not divisible, one deduces from (c) that H is not ν_G -dense. So there exists $m \in \mathbb{N}_+$ such that $H + mG \neq G$. Hence $m(G/H) \neq G/H$.

(d) \Rightarrow (a) We prove first that (d) implies $G^1 = 0$. Let $0 \neq x \in G$. Using Zorn's Lemma one can find a subgroup H of G with $x \notin H$ and maximal with respect to this property. Then the non-zero element $\bar{x} = x + H$ of the quotient G/H is contained in every non-trivial subgroup of G/H . Hence H is quasicyclic, i.e., isomorphic to a subgroup of Prüfer's group $\mathbb{Z}(p^\infty)$ for some prime p . Since G is strongly non-divisible, G/H cannot be divisible, so it must be finite. Hence $H \in \mathcal{C}(G)$. This proves that $x \notin \bigcap_{H \in \mathcal{C}(G)} H = G^1$ by (2.1). Therefore, $G^1 = 0$.

Now suppose that H is a subgroup of G . Since the property (d) is obviously preserved by quotients, we conclude that G/H is strongly non-divisible as well, so we can apply the first part of this argument to conclude that G/H is residually finite.

Our next aim is to prove the chain of implications (d) \Rightarrow (e) \Rightarrow (f) \Rightarrow (g) \Rightarrow (d) that will establish the equivalence of (d), (e), (f) and (g) and will conclude the proof of the theorem.

(d) \Leftrightarrow (e) Was proved in [15]. Here we need only the implication (d) \Rightarrow (e). Since the reader may have no easy access to [15], we provide a complete proof of this implication. Assume for a contradiction that $r_0(G)$ is infinite and fix a free subgroup L of G of infinite rank. Then there exists a surjective homomorphism $f : L \rightarrow \mathbb{Q}$. Since \mathbb{Q} is divisible, one can extend f to a surjective homomorphism $G \rightarrow \mathbb{Q}$, a contradiction since G is strongly non-divisible. Fix any subgroup $F \cong \mathbb{Z}^n$ of G . Since $r_0(G) = r_0(F)$, the quotient G/F is torsion. Let $G/F \cong \bigoplus_p B_p$, where each F_p is an abelian p -group. Assume for a contradiction that B_p is not bounded for some prime p . Since G is strongly non-divisible, B_p is reduced. Fix a basic subgroup B of B_p . Since B_p is unbounded and reduced, this implies that the subgroup B is unbounded too. Since B is a direct sum of cyclic subgroups, we deduce that B contains a subgroup $H \cong \bigoplus_n \mathbb{Z}(p^n)$. Fix a surjective homomorphism $h : H \rightarrow \mathbb{Z}(p^\infty)$. Since $\mathbb{Z}(p^\infty)$ is divisible, one can extend h to a surjective homomorphism $G \rightarrow \mathbb{Z}(p^\infty)$, a contradiction since G is strongly non-divisible.

(e) \Rightarrow (f) It suffices to note that every subgroup F of G of rank n contains a subgroup isomorphic to \mathbb{Z}^n and that a quotient of a group of the form $\bigoplus_p B_p$ (with each B_p a bounded abelian p -group) has the same form.

(f) \Rightarrow (g) We shall check below that the property (f) is inherited by all quotients of G . Moreover, there is an (order preserving) bijection between the proper finite-index subgroups of G containing a given proper subgroup H of G and the poset $\mathcal{C}(G/H) \setminus \{G/H\}$. Therefore, to prove (f) \Rightarrow (g) it suffices to show that (f) implies $\mathcal{C}(G) \neq \{G\}$, i.e., G has a proper finite-index subgroup. Fix any subgroup $F \cong \mathbb{Z}^n$ of G such that the quotient G/F has the form $G/F = \bigoplus_p B_p$, where each B_p is a bounded abelian p -group. If $G/F \neq 0$, then at least one $B_p \neq 0$, so B_p has a proper finite-index subgroup N_p . Then the inverse image of the subgroup $N_p \oplus \bigoplus_{q \neq p} B_q$ of G/F under the canonical projection $G \rightarrow G/F$ is a proper finite-index subgroup of G . Now assume that $G/F = 0$. But this means that $G = F \cong \mathbb{Z}^n$, so G has plenty of proper finite-index subgroups.

Let us check now that the property (f) is inherited by all quotients of G . Let G/H be a quotient of G with $r_0(G/H) = m \leq n = r_0(G)$. Fix any subgroup $F_1 \cong \mathbb{Z}^m$ of G/H . We have to see that the quotient $(G/H)/F_1$ has the form $\bigoplus_p B_p^*$, where each B_p^* is a bounded abelian p -group. Let a_1, \dots, a_m be a set of free generators of F_1 . Pick $x_1, \dots, x_m \in G$ such that $x_i + H = a_i$ in G/H for $i = 1, \dots, m$. Then x_1, \dots, x_m form an independent subset of G of size m . Since $r_0(G) = n \geq m$, we can complete this independent set to a maximal independent subset $X = \{x_1, \dots, x_n\}$ of G and let $F = \langle x_1, \dots, x_n \rangle$. Then $F \cong \mathbb{Z}^n$ and the canonical homomorphism $G \rightarrow G/H$ takes F onto F_1 . Therefore $(G/H)/F_1$ is isomorphic to a quotient of G/F . By (f), $G/F = \bigoplus_p B_p$, where each B_p is a bounded abelian p -group. Hence, its quotients have the same form.

(g) \Rightarrow (d) Let H be a proper subgroup of G . Then there exists $N \in \mathcal{C}(G)$ such that $H \subseteq N$. Consequently, $N/H \in \mathcal{C}(G/H)$ and so G/H is not divisible. \square

4 Functorial topologies vs subcategories of Ab

We enrich here our supply of examples of functorial topologies.

Example 4.1. Let G be an abelian group.


- (a) The p -adic topology ν_G^p of G has the family of subgroups $\{p^n G : n \in \mathbb{N}_+\}$ as a base of the neighborhoods of 0.
- (b) The pro - p -topology γ_G^p has the family of all subgroups H of G with G/H a finite abelian p -group as a base of the neighborhoods of 0. Obviously, $\gamma_G = \sup_p \gamma_G^p$.

- (c) The *p-Bohr topology* \mathcal{P}_G^p is the initial topology of all homomorphisms $G \rightarrow \mathbb{Z}(p^\infty)$, where $\mathbb{Z}(p^\infty)$ is equipped with the topology inherited from \mathbb{T} .
- (d) A topological group G is called \aleph_0 -*bounded*, if for every non-empty open subset U of G there exists a countable subset A of G with $G = U + A$. The \aleph_0 -bounded groups were introduced and studied by Guran [23].

The class \mathcal{G} of all \aleph_0 -bounded groups contains the class of totally bounded groups and \mathcal{G} is stable under taking products, subgroups and quotients. In particular, every abelian group G admits a maximal \aleph_0 -bounded topology that we shall denote by \mathfrak{G}_G (for more properties of these topologies see [8] or [9]).

- (e) The *pro-countable topology* ϱ_G has all subgroups of countable index as a base of the neighborhoods of 0. Obviously, $\varrho \leq \mathfrak{G}$.

In the large lattice \mathfrak{Ft} of functorial topologies one has the following diagram.



For every prime p and an abelian group G consider the subset

$$\mathcal{C}_p(G) = \{N \in \mathcal{C}(G) : |G/N| \text{ is a power of } p\}$$

of $\mathcal{C}(G)$. In analogy to narrow abelian groups one can introduce the following notion.

Definition 4.2. For a prime p , an abelian group G is said to be *p-narrow* if $\mathcal{C}_p(G)$ is countable.

Clearly, the subsets $\mathcal{C}_p(G)$ generate $\mathcal{C}(G)$ in the sense that every $N \in \mathcal{C}(G)$ is a finite intersection of subgroups $N_p \in \mathcal{C}_p(G)$.

In analogy to the first Ulm subgroup, we introduce

$$G_p^1 = \bigcap_{n \in \mathbb{N}_+} p^n G.$$

Definition 4.3. Let p be a prime. An abelian group G is *residually p-finite* if G is isomorphic to a subgroup of a direct product of finite abelian p -groups.

It is easy to see that $G_p^1 = \bigcap_{N \in \mathcal{C}_p(G)} N$, so G is residually p -finite if and only if $G_p^1 = 0$.

Remark 4.4. The results of the previous sections can be “localized at p ”, i.e., for any abelian group G ,

- $\gamma_G^p = \mathcal{P}_G^p$ if and only if G is a bounded abelian p -group, if and only if $\nu_G^p = \delta_G$;
- $\gamma^p = \inf\{\nu^p, \mathcal{P}^p\} = \inf\{\nu^p, \gamma\} = \inf\{\mathcal{P}^p, \gamma\}$;
- G is p -narrow if and only if G/pG is finite, if and only if $\gamma_G^p = \nu_G^p$;
- if $\mathcal{C}_p(G)$ is infinite, then $|\mathcal{C}_p(G)| = \chi(G, \gamma_G^p) = w(G, \gamma_G^p) = 2^{|G/pG|}$;
- if G is infinite and residually p -finite, then $\log |G| \leq w(G, \nu_G^p) = d(G, \gamma_G^p) = d(G, \nu_G^p) = |G/pG|$.

4.1 The equalizer of two functorial topologies

Inspired by the results of the previous sections, we consider now functorial topologies in general. In particular, we construct classes of abelian groups starting from functorial topologies. The starting example is that of narrow abelian groups, described by Theorem 3.1 as those abelian groups for which the profinite topology coincides with the natural topology (see Example 4.6 below for more details).

Let \mathcal{T} and \mathcal{S} be functorial topologies. Define the *equalizer* of \mathcal{T} and \mathcal{S} as

$$\mathcal{E}(\mathcal{T}, \mathcal{S}) = \{G \in \mathbf{Ab} : \mathcal{T}_G = \mathcal{S}_G\}.$$

Obviously, $0 \in \mathcal{E}(\mathcal{T}, \mathcal{S})$.

The following are the basic properties of the equalizer.

Lemma 4.5. *Let \mathcal{T} and \mathcal{S} be functorial topologies. Then:*

- (a) $\mathcal{E}(\mathcal{T}, \mathcal{S})$ is stable under taking finite products;
- (b) if \mathcal{T} and \mathcal{S} are hereditary, then $\mathcal{E}(\mathcal{T}, \mathcal{S})$ is stable under taking subgroups;
- (c) if \mathcal{T} and \mathcal{S} are ideal, then $\mathcal{E}(\mathcal{T}, \mathcal{S})$ is stable under taking quotients.

Proof. (a), (b) and (c) follow directly respectively from Lemma 2.1(a), (c) and (d). □

So the equalizer of two functorial topologies is always stable under taking finite products, but it may fail to be stable under taking arbitrary products, as item (d) of Example 4.6 shows.

Note that the class $\mathcal{E}(\mathcal{T}, \delta)$ is precisely the class $\mathcal{C}(\mathcal{T})$ introduced and studied in [1]. Following the terminology used in [1], we say that a class of abelian groups is a *discrete class* if it is stable under isomorphic groups, finite products and subgroups; moreover, it is an *ideal discrete class* if it is stable also under quotients.

So Lemma 4.5 says that $\mathcal{E}(\mathcal{T}, \mathcal{S})$ is a discrete class if \mathcal{T} and \mathcal{S} are hereditary, and $\mathcal{E}(\mathcal{T}, \mathcal{S})$ is an ideal discrete class if \mathcal{T} and \mathcal{S} are hereditary and ideal.

In the sequel we see various examples of classes of abelian groups described as equalizers of pairs of functorial topologies. To this end we need some more notation. Let \mathcal{N} be the class of all narrow abelian groups and for every prime p let \mathcal{N}_p be the class of all p -narrow abelian groups. Since every $N \in \mathcal{C}(G)$ is a finite intersection of subgroups $N_p \in \mathcal{C}_p(G)$, we have that

$$\mathcal{N} = \bigcap_p \mathcal{N}_p.$$

Example 4.6. (a) By Theorems 3.1 and 2.13,

$$\mathcal{N} = \mathcal{E}(\gamma, \nu) = \mathcal{E}(\nu, \mathcal{P}).$$

Hence, the class \mathcal{N} is closed under taking quotients; indeed, both the profinite and the natural topology are ideal, and so Lemma 4.5(c) applies.

- (b) The class \mathcal{N} contains all divisible abelian groups D , since both γ_D and ν_D coincide with the indiscrete topology of D as noted in Example 2.7(b).
- (c) In view of item (b), \mathcal{N} is not closed under taking subgroups, as every abelian group is subgroup of a divisible abelian group. So it suffices to consider $G = \mathbb{Z}(p)^{(\mathbb{N})}$, where $\gamma_G < \nu_G = \delta_G$, and the divisible hull of G .
- (d) Moreover, \mathcal{N} is not stable under taking infinite products, as $\mathbb{Z} \in \mathcal{N}$, while $\mathbb{Z}^{\mathbb{N}} \notin \mathcal{N}$.
- (e) According to Proposition 2.11, $\mathcal{E}(\gamma, \mathcal{P}) = \{\text{bounded abelian groups}\}$.
- (f) By Remark 4.4, $\mathcal{N}_p = \mathcal{E}(\gamma^p, \nu^p) = \mathcal{E}(\nu^p, \mathcal{P}^p)$, and
- (g) $\mathcal{E}(\gamma^p, \mathcal{P}) = \{\text{bounded abelian } p\text{-groups}\}$.

Item (a) of the next corollary is precisely [1, Theorem 2.3].

Corollary 4.7. *Let \mathcal{T} be a functorial topology. Then the classes $\mathcal{E}(\mathcal{T}, \delta)$ and $\mathcal{E}(\mathcal{T}, \iota)$ are stable under isomorphisms and finite products. Moreover:*

- (a) $\mathcal{E}(\mathcal{T}, \delta)$ is stable under taking subgroups (i.e., it is a discrete class); if \mathcal{T} is ideal, then $\mathcal{E}(\mathcal{T}, \delta)$ is stable also under taking quotients (i.e., it is an ideal discrete class);

(b) $\mathcal{E}(\mathcal{T}, \iota)$ is stable under taking quotients; if \mathcal{T} is hereditary, then $\mathcal{E}(\mathcal{T}, \iota)$ is stable also under taking subgroups (i.e., it is an ideal discrete class).

Proof. The stability under taking finite products is given by Lemma 4.5(a), while the stability under isomorphisms is obvious.

(a) Let $G \in \mathcal{E}(\mathcal{T}, \delta)$ and let H be a subgroup of G . Then $\mathcal{T}_H \geq \mathcal{T}_G \upharpoonright_H$ by Lemma 2.1(c). Since $\mathcal{T}_G \upharpoonright_H = \delta_G \upharpoonright_H = \delta_H$, we can conclude that $\mathcal{T}_H = \delta_H$, and hence $H \in \mathcal{E}(\mathcal{T}, \delta)$. If \mathcal{T} is ideal, then $\mathcal{E}(\mathcal{T}, \delta)$ is stable under taking quotients by Lemma 4.5(c).

(b) Let $G \in \mathcal{E}(\mathcal{T}, \iota)$ and let H be a subgroup of G . Then $\mathcal{T}_{G/H} \leq (\mathcal{T}_G)_q$ by Lemma 2.1(d). Since $(\mathcal{T}_G)_q = (\iota_G)_q = \iota_H$, we have $\mathcal{T}_{G/H} = \iota_{G/H}$, that is $G/H \in \mathcal{E}(\mathcal{T}, \iota)$. If \mathcal{T} is hereditary, then $\mathcal{E}(\mathcal{T}, \iota)$ is stable under taking subgroups by Lemma 4.5(b). \square

Example 4.8. (a) $\mathcal{E}(\delta, \iota) = \mathcal{E}(\varrho, \iota) = \mathcal{E}(\mathfrak{G}, \iota)$ is the singleton class consisting of the trivial group $\{0\}$;

(b) $\mathcal{E}(\nu, \delta) = \{\text{bounded abelian groups}\}$;

(c) $\mathcal{E}(\gamma, \delta) = \mathcal{E}(\mathcal{P}, \delta) = \{\text{finite abelian groups}\}$;

(d) $\mathcal{E}(\nu, \iota) = \mathcal{E}(\gamma, \iota) = \{\text{divisible abelian groups}\}$;

(e) $\mathcal{E}(\mathfrak{G}, \delta) = \mathcal{E}(\varrho, \delta) = \{\text{countable abelian groups}\}$;

(f) $\mathcal{E}(\nu^p, \delta) = \{\text{bounded abelian } p\text{-groups}\}$;

(g) $\mathcal{E}(\gamma^p, \delta) = \mathcal{E}(\mathcal{P}^p, \delta) = \{\text{finite abelian } p\text{-groups}\}$;

(i) $\mathcal{E}(\nu^p, \iota) = \mathcal{E}(\gamma^p, \iota) = \{p\text{-divisible abelian groups}\}$.

(j) $\mathcal{E}(\mathcal{P}^p, \iota) = \{\text{torsion abelian groups without non-trivial } p\text{-torsion elements}\}$.

A careful look at the examples reveals the following connection between topologies and classes of abelian groups. Given the class \mathcal{B} of all bounded abelian groups, one can obtain the natural topology ν_G of an abelian group G as the group topology on G having as base of the neighbourhoods of 0 the family $\{N \leq G : G/N \in \mathcal{B}\}$. Analogously, if \mathcal{F} is the class of all finite abelian groups, the profinite topology γ_G of any abelian group G is the group topology on G which has as a base of the neighbourhoods of 0 the family $\{N \leq G : G/N \in \mathcal{F}\}$.

This remark is generalized in the next theorem. According to Corollary 4.7, the properties required for the class \mathcal{C} are necessary.

Theorem 4.9. *Let \mathcal{C} be a discrete class.*

(a) *The family $\mathcal{C}_G = \{N \leq G : G/N \in \mathcal{C}\}$ is a base of the neighborhoods of 0 of a linear group topology $\mathcal{T}_G^{\mathcal{C}}$ on G .*

(b) *Moreover, $\mathcal{T}^{\mathcal{C}}$ is a linear hereditary functorial topology with $\mathcal{E}(\mathcal{T}^{\mathcal{C}}, \delta) = \mathcal{C}$; and*

(c) *\mathcal{C} is an ideal discrete class if and only if $\mathcal{T}^{\mathcal{C}}$ is ideal.*

Proof. (a) Let $N_1, N_2 \in \mathcal{C}_G$. Then $G/N_1, G/N_2 \in \mathcal{C}$. Since \mathcal{C} is stable under taking finite products, $G/N_1 \times G/N_2 \in \mathcal{C}$. Consider now the embedding $G/N_1 \cap N_2 \hookrightarrow G/N_1 \times G/N_2$. Since \mathcal{C} is stable under taking subgroups, $G/N_1 \cap N_2 \in \mathcal{C}$, and hence $N_1 \cap N_2 \in \mathcal{C}_G$. This proves that \mathcal{C}_G is a local base of a group topology $\mathcal{T}_G^{\mathcal{C}}$ on G . Clearly, $\mathcal{T}_G^{\mathcal{C}}$ is linear.

(b) To verify that $\mathcal{T}^{\mathcal{C}}$ is a functorial topology, consider a homomorphism $\phi : (G, \mathcal{T}_G^{\mathcal{C}}) \rightarrow (G', \mathcal{T}_{G'}^{\mathcal{C}})$ of abelian groups. Then ϕ is continuous; in fact, if $N' \in \mathcal{C}_{G'}$, then $G'/N' \in \mathcal{C}$. For $N = \phi^{-1}(N')$ we have $G/N \cong \phi(G)/\phi(N) = \phi(G)/N' \cap \phi(G) \cong \phi(G) + N'/N'$. Since \mathcal{C} is stable under taking subgroups, $G/N \in \mathcal{C}$ and so $N \in \mathcal{C}_G$. The equality $\mathcal{E}(\mathcal{T}^{\mathcal{C}}, \delta) = \mathcal{C}$ follows from the definitions.

We show now that $\mathcal{T}^{\mathcal{C}}$ is hereditary. To this end let G be an abelian group and H a subgroup of G . Let $N \in \mathcal{C}_G$, that is, $G/N \in \mathcal{C}$. We have $H/H \cap N \cong H + N/N \leq G/N$. Since \mathcal{C} is stable under taking subgroups, $H/H \cap N \in \mathcal{C}$, that is, $H \cap N \in \mathcal{C}_H$. This proves that $\mathcal{T}_G^{\mathcal{C}} \upharpoonright_H = \mathcal{T}_H^{\mathcal{C}}$, i.e., $\mathcal{T}^{\mathcal{C}}$ is hereditary.

(c) Assume now that \mathcal{C} is stable under taking quotients. Let G be an abelian group, H a subgroup of G and consider the canonical projection $\pi : G \rightarrow G/H$. Let $N \in \mathcal{C}_G$, that is, $G/N \in \mathcal{C}$. We have that $(G/H)/\pi(N) = (G/H)/(N + H/H) \cong G/N + H$. Since \mathcal{C} is stable under taking quotients, $(G/H)/\pi(N) \in \mathcal{C}$, i.e., $\pi(N) \in \mathcal{C}_{G/H}$. This proves that $(\mathcal{T}_G^{\mathcal{C}})_q = \mathcal{T}_{G/H}^{\mathcal{C}}$, hence $\mathcal{T}^{\mathcal{C}}$ is ideal.

Now suppose that $\mathcal{T}^{\mathcal{C}}$ is ideal. Then $\mathcal{C} = \mathcal{E}(\mathcal{T}^{\mathcal{C}}, \delta)$ is stable under taking quotients by Lemma 4.5(c). \square

This procedure is standard in the field of functorial topologies, and it completely describes a large class of functorial topologies (for further details see [1, Theorems 2.5 and 2.6]).

In the next example we exhibit a linear functorial topology that is not ideal.

Example 4.10. Consider the class $\mathcal{C} = \{\mathbb{Z}^n : n \in \mathbb{N}\}$. More precisely, let \mathcal{C} be the class of all finite-rank free abelian groups. Obviously, \mathcal{C} is stable under isomorphic groups, finite products and subgroups. But \mathcal{C} is not stable under taking quotients. By Theorem 4.9, $\mathcal{T}^{\mathcal{C}}$ is a linear hereditary functorial topology which is not ideal.

4.2 Subcategories of \mathbf{Ab} as inverse images via functorial topologies

We showed above that many subcategories of \mathbf{Ab} arise as equalizers of two functorial topologies. Here we consider other ways to generate subcategories of \mathbf{Ab} using functorial topologies (e.g., via inverse images). To start with, let us note that the classes $\mathcal{E}(\mathcal{T}, \delta)$ and $\mathcal{E}(\mathcal{T}, \iota)$ can be obtained also in a different way. Namely, let \mathcal{A} be a class of topological abelian groups and let

$$\mathcal{A}(\mathcal{T}) = \{G \in \mathbf{Ab} : (G, \mathcal{T}_G) \in \mathcal{A}\}.$$

In other words, $\mathcal{A}(\mathcal{T})$ is the inverse image of the subcategory \mathcal{A} along the functor $\mathcal{T} : \mathbf{Ab} \rightarrow \mathbf{TopAb}$, hence it is natural to expect that nice stability properties of \mathcal{A} will entail nice stability properties of $\mathcal{A}(\mathcal{T})$:

Lemma 4.11. *Let \mathcal{T} be a functorial topology and let \mathcal{A} be a class of topological abelian groups stable under taking products, subgroups and finer group topologies. Then:*

- (a) if $G_i \in \mathcal{A}(\mathcal{T})$ for every $i \in I$, then $G = \prod_{i \in I} G_i \in \mathcal{A}(\mathcal{T})$;
- (b) if $G \in \mathcal{A}(\mathcal{T})$, then $H \in \mathcal{A}(\mathcal{T})$ for every subgroup H of G .

Proof. (a) Since $\mathcal{T}_G \geq \prod_{i \in I} \mathcal{T}_{G_i}$ by Lemma 2.1(b), and since $(G, \prod_{i \in I} \mathcal{T}_{G_i}) \in \mathcal{A}$ according to the hypothesis, it follows that $(G, \mathcal{T}_G) \in \mathcal{A}$.

(b) Since $\mathcal{T}_H \geq \mathcal{T}_G \upharpoonright_H$ by Lemma 2.1(c), we deduce that $(H, \mathcal{T}_H) \in \mathcal{A}$ in view of the hypothesis. \square

Moreover, this construction produces a reflection functor $A_{\mathcal{T}} : \mathbf{Ab} \rightarrow \mathcal{A}(\mathcal{T})$ associating to an abelian group G , its biggest quotient G/H in $\mathcal{A}(\mathcal{T})$ (i.e., such that $(G/H, \mathcal{T}_{G/H}) \in \mathcal{A}$).

We can apply this approach for example for

- (a) $\mathcal{A} = \mathcal{H}$ the class of Hausdorff topological groups, so $\mathcal{H}(\mathcal{T}) = \{G \in \mathbf{Ab} : \mathcal{T}_G \text{ is Hausdorff}\}$;
- (b) $\mathcal{A} = \Delta$ the class of all discrete groups, so $\Delta(\mathcal{T}) = \mathcal{E}(\mathcal{T}, \delta)$;
- (c) $\mathcal{A} = \mathfrak{I}$ the class of all indiscrete groups, so $\mathfrak{I}(\mathcal{T}) = \mathcal{E}(\mathcal{T}, \iota)$.

Example 4.12. Here we consider examples concerning only ν .

- (a) For the class \mathfrak{T} of all totally bounded abelian groups, $\mathfrak{T}(\nu) = \mathcal{N}$.
- (b) For the class \mathfrak{C} (respectively, \mathfrak{L}) of all compact (respectively, locally compact) abelian groups, Orsatti [29] characterized the classes $\mathfrak{C}(\nu)$ and $\mathfrak{L}(\nu)$.

In the case $\mathcal{A} = \mathcal{H}$ we give the following

Example 4.13. For $\mathcal{A} = \mathcal{H}$, we have:

- (a) $\mathcal{H}(\mathcal{P}) = \mathcal{H}(\varrho) = \mathcal{H}(\mathfrak{G}) = \mathbf{Ab}$;
- (b) $\mathcal{H}(\gamma) = \mathcal{H}(\nu) = \{\text{residually finite abelian groups}\}$ by Lemma 2.3;
- (c) $\mathcal{H}(\gamma^p) = \mathcal{H}(\nu^p) = \{\text{residually } p\text{-finite abelian groups}\}$ by Remark 4.4.

In this case the reflection functor $H_{\mathcal{T}} : \mathbf{Ab} \rightarrow \mathcal{H}(\mathcal{T})$ associates to an abelian group G , its biggest quotient G/H such that $(G/H, \mathcal{T}_{G/H})$ is Hausdorff. For example,

- (a') $H_{\mathcal{P}}(G) = G$;
- (b') $H_{\gamma} = H_{\nu} = G/G^1$;
- (c') $H_{\gamma^p} = H_{\nu^p} = G/G_p^1$.

5 Open problems and final remarks

The next question should be compared with item (a) of Remark 2.14.

Question 5.1. *Let $G = \bigoplus_{\omega_1} \mathbb{Z}(2)$ and $H = \bigoplus_{\omega_1} \mathbb{Z}(3)$. Are (G, \mathfrak{G}_G) and (H, \mathfrak{G}_H) homeomorphic as topological spaces? What about (G, ϱ_G) and (H, ϱ_H) ?*

Problem 5.2. *Find a sufficient condition for a functorial topology \mathcal{T} and a pair of infinite abelian groups G, H , so that whenever $f : (G, \mathcal{T}_G) \rightarrow (H, \mathcal{T}_H)$ is a homeomorphism with $f(0) = 0$ there exist a homomorphism $\phi : G \rightarrow H$ and an infinite subset A of G containing 0, such that $f \upharpoonright_A = \phi \upharpoonright_A$.*

According to item (a) of Remark 2.14, this is the case of the functorial topologies γ and \mathcal{P} when G and H are bounded abelian groups (see [7] for more details). Note that if the pair G, H from Question 5.1 has the “linearization property” described above with respect to \mathfrak{G} or ϱ , we obtain a negative answer to Question 5.1.

Question 5.3. *Is it true that for every functorial topology \mathcal{T} every infinite abelian group G such that (G, \mathcal{T}_G) is Hausdorff is necessarily zero-dimensional?*

The answer is obviously “Yes” for all linear topologies. Moreover, it is “Yes” for $\mathcal{T} = \mathcal{P}$ and this follows from results due to van Douwen [19] and Shakhmatov [31].

The next problem concerns compactness-like properties of the functorial topologies. The weakest possible one, namely total boundedness is present in the case of the Bohr topology \mathcal{P} . Nevertheless, it is known that (G, \mathcal{P}_G) is pseudocompact for no infinite abelian group G [6]. Motivated by this fact and item (b) of Example 4.12, we propose the following general question in the line of Example 4.12(b):

Problem 5.4. *Study the class $\mathcal{A}(\mathcal{T})$ for a functorial topology \mathcal{T} when \mathcal{A} is a class of compact-like abelian groups.*

Question 5.5. *Do there exist a functorial topology \mathcal{T} and an infinite abelian group G such that (G, \mathcal{T}_G) is Hausdorff and $|G| > 2^{d(G, \mathcal{T}_G)}$?*

Let us recall that there exist Hausdorff topological groups with $|G| = 2^{2^{d(G)}}$.

Motivated by item (e) of Example 4.8, one can ask whether the pair \mathfrak{G}, ϱ has an analogous behavior as the pair \mathcal{P}, γ . So we leave open the following

Problem 5.6. *Describe the precise relation between \mathfrak{G} and ϱ , and compute $\mathcal{E}(\mathfrak{G}, \varrho)$.*

According to Lemma 2.9, one has

$$|d(G, \mathcal{P}_G)| = |G| \text{ and } |w(G, \mathcal{P}_G)| = 2^{|G|} \quad (5.1)$$

for every abelian group G . These relations depend only on the cardinality of the abelian group G . Theorems 3.3 and 3.6 show that things change for the profinite topology, where the algebraic structure of the group starts to have some impact through the following condition, relevant for the computation of $d(G, \nu_G)$:

$$\text{the set of cardinals } \{|G/pG| : p \in P\} \text{ has no top element.} \quad (5.2)$$

Obviously, (5.2) implies that $d(G, \nu_G)$ is a limit cardinal with $\text{cf}(d(G, \nu_G)) = \omega$. When (5.2) fails, then $d(G, \nu_G) = |G/pG|$ for some prime p , and consequently $w(G, \nu_G) = 2^{d(G, \nu_G)}$ by Theorems 3.3 and 3.6. Since this may occur independently on the cofinality of $|G/pG|$ (that may also be countable), this shows that $\text{cf}(d(G, \nu_G)) = \omega$ does not imply (5.2). Therefore, $w(G, \gamma_G) = 2^{d(G, \gamma_G)}$ may occur quite often (for example, when (5.2) fails, but not only in that case). To give a more precise account on this, let us recall that for a limit cardinal λ one puts $2^{<\lambda} = \sup\{2^\mu : \mu < \lambda\}$. In these terms, $w(G, \gamma_G) = 2^{<d(G, \nu_G)}$ when (5.2) holds (so $d(G, \nu_G)$ is a limit cardinal with $\text{cf}(d(G, \nu_G)) = \omega$). Since the cardinal function $2^{<\lambda}$ is strongly dependent on the cardinal arithmetics, this leaves open the following general problem:

Problem 5.7. *Describe the precise relation between the cardinals $|G|$, $w(G, \gamma_G)$ and $d(G, \gamma_G)$ for an abelian group G .*

To give a more precise form of this problem, one can present it as a “realization problem”:

Problem 5.8. *Characterize all triples of infinite cardinals (λ, κ, μ) such that $\lambda = d(G, \gamma_G)$, $\kappa = w(G, \gamma_G)$ and $\mu = |G|$ for some abelian group G .*

Obviously, one has to impose $\lambda \leq \mu \leq 2^\lambda$ and $\lambda \leq \kappa \leq 2^\lambda$ on the triples (λ, κ, μ) .

As a first step, one may consider the problem of realization of single cardinals. While every infinite cardinal λ can be of the form $d(G, \gamma_G)$ for some abelian group G of size λ (take the free abelian group G of size λ , then $d(G, \gamma_G) = \lambda$ and $w(G, \gamma_G) = 2^\lambda$), a successor non-exponential cardinal cannot be realized as $w(G, \gamma_G)$, according to Theorem 3.3.

Next comes the realization of a pair of infinite cardinals (λ, κ) as a pair $(d(G, \gamma_G), w(G, \gamma_G))$ for some abelian group G . As already seen above, all pairs $(\lambda, 2^\lambda)$ with an arbitrary infinite cardinal λ are realizable. The ‘‘antipodal’’ condition $\kappa = \lambda$ is discussed in the next remark (the equality $w(G, \gamma_G) = d(G, \gamma_G)$ should be compared with Question 5.1).

Remark 5.9. Recall that a cardinal κ is a *strong limit cardinal* if $2^\lambda < \kappa$ for every cardinal $\lambda < \kappa$; obviously, under the assumption of the Generalized Continuum Hypothesis (GCH), all limit cardinals are strong limit cardinals. If κ is a limit cardinal, then κ is a strong limit cardinal if and only if $\kappa = 2^{<\lambda}$.

- (a) One can show that $w(G, \gamma_G)$ is a strong limit cardinal for a residually finite abelian group G if and only if $w(G, \gamma_G) = d(G, \gamma_G)$. In such a case its cofinality is countable. Similarly, if (5.2) holds true, then $d(G, \gamma_G)$ is an uncountable strong limit cardinal if and only if $w(G, \gamma_G) = d(G, \gamma_G)$. In particular, no strong limit cardinal κ of uncountable cofinality can be realized as $w(G, \gamma_G)$.
- (b) One can realize the pair (κ, κ) for every strong limit cardinal κ of countable cofinality.

We end this section and the paper noting, as suggested by the referee, that the notion of functorial topology need not be confined to discrete abelian groups. Indeed, a functorial topology on \mathbf{TopAb} can be defined as a functor $\mathcal{T} : \mathbf{TopAb} \rightarrow \mathbf{TopAb}$ such that $U\mathcal{T} = U$.

In other words, a functorial topology on \mathbf{TopAb} is a functor $\mathcal{T} : \mathbf{TopAb} \rightarrow \mathbf{TopAb}$ such that $\mathcal{T}(G, \tau) = (G, \mathcal{T}_{(G, \tau)})$ for any $(G, \tau) \in \mathbf{TopAb}$, where $\mathcal{T}_{(G, \tau)}$ denotes the topology on G , and $\mathcal{T}(\phi) = \phi$ for any continuous homomorphism ϕ in \mathbf{TopAb} .

Example 5.10. (a) If (G, τ) is a topological abelian group, the *Bohr modification* of τ is the topology $\tau^+ = \sup\{\tau' : \tau' \leq \tau, \tau' \text{ totally bounded}\}$; this topology is the finest totally bounded group topology on G coarser than τ . Actually, $\tau^+ = \inf\{\tau, \mathcal{P}_G\}$. So the functor $\mathcal{P} : \mathbf{TopAb} \rightarrow \mathbf{TopAb}$, defined on the objects of \mathbf{TopAb} by $\mathcal{P}(G, \tau) = (G, \tau^+)$, is a functorial topology. Since $\delta_G^+ = \mathcal{P}_G$, this functorial topology extends the functor of the Bohr topology from \mathbf{Ab} to \mathbf{TopAb} .

- (b) For a topological group (G, τ) , let τ_γ be the group topology on G having as a local base the τ -open finite-index subgroups of G . So the functor $\gamma : \mathbf{TopAb} \rightarrow \mathbf{TopAb}$, defined on the objects of \mathbf{TopAb} by $\gamma(G, \tau) = (G, \tau_\gamma)$, is a functorial topology. Since $(\delta_G)_\gamma = \gamma_G$, this functorial topology extends the functor of the profinite topology from \mathbf{Ab} to \mathbf{TopAb} .

Following the verifications in Section 2.2, it is possible to prove that the inequality $\gamma \leq \mathcal{P}$ still holds in this more general setting.

On the other hand, for a topological group (G, τ) , let τ_ν be the group topology on G having as a local base the τ -open subgroups of G of the form mG for some $m \in \mathbb{N}_+$. Note that $(\delta_G)_\nu = \nu_G$. The next example shows that there exists a topological abelian group (G, τ) such that the inequality $\tau_\gamma \leq \tau_\nu$ fails.

Example 5.11. Let p be a prime, $G = \mathbb{Z}(p)^\omega$ and let τ be the product topology on G . Then $\tau_\gamma = \tau$, while $\tau_\nu = \iota_G$.

Moreover, the following example shows that the map $\nu : \mathbf{TopAb} \rightarrow \mathbf{TopAb}$, defined on the objects of \mathbf{TopAb} by $\nu(G, \tau) = (G, \tau_\nu)$ is not a functorial topology, as it does not send continuous homomorphisms to continuous homomorphisms.

Example 5.12. Let p be a prime, $G = \bigoplus_{\mathbb{N}} \mathbb{Z}(p)$ and $H = \bigoplus_{2\mathbb{N}} \mathbb{Z}(p) \oplus \bigoplus_{2\mathbb{N}+1} \mathbb{Z}(p^2)$, so that $G \subseteq H$. Moreover, let $N = 0 \oplus \bigoplus_{2\mathbb{N}+1} \mathbb{Z}(p) \subseteq G$ and note that $N = pH$. Let τ be a non-discrete group topology on G such that N is τ -open in G . Let σ be the group topology on H defined imposing that the embedding $i : (G, \tau) \rightarrow (H, \sigma)$ is continuous and open, so that N is σ -open in H . We verify that $i : (G, \tau_\nu) \rightarrow (H, \sigma_\nu)$ is not continuous. In fact, τ_ν is indiscrete as τ is not discrete and mG is either 0 or G for every $m \in \mathbb{N}_+$, while $pH = N$ is a non-trivial proper subgroup of G and it is σ_ν -open.

Finally, analogous considerations can be done about the p -Bohr topology, the pro- p -topology and the p -adic topology.

Acknowledgements

It is a pleasure to thank the referee for her/his unusually careful and competent reading, for the numerous useful comments and for the constructive criticism.

References

- [1] D. Boyer and A. Mader, *Functorial topologies on abelian groups*, Rocky Mountain J. Math. **10** (1980) no. 4, 695–708.
- [2] B. Charles, *Méthodes topologiques en théorie des groupes abéliens*, Proceedings of the Colloquium on Abelian Groups (Tihany, 1963), pp. 29–42, Akad. Kiado, Budapest, 1964.
- [3] M. M. Clementino, D. Dikranjan and W. Tholen, *Torsion theories and radicals in normal categories*, J. Algebra **305** (2006) no. 1, 98–129
- [4] W. W. Comfort, *Topological groups*, In: Handbook of Set-theoretic Topology, K. Kunen and J. E. Vaughan, editors, Chapter 24, pages 1143–1263. North-Holland, Amsterdam 1984.
- [5] W. W. Comfort and K. A. Ross, *Topologies induced by groups of characters*, Fund. Math. **55** (1964), 283–291.
- [6] W. Comfort and V. Saks, *Countably compact groups and finest totally bounded topologies*, Pacific J. Math. **49** (1973), 33–44.
- [7] L. de Leo and D. Dikranjan, *Straightening theorem for bounded abelian groups*, Topol. And Its Appl. **155** (2008), 2158–2176.
- [8] L. de Leo and M. Tkachenko, *The maximal ω -narrow group topology on abelian groups*, Houston J. Math. **36** (2010) no. 1, 215–227.
- [9] D. Dikranjan, *Functorial topologies on abelian groups*, in: Proceedings SUNLAG 2000, pp. 81–95, Caserta, 2001.
- [10] D. Dikranjan, *The lattice of group topologies and compact representations*, Nuclear groups and Lie groups. Current result and problems, Proc. Workshop on Lie groups and topological groups, Madrid, 1999, (Univ. Complutense), pp. 105–126.
- [11] D. Dikranjan, A. Giordano Bruno and L. Salce, *Adjoint algebraic entropy*, J. of Algebra **324** (2010), 442–463.
- [12] D. Dikranjan and E. Giuli, *Closure operators I*, Topology Appl. **27** (1987), 129–143.
- [13] D. Dikranjan and E. Giuli, *Factorizations, injectivity and compactness in categories of modules*, Comm. Algebra **19** (1) (1991), 45–83.
- [14] D. Dikranjan, B. Goldsmith, L. Salce and P. Zanardo, *Algebraic entropy for abelian groups*, Trans. Amer. Math. Soc. **361** (2009), 3401–3434.
- [15] D. Dikranjan and I. Prodanov, *A class of compact abelian groups*, Annuaire Univ. Sofia, Fac. Math. Méc. **70** (1975/76), 191–206.
- [16] D. Dikranjan, I. Prodanov and L. Stoyanov, *Topological Groups: Characters, Dualities and Minimal Group Topologies*, Pure and Applied Mathematics, Vol. **130**, Marcel Dekker Inc., New York-Basel, 1989.
- [17] D. Dikranjan and W. Tholen *Categorical Structure of Closure Operators with Applications to Topology, Algebra and Discrete Mathematics, Mathematics and its Applications*, vol. 346, Kluwer Academic Publishers, Dordrecht-Boston-London 1995.
- [18] D. Dikranjan and S. Watson, *Solution to van Douwen’s problem on Bohr topologies*, Jour. Pure Appl. Algebra **163** (2001) no. 2, 147–158.
- [19] E. K. van Douwen, *The maximal totally bounded group topology on G and the biggest minimal G -space for Abelian groups G* , Topology Appl. **34** (1990), 69–91.
- [20] L. Fuchs, *Infinite abelian Groups*, Vol. I and II, Academic Press, 1970 and 1973.
- [21] A. Giordano Bruno, *Adjoint entropy vs Topological entropy*, submitted.
- [22] I. Glicksberg, *Uniform boundedness for groups*, Canad. J. Math. **14** (1962), 269–276.
- [23] I. Guran, *On topological groups close to being Lindelöf*, Soviet Math. Dokl. **23** (1981), 173–175.
- [24] S. Kakutani, *On cardinal numbers related with a compact abelian group*, Proc. Imperial Acad. Tokyo **19** (1943), 366–372.
- [25] K. Kunen, *Bohr topology and partition theorems for vector spaces*, Topology Appl. **90** (1998), 97–107.
- [26] K. Kunen and W. Rudin, *Lacunarity and the Bohr topology*, Math. Proc. Cambridge Philos. Soc. **126** (1999) no. 1, 117–137.

- [27] A. Mader, *Basic concepts of functorial topologies*, abelian group theory (Oberwolfach, 1981), pp. 251–271, Lecture Notes in Math. **874**, Springer, Berlin–New York, 1981.
- [28] A. Mader and R. Mines, *Functorial topologies with totally ordered neighborhood bases*, Arch. Math. (Basel) **34** (1980) no. 3, 272–275.
- [29] A. Orsatti, *Una caratterizzazione dei gruppi abeliani compatti o localmente compatti nella topologia naturale*, Rend. Sem. Mat. Univ. Padova **39** (1967), 219–225.
- [30] L. Salce and P. Zanardo, *Abelian groups of zero adjoint entropy*, submitted.
- [31] D. Shakhmatov, *On zero-dimensionality of subgroups of locally compact groups*, Comment. Math. Univ. Carolin. **32** (1991) no. 3, 581–582.
- [32] S. Shkarin, *An infinite-dimensional separable pre-Hilbert space that is not homeomorphic to a closed hyperplane of it*, Izv. Ross. Akad. Nauk Ser. Mat. **63** (1999) no. 6, 209–219.
- [33] A. Weil, *Sur les Espaces a Structure Uniforme et sur la Topologie Generale*, Publ. Math. Univ. Strasbourg. Hermann & Cie., Paris, 1937.
- [34] M. D. Weiss, *Algebraic and other entropies of group endomorphisms*, Math. Systems Theory **8** (1974/75) no. 3, 243–248.