

Article

Antimicrobial Resistance and Molecular Epidemiology of *Staphylococcus aureus* from Hunters and Hunting Dogs

Vanessa Silva ^{1,2,3,4,*} , Manuela Caniça ^{5,6} , Vera Manageiro ^{5,6} , Madalena Vieira-Pinto ^{7,8} , José Eduardo Pereira ^{1,7,8}, Luís Maltez ^{1,7,8} , Patrícia Poeta ^{1,4,7,8,*}  and Gilberto Igrejas ^{2,3,4,t} 

- ¹ Microbiology and Antibiotic Resistance Team (MicroART), Department of Veterinary Sciences, University of Trás-os-Montes and Alto Douro (UTAD), 5000-801 Vila Real, Portugal; jeduardo@utad.pt (J.E.P.); lmaltez@utad.pt (L.M.)
 - ² Department of Genetics and Biotechnology, University of Trás-os-Montes and Alto Douro (UTAD), 5000-801 Vila Real, Portugal; gigrejas@utad.pt
 - ³ Functional Genomics and Proteomics Unit, University of Trás-os-Montes and Alto Douro (UTAD), 5000-801 Vila Real, Portugal
 - ⁴ LAQV-REQUIMTE, Department of Chemistry, NOVA School of Science and Technology, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal
 - ⁵ National Reference Laboratory of Antibiotic Resistances and Healthcare Associated Infections (NRL-AMR/HAI), Department of Infectious Diseases, National Institute of Health Dr. Ricardo Jorge, Av. Padre Cruz, 1649-016 Lisbon, Portugal; manuela.canica@insa.min-saude.pt (M.C.); vera.manageiro@insa.min-saude.pt (V.M.)
 - ⁶ Centre for the Studies of Animal Science, Institute of Agrarian and Agri-Food Sciences and Technologies, Oporto University, 4051-401 Oporto, Portugal
 - ⁷ CECAV—Veterinary and Animal Research Centre, University of Trás-os-Montes and Alto Douro (UTAD), 5000-801 Vila Real, Portugal; mmvpinto@utad.pt
 - ⁸ Associate Laboratory for Animal and Veterinary Science (AL4Animals), University of Trás-os-Montes and Alto Douro (UTAD), 5000-801 Vila Real, Portugal
- * Correspondence: vanessasilva@utad.pt (V.S.); ppoeta@utad.pt (P.P.)
† These authors have contributed equally to this work.



Citation: Silva, V.; Caniça, M.; Manageiro, V.; Vieira-Pinto, M.; Pereira, J.E.; Maltez, L.; Poeta, P.; Igrejas, G. Antimicrobial Resistance and Molecular Epidemiology of *Staphylococcus aureus* from Hunters and Hunting Dogs. *Pathogens* **2022**, *11*, 548. <https://doi.org/10.3390/pathogens11050548>

Academic Editors: Francesca Paola Nocera and Patrizia Nebbia

Received: 27 March 2022

Accepted: 29 April 2022

Published: 6 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Several studies have showed that a dog-to-human transmission of *Staphylococcus aureus* occurs. Hunting dogs do not have as much contact with their owners as dogs that live in the same household as the owners; however, these dogs have contact with their owners during hunting activities as well as when hunting game; therefore, we aimed to isolate *S. aureus* from hunters and their hunting dogs to investigate a possible *S. aureus* transmission. Nose and mouth samples were collected from 30 hunters and their 78 hunting dogs for staphylococcal isolation. The species identification was performed using MALDI-TOF. The antimicrobial susceptibility profiles were accessed using the Kirby–Bauer method and respective antimicrobial resistance genes were investigated by PCR. Multilocus sequence typing (MLST) and *spa*- and *agr*-typing was performed in all *S. aureus* isolates. *S. aureus* were detected in 10 (30%) human samples and in 11 (15.4%) dog samples of which 11 and 5 were methicillin-resistant *S. aureus* (MRSA). Other staphylococci were identified, particularly, *S. pseudintermedius*. Most *S. aureus* isolates were resistant to penicillin, erythromycin, and tetracycline. Evidence of a possible transmission of *S. aureus* between human and dogs was detected in three hunters and their dogs. *S. aureus* isolates were ascribed to 10 STs and 9 *spa*-types. A moderate colonization of *S. aureus* in hunting dogs and their owners was detected in this study. A few dog-to-dog and dog-to-human possible transmissions were identified.

Keywords: *Staphylococcus aureus*; MRSA; transmission; dogs; human-to-dog

1. Introduction

Staphylococci are natural colonizers of humans and some animal species. *S. aureus* and *S. epidermidis* are the most frequent colonizers of human skin and mucous membranes [1]. Approximately 30% of the human population are asymptomatic carriers of *S. aureus* [2].

Humans colonized by *S. aureus* are at higher risk of subsequent infection, both nosocomial and community-acquired [3]. Indeed, although considered a commensal organism, *S. aureus* is an opportunist pathogen that can cause a wide range of diseases ranging from mild skin infections to severe and potentially fatal ones [4]. Methicillin-resistant *S. aureus* (MRSA) has been first described in 1961 and has become a priority pathogen causing infections increasingly difficult to treat [5]. Despite a downward trend in the prevalence of MRSA in the EU, 25% of European countries continue to have a rate of invasive isolates above 25% [6,7]. Methicillin-resistance is driven by the acquisition of the *mec* genes (*mecA*, *mecB* or *mecC*) which encodes the penicillin-binding protein 2a (PBP2a) with a low affinity for β -lactam antibiotics [8,9]. Healthcare-associated (HA) and community-associated (CA)-MRSA strains have emerged and spread widely [10]. Although other studies have reported a linkage between animal and human MRSA, it was not until 2005 that the first case of transmission between humans and animals was demonstrated [11,12]. In that study, the farmers and one pig were colonized by a MRSA strain different from those usually found in HA-MRSA and CA-MRSA [12].

The role of animals in the spread and transmission of MRSA strains in the human community is not well understood yet; however, several studies conducted with farm workers, pet owners, and veterinarians, who are at greater risk of being colonized or infected by MRSA, show epidemiological evidence that suggests MRSA transmission between human and animal hosts occur in both directions [13–17]. MRSA transmission between humans and their pets may be more favored due to intimate contact and sharing of the same household [18]. Furthermore, studies have reported infections in pet owners caused by methicillin-resistant staphylococci from pets and vice versa, which indicates that MRSA colonization might also represent a potential health risk for both humans and animals [19,20]. *S. aureus* is not generally considered part of the normal flora of dogs but it can be found in dogs at rates between 5% and 10% [21–23]. Instead, *S. pseudintermedius* predominates in dogs. It can also colonize humans at very low frequencies and usually dog owners [19]. Nevertheless, HA-MRSA and CA-MRSA lineages have been increasingly identified in dogs and cats [24,25]. Close interactions between dogs and their owners creates favorable conditions for MRSA transmission; however, unlike other dogs, hunting dogs do not live indoors with the owners, as they are primarily used in hunting activities. Nevertheless, hunting dogs have direct contact with the natural environment and with game species. Thus, in this study, we intend to analyze the possible transmission of *S. aureus* between hunting dogs and their hunting owners, as well as to verify if the clonal lineages of the isolates are related to strains frequently found in the environment and wild animals. For this, *S. aureus*, MRSA, and other methicillin-resistant staphylococci were isolated from hunting dogs and hunters, and the isolates were analyzed for their antibiotic resistance profile, virulence, and clonal lineages.

2. Material and Methods

2.1. Samples and Bacterial Isolates

From August to December 2019, a total of 108 samples were collected from 30 hunters and their 78 hunting dogs. Samples were collected using a nasal and oral swab (one swab per individual). All hunters were males, and the dogs' ages, sex and breed were variable and are shown in Table S1. Swabs were inoculated into Brain Heart Infusion (BHI) broth containing 6.5% NaCl and incubated at 37 °C for 24 h. An aliquot of 100 μ L was then seeded onto Baird–Parker agar and oxacillin resistance screening agar base (ORSAB) plates supplemented with oxacillin (2 mg/L) and incubated at 37 °C for 24–48 h. Presumptive *S. aureus* and MRSA colonies were selected and further identified. In cases where MRSA was found, samples from dogs and/or dog owners that shared the same household were screened for the presence of other methicillin-resistant staphylococci species. Confirmation of staphylococci species was performed using Matrix-Assisted Laser Desorption/Ionization Time-of-Flight (MALDI-TOF).

2.2. Antimicrobial Susceptibility Testing and Resistance Genes

For all staphylococci identified, susceptibility to penicillin (1 U), cefoxitin (30 µg), gentamicin (10 µg), tobramycin (10 µg), tetracycline (30 µg), chloramphenicol (30 µg), erythromycin (15 µg), clindamycin (2 µg), ciprofloxacin (5 µg), linezolid (10), and trimethoprim-sulfamethoxazole (1.25 + 23.75 µg) was examined using the Kirby–Bauer disk-diffusion method. The results were analyzed according to the EUCAST 2018 guidelines, except for kanamycin, which followed the CLSI 2017 guidelines [26,27]. The reference strain *S. aureus* ATCC 25923 was used for quality control.

According to the phenotypic resistance profile, each isolate was screened for the presence of the following antimicrobial resistance genes by PCR, as previously described [28]: *bla_Z* and *mecA* (β-lactam resistance), *aac(6′)-aph(2′′)*, *aph(3′)-IIIa*, *ant(4′)-Ia* and *str* (aminoglycosides), *ermA*, *ermB*, *ermC*, *ermT*, *msr(A/B)*, *mphC*, *lnuA*, *lnuB*, *vgaA* and *vgaB* (macrolides and lincosamide), *tetK*, *tetM*, *tetL* and *tetO* (tetracycline) and, *fexA*, *fexB*, *cat_{pC194}*, *cat_{pC221}* and *cat_{pC223}* (chloramphenicol).

S. aureus isolates were screened for genes encoding virulence factors: hemolysins (*hla*, *hlb* and *hld*), exfoliative toxins (*eta* and *etb*), the leucocidin *lukS/F-PV*, and the toxic-shock syndrome toxin (*tst*) [29–31]. Additionally, the detection of the immune evasion cluster (IEC) system genes (*scn*, *chp*, *sak*, *sea* and *sep*) was also performed in *S. aureus* isolates which enabled the classification into different IEC types [32]. Finally, the presence of the virulence genes *lukS/F-I* and *siet* was investigated in all *S. pseudintermedius* isolates [33,34].

2.3. Molecular Typing in *S. aureus* Isolates

S. aureus isolates were typed by multilocus sequence typing (MLST) and *spa*-, *agr*- and *SCCmec*-typing. MLST was performed by amplifying and sequencing the amplicons of 7 housekeeping genes as previously described [35]. Isolates were subjected to *spa*-typing as previously described, and sequences were analyzed using Ridom Staph-Type software (version 1.5, Ridom GmbH, Wurzburg, Germany) [36]. The *agr* type of all *S. aureus* isolates was determined by the PCR as described in other studies [37]. Finally, all MRSA were characterized by *SCCmec* typing (I–V) using specific primers [38].

3. Results

A total of 108 samples (30 hunters and 78 hunting dogs) were analyzed in this study. From these samples, 21 (19.4%) *S. aureus* were isolated, of which 11 (52.4%) were MRSA. *S. aureus* were detected in 10 (30%) human samples and in 11 (15.4%) dog samples. Regarding the 11 MRSA isolates, 4 were isolated from humans and 7 from dogs. In cases when a hunter or a dog tested positive for *S. aureus* or MRSA, the remaining dogs and/or hunter living in the same household were screened for methicillin-resistant staphylococci. A total of 15 MRS were isolated from humans and dogs, namely, *S. pseudintermedius* (n = 5), *S. lentus* (n = 6), *S. sciuri* (n = 2), *S. cohnii*, and *S. vitulinus*. *S. lentus* were isolated from five dogs and one hunter who co-carried a MRSA strain (Table 1). *S. pseudintermedius* were isolated only from dogs, of which three co-carried MRSA and one co-carried *S. aureus*. *S. sciuri* and *S. cohnii* were isolated only from dog samples.

Table 1. Genetic characterization of the *S. aureus*, MRSA and MRS isolates grouped by hunter and the respective dogs.

	Isolate	Host	Species	Molecular Typing				Antimicrobial Resistance		Virulence Factors	
				ST (CC)	<i>spa</i>	SCC <i>mec</i>	<i>agr</i>	Phenotype	Genotype	IEC system	Other genes
Case 1	VS3182	Hunter 1	<i>S. aureus</i>	7353	t10042	-	NT	PEN	<i>blaZ</i>	-	<i>hla, hlb, hld</i>
	VS3183	Dog 1	<i>S. aureus</i>	7353	t10042	-	NT	PEN	<i>blaZ</i>	E	<i>hla, hld</i>
	VS3184	Dog 2	<i>S. aureus</i>	7353	t10042	-	NT	PEN	<i>blaZ</i>	E	<i>hla, hld</i>
	VS3185	Dog 2	<i>S. pseudintermedius</i>	-	-	-	-	PEN, FOX	<i>mecA</i>		<i>lukS/F-I, siet</i>
	VS3186	Dog 3	<i>S. cohnii</i>	-	-	-	-	PEN	<i>mecA</i>		
Case 2	VS3187	Hunter 2	<i>S. aureus</i>	30 (30)	t012	-	III	Susceptible	-	E	<i>hla, hld, tst</i>
	VS3188	Dog 1	<i>S. aureus</i>	30 (30)	t012	-	III	Susceptible	-	-	<i>hla, hlb, hld, tst</i>
Case 3	VS3189	Hunter 3	<i>S. aureus</i>	9 (9)	t2922	N.T.	II	PEN, FOX, CN, TOB, KAN, CD, TET, FD	<i>mecA, blaZ, aac(6')-Ie-aph(2'')-Ia, aph(3')-IIIa, lnuB, vgaA, tetM, tetK</i>	-	<i>hla, hlb, hld</i>
	VS3190	Dog 1	<i>S. aureus</i>	9 (9)	t2922	N.T.	II	PEN, FOX, CN, TOB, KAN, CD, TET, FD, C	<i>mecA, blaZ, aac(6')-Ie-aph(2'')-Ia, aph(3')-IIIa, vgaA, lnuB, tetM, tetK, cat_{p221}</i>	-	<i>hla, hlb, hld</i>
	VS3191	Dog 2	<i>S. aureus</i>	9 (9)	t2922	N.T.	II	PEN, FOX, CN, TOB, KAN, CD, TET, FD, C	<i>mecA, blaZ, aac(6')-Ie-aph(2'')-Ia, aph(3')-IIIa, vgaA, lnuB, tetM, tetK, cat_{p221}</i>	-	<i>hla, hlb, hld</i>
	VS3192	Dog 5	<i>S. lentus</i>	-	-	-	-	PEN	<i>mecA</i>	-	
Case 4	VS3193	Hunter 4	<i>S. aureus</i>	8 (8)	t121	IV	I	PEN, FOX	<i>mecA, blaZ,</i>	-	<i>hla, hlb, hld</i>
	VS3194		<i>S. lentus</i>	-	-			PEN	<i>mecA</i>		
	VS3195	Dog 1	<i>S. aureus</i>	5 (5)	t179	IV	II	PEN, FOX, ERY	<i>mecA, blaZ ermC,</i>	E	<i>hla, hld</i>
	VS3196		<i>S. pseudintermedius</i>	-	-	-	-	PEN, FOX	<i>mecA, blaZ</i>	-	<i>lukS/F-I</i>
	VS3197	Dog 2	<i>S. aureus</i>	5 (5)	t179	IV	II	PEN, FOX, ERY	<i>mecA, blaZ, ermC,</i>	E	<i>hla, hld</i>
	VS3198		<i>S. pseudintermedius</i>	-	-	-	-	PEN, FOX, CN, TOB, KAN	<i>mecA, blaZ, aac(6')-Ie-aph(2'')-Ia, aph(3')-IIIa, str</i>	-	<i>lukS/F-I, siet</i>

Table 1. Cont.

	Isolate	Host	Species	Molecular Typing				Antimicrobial Resistance		Virulence Factors	
				ST (CC)	<i>spa</i>	SCC <i>mec</i>	<i>agr</i>	Phenotype	Genotype	IEC system	Other genes
	VS3199	Dog 3	<i>S. aureus</i>	5 (5)	t179	IV	II	PEN, FOX, ERY	<i>mecA, blaZ</i>	E	<i>hla, hld</i>
	VS3200		<i>S. pseudintermedius</i>	-	-	-	-	PEN, FOX	<i>mecA, blaZ</i>	-	<i>lukS/F-I, siet</i>
Case 5	VS3201	Dog 2	<i>S. aureus</i>	718	t11333	IV	II	PEN, FOX, ERY, CD	<i>mecA, blaZ, ermC,</i>	E	<i>hld</i>
	VS3202	Dog 3	<i>S. aureus</i>	718	t11333	IV	II	PEN, FOX, ERY	<i>mecA, blaZ, ermC,</i>	E	<i>hld</i>
Case 6	VS3203	Dog1	<i>S. aureus</i>	398 (398)	t5635	-	I	Susceptible	-	-	<i>hla, hld</i>
	VS3204	Dog 3	<i>S. sciuri</i>	-	-	-	-	PEN	<i>mecA, blaZ</i>	-	
	VS3205	Hunter 5	<i>S. aureus</i>	34 (30)	t166	-	III	PEN, ERY	<i>blaZ, ermC</i>	E	<i>hld</i>
Case 7	VS3206	Dog1	<i>S. pseudintermedius</i>	-	-	-	-	PEN, FOX	<i>mecA, blaZ</i>	-	<i>lukS/F-I, siet</i>
	VS3207	Dog2	<i>S. lentus</i>	-	-	-	-	PEN	<i>mecA</i>	-	
Case 8	VS3208	Hunter 6	<i>S. aureus</i>	718	t11333		II	Susceptible	-	E	<i>hla, hld</i>
	VS3209	Dog 1	<i>S. vitulinus</i>	-	-			PEN	<i>mecA</i>		
Case 9	VS3210	Hunter 7	<i>S. aureus</i>	398 (398)	t5635	-	I	PEN, ERY	<i>blaZ, ermT</i>	<i>scn</i>	<i>hla, hld</i>
	VS3211	Dog1	<i>S. lentus</i>	-	-	-		PEN, CN, KAN, CD, C	<i>mecA, aac(6')-Ie-aph(2'')-Ia, aph(3')-IIIa, mphC, cat_{p221}</i>	-	<i>hla</i>
Case 10	VS3212	Hunter 8	<i>S. aureus</i>	7343	t012	N.T.	III	PEN, FOX	<i>mecA, blaZ</i>	-	<i>hla, hlb, hld</i>
	VS3213	Dog	<i>S. sciuri</i>	-	-			PEN	<i>mecA</i>		
Case 11	VS3214	Hunter 9	<i>S. aureus</i>	5 (5)	t179	IV	II	PEN, FOX, ERY,	<i>mecA, blaZ, ermA</i>	E	<i>hla, hld</i>
	VS3215	Dog 1	<i>S. lentus</i>	-	-	-	-	PEN	<i>mecA, blaZ</i>	-	
	VS3216	Dog 2	<i>S. lentus</i>	-	-	-	-	PEN	<i>mecA, blaZ,</i>	-	
Case 12	VS3217	Hunter 10	<i>S. aureus</i>	7	t091	-	I	PEN	<i>blaZ</i>	G	<i>hla, hld</i>

Abbreviation: NT: not typeable; PEN: penicillin; FOX: ceftiofur; CN: gentamicin; TOB: tobramycin; KAN: kanamycin; ERY: erythromycin; CD: clindamycin; TET: tetracycline; FD: fusidic acid; C: chloramphenicol; IEC: immune evasion cluster; ST: sequence type; CC: clonal complex. MRSA isolates are presented in bold.

Table 1 is divided into the cases of a hunter and their respective hunting dogs. In general, MRSA isolates were ascribed to five STs (ST9, ST8, ST5, ST718, and ST7343) and five *spa*-types (t2922, t121, t11333, t012 and t179) (Figure 1). Isolates belonging to ST5, ST8, and ST718 were typed as SCC*mec* IV, whereas isolates ascribed to ST9 and ST7343 were not typeable by SCC*mec*-typing. MRSA isolates were ascribed to *agr* type II (n = 9), I and III. Only three MRSA isolates displayed a multidrug-resistant profile, all from Case 3, showing resistances to penicillin, ceftazidime, gentamicin, tobramycin, kanamycin, clindamycin, fusidic acid, and chloramphenicol encoded by the *mecA*, *blaZ*, *aac(6′)-Ie-aph(2′′)-Ia*, *aph(3′)-IIIa*, *vgaA*, *lnuB*, and *cat_{p221}* genes. The remaining MRSA isolates showed resistance mainly to penicillin, ceftazidime, and erythromycin conferred by the *mecA*, *blaZ*, *ermA*, and *ermC* genes. Six out of the eleven MRSA isolates harbored the *scn* and *sak* genes of the IEC system as were categorized as type E. Among the other virulence genes, *hla* (n = 9), *hly* (n = 5), *hld* (n = 11), and *tst* (n = 2) were detected among MRSA isolates.




Figure 1. Phylogenetic tree inferred from the analysis of the MLST analysis. The tree was created using the Complete-Linkage method of Hierarchical Clustering. Hamming distance was used to measure genetic distance.

Methicillin-susceptible *S. aureus* (MSSA) isolates were detected in 5 hunters and 5 dogs. These isolates were ascribed to ST7353-t10042 (n = 3), ST30-t012 (n = 2), ST398-t5635 (n = 2), ST34-t166, ST718-t11333 and ST7-t091. MSSA isolates were typed as *agr* type III (n = 3), type I (n = 3) and type II (n = 1), and the ST7353-t10042 isolates were not typeable. From the ten MSSA isolates, four were susceptible to all antibiotics tested. Resistance to penicillin encoded by the *blaZ* gene was identified in five isolates and resistance to erythromycin was identified in two isolates harboring the *ermC* and *ermT* genes. MSSA isolates were grouped in IEC type E (n = 5) and G (n = 1), and one isolate carried only the *scn* gene.

As previously mentioned, in cases where MRSA was found, samples from dogs and/or dog owners that shared the same household were screened for the presence of other methicillin-resistant staphylococci species. Among the non-*aureus* staphylococci, five *S. pseudintermedius* were isolated from five dogs. All *S. pseudintermedius* isolates were resistant to penicillin and ceftazidime and carried the *mecA* gene, and therefore, they were considered MRSP. One *S. pseudintermedius* also showed resistance aminoglycosides conferred by the *aph(2′′)-Ia*, *aph(3′)-IIIa* and *str* genes. All *S. pseudintermedius* harbored the *lukS/F-I* genes and four also carried the virulence gene *siet*. All the coagulase-negative staphylococci (CoNS) isolates were recovered from dogs except for one *S. lentus*. CoNS were positive for the *mecA* gene and were resistant to penicillin. Only one CoNS (*S. lentus* VS3211) had a multidrug-resistance profile showing resistance to penicillin, gentamicin, kanamycin, clindamycin, and chloramphenicol encoded by the *mecA*, *aac(6′)-Ie-aph(2′′)-Ia*,

aph(3')-IIIa, *mphC* and *cat*_{p221} genes. Moreover, this isolate was also the only CoNS to carry a virulence gene, the *hla* gene.

4. Discussion

S. aureus is part of the normal human microbiome in approximately 30% of the human population [39]. In fact, 20% of humans are permanent carriers of *S. aureus* and about 60% are intermittent carriers [39]. This data is in accordance with the frequency of *S. aureus* obtained in our study (30%) in healthy hunters. Half of the *S. aureus* isolates were MRSA, which corresponds to 16.6% of the 30 hunters tested. MRSA frequency in healthy community humans is very variable and is influenced by geographic location, demographic characteristics, sampling years, among others. Even so, several studies have reported similar frequencies to those obtained in our study. In the study by Velasco et al. none of the 550 undergraduate students carried MRSA and only 7.6% were positive for *S. aureus* [40]. Other studies have reported a frequency of MRSA in healthy humans between 14.64% and 24.7% [22,41,42]. In our study, the frequency of *S. aureus* and MRSA in hunting dogs was 15.4% and 9%, respectively. János et al. also reported similar results with a frequency of *S. aureus* and MRSA of 11.62% and 9.30%, respectively, in kennel dogs from Romania [43]. Other studies have reported a lower prevalence of MRSA in dogs [44,45]. In contrast, in a previous study from Portugal, a higher prevalence of MRSA (30%) was detected in dogs [46]. In the only available study of *S. aureus* in hunting dogs, 36.9% and 23.7% of the dogs were nasal carriers of *S. aureus* and MRSA, respectively, which is a much higher frequency than that found in our study [47].

A few studies have reported transmission of *S. aureus* and MRSA between dogs and humans in the same household and people working in close contact with dogs [45,48–51]. Most studies rely on molecular typing techniques, such as whole genome sequencing, rep-PCR, and pulsed field gel electrophoresis, to verify dog-to-human transmission. In our study, three cases (Cases 1, 2, and 3) concerning a possible transmission between dogs and owners were identified. Our results demonstrate that hunting dogs and their owners carried *S. aureus* and MRSA strains with clonal similarity, indicating a possible transmission via direct transfer from animals to humans or vice versa. Nevertheless, it is important to point out that whole genome sequencing should have been performed to confirm the bacterial transmission. In Case 1, the hunter and his dogs share the same clonal lineage ST7353-t10042. *S. aureus* ST7353 was first reported in this study and is a single locus variant of the ST45 (CC45) with a single-point mutation in the *pta* gene. *S. aureus* belonging to CC45 is primarily known as a human-associated clone. Nevertheless, the CC45 is characterized by its diversity since it has been associated with MRSA, MSSA, HA-MRSA, CA-MRSA, commensal clones, and it has also been isolated from pets, livestock, wild animals, and the environment [52–57]. Furthermore, *S. aureus* ST45 have unique genetic differences from other *S. aureus* clades since it has been shown that ST45 branches off near the root of the *S. aureus* population [58]. *spa*-type t10042 is a rare *spa*-type and it has only been detected once since it was first reported in 2012, and is associated with human isolates in Europe [59].

A transmission of MSSA might have also occurred in Case 2 since *S. aureus* ST30-t012 was isolated from both the hunter and his hunting dog. Furthermore, the pattern of antimicrobial resistance was similar in both isolates. MSSA-ST30 is an ancestral strain of an epidemic MRSA clone which evolved into MSSA, and is also known as the Southwest Pacific clone [60]. ST30 is an international successful clone since it has been found in Australia, Europe and Asia [61]. This lineage is primarily associated with humans, but it has been also found among animals and in the environment [55,62]. Moreover, ST30-MSSA-t012 has been detected in hospitalized humans and livestock workers in Portugal, in the same region where the samples of this study were collected [63,64]. Both ST30-MSSA-t012 isolates carried the virulence gene *tst*, in addition to the hemolysins genes, which is in accordance with other studies that have shown that *S. aureus* ST30 often carries pathogenicity islands including the *tst* gene [65].

In Case 3, MRSA strains with the same clonal lineage, ST9-t2922, were isolated from the hunter and two hunting dogs. ST9 is a livestock-associated MRSA (LA-MRSA) lineage that is predominant in Asia. As in our study, this lineage lacks several important virulence genes, such as *lukF/S-PVL*. Furthermore, ST9 MRSA strains are usually multidrug resistant [66]. In our study, all ST9 isolates were resistant to penicillin, cefoxitin, aminoglycosides, clindamycin, and tetracycline encoded by the *mecA*, *blaZ*, *aac(6′)-Ie-aph(2′′)-Ia*, *aph(3′)-IIIa*, *lnuB*, *vgaA*, *tetK*, and *tetM* genes. In addition, MRSA isolates from dogs were also resistant to chloramphenicol and harbored the *cat_{p221}* gene. These minor genetic variations between the dogs and owner isolates may have evolved in the two different hosts after interspecies transmission, as was the case in other studies in which *S. aureus* was transmitted between dogs and humans [52]. MRSA *spa*-type t2922 is often associated with LA-MRSA, not so much with ST9, but rather with ST398 which is the predominant LA-MRSA lineage in Europe [67]. Even so, ST9-t2922 has been detected among pigs in China and Taiwan [68,69]. ST9-t2922 isolates were not typeable by SCC*mec* typing which may be due to the high diversity of SCC*mec* types reported among ST9 MRSA strains [66,70,71].

Dog-to-dog staphylococci transmission has been documented, particularly dogs living together in the same household [72,73]. Transmission between dogs may occur due to several factors, such as dog-to-dog contact, sharing the same water and food, and sharing the same environment. In our study, in Case 4, the hunter and three hunting dogs were colonized by MRSA strains. All dogs' isolates belonged to ST5-t179, which suggests a possible dog-to-dog transmission. The MRSA isolates from dogs were assigned to IEC type E which indicates a possible human origin [74]; however, transmission between the hunter owner and the dogs does not seem to have occurred as the clonal lineages differ. MRSA ST5-t179 SCC*mec* type IV, also known as the Pediatric clone, is a classic human pathogen predominant in HA-MRSA, and has been repeatedly isolated from human infections in Portugal [63,75]. Nevertheless, we believe that this is the first study reporting ST5-MRSA-t179 in dogs. As for MRSA isolated from the hunter-owner, it belongs to ST8, *spa*-type t121, and SCC*mec* IV. MRSA ST8 is a common CA-MRSA clone frequently detected in the USA, which is often related with the USA300 clone; however, the presence of PVL encoding genes is a marker of the USA300 clone, and in our study, the ST8 MRSA isolate lacked this gene [76]. In fact, the epidemiology of the ST8 MRSA clone differs remarkably among world regions. For instance, in Europe, ST8 is commonly detected in community humans, but most of them are non-USA300 [76]. Nevertheless, in Case 4, a transmission of MRSP may have occurred between dogs. Two MRSP showed resistance to penicillin and cefoxitin and carried the *blaZ* and *mecA* genes; however, one MRSP isolate (VS3198) also showed resistance to aminoglycosides conferred by the *aac(6′)-Ie-aph(2′′)-Ia*, *aph(3′)-IIIa*, and *str* which may have been acquired after the transmission. The MRSP isolates also carried the virulence genes *lukS/F-I* and *siet*, which have been previously reported in both commensal and clinical strains, indicating that these genes may be ubiquitous in this *S. pseudintermedius* [77].

Another possible dog-to-dog transmission can be observed in Case 5. Both dogs sharing the same household carried MRSA strains belonging to ST718, t11333, and SCC*mec* IV, and they harbored the same resistance and IEC genes. *S. aureus* ST718 is a rare human-associated clone that has been reported in a few countries with regard to human infections and community humans [78–81]. Although this clonal lineage is most often associated with MSSA strains, it has also been identified among MRSA isolates that are ascribed to SCC*mec* type IV, similarly to the one obtained in this study [78]. Interestingly, in our study, one ST718-t11333 MSSA strain was also isolated from a hunter (Case 8) which shows the adaptability of this lineage. *S. aureus* ST718-t11333 has been identified among animals in only one study that was conducted with owl samples in Portugal [62]. Since hunters and their hunting dogs are in direct contact with wild game animals, transmission may also occur, especially between dogs and wild game animals.

In no other Case does the transmission of *S. aureus* between dogs and humans appear to have occurred; however, *S. aureus* was isolated from five more hunters and one dog.

One hunter (Case 9) and one hunting dog (Case 6) shared the same *S. aureus* clone, ST398-t5635, but with no apparent relationship or contact between them. Animals are considered the main reservoir of *S. aureus* CC398; however, this lineage is divided into two clades: the classical LA clade and the human clade [82]. It is believed that ST398 was originally a human-associated clone, and it has adapted to animals through the loss of integrase group 3 prophages containing the IEC system genes, and with it, they acquired tetracycline resistance [83,84]; however, it has been shown that a re-adaptation of *S. aureus* CC398 to humans may occur with the acquisition of IEC [83–85]. Both of our CC398 isolates lacked the tetracycline resistance, which is a marker of animal adaptation; however, the human isolate was resistant to penicillin and erythromycin carrying the *blaZ* and *ermT* genes. Furthermore, this isolate also carried the *scn* gene of the IEC system. Studies have shown that CC398 related to humans and human infections often carry the *ermT* and the *chp* and *scn* genes, which may indicate a human adaptation [85,86]. It is important to point out that most MRSA and MRSP isolates did not present a multidrug-resistant profile that is common in methicillin-resistant isolates; therefore, MRSA that is simply β -lactam resistant, as many in this study are, would be of lesser concern, as it is an opportunistic pathogen, and thus, there are still a multitude of appropriate therapeutic options.

Methicillin-resistant CoNS (MRCoNS) were also isolated from both hunters and dogs in this study. A few studies have been conducted investigating the frequency of CoNS and MRCoNS in healthy dogs, and the CoNS species detected in those studies are very variable. Ma et al. reported that the most frequent CoNS among dogs in Australia was *S. sciuri*, whereas in Brazil and the United Kingdom, it was *S. epidermidis* followed by *S. simulans* and *S. epidermidis*, followed by *S. warneri*, respectively [44,87,88]. In Thailand, the most common CoNS species was *S. chromogenes* [89]. In our study, the most prevalent CoNS was *S. lentus*, which was isolated from dogs and one human. Furthermore, among all MRCoNS only one (*S. lentus* VS3211) was multidrug-resistant, and showed resistance to penicillin, aminoglycosides, clindamycin, and chloramphenicol which was conferred by the *mecA*, *aac(6′)-Ie-aph(2′′)-Ia*, *aph(3′)-IIIa*, *mphC*, and *cat_{p221}* genes. *S. lentus* is considered to be an animal commensal and pathogen species [28]. Nevertheless, it has also been identified as the etiological agent of human infections [90]. As expected, none of the MRCoNS isolates showed phenotypic resistance to ceftiofur despite carrying the *mecA* gene. Studies have shown that the *mecA* gene may have originated from the CoNS species belonging to the *S. sciuri* group, which includes *S. sciuri*, *S. viutlinus*, and *S. lentus*, and these species often carry *mecA* homologues that do not confer phenotypic resistance [91,92]. Transmission of MRCoNS is harder to confirm since molecular typing methods are not available for all species. Nevertheless, we can hypothesize that in Case 11, the hunting dogs may be sharing the same *S. lentus* clone, since both isolates showed the same resistance pattern.

5. Conclusions

Genetic similarity was observed between *S. aureus* and MRSA isolates from hunters and their hunting dogs, suggesting possible human-to-dog and dog-to-dog transmissions which could pose a public health risk. *S. aureus* isolated from hunters and their hunting dogs living in the same household showed identical STs, *spa*-, *SCCmec*-, and *agr*-types, as well as similar resistance and virulence patterns. Most *S. aureus* isolates were classical human-associated clones which may point to one-way transmission from humans to dogs. Furthermore, several *S. aureus* isolates carried the genes encoding the IEC system, which reinforces a possible human origin; therefore, the role of *S. aureus* as a zoonotic pathogen is potentiated; however, although this study points to a possible *S. aureus* transmission, whole genome sequencing should be carried to confirm the human-to-dog and dog-to-dog transmissions.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/pathogens11050548/s1>, Table S1: Breed, gender, and age of all dogs sampled in this study.

Author Contributions: Conceptualization, V.S. and P.P.; methodology, V.S.; validation, M.C., M.V.-P. and P.P.; investigation, V.S.; resources, J.E.P. and L.M.; data curation, V.S. and V.M.; writing—original draft preparation, V.S.; writing—review and editing, V.S.; supervision, G.I. and P.P. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by the R&D Project CAREBIO2: Comparative assessment of antimicrobial resistance in environmental biofilms through proteomics—towards innovative therapeutic biomarkers, with reference NORTE-01-0145-FEDER-030101 and PTDC/SAU-INF/30101/2017, financed by the European Regional Development Fund (ERDF) through the Northern Regional Operational Program (NORTE 2020) and the Foundation for Science and Technology (FCT). This work was supported by the Associate Laboratory for Green Chemistry-LAQV, which is financed by national funds from FCT/MCTES (UIDB/50006/2020 and UIDP/50006/2020) and by the projects UIDB/CVT/00772/2020 and LA/P/0059/2020 funded by the Portuguese Foundation for Science and Technology (FCT). Vanessa Silva is grateful to FCT (Fundação para a Ciência e a Tecnologia) for financial support through the PhD grant SFRH/BD/137947/2018.

Institutional Review Board Statement: The study was conducted according to the Helsinki Declaration (ICH-GCP principles), in compliance with Schedule Y/ICMR Guidelines, the Oviedo Convention, and was approved by the Ethics Committee of University of Trás-os-Montes e Alto Douro (EC-UTAD, 8 November 2019).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Not applicable.

Acknowledgments: The authors acknowledge the Federation of Hunters Associations of the 1st Game Region (Federação das Associações de Caçadores da 1ª Região Cinegética) (FACIRC) for the collaboration and contribution to the collection of samples.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lee, D.C.; Kananurak, A.; Tran, M.T.; Connolly, P.A.; Polage, C.R.; Iwase, T.; Bevins, C.L.; Underwood, M.A. Bacterial Colonization of the Hospitalized Newborn: Competition between *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Pediatr. Infect. Dis. J.* **2019**, *38*, 682–686. [[CrossRef](#)] [[PubMed](#)]
2. González-García, S.; Hamdan-Partida, A.; Bustos-Hamdan, A.; Bustos-Martínez, J. Factors of Nasopharynx that Favor the Colonization and Persistence of *Staphylococcus aureus*. In *Pharynx-Diagnosis and Treatment*; IntechOpen: Rijeka, Croatia, 2021; ISBN 1789856094.
3. Sakr, A.; Brégeon, F.; Mège, J.-L.; Rolain, J.-M.; Blin, O. *Staphylococcus aureus* Nasal Colonization: An Update on Mechanisms, Epidemiology, Risk Factors, and Subsequent Infections. *Front. Microbiol.* **2018**, *9*, 2419. [[CrossRef](#)] [[PubMed](#)]
4. Ahmad-Mansour, N.; Loubet, P.; Pouget, C.; Dunyach-Remy, C.; Sotto, A.; Lavigne, J.-P.; Molle, V. *Staphylococcus aureus* toxins: An update on their pathogenic properties and potential treatments. *Toxins* **2021**, *13*, 677. [[CrossRef](#)] [[PubMed](#)]
5. Jevons, M.P. “Celbenin”-resistant staphylococci. *Br. Med. J.* **1961**, *1*, 124. [[CrossRef](#)]
6. European Centre for Disease Prevention and Control (ECDC). *Antimicrobial resistance in the EU/EEA (EARS-Net) - Annual Epidemiological Report for 2019*; European Centre for Disease Prevention and Control (ECDC): Stockholm, Sweden, 2020.
7. WHO Regional Office for Europe and European Centre for Disease Prevention and Control. *Surveillance of Antimicrobial Resistance in Europe, 2020 Data*; WHO Regional Office for Europe and European Centre for Disease Prevention and Control: Copenhagen, Denmark, 2021.
8. Shalaby, M.-A.W.; Dokla, E.M.E.; Serya, R.A.T.; Abouzid, K.A.M. Penicillin binding protein 2a: An overview and a medicinal chemistry perspective. *Eur. J. Med. Chem.* **2020**, *199*, 112312. [[CrossRef](#)]
9. Schwendener, S.; Perreten, V. The bla and mec families of β -lactam resistance genes in the genera *Micrococcus*, *Mammaliococcus* and *Staphylococcus*: An in-depth analysis with emphasis on *Micrococcus*. *J. Antimicrob. Chemother.* **2022**, dkac107. [[CrossRef](#)]
10. Boswihi, S.S.; Udo, E.E. Methicillin-resistant *Staphylococcus aureus*: An update on the epidemiology, treatment options and infection control. *Curr. Med. Res. Pract.* **2018**, *8*, 18–24. [[CrossRef](#)]
11. Crespo-Piazuelo, D.; Lawlor, P.G. Livestock-associated methicillin-resistant *Staphylococcus aureus* (LA-MRSA) prevalence in humans in close contact with animals and measures to reduce on-farm colonisation. *Ir. Vet. J.* **2021**, *74*, 21. [[CrossRef](#)]
12. Voss, A.; Loeffen, F.; Bakker, J.; Klaassen, C.; Wulf, M. Methicillin-resistant *Staphylococcus aureus* in Pig Farming. *Emerg. Infect. Dis.* **2005**, *11*, 1965–1966. [[CrossRef](#)]
13. Loeffler, A.; Pfeiffer, D.U.; Lindsay, J.A.; Soares-Magalhaes, R.; Lloyd, D.H. Lack of transmission of methicillin-resistant *Staphylococcus aureus* (MRSA) between apparently healthy dogs in a rescue kennel. *Vet. Microbiol.* **2010**, *141*, 178–181. [[CrossRef](#)]

14. Van Duijkeren, E.; Hengeveld, P.; Zomer, T.P.; Landman, F.; Bosch, T.; Haenen, A.; van de Giessen, A. Transmission of MRSA between humans and animals on duck and turkey farms. *J. Antimicrob. Chemother.* **2016**, *71*, 58–62. [[CrossRef](#)] [[PubMed](#)]
15. Sahibzada, S.; Abraham, S.; Coombs, G.W.; Pang, S.; Hernández-Jover, M.; Jordan, D.; Heller, J. Transmission of highly virulent community-associated MRSA ST93 and livestock-associated MRSA ST398 between humans and pigs in Australia. *Sci. Rep.* **2017**, *7*, 5273. [[CrossRef](#)] [[PubMed](#)]
16. Van Balen, J.C.; Landers, T.; Nutt, E.; Dent, A.; Hoet, A.E. Molecular epidemiological analysis to assess the influence of pet-ownership in the biodiversity of Staphylococcus aureus and MRSA in dog- and non-dog-owning healthy households. *Epidemiol. Infect.* **2017**, *145*, 1135–1147. [[CrossRef](#)] [[PubMed](#)]
17. Kuroda, T.; Kinoshita, Y.; Niwa, H.; Shinzaki, Y.; Tamura, N.; Hobo, S.; Kuwano, A. Methicillin-resistant Staphylococcus aureus colonisation and infection in Thoroughbred racehorses and veterinarians in Japan. *Vet. Rec.* **2016**, *178*, 473. [[CrossRef](#)]
18. Bramble, M.; Morris, D.; Tolomeo, P.; Lautenbach, E. Potential Role of Pet Animals in Household Transmission of Methicillin-Resistant Staphylococcus aureus: A Narrative Review. *Vector-Borne Zoonotic Dis.* **2010**, *11*, 617–620. [[CrossRef](#)]
19. Gómez-Sanz, E.; Torres, C.; Ceballos, S.; Lozano, C.; Zarazaga, M. Clonal Dynamics of Nasal Staphylococcus aureus and Staphylococcus pseudintermedius in Dog-Owning Household Members. Detection of MSSA ST398. *PLoS ONE* **2013**, *8*, e69337. [[CrossRef](#)]
20. Lozano, C.; Rezusta, A.; Ferrer, I.; Pérez-Laguna, V.; Zarazaga, M.; Ruiz-Ripa, L.; Revillo, M.J.; Torres, C. Staphylococcus pseudintermedius Human Infection Cases in Spain: Dog-to-Human Transmission. *Vector-Borne Zoonotic Dis.* **2017**, *17*, 268–270. [[CrossRef](#)]
21. Boost, M.V.; O'Donoghue, M.M.; James, A. Prevalence of Staphylococcus aureus carriage among dogs and their owners. *Epidemiol. Infect.* **2008**, *136*, 953–964. [[CrossRef](#)]
22. Daley, P.; Bajgai, J.; Penney, C.; Williams, K.; Whitney, H.; Golding, G.R.; Weese, S. A cross sectional study of animal and human colonization with Methicillin-Resistant Staphylococcus aureus (MRSA) in an Aboriginal community. *BMC Public Health* **2016**, *16*, 595. [[CrossRef](#)]
23. Ma, G.C.; Worthing, K.A.; Gottlieb, T.; Ward, M.P.; Norris, J.M. Molecular characterization of community-associated methicillin-resistant Staphylococcus aureus from pet dogs. *Zoonoses Public Health* **2020**, *67*, 222–230. [[CrossRef](#)]
24. Penna, B.; Silva, M.B.; Soares, A.E.R.; Vasconcelos, A.T.R.; Ramundo, M.S.; Ferreira, F.A.; Silva-Carvalho, M.C.; de Sousa, V.S.; Rabello, R.F.; Bandeira, P.T.; et al. Comparative genomics of MRSA strains from human and canine origins reveals similar virulence gene repertoire. *Sci. Rep.* **2021**, *11*, 4724. [[CrossRef](#)] [[PubMed](#)]
25. Taniguchi, Y.; Koide, S.; Maeyama, Y.; Tamai, K.; Hayashi, W.; Tanaka, H.; Iimura, M.; Suzuki, M.; Nagano, Y.; Arakawa, Y.; et al. Predominance of methicillin-resistant Staphylococcus aureus SCCmec type II-CC5 and SCCmec type IV-CC1/CC8 among companion animal clinical isolates in Japan: Findings from phylogenetic comparison with human clinical isolates. *J. Glob. Antimicrob. Resist.* **2020**, *20*, 253–259. [[CrossRef](#)] [[PubMed](#)]
26. CLSI Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing; Clinical and Laboratory Standards Institute*: Wayne, PA, USA, 2017.
27. European Committee on Antimicrobial Susceptibility Testing (EUCAST). *Breakpoint Tables for Interpretation of MICs and Zone Diameters Version 8.0.*; EUCAST: Växjö, Sweden, 2018.
28. Silva, V.; Caniça, M.; Ferreira, E.; Vieira-Pinto, M.; Saraiva, C.; Pereira, J.E.; Capelo, J.L.; Igrejas, G.; Poeta, P. Multidrug-Resistant Methicillin-Resistant Coagulase-Negative Staphylococci in Healthy Poultry Slaughtered for Human Consumption. *Antibiotics* **2022**, *11*, 365. [[CrossRef](#)] [[PubMed](#)]
29. Jarraud, S.; Mougel, C.; Thioulouse, J.; Lina, G.; Meugnier, H.; Forey, F.; Etienne, J.; Vandenesch, F.; Nesme, X. Relationships between Staphylococcus aureus Genetic Background, Virulence Factors, agr Groups (Alleles), and Human Disease. *Infect. Immun.* **2002**, *70*, 631–641. [[CrossRef](#)] [[PubMed](#)]
30. Lina, G.; Piemont, Y.; Godail-Gamot, F.; Bes, M.; Peter, M.-O.; Gauduchon, V.; Vandenesch, F.; Etienne, J. Involvement of Panton-Valentine Leukocidin—Producing Staphylococcus aureus in Primary Skin Infections and Pneumonia. *Clin. Infect. Dis.* **1999**, *29*, 1128–1132. [[CrossRef](#)]
31. Yamaguchi, T.; Nishifuji, K.; Sasaki, M.; Fudaba, Y.; Aepfelbacher, M.; Takata, T.; Ohara, M.; Komatsuzawa, H.; Amagai, M.; Sugai, M. Identification of the Staphylococcus aureus etd pathogenicity island which encodes a novel exfoliative toxin, ETD, and EDIN-B. *Infect. Immun.* **2002**, *70*, 5835–5845. [[CrossRef](#)]
32. Van Wamel, W.J.B.; Rooijackers, S.H.M.; Ruyken, M.; van Kessel, K.P.M.; van Strijp, J.A.G. The innate immune modulators staphylococcal complement inhibitor and chemotaxis inhibitory protein of Staphylococcus aureus are located on beta-hemolysin-converting bacteriophages. *J. Bacteriol.* **2006**, *188*, 1310–1315. [[CrossRef](#)]
33. Iyori, K.; Futagawa-Saito, K.; Hisatsune, J.; Yamamoto, M.; Sekiguchi, M.; Ide, K.; Son, W.; Olivry, T.; Sugai, M.; Fukuyasu, T. Staphylococcus pseudintermedius exfoliative toxin EXI selectively digests canine desmoglein 1 and causes subcorneal clefts in canine epidermis. *Vet. Dermatol.* **2011**, *22*, 319–326. [[CrossRef](#)]
34. Futagawa-Saito, K.; Sugiyama, T.; Karube, S.; Sakurai, N.; Ba-Thein, W.; Fukuyasu, T. Prevalence and Characterization of Leukotoxin-Producing Staphylococcus intermedius in Isolates from Dogs and Pigeons. *J. Clin. Microbiol.* **2004**, *42*, 5324–5326. [[CrossRef](#)]
35. Enright, M.C.; Day, N.P.; Davies, C.E.; Peacock, S.J.; Spratt, B.G. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of Staphylococcus aureus. *J. Clin. Microbiol.* **2000**, *38*, 1008–1015. [[CrossRef](#)]

36. Harmsen, D.; Claus, H.; Witte, W.; Rothgänger, J.; Claus, H.; Turnwald, D.; Vogel, U. Typing of Methicillin-Resistant *Staphylococcus aureus* in a University Hospital Setting by Using Novel Software for spa Repeat Determination and Database Management. *J. Clin. Microbiol.* **2003**, *41*, 5442–5448. [[CrossRef](#)] [[PubMed](#)]
37. Shopsin, B.; Mathema, B.; Alcabes, P.; Said-Salim, B.; Lina, G.; Matsuka, A.; Martinez, J.; Kreiswirth, B.N. Prevalence of agr Specificity Groups among *Staphylococcus aureus* Strains Colonizing Children and Their Guardians. *J. Clin. Microbiol.* **2003**, *57*, 456–459. [[CrossRef](#)] [[PubMed](#)]
38. Zhang, K.; McClure, J.-A.; Elsayed, S.; Louie, T.; Conly, J.M. Novel Multiplex PCR Assay for Characterization and Concomitant Subtyping of Staphylococcal Cassette Chromosome *mec* Types I to V in Methicillin-Resistant *Staphylococcus aureus*. *J. Clin. Microbiol.* **2005**, *43*, 5026–5033. [[CrossRef](#)]
39. Laux, C.; Peschel, A.; Krismer, B. *Staphylococcus aureus* Colonization of the Human Nose and Interaction with Other Microbiome Members. *Microbiol. Spectr.* **2019**, *7*, 2. [[CrossRef](#)]
40. Velasco, V.; Buyukcangaz, E.; Sherwood, J.S.; Stepan, R.M.; Koslofsky, R.J.; Logue, C.M. Characterization of *Staphylococcus aureus* from Humans and a Comparison with Isolates of Animal Origin, in North Dakota, United States. *PLoS ONE* **2015**, *10*, e0140497. [[CrossRef](#)]
41. Bektas, S.; Obradovic, A.; Aljicevic, M.; Numanovic, F.; Hodzic, D.; Sporisevic, L. The frequency of community-acquired methicillin-resistant *Staphylococcus aureus* (CA-mrsa) among samples in institute for public health in canton Sarajevo. *Mater. Sociomed.* **2016**, *28*, 61–65. [[CrossRef](#)]
42. Kasela, M.; Grzegorzcyk, A.; Nowakowicz-Dębek, B.; Malm, A. The Prevalence of Virulence Determinants and Antibiotic Resistance Patterns in Methicillin—Resistant *Staphylococcus aureus* in a Nursing Home in Poland. *Pathogens* **2021**, *10*, 427. [[CrossRef](#)]
43. János, D.; Viorel, H.; Ionica, I.; Corina, P.; Tiana, F.; Roxana, D. Carriage of Multidrug Resistance Staphylococci in Shelter Dogs in Timisoara, Romania. *Antibiotics* **2021**, *10*, 801. [[CrossRef](#)]
44. Ma, G.C.; Worthing, K.A.; Ward, M.P.; Norris, J.M. Commensal Staphylococci Including Methicillin-Resistant *Staphylococcus aureus* from Dogs and Cats in Remote New South Wales, Australia. *Microb. Ecol.* **2020**, *79*, 164–174. [[CrossRef](#)]
45. Huang, T.; Chou, C. Methicillin-sensitive and methicillin-resistant *Staphylococcus aureus* strains and their toxin genes in the nostrils of dogs and workers at an animal shelter. *J. Appl. Microbiol.* **2019**, *126*, 1899–1909. [[CrossRef](#)]
46. Coelho, C.; Torres, C.; Radhouani, H.; Pinto, L.; Lozano, C.; Gómez-Sanz, E.; Zaragaza, M.; Igrejas, G.; Poeta, P. Molecular Detection and Characterization of Methicillin-Resistant *Staphylococcus aureus* (MRSA) Isolates from Dogs in Portugal. *Microb. Drug Resist.* **2011**, *17*, 333–337. [[CrossRef](#)] [[PubMed](#)]
47. Mustapha, M.; Bukar-Kolo, Y.M.; Geidam, Y.A.; Gulani, I.A. Phenotypic and genotypic detection of methicillin-resistant *Staphylococcus aureus* in hunting dogs in Maiduguri metropolitan, Borno State, Nigeria. *Vet. World* **2016**, *9*, 501–506. [[CrossRef](#)] [[PubMed](#)]
48. Fabri, F.V.; Pinto, N.B.; de Mattos, M.d.S.F.; Rodrigues, R.F.; Shinohara, D.R.; Pereira, P.M.; Nishiyama, S.A.B.; Tognim, M.C.B. First report of oxacillin-susceptible *mecA*-positive *Staphylococcus aureus* in healthy dogs and their owners in southern Brazil. *Prev. Vet. Med.* **2021**, *189*, 105286. [[CrossRef](#)] [[PubMed](#)]
49. Ferreira, J.P.; Anderson, K.L.; Correa, M.T.; Lyman, R.; Ruffin, F.; Reller, L.B.; Fowler, V.G., Jr. Transmission of MRSA between Companion Animals and Infected Human Patients Presenting to Outpatient Medical Care Facilities. *PLoS ONE* **2011**, *6*, e26978. [[CrossRef](#)]
50. Davis, M.F.; Misic, A.M.; Morris, D.O.; Moss, J.T.; Tolomeo, P.; Beiting, D.P.; Nachamkin, I.; Lautenbach, E.; Rankin, S.C. Genome sequencing reveals strain dynamics of methicillin-resistant *Staphylococcus aureus* in the same household in the context of clinical disease in a person and a dog. *Vet. Microbiol.* **2015**, *180*, 304–307. [[CrossRef](#)]
51. Oh, J.-Y.; Chae, J.-C.; Han, J.-I.; Song, W.-K.; Lee, C.-M.; Park, H.-M. Distribution and epidemiological relatedness of methicillin-resistant *Staphylococcus aureus* isolated from companion dogs, owners, and environments. *J. Vet. Med. Sci.* **2020**, *82*, 1379–1386. [[CrossRef](#)] [[PubMed](#)]
52. Sahin-Tóth, J.; Kovács, E.; Tóthpál, A.; Juhász, J.; Forró, B.; Bányai, K.; Havril, K.; Horváth, A.; Ghidán, Á.; Dobay, O. Whole genome sequencing of coagulase positive staphylococci from a dog-and-owner screening survey. *PLoS ONE* **2021**, *16*, e0245351. [[CrossRef](#)]
53. Effelsberg, N.; Stegger, M.; Peitzmann, L.; Altinok, O.; Coombs, G.W.; Pichon, B.; Kearns, A.; Randad, P.R.; Heaney, C.D.; Bletz, S. Global Epidemiology and Evolutionary History of *Staphylococcus aureus* ST45. *J. Clin. Microbiol.* **2022**, *59*, e02198-20. [[CrossRef](#)]
54. Asanin, J.; Misic, D.; Aksentijevic, K.; Tambur, Z.; Rakonjac, B.; Kovacevic, I.; Sperser, J.; Loncaric, I. Genetic Profiling and Comparison of Human and Animal Methicillin-Resistant *Staphylococcus aureus* (MRSA) Isolates from Serbia. *Antibiotics* **2019**, *8*, 26. [[CrossRef](#)]
55. Silva, V.; Ferreira, E.; Manageiro, V.; Reis, L.; Tejedor-Junco, M.T.; Sampaio, A.; Capelo, J.L.; Caniça, M.; Igrejas, G.; Poeta, P. Distribution and Clonal Diversity of *Staphylococcus aureus* and Other Staphylococci in Surface Waters: Detection of ST425-t742 and ST130-t843 *mecC*-Positive MRSA Strains. *Antibiotics* **2021**, *10*, 1416. [[CrossRef](#)]
56. Luzzago, C.; Locatelli, C.; Franco, A.; Scaccabarozzi, L.; Gualdi, V.; Viganò, R.; Sironi, G.; Besozzi, M.; Castiglioni, B.; Lanfranchi, P.; et al. Clonal diversity, virulence-associated genes and antimicrobial resistance profile of *Staphylococcus aureus* isolates from nasal cavities and soft tissue infections in wild ruminants in Italian Alps. *Vet. Microbiol.* **2014**, *170*, 157–161. [[CrossRef](#)] [[PubMed](#)]

57. Zou, G.; Matuszewska, M.; Jia, M.; Zhou, J.; Ba, X.; Duan, J.; Zhang, C.; Zhao, J.; Tao, M.; Fan, J. A Survey of Chinese Pig Farms and Human Healthcare Isolates Reveals Separate Human and Animal Methicillin-Resistant *Staphylococcus aureus* Populations. *Adv. Sci.* **2021**, *9*, 2103388. [[CrossRef](#)] [[PubMed](#)]
58. Driebe, E.M.; Sahl, J.W.; Roe, C.; Bowers, J.R.; Schupp, J.M.; Gillece, J.D.; Kelley, E.; Price, L.B.; Pearson, T.R.; Hepp, C.M. Using whole genome analysis to examine recombination across diverse sequence types of *Staphylococcus aureus*. *PLoS ONE* **2015**, *10*, e0130955. [[CrossRef](#)] [[PubMed](#)]
59. Michiels, B.; Appelen, L.; Franck, B.; den Heijer, C.D.J.; Bartholomeeusen, S.; Coenen, S. *Staphylococcus aureus*, including methicillin-resistant *Staphylococcus aureus*, among general practitioners and their patients: A cross-sectional study. *PLoS ONE* **2015**, *10*, e0140045. [[CrossRef](#)] [[PubMed](#)]
60. Zurita, J.; Barba, P.; Ortega-Paredes, D.; Mora, M.; Rivadeneira, S. Local circulating clones of *Staphylococcus aureus* in Ecuador. *Braz. J. Infect. Dis.* **2016**, *20*, 525–533. [[CrossRef](#)]
61. Goudarzi, H.; Goudarzi, M.; Sabzehali, F.; Fazeli, M.; Salimi Chirani, A. Genetic analysis of methicillin-susceptible *Staphylococcus aureus* clinical isolates: High prevalence of multidrug-resistant ST239 with strong biofilm-production ability. *J. Clin. Lab. Anal.* **2020**, *34*, e23494. [[CrossRef](#)]
62. Silva, V.; Lopes, A.F.; Soeiro, V.; Caniça, M.; Manageiro, V.; Pereira, J.E.; Maltez, L.; Capelo, J.L.; Igrejas, G.; Poeta, P. Nocturnal Birds of Prey as Carriers of *Staphylococcus aureus* and Other *Staphylococci*: Diversity, Antimicrobial Resistance and Clonal Lineages. *Antibiotics* **2022**, *11*, 240. [[CrossRef](#)]
63. Silva, V.; Almeida, F.; Carvalho, J.A.; Castro, A.P.; Ferreira, E.; Manageiro, V.; Tejedor-Junco, M.T.; Caniça, M.; Igrejas, G.; Poeta, P. Emergence of community-acquired methicillin-resistant *Staphylococcus aureus* EMRSA-15 clone as the predominant cause of diabetic foot ulcer infections in Portugal. *Eur. J. Clin. Microbiol. Infect. Dis.* **2020**, *39*, 179–186. [[CrossRef](#)]
64. Silva, V.; Alfarela, C.; Caniça, M.; Manageiro, V.; Nóvoa, M.; Leiva, B.; Kress, M.; Capelo, J.L.; Poeta, P.; Igrejas, G. A One Health Approach Molecular Analysis of *Staphylococcus aureus* Reveals Distinct Lineages in Isolates from Miranda Donkeys (*Equus asinus*) and Their Handlers. *Antibiotics* **2022**, *11*, 374. [[CrossRef](#)]
65. Papadimitriou-Olivgeris, M.; Drougka, E.; Fligou, F.; Dodou, V.; Kolonitsiou, F.; Filos, K.S.; Anastassiou, E.D.; Petinaki, E.; Marangos, M.; Spiliopoulou, I. Spread of Tst-Positive *Staphylococcus aureus* Strains Belonging to ST30 Clone among Patients and Healthcare Workers in Two Intensive Care Units. *Toxins* **2017**, *9*, 270. [[CrossRef](#)]
66. Chuang, Y.-Y.; Huang, Y.-C. Livestock-associated methicillin-resistant *Staphylococcus aureus* in Asia: An emerging issue? *Int. J. Antimicrob. Agents* **2015**, *45*, 334–340. [[CrossRef](#)] [[PubMed](#)]
67. Peeters, L.E.J.; Argudín, M.A.; Azadikhah, S.; Butaye, P. Antimicrobial resistance and population structure of *Staphylococcus aureus* recovered from pigs farms. *Vet. Microbiol.* **2015**, *180*, 151–156. [[CrossRef](#)] [[PubMed](#)]
68. Yan, X.; Yu, X.; Tao, X.; Zhang, J.; Zhang, B.; Dong, R.; Xue, C.; Grundmann, H.; Zhang, J. *Staphylococcus aureus* ST398 from slaughter pigs in northeast China. *Int. J. Med. Microbiol.* **2014**, *304*, 379–383. [[CrossRef](#)] [[PubMed](#)]
69. Lo, Y.P.; Wan, M.T.; Chen, M.M.; Su, H.Y.; Lauderdale, T.L.; Chou, C.C. Molecular characterization and clonal genetic diversity of methicillin-resistant *Staphylococcus aureus* of pig origin in Taiwan. *Comp. Immunol. Microbiol. Infect. Dis.* **2012**, *35*, 513–521. [[CrossRef](#)] [[PubMed](#)]
70. Lee, C.-Y.; Fang, Y.-P.; Chang, Y.-F.; Wu, T.-H.; Yang, Y.-Y.; Huang, Y.-C. Comparison of molecular epidemiology of bloodstream methicillin-resistant *Staphylococcus aureus* isolates between a new and an old hospital in central Taiwan. *Int. J. Infect. Dis.* **2019**, *79*, 162–168. [[CrossRef](#)]
71. Sinlapasorn, S.; Lulitanond, A.; Angkititrakul, S.; Chanawong, A.; Wilailuckana, C.; Tavichakorntrakool, R.; Chindawong, K.; Seelaget, C.; Krasaesom, M.; Chartchai, S.; et al. SCCmec IX in methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative staphylococci from pigs and workers at pig farms in Khon Kaen, Thailand. *J. Med. Microbiol.* **2015**, *64*, 1087–1093. [[CrossRef](#)]
72. Windahl, U.; Ågren, J.; Holst, B.S.; Börjesson, S. Colonization with methicillin-resistant *Staphylococcus pseudintermedius* in multi-dog households: A longitudinal study using whole genome sequencing. *Vet. Microbiol.* **2016**, *189*, 8–14. [[CrossRef](#)]
73. Yarbrough, M.L.; Lainhart, W.; Burnham, C.A.D. Epidemiology, Clinical Characteristics, and Antimicrobial Susceptibility Profiles of Human Clinical Isolates of *Staphylococcus intermedius* Group. *J. Clin. Microbiol.* **2022**, *56*, e01788–e01817. [[CrossRef](#)]
74. Cuny, C.; Layer, F.; Hansen, S.; Werner, G.; Witte, W. Nasal Colonization of Humans with Occupational Exposure to Raw Meat and to Raw Meat Products with Methicillin-Susceptible and Methicillin-Resistant *Staphylococcus aureus*. *Toxins* **2019**, *11*, 190. [[CrossRef](#)]
75. Tavares, A.; Miragaia, M.; Rolo, J.; Coelho, C.; de Lencastre, H. High prevalence of hospital-associated methicillin-resistant *Staphylococcus aureus* in the community in Portugal: Evidence for the blurring of community–hospital boundaries. *Eur. J. Clin. Microbiol. Infect. Dis.* **2013**, *32*, 1269–1283. [[CrossRef](#)]
76. Strauß, L.; Stegger, M.; Akpaka, P.E.; Alabi, A.; Breurec, S.; Coombs, G.; Egyir, B.; Larsen, A.R.; Laurent, F.; Monecke, S.; et al. Origin, evolution, and global transmission of community-acquired *Staphylococcus aureus* ST8. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E10596–E10604. [[CrossRef](#)] [[PubMed](#)]
77. Ruiz-Ripa, L.; Simón, C.; Ceballos, S.; Ortega, C.; Zarazaga, M.; Torres, C.; Gómez-Sanz, E.S. *pseudintermedius* and *S. aureus* lineages with transmission ability circulate as causative agents of infections in pets for years. *BMC Vet. Res.* **2021**, *17*, 42. [[CrossRef](#)] [[PubMed](#)]

78. Lahiri, S.D.; McLaughlin, R.E.; Whiteaker, J.D.; Ambler, J.E.; Alm, R.A. Molecular characterization of MRSA isolates bracketing the current EUCAST ceftaroline-susceptible breakpoint for *Staphylococcus aureus*: The role of PBP2a in the activity of ceftaroline. *J. Antimicrob. Chemother.* **2015**, *70*, 2488–2498. [[CrossRef](#)] [[PubMed](#)]
79. Cavalcante, F.S.; Abad, E.D.; Lyra, Y.C.; Saintive, S.B.; Ribeiro, M.; Ferreira, D.C.; dos Santos, K.R.N. High prevalence of methicillin resistance and PVL genes among *Staphylococcus aureus* isolates from the nares and skin lesions of pediatric patients with atopic dermatitis. *Braz. J. Med. Biol. Res.* **2015**, *48*, 588–594. [[CrossRef](#)]
80. Hirose, M.; Aung, M.S.; Fukuda, A.; Yahata, S.; Fujita, Y.; Saitoh, M.; Hirose, Y.; Urushibara, N.; Kobayashi, N. Antimicrobial Resistance and Molecular Epidemiological Characteristics of Methicillin-Resistant and Susceptible *Staphylococcal* Isolates from Oral Cavity of Dental Patients and Staff in Northern Japan. *Antibiotics* **2021**, *10*, 1316. [[CrossRef](#)]
81. Ruimy, R.; Angebault, C.; Djossou, F.; Dupont, C.; Epelboin, L.; Jarraud, S.; Armand Lefevre, L.; Bes, M.; Elena Lixandru, B.; Bertine, M.; et al. Are host genetics the predominant determinant of persistent nasal *Staphylococcus aureus* carriage in humans? *J. Infect. Dis.* **2010**, *202*, 924–934. [[CrossRef](#)]
82. Van der Mee-Marquet, N.; Corvaglia, A.-R.; Valentin, A.-S.; Hernandez, D.; Bertrand, X.; Girard, M.; Kluytmans, J.; Donnio, P.-Y.; Quentin, R.; François, P. Analysis of prophages harbored by the human-adapted subpopulation of *Staphylococcus aureus* CC398. *Infect. Genet. Evol.* **2013**, *18*, 299–308. [[CrossRef](#)]
83. Cuny, C.; Abdelbary, M.; Layer, F.; Werner, G.; Witte, W. Prevalence of the immune evasion gene cluster in *Staphylococcus aureus* CC398. *Vet. Microbiol.* **2015**, *177*, 219–223. [[CrossRef](#)]
84. Stegger, M.; Liu, C.M.; Larsen, J.; Soldanova, K.; Aziz, M.; Contente-Cuomo, T.; Petersen, A.; Vandendriessche, S.; Jiménez, J.N.; Mammina, C.; et al. Rapid Differentiation between Livestock-Associated and Livestock-Independent *Staphylococcus aureus* CC398 Clades. *PLoS ONE* **2013**, *8*, e79645. [[CrossRef](#)]
85. Argudín, M.A.; Deplano, A.; Vandendriessche, S.; Dodémont, M.; Nonhoff, C.; Denis, O.; Roisin, S. CC398 *Staphylococcus aureus* subpopulations in Belgian patients. *Eur. J. Clin. Microbiol. Infect. Dis.* **2018**, *37*, 911–916. [[CrossRef](#)]
86. Lekkerkerk, W.S.N.; Van Wamel, W.J.B.; Sniijders, S.V.; Willems, R.J.; Van Duijkeren, E.; Broens, E.M.; Wagenaar, J.A.; Lindsay, J.A.; Vos, M.C. What Is the Origin of Livestock-Associated Methicillin-Resistant *Staphylococcus aureus* Clonal Complex 398 Isolates from Humans without Livestock Contact? An Epidemiological and Genetic Analysis. *J. Clin. Microbiol.* **2015**, *53*, 1836–1841. [[CrossRef](#)] [[PubMed](#)]
87. Teixeira, I.M.; de Oliveira Ferreira, E.; de Araújo Penna, B. Dogs as reservoir of methicillin resistant coagulase negative staphylococci strains – A possible neglected risk. *Microb. Pathog.* **2019**, *135*, 103616. [[CrossRef](#)] [[PubMed](#)]
88. Schmidt, V.M.; Williams, N.J.; Pinchbeck, G.; Corless, C.E.; Shaw, S.; McEwan, N.; Dawson, S.; Nuttall, T. Antimicrobial resistance and characterisation of staphylococci isolated from healthy Labrador retrievers in the United Kingdom. *BMC Vet. Res.* **2014**, *10*, 17. [[CrossRef](#)] [[PubMed](#)]
89. Phumthanakorn, N.; Prapasarakul, N.; Yindee, J.; Gronsang, D. Frequency, Distribution, and Antimicrobial Resistance of Coagulase-Negative Staphylococci Isolated from Clinical Samples in Dogs and Cats. *Microb. Drug Resist.* **2021**, *28*, 236–243. [[CrossRef](#)] [[PubMed](#)]
90. Wu, C.; Zhang, X.; Liang, J.; Li, Q.; Lin, H.; Lin, C.; Liu, H.; Zhou, D.; Lu, W.; Sun, Z.; et al. Characterization of florfenicol resistance genes in the coagulase-negative *Staphylococcus* (CoNS) isolates and genomic features of a multidrug-resistant *Staphylococcus lentus* strain H29. *Antimicrob. Resist. Infect. Control* **2021**, *10*, 9. [[CrossRef](#)]
91. Monecke, S.; Müller, E.; Schwarz, S.; Hotzel, H.; Ehrlich, R. Rapid microarray-based identification of different *mecA* alleles in staphylococci. *Antimicrob. Agents Chemother.* **2012**, *56*, 5547–5554. [[CrossRef](#)]
92. Nemeghaire, S.; Argudín, M.A.; Feßler, A.T.; Hauschild, T.; Schwarz, S.; Butaye, P. The ecological importance of the *Staphylococcus sciuri* species group as a reservoir for resistance and virulence genes. *Vet. Microbiol.* **2014**, *171*, 342–356. [[CrossRef](#)]