

Kent Academic Repository

Full text document (pdf)

Citation for published version

Oakley, Joseph L. and Bicknell, Jake Emmerson (2022) The impacts of tropical agriculture on biodiversity: a meta-analysis. *Journal of Applied Ecology*. (In press)

DOI

Link to record in KAR

<https://kar.kent.ac.uk/97269/>

Document Version

Author's Accepted Manuscript

Copyright & reuse

Content in the Kent Academic Repository is made available for research purposes. Unless otherwise stated all content is protected by copyright and in the absence of an open licence (eg Creative Commons), permissions for further reuse of content should be sought from the publisher, author or other copyright holder.

Versions of research

The version in the Kent Academic Repository may differ from the final published version.

Users are advised to check <http://kar.kent.ac.uk> for the status of the paper. **Users should always cite the published version of record.**

Enquiries

For any further enquiries regarding the licence status of this document, please contact:

researchsupport@kent.ac.uk

If you believe this document infringes copyright then please contact the KAR admin team with the take-down information provided at <http://kar.kent.ac.uk/contact.html>

The impacts of tropical agriculture on biodiversity: a meta-analysis

Journal:	<i>Journal of Applied Ecology</i>
Manuscript ID	JAPPL-2022-00125.R1
Manuscript Type:	Research Article
Date Submitted by the Author:	17-Aug-2022
Complete List of Authors:	Oakley, Joseph; University of Kent Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation Bicknell, Jake; University of Kent Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation
Key-words:	Agriculture, Biodiversity, Food systems, Tropical, Meta-Analysis, Crops, Ecosystem services, Intensification

SCHOLARONE™
Manuscripts

12 **Abstract**

13 1. Biodiversity underpins all food production and strengthens agricultural resilience to crop
14 failure. However, agricultural expansion is the primary driver of biodiversity loss,
15 particularly in the tropics where crop production is increasing and intensifying rapidly to
16 meet a growing global food demand. It is therefore crucial to ask, how do different crops
17 and crop production systems impact biodiversity?

18 2. We first use the FAO database of harvested crop area to explore temporal changes in crop
19 area and intensification across the entire tropical realm. We show that the harvested area of
20 tropical crops has more than doubled since 1961, with ever-increasing intensification. The
21 harvested area in 2019 was 7.21 million km², equivalent to 5.5% of global ice-free land area,
22 or 11.5% of land area in the tropics.

23 3. Second, we conducted a meta-analysis of 194 studies and 1,368 pairwise comparisons to
24 assess the impact of tropical agriculture on biodiversity, comparing biodiversity values in
25 food crop sites versus natural reference habitats.

26 4. Our meta-analysis shows that crop type, rotation time and level of shading are important
27 determinants of biodiversity assemblages. Perennial tropical crops that are grown in shaded
28 plantations or agroforests (e.g., banana and coffee) support higher biodiversity, while crops
29 cultivated in unshaded and often homogeneous croplands (e.g., maize, sugarcane, and oil
30 palm), and particularly annual crops, have impoverished biodiversity communities.

31 5. *Policy implications:* These findings inform our understanding of how different crops and
32 crop production systems impact biodiversity, and may serve as a warning sign for
33 agricultural systems that rely on the ecological functions provided by biodiversity.

34 **Keywords:** agriculture, biodiversity, crops, ecosystem services, food systems, intensification,
35 meta-analysis, tropical

36 **Introduction**

37 Biodiversity underpins all food production and strengthens agricultural resilience to crop
38 failure due to the ecological functions that animals provide (Bélanger & Pilling 2019).
39 However, many of the species that perform these functions are disappearing, in part due to the
40 intensification of agricultural systems (Bélanger & Pilling 2019; Foley *et al.* 2005; Figure S1).
41 With the demand for food predicted to double by 2050 from 2010 levels (Springmann *et al.*
42 2018), food security is an increasingly important global issue (Rosegrant & Cline 2003). It is
43 therefore important to consider how different crop production systems impact biodiversity
44 communities.

45

46 Agricultural expansion is a major driver of habitat loss (Curtis *et al.* 2018; Foley *et al.* 2005;
47 Phalan *et al.* 2013) and one of the most detrimental disturbances to biodiversity assemblages
48 (Gibson *et al.* 2011; Green, 2005; Newbold *et al.* 2014). Over the last sixty years, the
49 production of different tropical crops has increased by varying degrees (Phalan *et al.* 2013). In
50 the next three decades, to meet a growing demand for food, it is predicted that agricultural
51 expansion will continue to increase. This is expected to occur mostly in poorer countries
52 throughout the tropics, where land for crop production often comes at the expense of natural
53 habitats (Tilman *et al.* 2011).

54

55 The tropics are extremely biodiverse, with tropical forests alone containing more than two-
56 thirds of the world's terrestrial biodiversity (Giam 2017). The presence of wild animals in
57 ecosystems is important due to the ecological functions and ecosystem services that they
58 provide, such as pollination, seed dispersal, nutrient cycling, energy flow through trophic
59 levels, and pest control (Bélanger & Pilling 2019; Mathieu *et al.* 2005; Valencia-Aguilar *et al.*
60 2013; Willig *et al.* 2007). Therefore, the promotion of biodiversity in agricultural systems,

61 alongside appropriate management, can provide these benefits in addition to high crop yields
62 (Bélanger & Pilling 2019; Clough *et al.* 2011). In some taxa, particularly birds and bats,
63 agricultural conversion affects the relative composition of functional groups. Insectivorous and
64 carnivorous species that provide pest control services often decline, whilst the proportion of
65 frugivores, nectarivores and granivores may increase, depending on food availability within
66 the cropland (Mtsetfwa *et al.* 2018; Tschardtke *et al.* 2008; Willig *et al.* 2007). These changes
67 affect the ability of biodiversity communities to perform functions important to food
68 production, particularly pollination and pest control (Bélanger & Pilling 2019).

69

70 The magnitude to which agriculture affects biodiversity varies greatly between different crops
71 and agricultural management practices. For example, rice fields are generally less biodiverse
72 than the natural forests or wetlands that they replace (Mathieu *et al.* 2005; Tschardtke *et al.*
73 2008). However, well-managed rice fields can maintain biodiversity and provide important
74 foraging and breeding grounds for some birds, including rare species (Elphick *et al.*, 2010).
75 Forest conversion for oil palm is the one of the greatest threats to biodiversity in Southeast
76 Asia, characterised by the loss of high conservation value species, and overall, harbouring
77 fewer species than natural forests (Fitzherbert *et al.* 2008; Wilcove & Koh 2010). Crops such
78 as coffee and cacao, when grown in shaded plantations, support a greater diversity than those
79 grown in open monocultures, since they provide arboreal habitats and are more structurally
80 similar to natural forests (Estrada, *et al.* 1997; Zermeño-Hernández *et al.* 2016). In addition to
81 the ecological conditions of croplands, crop rotation times (e.g., perennial or annual), proximity
82 to natural habitats, fragmentation, and connectivity are other major factors that influence the
83 capacity for agricultural areas to support biodiversity (Haddad *et al.* 2015; Şekercioğlu *et al.*
84 2019).

85

86 Despite numerous studies on the impacts of tropical food crops on biodiversity, most are
87 limited to certain crops, taxa, and geographic regions. Therefore, a global analysis to identify
88 and compare the impacts of individual tropical food crops on biodiversity assemblages is
89 needed. Here we explore trends in crop production in the tropics between 1961 and 2019,
90 identifying the crops which have expanded the most. We then present a meta-analysis to assess
91 the impacts of tropical agriculture on animal diversity. We investigate whether biodiversity
92 impacts vary between different crops, shading levels, crop rotation times, taxonomic groups,
93 and geographic regions. We expected that agricultural systems that are structurally complex,
94 or similar to natural counterparts (e.g. shaded crops), would maintain biodiversity closer to
95 natural levels, whilst crop sites that are homogeneous and structurally simple (e.g. unshaded
96 crops) would harbour impoverished biodiversity assemblages. Furthermore, we hypothesised
97 that perennial crops such as coffee, cacao and banana would better support biodiversity than
98 annual crops such as maize and sugarcane. We also expected to see differences in agricultural
99 impacts between different geographic regions, due to the variation in crop species and
100 agricultural practices in different parts of the world. Quantifying the impacts of different food
101 crops and their cultivation approaches on biodiversity can inform our understanding of changes
102 to the ecological contribution of biodiversity in tropical agricultural landscapes. In turn, this
103 may inform potential improvements to agricultural practices, and the long-term sustainability
104 of tropical food production.

105

106 **Materials and Methods**

107 **Quantifying tropical crop expansion**

108 In order to quantify crop expansion in the tropics, following Phalan *et al.* (2013), we defined
109 tropical countries as those with at least one-third of their land area situated within the tropics.

110 We used this definition because data on crop harvesting were only available as totals per

111 country for each crop. We used data from FAOSTAT (fao.org/faostat/) on the production and
112 area harvested for all food crops in 115 tropical countries for the years 1961-2019. The
113 harvested area of each of the 137 crops was totalled in each year to compute pan-tropical
114 estimates for each crop's total harvested area per year, and changes in harvested area.

115

116 While the FAO provides some of the best available data on crop harvesting, it must be
117 acknowledged that it has some limitations, so caution must be taken when interpreting the data.
118 Where annual crops are harvested in rotation on the same land multiple times a year, they are
119 all counted towards crop harvesting data, so may lead to overestimations of the true harvested
120 land area. Conversely, underestimations may also occur since crop harvesting data excludes
121 areas where crops were planted but not harvested due to natural calamities or economic reasons.
122 Additionally, there are discrepancies in the reporting of data between countries, with some
123 reporting the entire cultivated area of perennial crops, while others report only the productive
124 area (FAO 2011). Therefore, there may be some discrepancies between the reported and true
125 harvested areas of crops, but the results are likely to be indicative of trends.

126

127 **Literature search to quantify agricultural impacts on biodiversity**

128 To quantify the relative impacts of different tropical crops on biodiversity, we first conducted
129 a rapid evidence assessment (REA) to search for peer-reviewed studies measuring biodiversity
130 in both food crops sites and natural reference sites, based on inclusion and exclusion criteria
131 (described below). We used Web of Science to search for studies published prior to 9th June
132 2020.

133

134 After trialling various search strings, we finally conducted our search using the query: TS =
135 (*tropic* AND (agricultur* OR farm* OR plantation* OR crop* OR agroforest*)) AND

136 (biodiversity OR wildlife OR *fauna* OR bird* OR mammal* OR bat* OR reptil* OR
137 amphibia* OR insect* OR invertebrate*) AND (abundance* OR *diversit* OR richness* OR
138 communit*). We restricted search results to journals within the subject areas: ecology,
139 environmental sciences, biodiversity conservation, entomology, forestry, multidisciplinary
140 sciences, agriculture multidisciplinary, zoology, and ornithology. We limited our search to
141 English language studies, with no restrictions on the date of publication. This search returned
142 3,900 results (Figure 1).

143

144 The lead author (JLO) subsequently screened the retrieved studies for relevance based on the
145 title, abstract, and full text of the articles. A conservative approach was taken in the inclusion
146 of papers during the title and abstract screening to reduce errors of omission. Both authors
147 screened a subset of studies independently and assessed the level of agreement using Cohen's
148 kappa statistic (Cohen 1960), scoring 0.8, to ensure the inclusion criteria were applied
149 consistently. Studies that met our inclusion criteria: (a) reported vertebrate or
150 macroinvertebrate species richness, density, or abundance within both an area cultivated for
151 food crops and a paired natural landscape of any size with little or no disturbance - yielding us
152 a pairwise comparison for the calculation of effect sizes in the meta-analysis, (b) were located
153 within the tropics, and (c) provided or allowed us to calculate the mean, standard deviation,
154 and sample size, from which we could compute an effect size. We were unable to calculate
155 effect sizes for pairwise comparisons where the standard deviation was zero or the sample size
156 was one, therefore they were excluded. We also excluded pairwise comparisons where food
157 crops were mixed with other anthropogenic land uses (e.g., pasture). Studies that measured
158 biodiversity in aquatic ecosystems within agricultural and reference sites (e.g., streams,
159 irrigated croplands or wetlands) were included.

160

161 Our screening process resulted in 194 studies (Figure 1; Table S1) which contributed to our
162 final dataset, amounting to a total of 1,364 pairwise comparisons for 13 crop categories (Table
163 S2), from 34 countries (Figure S2), spanning five geographic regions (Tables S1-S2): Africa
164 ($N_{\text{studies}}=38$, $N_{\text{comparisons}}=281$), Asia ($N_{\text{studies}}=55$, $N_{\text{comparisons}}=432$), Central America ($N_{\text{studies}}=48$,
165 $N_{\text{comparisons}}=371$), South America ($N_{\text{studies}}=52$, $N_{\text{comparisons}}=278$), and Oceania ($N_{\text{studies}}=1$,
166 $N_{\text{comparisons}}=2$). Brazil, Malaysia, Mexico, and Indonesia were the most well-studied countries,
167 comprising more than 50% of all studies (Figure S2). Macroinvertebrates were the most well-
168 represented group ($N_{\text{comparisons}}=613$), followed by birds ($N_{\text{comparisons}}=428$), mammals
169 ($N_{\text{comparisons}}=248$), herpetofauna ($N_{\text{comparisons}}=65$), and fish ($N_{\text{comparisons}}=10$).

170

171 **Data extraction and meta-analysis**

172 The data that met the inclusion criteria were extracted by JLO with a second opinion from JEB
173 where necessary. For each pairwise comparison, we extracted the mean and standard deviation
174 of the biodiversity data. Where studies reported median values, we used these directly (Higgins
175 *et al.*, 2019). We converted standard error, interquartile ranges and confidence intervals to
176 standard deviation. Data were extracted from tables, figures or the text of each study. For those
177 that presented data graphically, we used WebPlotDigitiser (<https://apps.automeris.io/wpd/>) to
178 extract the data. Where studies provided multiple pairwise comparisons (e.g., different crops,
179 taxonomic groups, or geographic locations) we recorded each separately. For those that
180 provided separate pairwise comparisons for food crops and other agricultural or anthropogenic
181 habitats with reference sites, we only extracted the food crop comparison. We considered
182 sample sizes as the number of independent sites within a study. For each pairwise comparison,
183 we also recorded the taxonomic group (birds, fish, herpetofauna, invertebrates, or mammals),
184 geographic region (Africa, Asia, Central America, South America, or Oceania), crop rotation
185 time, and level of shading. We divided shading into three categories: ‘Shaded crops’ were those

186 characterised by natural or planted shade trees above the crop in question. This was most
187 common in cacao and coffee; ‘Unshaded crops’ contained crops grown in open land with sparse
188 or no shade trees. In the case of large crops, e.g. oil palm, where the mature trees create shade
189 we considered these unshaded crops because they were not being shaded by a second
190 vegetation type; finally, ‘Crops with some vegetation’ included those which the authors stated
191 had moderate levels of shade trees, understory vegetation, or something to a similar effect.
192 These were classified to the best of our ability with the information available in the papers. We
193 calculated an effect size for individual crops if there were at least four studies reporting data
194 for that crop. For single crops represented by fewer than four studies, we grouped these and
195 reported them as ‘all other tropical crops’ (e.g., ‘brazil nut’, or ‘pineapple’). When biodiversity
196 values were provided for sites that did not distinguish between multiple different crops, we
197 reported them as ‘mixed tropical crops’ (e.g., ‘annual crops’, or ‘sugarcane, pineapple, and
198 banana’). We divided data into four categories for crop rotation time, classified as ‘annual’,
199 ‘perennial’, ‘mixed’, or ‘unknown’ if the crops were not specified.

200

201 To assess the magnitude of the impact of tropical agriculture upon biodiversity, we calculated
202 the Hedges’ g effect size of the standardized mean difference between agricultural and natural
203 reference sites. Some studies provided multiple pairwise comparisons with a common control
204 (natural reference) site, so we accounted for the potential non-independence of these by nesting
205 them within study, computing a mean for each study (Borenstein *et al.*, 2009). We used a
206 random-effects model, which weighted each comparison by the inverse of within-study
207 variance and between-study variance (Borenstein *et al.* 2009; Koricheva *et al.* 2013).

208

209 In cases where data were extracted from figures, and the variance was so small that it was
210 indiscernible from the mean, we recorded the variance as 0.001 so that an effect size could be

211 computed. The effect direction was reported as positive for cases where the biodiversity value
212 was more favourable in the reference site than the agricultural site, and negative for cases where
213 the biodiversity value was less favourable in the reference site than agricultural site. In cases
214 where there was a greater abundance and/or diversity of invasive species in the agricultural
215 site, this was deemed negative. Therefore, a negative effect size indicates that the agricultural
216 site had an impoverished biodiversity community, and a positive effect size indicates that the
217 agricultural site supported higher levels of biodiversity than the reference site. We considered
218 effect sizes to be significant if the confidence interval did not overlap zero (Koricheva *et al.*
219 2013).

220

221 We calculated the mean effect size for the overall dataset, and the mean effect size for each of
222 the moderator variables (crop type, shading, crop rotation time, taxonomic group, geographic
223 region, and biodiversity metric – richness or abundance). Where fewer than four studies were
224 used for each category, they contributed to the calculation of the overall effect size, but were
225 otherwise not displayed separately in Figure 3.

226

227 To test for publication bias, we followed Nakagawa *et al.* (2017). As such, we plotted funnel
228 plots of standard error and precision for Hedges' g (Figure S3), and calculated the Classic Fail-
229 safe N. The Classic Fail-safe N was 5,151, which means that we would need to locate and
230 include 5,151 null studies in order to overturn the significance of our results (Borenstein *et al.*
231 2009; Koricheva *et al.* 2013). The symmetry of the funnel plots and high Fail-safe N suggest
232 that publication bias is minimal or non-existent in our dataset. We conducted all meta-analyses
233 in the Comprehensive Meta-analysis v3.0 software (Borenstein *et al.* 2013).

234

235 **Results**

236 **Crop expansion**

237 According to the FAO data, the summed harvested area of crops in tropical countries in 2019
238 was 7.21 million km² (Figure 2), equivalent to 5.5% of global ice-free land area, or 11.5% of
239 land area in the tropics (i.e., approximately equivalent to the size of the Australian continent).
240 The top ten crops by harvested area in tropical countries in 2019 were rice, maize, soybeans,
241 wheat, sorghum, beans, millet, oil palm, cassava, and groundnuts, which together accounted
242 for two-thirds (67%) of total harvested area (Figure 2a). Across the tropics, the total area of
243 harvested land has more than doubled between 1961 and 2019 (Figure 2b). The mean annual
244 rate of expansion has accelerated in the past two decades, almost doubling in 2000-2019
245 compared to that of 1980-1999. Production has increased at a greater rate than harvested area
246 (Figure S1), showing the overall increasing intensification of tropical food production.

247

248 Between 1961 and 2019, soybeans were the most rapidly expanding crop both in terms of
249 absolute area, increasing by 0.54 million km² (Figure 2c), and percentage, increasing by
250 4,597% (Figure 2d). After soybeans, maize, rice, and oil palm expanded most in absolute area,
251 while oil palm, cow peas, and sugarcane increased by the greatest percentage.

252

253 **Biodiversity impacts**

254 Our results suggest that, overall, food crop expansion has contributed towards biodiversity loss
255 in tropical regions, although the direction and magnitude of the impact depends on the crop,
256 level of shading, rotation time, taxonomic group, and geographic region. The overall effect of
257 tropical agriculture upon biodiversity is negative and significantly different from zero (Figure
258 3; mean Hedges' g [\pm 95% CI] = -0.59 [-0.67 to -0.51], $p < 0.001$; Table S3).

259

260 Exploring the data by crops, we found that effect sizes were negative and significantly different
261 from zero in maize, oil palm, sugarcane, ‘all other tropical crops’, tea, rice, cacao, and ‘mixed
262 tropical crop’ sites, compared with natural habitats (Figure 3a; Table S3). Biodiversity
263 responses were in general negative but not significant in citrus, allspice, and coffee plantations,
264 while banana and mixed cacao and coffee plantations showed a positive effect size, though not
265 significantly different from zero. Examining our results by level of shading, we found that for
266 shaded and unshaded crops, biodiversity showed a negative and significant difference from
267 zero, with unshaded crops having a considerably greater negative effect size than shaded crops
268 (Figure 3b; Table S3). However, we do not find a significant affect for crops with some
269 vegetation where the confidence intervals were particularly wide and overlapped zero. We find
270 that crop rotation time is an important determinant of impacts, with annual crops showing a
271 greater negative response than perennial crops that have longer rotation periods, though both
272 categories had a significantly negative effect size (Figure 3c). Effect sizes were negative but
273 not significantly different from zero for croplands with ‘mixed’ annual and perennial crops,
274 and for ‘unknown’ where studies didn’t specify whether crops were annual or perennial.
275 Exploring the results by taxonomic group, we found that bird, herpetofauna, and invertebrate
276 assemblages showed significantly negative effect sizes of similar magnitudes in response to
277 agricultural treatments, while mammal responses were negative but not significant (Figure 3d;
278 Table S3). Examining our results by geographic region, we found there was a significantly
279 negative effect of agriculture on biodiversity in all tropical regions (Figure 3e; Table S3). Asia
280 showed the greatest negative response, followed by South America, Africa, and lastly, Central
281 America. Finally, comparing by biodiversity metric, effect sizes for both richness and
282 abundance were negative and significantly different from zero, with richness showing the
283 strongest response to agriculture (Figure S4). Breaking this down by crop, in all cases, the
284 effect sizes for richness were consistently more negative than those for abundance, more likely

285 to be significant, and in the case of coffee, there was evidence of a positive effect on abundance
286 (Figure S4).

287

288 **Discussion**

289 Our study supports the existing literature highlighting the adverse impacts of tropical
290 agriculture upon animal assemblages (Chapman *et al.* 2019; Gibson *et al.*, 2011; Ocampo-
291 Ariza *et al.* 2019; Ramamonjisoa *et al.* 2020). Adding to this, our meta-analysis is the first to
292 compare the magnitude and direction of the impacts of different food crops across the whole
293 of the tropics, and demonstrates that agricultural conversion across a range of ecosystems has
294 an effect on biodiversity, depending on the type of crop and intensity of land use. We also
295 demonstrate the sheer scale of tropical crop expansion (Figure 2), with our findings of increased
296 acceleration of crop expansion over the past two decades corroborating those of Potapov *et al.*
297 (2022), which are based on remote sensing data. Potapov *et al.* (2022) further emphasises the
298 magnitude of crop expansion in the tropics, showing that globally, conversion of natural
299 vegetation to croplands was proportionately largest in Africa, Southeast Asia, and South
300 America. Our results demonstrate that intensification is increasing year-on-year due to
301 production increases out-accelerating area increases (Figure S1). Intensification is particularly
302 concerning because there is increasing evidence that croplands with impoverished biodiversity
303 communities can produce lower yields, and require higher levels of chemical inputs (Bélanger
304 & Pilling 2019). This is therefore due in part to intensification undermining the pollination and
305 other services provided by biodiversity, because of the impact intensification has on
306 biodiversity assemblages as illustrated herein. Indeed, in general, it is known that crop systems
307 support widespread, common, and generalist species, while more specialist, disturbance-
308 sensitive, endemic, and threatened species are likely to be absent (Gallmetzer & Schulze 2015;
309 Şekercioğlu 2012), and along with them, their specific functions lost.

310

311 A particularly important finding from our study is the relative impacts from different crop
312 production systems. We show that unshaded crops result in the most impoverished biodiversity
313 communities, however, the effects varied greatly depending on the crop species. Impoverished
314 biodiversity in agricultural sites could be associated with reduced structural complexity, the
315 removal of understory vegetation, destructive land management practices (Bohada-Murillo *et al.*
316 *al.* 2020; Castaño-Villa *et al.* 2014; Zermeño-Hernández *et al.* 2016), use of agrochemicals
317 (Smith *et al.*, 2016; Zermeño-Hernández *et al.* 2016), reduced resource availability (Mang &
318 Brodie, 2015), changes in soil quality and communities (Franco *et al.* 2019; Smith *et al.* 2016),
319 and an increase in pest or invasive species (Paini *et al.* 2016; Suzán *et al.* 2008). Crops grown
320 in systems that are structurally complex or similar to natural ecosystems, such as agroforests
321 (e.g., some cacao, coffee, and banana plantations), harbour biodiversity closer to natural levels
322 (Estrada *et al.* 1997; Zermeño-Hernández *et al.* 2016). The substantially smaller impact of
323 shaded crops than unshaded crops highlights the potential for improving agricultural practices
324 to reduce biodiversity loss, and this may explain why abundance in some coffee plantations
325 can increase. The wide confidence intervals for crops with some vegetation could be due to
326 fewer studies, or variation in the capacity for different types of vegetation to support
327 biodiversity (e.g., croplands with scattered shade trees provide a different habitat from those
328 with an intact understory). Better measures of agricultural intensity could include chemical
329 inputs, monoculture vs polyculture, and weed richness and cover, however, these details were
330 often not reported in studies. We also show that crops that are harvested on an annual basis,
331 such as maize, sugarcane and rice, result in greater biodiversity impacts when compared with
332 crops that have longer rotation periods, such as coffee, tea, citrus, allspice, cacao and banana.
333 However, oil palm (a perennial with ~25-year rotation cycles) which has significant impacts
334 on biodiversity, does not follow this trend. This may be due to oil palm often being planted

335 within large-scale, high-yield monocultures, but also the fact that 80% of oil palm is produced
336 in the highly biodiverse Southeast Asia biodiversity hotspot, much of this replacing tropical
337 forests (Fitzherbert *et al.* 2008).

338

339 While our findings provide insights into the impacts of different crops on biodiversity, there is
340 a distinct lack of data for most crops. Of the top ten crops in terms of harvested area in the
341 tropics, our REA only returned enough studies for rice, maize, and oil palm to be analysed
342 individually. The large negative effect size of the ‘all other tropical crops’ category highlights
343 the need for more research on understudied crops to identify their individual impacts. Despite
344 soybeans being the most rapidly expanding crop in recent decades, we only found one study
345 reporting biodiversity in soybean sites with data that met our criteria for the meta-analysis
346 (Moura *et al.* 2013). Soybean expansion is well documented, particularly in Brazil. It has been
347 responsible for large areas of deforestation of the Amazon and habitat loss in the globally
348 important Cerrado biome (Kastens *et al.* 2017; Soterroni *et al.* 2019). Nonetheless, the
349 biodiversity impacts of soybeans are understudied compared with other tropical crops such as
350 cacao, coffee, and oil palm, which account for considerably less harvested land area
351 (fao.org/faostat/). Many lesser-known crops are grown by small-scale subsistence farmers and
352 are less likely to gain attention from conservationists than industrially produced crops that are
353 traded internationally (Balmford *et al.* 2012). Our REA also showed some geographic bias in
354 the papers we found. In the Neotropics, research is concentrated in Brazil, Mexico, Costa Rica,
355 Colombia, and Peru, and in Asia the majority of studies come from Malaysia, Indonesia, and
356 India (Figure S2). Most other countries provided few or no studies; research in tropical Oceania
357 is particularly limited. Gaps in our dataset may be due in part to our restricting of the literature
358 search to English language studies. It must therefore be acknowledged that language bias
359 presents a limitation to our study. The inclusion of non-English languages could provide further

360 data and consequently potentially alter effect sizes (Konno *et al.* 2020). However, as
361 demonstrated by our assessments of publication bias (described in the Methods), any changes
362 to effect sizes as a result of missing articles, are highly unlikely to overturn the conclusions of
363 the study.

364

365 In our analysis, richness metrics declined more than abundance metrics. Both are concerning,
366 because there is an abundant literature to show that, in general, the first species to be lost under
367 habitat conversion (and therefore reduce richness), are the most sensitive species that are
368 typically of conservation concern (e.g. Newbold *et al.* 2015 and studies therein). On the other
369 hand, reductions in abundance metrics (and richness metrics) indicate potential declines in the
370 provisioning of ecosystem functions performed by key groups. As such, birds which are
371 important mobile seed dispersers and pest controllers, showed the greatest negative response
372 to agricultural conversion while mammals displayed the most tolerance, reflecting the findings
373 of Gibson *et al.* (2011). It has been suggested that large-bodied mammals are often extirpated
374 due to habitat loss, whereas small nonflying mammal and bat populations can thrive in
375 agricultural habitats (Daily *et al.* 2003; Gibson *et al.* 2011; Wearn *et al.* 2017).

376

377 In many studies used in our meta-analysis, reference sites were fragmented landscapes.
378 Evidence suggests that due to fragmentation, 70% of global forest lies within 1 km of the forest
379 edge (Haddad *et al.* 2015). Agricultural land can have adverse impacts upon biodiversity at
380 considerable distances into natural habitats (Hurst *et al.* 2013; Scriven *et al.* 2018). Therefore,
381 biodiversity levels in reference sites would be influenced by factors such as proximity to
382 agricultural land, patch size, connectivity, edge effects, and the intensity of land use in the
383 surrounding matrix (Prugh *et al.* 2008). Consequently, the true effects of agricultural
384 conversion are likely to be greater than our estimates, when considering the additional impacts

385 of fragmentation (Haddad *et al.* 2015). Nonetheless, the relative differences between the
386 impacts of different crops are likely to remain largely the same.

387

388 Understanding the consequences of food cultivation on biodiversity can help to identify
389 improvements to agricultural practices and influence consumer choice. Since much of the food
390 produced in tropical regions is exported internationally, a large proportion of impacts on
391 tropical biodiversity are remotely driven by industrialised countries (Green *et al.* 2019; Lenzen
392 *et al.* 2012). This study therefore provides us with food for thought regarding the positive and
393 negative environmental impacts caused by our food choices. It is particularly pertinent as we
394 are trying to improve the transparency of food supply chains and connecting consumer markets
395 to habitat destruction and biodiversity loss through projects such as Trase
396 (<http://www.trase.earth>). The knowledge gained from this study could also be incorporated into
397 the modelling of future agricultural expansion scenarios (e.g., Chaplin-Kramer *et al.* 2015),
398 helping to identify areas for crop expansion with minimal adverse impacts on biodiversity.
399 Most of all though, our findings may serve as a warning sign for agricultural systems that rely
400 on the ecological functions provided by biodiversity to maximise their yields. This is crucial,
401 because with an ever-increasing global food demand, yield deficits could result in further
402 expansion to the area footprint of tropical agriculture.

403

404

405 **Author contributions**

406 JLO led the manuscript writing, conducted the REA and analysed the data. JEB conceived the
407 study, co-wrote the manuscript and assisted with data analysis.

408

409 **Acknowledgments**

410 We would like to thank C. Gardner for initial discussions regarding the study concept.

411

412 **Conflicts of interests**

413 The authors declare no conflicts of interests.

414

415 **Data availability**

416 Should the manuscript be accepted, the dataset will be archived in the repository Dryad, and
417 the data DOI will be included at the end of the article.

418 **References**

- 419 Balmford, A., Green, R., & Phalan, B. (2012). What conservationists need to know about
420 farming. *Proceedings of the Royal Society B: Biological Sciences*, 279(1739), 2714–
421 2724. doi: 10.1098/rspb.2012.0515
- 422 Bélanger, J., & Pilling, D. (Eds.). (2019). *The State of the World's Biodiversity for Food and*
423 *Agriculture*. Rome: FAO. doi: 10.4060/CA3129EN
- 424 Bohada-Murillo, M., Castaño-Villa, G. J., & Fontúrbel, F. E. (2020). The effects of forestry
425 and agroforestry plantations on bird diversity: A global synthesis. *Land Degradation*
426 *& Development*, 31(5), 646–654. doi: 10.1002/ldr.3478
- 427 Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009). *Introduction to*
428 *Meta-Analysis*. Chichester, U.K: John Wiley & Sons.
- 429 Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2013). *Comprehensive*
430 *Meta-Analysis (Version 2)*. Engelwood, NJ: Biostat.
- 431 Castaño-Villa, G. J., Ramos-Valencia, S. A., & Fontúrbel, F. E. (2014). Fine-scale habitat
432 structure complexity determines insectivorous bird diversity in a tropical forest. *Acta*
433 *Oecologica*, 61, 19–23. doi: 10.1016/j.actao.2014.10.002
- 434 Chaplin-Kramer, R., Sharp, R. P., Mandle, L., Sim, S., Johnson, J., Butnar, I., ... Kareiva, P.
435 M. (2015). Spatial patterns of agricultural expansion determine impacts on
436 biodiversity and carbon storage. *Proceedings of the National Academy of Sciences*,
437 112(24), 7402–7407. doi: 10.1073/pnas.1406485112
- 438 Chapman, P. M., Loveridge, R., Rowcliffe, J. M., Carbone, C., Bernard, H., Davison, C. W.,
439 & Ewers, R. M. (2019). Minimal Spillover of Native Small Mammals From Bornean
440 Tropical Forests Into Adjacent Oil Palm Plantations. *Frontiers in Forests and Global*
441 *Change*, 2, 2. doi: 10.3389/ffgc.2019.00002

- 442 Clough, Y., Barkmann, J., Juhbandt, J., Kessler, M., Wanger, T. C., Anshary, A., ...
443 Tschardtke, T. (2011). Combining high biodiversity with high yields in tropical
444 agroforests. *Proceedings of the National Academy of Sciences*, *108*(20), 8311–8316.
445 doi: 10.1073/pnas.1016799108
- 446 Cohen, J. (1960). A coefficient of agreement for nominal scales. *Educational and*
447 *Psychological Measurement*, *20*(1), 37–46.
448 <https://doi.org/10.1177/001316446002000104>
- 449 Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A., & Hansen, M. C. (2018). Classifying
450 drivers of global forest loss. *Science (New York, N.Y.)*, *361*(6407), 1108–1111. doi:
451 10.1126/science.aau3445
- 452 Daily, G. C., Ceballos, G., Pacheco, J., Suzán, G., & Sánchez-Azofeifa, A. (2003).
453 Countryside Biogeography of Neotropical Mammals: Conservation Opportunities in
454 Agricultural Landscapes of Costa Rica. *Conservation Biology*, *17*(6), 1814–1826. doi:
455 10.1111/j.1523-1739.2003.00298.x
- 456 Elphick, C. S., Baicich, P., Parsons, K. C., Fasola, M., & Mugica, L. (2010). The Future for
457 Research on Waterbirds in Rice Fields. *Waterbirds*, *33*(sp1), 231. doi:
458 10.1675/063.033.s117
- 459 Estrada, A., Coates-Estrada, R., & Meritt, D. A. (1997). Anthropogenic landscape changes
460 and avian diversity at Los Tuxtlas, Mexico. *Biodiversity & Conservation*, *6*(1), 19–43.
461 doi: 10.1023/A:1018328930981
- 462 FAO (2011). *Tree crops. Guidelines for estimating area data*. Rome: Food and Agriculture
463 Organization of the United Nations: <https://www.fao.org/3/cb2463en/cb2463en.pdf>.
464 Accessed April 2022
- 465 FAOSTAT. Food and Agriculture Organization of the United Nations:
466 <http://www.fao.org/faostat/>. Accessed June 2021.

- 467 Fitzherbert, E., Struebig, M., Morel, A., Danielsen, F., Bruhl, C., Donald, P., & Phalan, B.
468 (2008). How will oil palm expansion affect biodiversity? *Trends in Ecology &*
469 *Evolution*, 23(10), 538–545. doi: 10.1016/j.tree.2008.06.012
- 470 Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., ... Snyder, P.
471 K. (2005). Global Consequences of Land Use. *Science*, 309(5734), 570–574. doi:
472 10.1126/science.1111772
- 473 Franco, A. L. C., Sobral, B. W., Silva, A. L. C., & Wall, D. H. (2019). Amazonian
474 deforestation and soil biodiversity. *Conservation Biology*, 33(3), 590–600. doi:
475 10.1111/cobi.13234
- 476 Gallmetzer, N., & Schulze, C. H. (2015). Impact of oil palm agriculture on understory
477 amphibians and reptiles: A Mesoamerican perspective. *Global Ecology and*
478 *Conservation*, 4, 95–109. doi: 10.1016/j.gecco.2015.05.008
- 479 Giam, X. (2017). Global biodiversity loss from tropical deforestation. *Proceedings of the*
480 *National Academy of Sciences*, 114(23), 5775–5777. doi: 10.1073/pnas.1706264114
- 481 Gibson, L., Lee, T. M., Koh, L. P., Brook, B. W., Gardner, T. A., Barlow, J., ... Sodhi, N. S.
482 (2011). Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature*,
483 478(7369), 378–381. doi: 10.1038/nature10425
- 484 Green, J. M. H., Croft, S. A., Durán, A. P., Balmford, A. P., Burgess, N. D., Fick, S., ...
485 West, C. D. (2019). Linking global drivers of agricultural trade to on-the-ground
486 impacts on biodiversity. *Proceedings of the National Academy of Sciences*, 116(46),
487 23202–23208. doi: 10.1073/pnas.1905618116
- 488 Green, R. E. (2005). Farming and the Fate of Wild Nature. *Science*, 307(5709), 550–555. doi:
489 10.1126/science.1106049


- 490 Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., ...
491 Townshend, J. R. (2015). Habitat fragmentation and its lasting impact on Earth's
492 ecosystems. *Science Advances*, *1*(2), e1500052. doi: 10.1126/sciadv.1500052
- 493 Higgins, J. P. T., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., & Welch, V. A.
494 (Eds.). (2019). *Cochrane Handbook for Systematic Reviews of Interventions* (2nd
495 Edition). Chichester: John Wiley & Sons.
- 496 Hurst, Z. M., McCleery, R. A., Collier, B. A., Fletcher, R. J., Silvy, N. J., Taylor, P. J., &
497 Monadjem, A. (2013). Dynamic Edge Effects in Small Mammal Communities across
498 a Conservation-Agricultural Interface in Swaziland. *PLoS ONE*, *8*(9), e74520. doi:
499 10.1371/journal.pone.0074520
- 500 Kastens, J. H., Brown, J. C., Coutinho, A. C., Bishop, C. R., & Esquerdo, J. C. D. M. (2017).
501 Soy moratorium impacts on soybean and deforestation dynamics in Mato Grosso,
502 Brazil. *PLOS ONE*, *12*(4), e0176168. doi: 10.1371/journal.pone.0176168
- 503 Konno, K., Akasaka, M., Koshida, C., Katayama, N., Osada, N., Spake, R., & Amano, T.
504 (2020). Ignoring non-English-language studies may bias ecological meta-analyses.
505 *Ecology and Evolution*, *10*(13), 6373–6384. <https://doi.org/10.1002/ece3.6368>
- 506 Koricheva, J., Gurevitch, J., & Mengersen, K. (2013). *Handbook of Meta-analysis in Ecology
507 and Evolution*. Princeton: Princeton University Press.
- 508 Lenzen, M., Moran, D., Kanemoto, K., Foran, B., Lobefaro, L., & Geschke, A. (2012).
509 International trade drives biodiversity threats in developing nations. *Nature*,
510 *486*(7401), 109–112. doi: 10.1038/nature11145
- 511 Mang, S. L., & Brodie, J. F. (2015). Impacts of non-oil tree plantations on biodiversity in
512 Southeast Asia. *Biodiversity and Conservation*, *24*(14), 3431–3447. doi:
513 10.1007/s10531-015-1022-5

- 514 Mathieu, J., Rossi, J.-P., Mora, P., Lavelle, P., Martins, P. F. da S., Rouland, C., & Grimaldi,
515 M. (2005). Recovery of Soil Macrofauna Communities after Forest Clearance in
516 Eastern Amazonia, Brazil. *Conservation Biology*, *19*(5), 1598–1605. doi:
517 10.1111/j.1523-1739.2005.00200.x
- 518 Moura, N. G., Lees, A. C., Andretti, C. B., Davis, B. J. W., Solar, R. R. C., Aleixo, A., ...
519 Gardner, T. A. (2013). Avian biodiversity in multiple-use landscapes of the Brazilian
520 Amazon. *Biological Conservation*, *167*, 339–348. doi: 10.1016/j.biocon.2013.08.023
- 521 Mtsetfwa, F., McCleery, R. A., & Monadjem, A. (2018). Changes in bat community
522 composition and activity patterns across a conservation-agriculture boundary. *African*
523 *Zoology*, *53*(3), 99–106. doi: 10.1080/15627020.2018.1531726
- 524 Nakagawa, S., Noble, D. W. A., Senior, A. M., & Lagisz, M. (2017). Meta-evaluation of
525 meta-analysis: Ten appraisal questions for biologists. *BMC Biology*, *15*(1), 18. doi:
526 10.1186/s12915-017-0357-7
- 527 Newbold, T., Hudson, L. N., Hill, S. L. L., Contu, S., Lysenko, I., Senior, R. A., ... Purvis,
528 A. (2015). Global effects of land use on local terrestrial biodiversity. *Nature*,
529 *520*(7545), 45–50. doi: 10.1038/nature14324
- 530 Newbold, T., Hudson, L. N., Phillips, H. R. P., Hill, S. L. L., Contu, S., Lysenko, I., ...
531 Purvis, A. (2014). A global model of the response of tropical and sub-tropical forest
532 biodiversity to anthropogenic pressures. *Proceedings of the Royal Society B:*
533 *Biological Sciences*, *281*(1792), 20141371. doi: 10.1098/rspb.2014.1371
- 534 Ocampo-Ariza, C., Denis, K., Njie Motombi, F., Bobo, K. S., Kreft, H., & Waltert, M.
535 (2019). Extinction thresholds and negative responses of Afrotropical ant-following
536 birds to forest cover loss in oil palm and agroforestry landscapes. *Basic and Applied*
537 *Ecology*, *39*, 26–37. doi: 10.1016/j.baae.2019.06.008

- 538 Paini, D. R., Sheppard, A. W., Cook, D. C., De Barro, P. J., Worner, S. P., & Thomas, M. B.
539 (2016). Global threat to agriculture from invasive species. *Proceedings of the*
540 *National Academy of Sciences*, *113*(27), 7575–7579. doi: 10.1073/pnas.1602205113
- 541 Phalan, B., Bertzky, M., Butchart, S. H. M., Donald, P. F., Scharlemann, J. P. W.,
542 Stattersfield, A. J., & Balmford, A. (2013). Crop Expansion and Conservation
543 Priorities in Tropical Countries. *PLOS ONE*, *8*(1), e51759. doi:
544 10.1371/journal.pone.0051759
- 545 Potapov, P., Turubanova, S., Hansen, M. C., Tyukavina, A., Zalles, V., Khan, A., Song, X.-
546 P., Pickens, A., Shen, Q., & Cortez, J. (2022). Global maps of cropland extent and
547 change show accelerated cropland expansion in the twenty-first century. *Nature Food*,
548 *3*(1), 19–28. <https://doi.org/10.1038/s43016-021-00429-z>
- 549 Prugh, L. R., Hodges, K. E., Sinclair, A. R. E., & Brashares, J. S. (2008). Effect of habitat
550 area and isolation on fragmented animal populations. *Proceedings of the National*
551 *Academy of Sciences*, *105*(52), 20770–20775. doi: 10.1073/pnas.0806080105
- 552 Ramamonjisoa, N., Sakai, M., Ndirantsoa, S. H., Kakehashi, R., Kurabayashi, A., Tomaru,
553 N., & Natuhara, Y. (2020). Hotspots of stream tadpole diversity in forest and
554 agricultural landscapes in Ranomafana, Madagascar. *Landscape and Ecological*
555 *Engineering*, *16*(3), 207–221. doi: 10.1007/s11355-020-00407-w
- 556 Rosegrant, M. W., & Cline, S. A. (2003). Global Food Security: Challenges and Policies.
557 *Science*, *302*(5652), 1917–1919. doi: 10.1126/science.1092958
- 558 Scriven, S. A., Gillespie, G. R., Laimun, S., & Goossens, B. (2018). Edge effects of oil palm
559 plantations on tropical anuran communities in Borneo. *Biological Conservation*, *220*,
560 37–49. doi: 10.1016/j.biocon.2018.02.006

- 561 Şekercioglu, Ç. H. (2012). Bird functional diversity and ecosystem services in tropical
562 forests, agroforests and agricultural areas. *Journal of Ornithology*, 153(1), 153–161.
563 doi: 10.1007/s10336-012-0869-4
- 564 Şekercioglu, Ç. H., Mendenhall, C. D., Oviedo-Brenes, F., Horns, J. J., Ehrlich, P. R., &
565 Daily, G. C. (2019). Long-term declines in bird populations in tropical agricultural
566 countryside. *Proceedings of the National Academy of Sciences*, 116(20), 9903–9912.
567 doi: 10.1073/pnas.1802732116
- 568 Smith, P., House, J. I., Bustamante, M., Sobocká, J., Harper, R., Pan, G., ... Pugh, T. A. M.
569 (2016). Global change pressures on soils from land use and management. *Global
570 Change Biology*, 22(3), 1008–1028. doi: 10.1111/gcb.13068
- 571 Soterroni, A. C., Ramos, F. M., Mosnier, A., Fargione, J., Andrade, P. R., Baumgarten, L., ...
572 Polasky, S. (2019). Expanding the Soy Moratorium to Brazil's Cerrado. *Science
573 Advances*, 5(7), eaav7336. doi: 10.1126/sciadv.aav7336
- 574 Springmann, M., Clark, M., Mason-D'Croz, D., Wiebe, K., Bodirsky, B. L., Lassaletta, L., ...
575 Willett, W. (2018). Options for keeping the food system within environmental limits.
576 *Nature*, 562(7728), 519–525. doi: 10.1038/s41586-018-0594-0
- 577 Suzán, G., Armién, A., Mills, J. N., Marcé, E., Ceballos, G., Ávila, M., ... Yates, T. L.
578 (2008). Epidemiological considerations of rodent community composition in
579 fragmented landscapes in Panama. *Journal of Mammalogy*, 89(3), 684–690. doi:
580 10.1644/07-MAMM-A-015R1.1
- 581 Tilman, D., Balzer, C., Hill, J., & Befort, B. L. (2011). Global food demand and the
582 sustainable intensification of agriculture. *Proceedings of the National Academy of
583 Sciences*, 108(50), 20260–20264. doi: 10.1073/pnas.1116437108


- 584 Tschardtke, T., Sekercioglu, C. H., Dietsch, T. V., Sodhi, N. S., Hoehn, P., & Tylianakis, J.
585 M. (2008). Landscape constraints on functional diversity of birds and insects in
586 tropical agroecosystems. *Ecology*, 89(4), 944–951. doi: 10.1890/07-0455.1
- 587 Valencia-Aguilar, A., Cortés-Gómez, A. M., & Ruiz-Agudelo, C. A. (2013). Ecosystem
588 services provided by amphibians and reptiles in Neotropical ecosystems. *International
589 Journal of Biodiversity Science, Ecosystem Services & Management*, 9(3), 257–272.
590 doi: 10.1080/21513732.2013.821168
- 591 Wearn, O. R., Rowcliffe, J. M., Carbone, C., Pfeifer, M., Bernard, H., & Ewers, R. M.
592 (2017). Mammalian species abundance across a gradient of tropical land-use
593 intensity: A hierarchical multi-species modelling approach. *Biological Conservation*,
594 212, 162–171. doi: 10.1016/j.biocon.2017.05.007
- 595 Wilcove, D. S., & Koh, L. P. (2010). Addressing the threats to biodiversity from oil-palm
596 agriculture. *Biodiversity and Conservation*, 19(4), 999–1007. doi: 10.1007/s10531-
597 009-9760-x
- 598 Willig, M. R., Presley, S. J., Bloch, C. P., Hice, C. L., Yanoviak, S. P., Díaz, M. M., ...
599 Weaver, S. C. (2007). Phyllostomid Bats of Lowland Amazonia: Effects of Habitat
600 Alteration on Abundance. *Biotropica*, 39(6), 737–746. doi: 10.1111/j.1744-
601 7429.2007.00322.x
- 602 Zermeño-Hernández, I., Pingarrón, A., & Martínez-Ramos, M. (2016). Agricultural land-use
603 diversity and forest regeneration potential in human- modified tropical landscapes.
604 *Agriculture, Ecosystems & Environment*, 230, 210–220. doi:
605 10.1016/j.agee.2016.06.007
606

607 **Figures**

608


609 **Figure 1.** PRISMA diagram of the number of studies included during each filtering stage of

610 the rapid evidence assessment. See methods for inclusion criteria.



611

612 **Figure 2.** Changes in harvested area of tropical crops from 1961-2019. (a) Harvested area of
 613 individual food crops. (b) Total harvested area of food crops. (c) Increase in harvested area of
 614 food crops by absolute area and (d) by percentage, in tropical countries from 1961-2019. The
 615 top ten tropical crops by area in 2019 are shown. Additionally, sugarcane and cow peas, which
 616 were in the top ten by area increase, are also shown. The harvested areas of ‘all other tropical
 617 crops’ were combined. Data: FAOSTAT.



618

← More impoverished biodiversity community in agricultural area compared to reference habitat

619 **Figure 3.** Effect sizes of agricultural impacts on biodiversity by (a) crop, (b) intensity, (c) crop
 620 rotation time, (d) taxonomic group (omitting fish $N_{\text{studies}}=3$), and (e) geographic region
 621 (omitting Oceania $N_{\text{studies}}=1$). The number of pairwise comparisons between agricultural and
 622 reference sites per category is reported in parentheses. The black vertical lines show the mean
 623 standardised effect size (Hedges' g), and 95% CI are indicated by the width of the boxes. Effect
 624 sizes are significant if the confidence intervals do not overlap zero. The tall vertical black lines
 625 and grey dashed lines represent an effect size of zero and mean overall effect size respectively.
 626 For single crops represented by fewer than four studies, we grouped these and reported them
 627 as 'all other tropical crops'. When biodiversity values were provided for sites that did not
 628 distinguish between multiple different crops, we reported them as 'mixed tropical crops'

1

2

3

4

5

**The impacts of tropical agriculture on biodiversity: a meta-
analysis**

6

7


Joseph L. Oakley and Jake E. Bicknell

8

9

10

Supporting Information



11

12 **Figure S1.** Changes in crop production by weight and harvested area from 1961 to 2019.


13 Changes are measured as the percentage changes for each year compared to 1961. The top ten

14 crops by harvested area in 2019 are shown, as well as sugarcane, cowpeas, ‘all other tropical

15 crops’ and the total. Data: FAOSTAT.

16

17





18

19 **Figure S2** Map of study sites by country. Country colour illustrates the number of studies per20 country ($N_{\text{countries}}=34$, $N_{\text{studies}}=194$).

21


22

a**b**

23

24 **Figure S3.** Publication bias in the meta-analysis. (a) Funnel plot of the relationship between
25 mean effect size and standard error for each study. (b) Funnel plot of the relationship between
26 mean effect size and the precision of each study.

27



28

29 **Figure S4.** Effect sizes of agricultural impacts on biodiversity, broken down by crop type and
 30 by biodiversity metric (richness or abundance). The black vertical lines show the mean
 31 standardised effect size (Hedges' g), and 95% CI are indicated by the width of the lines. Where
 32 these go beyond the scale of -4 to +4, this is indicated by an arrow. Effect sizes are significant
 33 if the confidence intervals do not overlap zero. For single crops represented by fewer than four
 34 studies, we grouped these and reported them as 'all other tropical crops'. When biodiversity
 35 values were provided for sites that did not distinguish between multiple different crops, we
 36 reported them as 'mixed tropical crops'. Overall effect sizes for abundance and richness are
 37 found in Table S3.

Table S1. Details of the studies included in the meta-analysis

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Ackerman IL, Constantino R, Gauch, Jr HG, Lehmann J, Riha SJ, Fernandes ECM. 2009. Termite (Insecta: Isoptera) Species Composition in a Primary Rain Forest and Agroforests in Central Amazonia. <i>Biotropica</i> 41:226–233.	Shaded crops	Mixed tropical crops	Invertebrates	South America	Unknown
Adedoja O, Kehinde T. 2018. Changes in interaction network topology and species composition of flower-visiting insects across three land use types. <i>African Journal of Ecology</i> 56:964–971.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Annual
Allen L, Reeve R, Nousek-McGregor A, Villacampa J, MacLeod R. 2019. Are orchid bees useful indicators of the impacts of human disturbance? <i>Ecological Indicators</i> 103:745–755.	Unshaded crops	Banana	Invertebrates	South America	Perennial
	Shaded crops	Banana	Invertebrates	South America	Perennial
Almeida SM, Silva LC, Cardoso MR, Cerqueira PV, Juen L, Santos MPD. 2016. The effects of oil palm plantations on the functional diversity of Amazonian birds. <i>Journal of Tropical Ecology</i> 32:510–525.	Unshaded crops	Oil palm	Birds	South America	Perennial
Alonso-Rodríguez AM, Finegan B, Fiedler K. 2017. Neotropical moth assemblages degrade due to oil palm expansion. <i>Biodiversity and Conservation</i> 26:2295–2326.	Unshaded crops	Oil palm	Invertebrates	Central America	Perennial
Alvarez-Alvarez EA, Corcuera P, Almazán-Núñez RC. 2018. Spatiotemporal variation in the structure and diet types of bird assemblages in tropical dry forest in southwestern Mexico. <i>The Wilson Journal of Ornithology</i> 130:457.	Unshaded crops	Mixed tropical crops	Birds	Central America	Mixed
Arellano L, Favila ME, Huerta C. 2005. Diversity of dung and carrion beetles in a disturbed Mexican tropical montane cloud forest and on shade coffee plantations. <i>Biodiversity and Conservation</i> 14:601–615.	Shaded crops	Coffee	Invertebrates	Central America	Perennial
Arenas-Clavijo A, Armbrrecht I. 2019. Soil ants (Hymenoptera: Formicidae) and ground beetles (Coleoptera: Carabidae) in a coffee agroforestry landscape during a severe-drought period. <i>Agroforestry Systems</i> 93:1781–1792.	Unshaded crops	Coffee	Invertebrates	South America	Perennial
	Shaded crops	Coffee	Invertebrates	South America	Perennial
Ashwini KM, Sridhar KR. 2006. Seasonal abundance and activity of pill millipedes (<i>Arthrosphaera magna</i>) in mixed plantation and semi-evergreen forest of southern India. <i>Acta Oecologica</i> 29:27–32.	Shaded crops	Mixed tropical crops	Invertebrates	Asia	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Azhar B, Lindenmayer DB, Wood J, Fischer J, Manning A, McElhinny C, Zakaria M. 2011. The conservation value of oil palm plantation estates, smallholdings and logged peat swamp forest for birds. <i>Forest Ecology and Management</i> 262:2306–2315.	Unshaded crops	Oil palm	Birds	Asia	Perennial
Bakermans MH, Vitz AC, Rodewald AD, Rengifo CG. 2009. Migratory songbird use of shade coffee in the Venezuelan Andes with implications for conservation of cerulean warbler. <i>Biological Conservation</i> 142:2476–2483.	Shaded crops	Coffee	Birds	South America	Perennial
Barnes AD et al. 2017. Direct and cascading impacts of tropical land-use change on multi-trophic biodiversity. <i>Nature Ecology & Evolution</i> 1:1511–1519.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
	Unshaded crops	Oil palm	Birds	Asia	Perennial
	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Basset Y et al. 2008. Changes in Arthropod Assemblages along a Wide Gradient of Disturbance in Gabon. <i>Conservation Biology</i> 22:1552–1563.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Mixed
Beck J, Schulze CH, Linsenmair KE, Fiedler K. 2002. From forest to farmland: diversity of geometrid moths along two habitat gradients on Borneo. <i>Journal of Tropical Ecology</i> 18:33–51.	Unshaded crops	Mixed tropical crops	Invertebrates	Asia	Mixed
Belshaw R, Bolton B. 1993. The effect of forest disturbance on the leaf litter ant fauna in Ghana. <i>Biodiversity and Conservation</i> 2:656–666.	Unshaded crops	Cacao	Invertebrates	Africa	Perennial
Bennett RE, Leuenberger W, Bosarreyes Leja BB, Sagone Cáceres A, Johnson K, Larkin J. 2018. Conservation of Neotropical migratory birds in tropical hardwood and oil palm plantations. <i>PLOS ONE</i> 13:e0210293.	Unshaded crops	Oil palm	Birds	Central America	Perennial
Benstead JP, Douglas MM, Pringle CM. 2003. Relationships of stream invertebrate communities to deforestation in eastern Madagascar. <i>Ecological Applications</i> 13:1473–1490.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Annual
Bobo KS, Waltert M, Fermon H, Njokagbor J, Mühlenberg M. 2006. From Forest to Farmland: Butterfly Diversity and Habitat Associations Along a Gradient of Forest Conversion in Southwestern Cameroon. <i>Journal of Insect Conservation</i> 10:29–42.	Shaded crops	Cacao and coffee	Invertebrates	Africa	Perennial
	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Annual
Bos MM, Steffan-Dewenter I, Tschardt T. 2007. The contribution of cacao agroforests to the conservation of lower canopy ant and beetle diversity in Indonesia. <i>Biodiversity and Conservation</i> 16:2429–2444.	Shaded crops	Cacao	Invertebrates	Asia	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Braga RF, Korasaki V, Audino LD, Louzada J. 2012. Are Dung Beetles Driving Dung-Fly Abundance in Traditional Agricultural Areas in the Amazon? <i>Ecosystems</i> 15:1173–1181.	Unshaded crops	Mixed tropical crops	Invertebrates	South America	Unknown
	Shaded crops	Mixed tropical crops	Invertebrates	South America	Perennial
Buechley ER, Şekercioglu ÇH, Atickem A, Gebremichael G, Ndungu JK, Mahamued BA, Beyene T, Mekonnen T, Lens L. 2015. Importance of Ethiopian shade coffee farms for forest bird conservation. <i>Biological Conservation</i> 188:50–60.	Shaded crops	Coffee	Birds	Africa	Perennial
Cabra-García J, Bermúdez-Rivas C, Osorio AM, Chacón P. 2012. Cross-taxon congruence of α and β diversity among five leaf litter arthropod groups in Colombia. <i>Biodiversity and Conservation</i> 21:1493–1508.	Unshaded crops	Sugarcane	Invertebrates	South America	Annual
Cajaiba RL, Périco E, Dalzochio MS, da Silva WB, Bastos R, Cabral JA, Santos M. 2017. Does the composition of Scarabaeidae (Coleoptera) communities reflect the extent of land use changes in the Brazilian Amazon? <i>Ecological Indicators</i> 74:285–294.	Unshaded crops	Cacao	Invertebrates	South America	Perennial
	Unshaded crops	Cacao	Invertebrates	South America	Perennial
	Unshaded crops	Cacao	Invertebrates	South America	Perennial
Carvalho RL, Andresen E, Barônio GJ, Oliveira VHF, Louzada J, Braga RF. 2020. Is dung removal a good proxy for other dung beetle functions when monitoring for conservation? A case study from the Brazilian Amazon. <i>Ecological Indicators</i> 109:105841.	Unshaded crops	Mixed tropical crops	Invertebrates	South America	Mixed
Castillo LE, Martínez E, Ruepert C, Savage C, Gilek M, Pinnock M, Solis E. 2006. Water quality and macroinvertebrate community response following pesticide applications in a banana plantation, Limon, Costa Rica. <i>Science of The Total Environment</i> 367:418–432.	Unshaded crops	Banana	Invertebrates	Central America	Perennial
Céspedes LN, Bayly NJ. 2019. Over-winter ecology and relative density of Canada Warbler <i>Cardellina canadensis</i> in Colombia: the basis for defining conservation priorities for a sharply declining long-distance migrant. <i>Bird Conservation International</i> 29:232–248.	Shaded crops	Coffee	Birds	South America	Perennial
Chapman PM, Loveridge R, Rowcliffe JM, Carbone C, Bernard H, Davison CW, Ewers RM. 2019. Minimal Spillover of Native Small Mammals From Bornean Tropical Forests Into Adjacent Oil Palm Plantations. <i>Frontiers in Forests and Global Change</i> 2:2.	Unshaded crops	Oil palm	Mammals	Asia	Perennial
Chellaiah D, Yule CM. 2018. Riparian buffers mitigate impacts of oil palm plantations on aquatic macroinvertebrate community structure in tropical streams of Borneo. <i>Ecological Indicators</i> 95:53–62.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
	Shaded crops	Oil palm	Invertebrates	Asia	Perennial
	Crops with some vegetation	Oil palm	Invertebrates	Asia	Perennial
Chiawo DO, Kombe WN, Craig AJFK. 2018. Bird responses to land use change: guild diversity in a Kenyan coastal forest and adjoining habitats. <i>Emu - Austral Ornithology</i> 118:281–292.	Unshaded crops	Mixed tropical crops	Birds	Africa	Annual
Clough Y et al. 2016. Land-use choices follow profitability at the expense of ecological functions in Indonesian smallholder landscapes. <i>Nature Communications</i> 7:13137.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
	Unshaded crops	Oil palm	Birds	Asia	Perennial
Costa C, Oliveira VHF, Maciel R, Beiroz W, Korasaki V, Louzada J. 2017. Variegated tropical landscapes conserve diverse dung beetle communities. <i>PeerJ</i> 5:e3125.	Shaded crops	Coffee	Invertebrates	South America	Perennial
Coulibaly T, Akpesse AAM, Boga J-P, Yapi A, Kouassi KP, Roisin Y. 2016. Change in termite communities along a chronosequence of mango tree orchards in the north of Côte d'Ivoire. <i>Journal of Insect Conservation</i> 20:1011–1019.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Mixed
	Crops with some vegetation	All other tropical crops	Invertebrates	Africa	Perennial
	Shaded crops	All other tropical crops	Invertebrates	Africa	Perennial
da Cunha Bitar YO, Juen L, Pinheiro LC, Santos-Costa MC dos. 2015. Anuran Beta Diversity in a Mosaic Anthropogenic Landscape in Transitional Amazon. <i>Journal of Herpetology</i> 49:75–82.	Unshaded crops	Mixed tropical crops	Herpetofauna	South America	Annual
Daily GC, Ceballos G, Pacheco J, Suzán G, Sánchez-Azofeifa A. 2003. Countryside Biogeography of Neotropical Mammals: Conservation Opportunities in Agricultural Landscapes of Costa Rica. <i>Conservation Biology</i> 17:1814–1826.	Unshaded crops	Coffee	Mammals	Central America	Perennial
DaRocha WD, Neves FS, Dáttilo W, Delabie JHC. 2016. Epiphytic bromeliads as key components for maintenance of ant diversity and ant–bromeliad interactions in agroforestry system canopies. <i>Forest Ecology and Management</i> 372:128–136.	Shaded crops	Cacao	Invertebrates	South America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Davies TE, Clarke RH, Ewen JG, Fazey IRA, Pettorelli N, Cresswell W. 2015. The effects of land-use change on the endemic avifauna of Makira, Solomon Islands: endemics avoid monoculture. <i>Emu - Austral Ornithology</i> 115:199–213.	Unshaded crops	Cacao	Birds	Asia	Perennial
	Unshaded crops	Mixed tropical crops	Birds	Asia	Perennial
	Shaded crops	Mixed tropical crops	Birds	Asia	Mixed
De Beenhouwer M, Geeraert L, Mertens J, Van Geel M, Aerts R, Vanderhaegen K, Honnay O. 2016. Biodiversity and carbon storage co-benefits of coffee agroforestry across a gradient of increasing management intensity in the SW Ethiopian highlands. <i>Agriculture, Ecosystems & Environment</i> 222:193–199.	Shaded crops	Coffee	Invertebrates	Africa	Perennial
De la Mora A, Murnen CJ, Philpott SM. 2013. Local and landscape drivers of biodiversity of four groups of ants in coffee landscapes. <i>Biodiversity and Conservation</i> 22:871–888.	Unshaded crops	Coffee	Invertebrates	Central America	Perennial
	Shaded crops	Coffee	Invertebrates	Central America	Perennial
de Lima RF, Dallimer M, Atkinson PW, Barlow J. 2013. Biodiversity and land-use change: understanding the complex responses of an endemic-rich bird assemblage. <i>Diversity and Distributions</i> 19:411–422.	Shaded crops	Mixed tropical crops	Birds	Africa	Mixed
Delabie JHC, Jahyny B, do Nascimento IC, Mariano CSF, Lacau S, Campiolo S, Philpott SM, Leponce M. 2007. Contribution of cocoa plantations to the conservation of native ants (Insecta: Hymenoptera: Formicidae) with a special emphasis on the Atlantic Forest fauna of southern Bahia, Brazil. <i>Biodiversity and Conservation</i> 16:2359–2384.	Shaded crops	Cacao	Invertebrates	South America	Perennial
Dosso K, Roisin Y, Tiho S, Konaté S, Yéo K. 2017. Short-term changes in the structure of termite assemblages associated with slash-and-burn agriculture in Côte d'Ivoire. <i>Biotropica</i> 49:856–861.	Shaded crops	Mixed tropical crops	Invertebrates	Africa	Mixed
Edwards DP et al. 2014a. Selective-logging and oil palm: multitaxon impacts, biodiversity indicators, and trade-offs for conservation planning. <i>Ecological Applications</i> 24:2029–2049.	Unshaded crops	Oil palm	Birds	Asia	Perennial
	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
	Unshaded crops	Oil palm	Mammals	Asia	Perennial
Edwards DP, Hodgson JA, Hamer KC, Mitchell SL, Ahmad AH, Cornell SJ, Wilcove DS. 2010. Wildlife-friendly oil palm plantations fail to protect biodiversity effectively: Farming and the fate of tropical biodiversity. <i>Conservation Letters</i> 3:236–242.	Unshaded crops	Oil palm	Birds	Asia	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Edwards FA, Edwards DP, Hamer KC, Davies RG. 2013. Impacts of logging and conversion of rainforest to oil palm on the functional diversity of birds in Sundaland. <i>Ibis</i> 155:313–326.	Unshaded crops	Oil palm	Birds	Asia	Perennial
Edwards FA, Edwards DP, Larsen TH, Hsu WW, Benedick S, Chung A, Vun Khen C, Wilcove DS, Hamer KC. 2014b. Does logging and forest conversion to oil palm agriculture alter functional diversity in a biodiversity hotspot?: Functional diversity and land-use change in Borneo. <i>Animal Conservation</i> 17:163–173.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Elisei T, Valadares E, Martins CF, Albuquerque FA. 2017. Diversity and Structure of Social Wasps Community (Hymenoptera: Vespidae, Polistinae) in Neotropical Dry Forest. <i>Sociobiology</i> 64:111.	Unshaded crops	Mixed tropical crops	Invertebrates	South America	Unknown
Estrada A, Coates-Estrada R, Dadda AA, Cammarano P. 1998. Dung and carrion beetles in tropical rain forest fragments and agricultural habitats at Los Tuxtlas, Mexico. <i>Journal of Tropical Ecology</i> 14:577–593.	Unshaded crops	Citrus	Invertebrates	Central America	Perennial
	Shaded crops	Coffee	Invertebrates	Central America	Perennial
	Unshaded crops	Allspice	Invertebrates	Central America	Perennial
	Shaded crops	Cacao and coffee	Invertebrates	Central America	Perennial
	Shaded crops	Cacao	Invertebrates	Central America	Perennial
Estrada A, Coates-Estrada R, Meritt D. 1994. Non flying mammals and landscape changes in the tropical rain forest region of Los Tuxtlas, Mexico. <i>Ecography</i> 17:229–241.	Unshaded crops	Citrus	Mammals	Central America	Perennial
	Unshaded crops	Allspice	Mammals	Central America	Perennial
	Shaded crops	Coffee	Mammals	Central America	Perennial
	Shaded crops	Cacao	Mammals	Central America	Perennial
	Shaded crops	Cacao and coffee	Mammals	Central America	Perennial
Estrada A, Coates-Estrada R, Meritt Jr DA. 1997. Anthropogenic changes and avian diversity at Los Tuxtlas, Mexico. <i>Biodiversity and Conservation</i> 6:19–43.	Unshaded crops	Banana	Birds	Central America	Perennial
	Unshaded crops	All other tropical crops	Birds	Central America	Annual

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
	Unshaded crops	Maize	Birds	Central America	Annual
	Unshaded crops	Allspice	Birds	Central America	Perennial
	Shaded crops	Cacao and coffee	Birds	Central America	Perennial
	Unshaded crops	Citrus	Birds	Central America	Perennial
	Shaded crops	Coffee	Birds	Central America	Perennial
	Shaded crops	Cacao	Birds	Central America	Perennial
Estrada A, Coates-Estrada R. 2005. Diversity of Neotropical migratory landbird species assemblages in forest fragments and man-made vegetation in Los Tuxtlas, Mexico. <i>Biodiversity and Conservation</i> 14:1719–1734.	Unshaded crops	Mixed tropical crops	Birds	Central America	Annual
	Unshaded crops	Mixed tropical crops	Birds	Central America	Perennial
	Shaded crops	Mixed tropical crops	Birds	Central America	Perennial
Estrada A, D AA, Coates-Estrada R. 1999. Tropical rain forest fragmentation, howler monkeys (<i>Alouatta palliata</i>), and dung beetles at Los Tuxtlas, Mexico. <i>American Journal of Primatology</i> 48:253–262.	Unshaded crops	Citrus	Invertebrates	Central America	Perennial
	Unshaded crops	Allspice	Invertebrates	Central America	Perennial
	Shaded crops	Coffee	Invertebrates	Central America	Perennial
	Shaded crops	Cacao and coffee	Invertebrates	Central America	Perennial
	Shaded crops	Cacao	Invertebrates	Central America	Perennial
Faria D, Baumgarten J. 2007. Shade cacao plantations (<i>Theobroma cacao</i>) and bat conservation in southern Bahia, Brazil. <i>Biodiversity and Conservation</i> 16:291–312.	Shaded crops	Cacao	Mammals	South America	Perennial
Faria D, Laps RR, Baumgarten J, Cetra M. 2006. Bat and Bird Assemblages from Forests and Shade Cacao Plantations in Two Contrasting Landscapes in the Atlantic Forest of Southern Bahia, Brazil. <i>Biodiversity and Conservation</i> 15:587–612.	Shaded crops	Cacao	Birds	South America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
	Shaded crops	Cacao	Mammals	South America	Perennial
Faria D, Paciencia MLB, Dixo M, Laps RR, Baumgarten J. 2007. Ferns, frogs, lizards, birds and bats in forest fragments and shade cacao plantations in two contrasting landscapes in the Atlantic forest, Brazil. <i>Biodiversity and Conservation</i> 16:2335–2357.	Shaded crops	Cacao	Herpetofauna	South America	Perennial
Faria D. 2006. Phyllostomid bats of a fragmented landscape in the north-eastern Atlantic forest, Brazil. <i>Journal of Tropical Ecology</i> 22:531–542.	Shaded crops	Cacao	Mammals	South America	Perennial
Faruk A, Belabut D, Ahmad N, Knell RJ, Garner TWJ. 2013. Effects of Oil-Palm Plantations on Diversity of Tropical Anurans: Effects of Oil-Palm Plantations on Anurans. <i>Conservation Biology</i> 27:615–624.	Unshaded crops	Oil palm	Herpetofauna	Asia	Perennial
Fayle TM, Turner EC, Snaddon JL, Chey VK, Chung AYC, Eggleton P, Foster WA. 2010. Oil palm expansion into rain forest greatly reduces ant biodiversity in canopy, epiphytes and leaf-litter. <i>Basic and Applied Ecology</i> 11:337–345.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Feijoo A, Carvajal AF, Zúñiga MC, Quintero H, Fragoso C. 2011. Diversity and abundance of earthworms in land use systems in central-western Colombia. <i>Pedobiologia</i> 54:S69–S75.	Unshaded crops	Mixed tropical crops	Invertebrates	South America	Mixed
	Shaded crops	Mixed tropical crops	Invertebrates	South America	Perennial
Filgueiras BKC, Tabarelli M, Leal IR, Vaz-de-Mello FZ, Iannuzzi L. 2015. Dung beetle persistence in human-modified landscapes: Combining indicator species with anthropogenic land use and fragmentation-related effects. <i>Ecological Indicators</i> 55:65–73.	Unshaded crops	Sugarcane	Invertebrates	South America	Annual
	Unshaded crops	Sugarcane	Invertebrates	South America	Annual
Fotso AK, Hanna R, Tindo M, Doumtsop A, Nagel P. 2015. How plants and honeydew-producing hemipterans affect ant species richness and structure in a tropical forest zone. <i>Insectes Sociaux</i> 62:443–453.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Unknown
Franco ALC, Bartz MLC, Cherubin MR, Baretta D, Cerri CEP, Feigl BJ, Wall DH, Davies CA, Cerri CC. 2016. Loss of soil (macro)fauna due to the expansion of Brazilian sugarcane acreage. <i>Science of The Total Environment</i> 563–564:160–168.	Unshaded crops	Sugarcane	Invertebrates	South America	Annual
Freudmann A, Mollik P, Tschapka M, Schulze CH. 2015. Impacts of oil palm agriculture on phyllostomid bat assemblages. <i>Biodiversity and Conservation</i> 24:3583–3599.	Unshaded crops	Oil palm	Mammals	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Frishkoff LO, Karp DS, M’Gonigle LK, Mendenhall CD, Zook J, Kremen C, Hadly EA, Daily GC. 2014. Loss of avian phylogenetic diversity in neotropical agricultural systems. <i>Science</i> 345:1343–1346.	Unshaded crops	Mixed tropical crops	Birds	Central America	Mixed
	Crops with some vegetation	Mixed tropical crops	Birds	Central America	Mixed
Furtado IS, Martins MB. 2018. The impacts of land use intensification on the assembly of drosophilidae (Diptera). <i>Global Ecology and Conservation</i> 16:e00432.	Unshaded crops	Mixed tropical crops	Invertebrates	South America	Perennial
	Unshaded crops	Mixed tropical crops	Invertebrates	South America	Annual
Gallmetzer N, Schulze CH. 2015. Impact of oil palm agriculture on understory amphibians and reptiles: A Mesoamerican perspective. <i>Global Ecology and Conservation</i> 4:95–109.	Unshaded crops	Oil palm	Herpetofauna	Central America	Perennial
Geissen V, Peña-Peña K, Huerta E. 2009. Effects of different land use on soil chemical properties, decomposition rate and earthworm communities in tropical Mexico. <i>Pedobiologia</i> 53:75–86.	Unshaded crops	Banana	Invertebrates	Central America	Perennial
	Shaded crops	Banana	Invertebrates	Central America	Perennial
Gillespie GR, Ahmad E, Elahan B, Evans A, Ancrenaz M, Goossens B, Scroggie MP. 2012. Conservation of amphibians in Borneo: Relative value of secondary tropical forest and non-forest habitats. <i>Biological Conservation</i> 152:136–144.	Unshaded crops	Oil palm	Herpetofauna	Asia	Perennial
Gilroy JJ, Prescott GW, Cardenas JS, Castañeda PG del P, Sánchez A, Rojas-Murcia LE, Medina Uribe CA, Haugaasen T, Edwards DP. 2015. Minimizing the biodiversity impact of Neotropical oil palm development. <i>Global Change Biology</i> 21:1531–1540.	Unshaded crops	Oil palm	Birds	South America	Perennial
	Unshaded crops	Oil palm	Invertebrates	South America	Perennial
	Unshaded crops	Oil palm	Herpetofauna	South America	Perennial
Glor RE, Flecker AS, Benard MF, Power AG. 2001. Lizard diversity and agricultural disturbance in a Caribbean forest landscape. <i>Biodiversity and Conservation</i> 10:711–723.	Unshaded crops	Oil palm	Herpetofauna	Central America	Perennial
	Shaded crops	Cacao	Herpetofauna	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Goodale E, Kotagama SW, Raman TRS, Sidhu S, Goodale U, Parker S, Chen J. 2014. The response of birds and mixed-species bird flocks to human-modified landscapes in Sri Lanka and southern India. <i>Forest Ecology and Management</i> 329:384–392.	Unshaded crops	Mixed tropical crops	Birds	Asia	Unknown
Gordon C, Manson R, Sundberg J, Cruz-Angón A. 2007. Biodiversity, profitability, and vegetation structure in a Mexican coffee agroecosystem. <i>Agriculture, Ecosystems & Environment</i> 118:256–266.	Shaded crops	Coffee	Mammals	Central America	Perennial
	Shaded crops	Coffee	Birds	Central America	Perennial
Gove AD, Hylander K, Nemomissa S, Shimelis A, Enkossa W. 2013. Structurally complex farms support high avian functional diversity in tropical montane Ethiopia. <i>Journal of Tropical Ecology</i> 29:87–97.	Unshaded crops	Mixed tropical crops	Birds	Africa	Unknown
Gray CL, Lewis OT, Chung AYC, Fayle TM. 2015. Riparian reserves within oil palm plantations conserve logged forest leaf litter ant communities and maintain associated scavenging rates. <i>Journal of Applied Ecology</i> 52:31–40.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Greenler SM, Ebersole JJ. 2015. Bird communities in tropical agroforestry ecosystems: an underappreciated conservation resource. <i>Agroforestry Systems</i> 89:691–704.	Shaded crops	Cacao	Birds	Central America	Perennial
Guéi AM, N'Dri JK, Zro FGB, Bakayoko S, Tondoh JE. 2019. Relationships between soil morpho-chemical parameters and earthworm community attributes in tropical agro-ecosystems in the Centre-West region of Côte d'Ivoire, Africa. <i>Tropical Ecology</i> 60:209–218.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Unknown
	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Mixed
	Unshaded crops	Cacao	Invertebrates	Africa	Perennial
Halffter G, Pineda E, Arellano L, Escobar F. 2007. Instability of Copronecrophagous Beetle Assemblages (Coleoptera: Scarabaeinae) in a Mountainous Tropical Landscape of Mexico. <i>Environmental Entomology</i> 36:1397–1407.	Shaded crops	Coffee	Invertebrates	Central America	Perennial
Harada LM, Araújo IS, Overal WL, Silva FAB. 2020. Comparison of dung beetle communities (Coleoptera: Scarabaeidae: Scarabaeinae) in oil palm plantations and native forest in the eastern Amazon, Brazil. <i>Revista Brasileira de Entomologia</i> 64:e2019102.	Unshaded crops	Oil palm	Invertebrates	South America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Harterreiten-Souza ES, Pujol-Luz JR, Sujii ER. 2016. Influence of Various Farmland Habitats on Abundance of Taeniaptera (Diptera: Micropezidae). Florida Entomologist 99:740–743.	Unshaded crops	Mixed tropical crops	Invertebrates	South America	Annual
Harvey CA, Gonzalez J, Somarriba E. 2006. Dung Beetle and Terrestrial Mammal Diversity in Forests, Indigenous Agroforestry Systems and Plantain Monocultures in Talamanca, Costa Rica. Biodiversity and Conservation 15:555–585.	Unshaded crops	All other tropical crops	Invertebrates	Central America	Perennial
	Unshaded crops	All other tropical crops	Mammals	Central America	Perennial
	Shaded crops	Banana	Invertebrates	Central America	Perennial
	Shaded crops	Cacao	Invertebrates	Central America	Perennial
	Shaded crops	Cacao	Mammals	Central America	Perennial
	Shaded crops	Banana	Mammals	Central America	Perennial
Harvey CA, González Villalobos JA. 2007. Agroforestry systems conserve species-rich but modified assemblages of tropical birds and bats. Biodiversity and Conservation 16:2257–2292.	Unshaded crops	All other tropical crops	Birds	Central America	Perennial
	Shaded crops	Banana	Birds	Central America	Perennial
	Shaded crops	Cacao	Birds	Central America	Perennial
	Unshaded crops	All other tropical crops	Mammals	Central America	Perennial
	Shaded crops	Banana	Mammals	Central America	Perennial
	Shaded crops	Cacao	Mammals	Central America	Perennial
Helbig-Bonitz M, Ferger SW, Böhning-Gaese K, Tschapka M, Howell K, Kalko EKV. 2015. Bats are Not Birds - Different Responses to Human Land-use on a Tropical Mountain. Biotropica 47:497–508.	Shaded crops	Coffee	Birds	Africa	Perennial
	Shaded crops	Mixed tropical crops	Birds	Africa	Unknown

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
	Shaded crops	Mixed tropical crops	Mammals	Africa	Unknown
	Shaded crops	Coffee	Mammals	Africa	Perennial
	Shaded crops	Mixed tropical crops	Mammals	Africa	Mixed
Hoehn P, Steffan-Dewenter I, Tschardt T. 2010. Relative contribution of agroforestry, rainforest and openland to local and regional bee diversity. <i>Biodiversity and Conservation</i> 19:2189–2200.	Unshaded crops	Mixed tropical crops	Invertebrates	Asia	Mixed
	Shaded crops	Mixed tropical crops	Invertebrates	Asia	Mixed
Horgan FG. 2009. Invasion and retreat: shifting assemblages of dung beetles amidst changing agricultural landscapes in central Peru. <i>Biodiversity and Conservation</i> 18:3519–3541.	Shaded crops	Coffee	Invertebrates	South America	Perennial
Horner-Devine MC, Daily GC, Ehrlich PR, Boggs CL. 2003. Countryside Biogeography of Tropical Butterflies. <i>Conservation Biology</i> 17:168–177.	Unshaded crops	Coffee	Invertebrates	Central America	Perennial
Huang JC, Rustiati EL, Nusalawo M, Kingston T. 2019. Echolocation and roosting ecology determine sensitivity of forest-dependent bats to coffee agriculture. <i>Biotropica</i> 51:757–768.	Shaded crops	Coffee	Mammals	Asia	Perennial
Huerta E, Kampichler C, Geissen V, Ochoa-Gaona S, Jong B de, Hernández-Daumás S. 2009. Towards an ecological index for tropical soil quality based on soil macrofauna. <i>Pesquisa Agropecuária Brasileira</i> 44:1056–1062.	Unshaded crops	Maize	Invertebrates	Central America	Annual
	Unshaded crops	Mixed tropical crops	Invertebrates	Central America	Mixed
	Unshaded crops	All other tropical crops	Invertebrates	Central America	Perennial
Kapoor V. 2008. Effects of rainforest fragmentation and shade-coffee plantations on spider communities in the Western Ghats, India. <i>Journal of Insect Conservation</i> 12:53–68.	Shaded crops	Coffee	Invertebrates	Asia	Perennial
Kessler M et al. 2009. Alpha and beta diversity of plants and animals along a tropical land-use gradient. <i>Ecological Applications</i> 19:2142–2156.	Shaded crops	Cacao	Birds	Asia	Perennial
	Shaded crops	Cacao	Invertebrates	Asia	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
King DI, Hernandez-Mayorga MD, Trubey R, Raudales R, Rappole JH. 2007. An Evaluation of the Contribution of Cultivated Allspice (<i>Pimenta Dioca</i>) to Vertebrate Biodiversity Conservation in Nicaragua. <i>Biodiversity and Conservation</i> 16:1299–1320.	Shaded crops	Allspice	Mammals	Central America	Perennial
	Shaded crops	Allspice	Herpetofauna	Central America	Perennial
	Shaded crops	Allspice	Birds	Central America	Perennial
Klarner B, Winkelmann H, Krashevskaya V, Maraun M, Widyastuti R, Scheu S. 2017. Trophic niches, diversity and community composition of invertebrate top predators (Chilopoda) as affected by conversion of tropical lowland rainforest in Sumatra (Indonesia). <i>PLOS ONE</i> 12:e0180915.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Knowlton JL, Mata Zayas EE, Ripley AJ, Valenzuela-Cordova B, Collado-Torres R. 2019. Mammal Diversity in Oil Palm Plantations and Forest Fragments in a Highly Modified Landscape in Southern Mexico. <i>Frontiers in Forests and Global Change</i> 2:67.	Unshaded crops	Oil palm	Mammals	Central America	Perennial
Kone M, Konate S, Yeo K, Kouassi PK, Linsenmair KE. 2012. Changes in ant communities along an age gradient of cocoa cultivation in the Oumé region, central Côte d'Ivoire: Ant communities in cocoa plantations. <i>Entomological Science</i> 15:324–339.	Unshaded crops	Cacao	Invertebrates	Africa	Perennial
	Shaded crops	Cacao	Invertebrates	Africa	Perennial
	Crops with some vegetation	Mixed tropical crops	Invertebrates	Africa	Mixed
	Unshaded crops	Cacao	Invertebrates	Africa	Perennial
	Crops with some vegetation	Mixed tropical crops	Invertebrates	Africa	Mixed
	Shaded crops	Cacao	Invertebrates	Africa	Perennial
	Konopik O, Steffan-Dewenter I, Grafe TU. 2015. Effects of Logging and Oil Palm Expansion on Stream Frog Communities on Borneo, Southeast Asia. <i>Biotropica</i> 47:636–643.	Shaded crops	Oil palm	Herpetofauna	Asia
Kudavidanage EP, Wanger TC, Alwis C, Sanjeewa S, Kotagama SW. 2012. Amphibian and butterfly diversity across a tropical land-use gradient in Sri Lanka; implications for conservation decision making: Land-use change affects amphibians and butterflies. <i>Animal Conservation</i> 15:253–265.	Crops with some vegetation	Mixed tropical crops	Herpetofauna	Asia	Unknown

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
	Crops with some vegetation	Mixed tropical crops	Invertebrates	Asia	Unknown
Kühnert K, Grass I, Waltert M. 2019. Sacred groves hold distinct bird assemblages within an Afrotropical savanna. <i>Global Ecology and Conservation</i> 18:e00656.	Unshaded crops	Mixed tropical crops	Birds	Africa	Annual
Kuppler J, Fricke J, Hemp C, Steffan-Dewenter I, Peters MK. 2015. Conversion of savannah habitats to small-scale agriculture affects grasshopper communities at Mt. Kilimanjaro, Tanzania. <i>Journal of Insect Conservation</i> 19:509–518.	Unshaded crops	Maize	Invertebrates	Africa	Annual
Lees AC, Moura NG, de Almeida AS, Vieira ICG. 2015. Poor Prospects for Avian Biodiversity in Amazonian Oil Palm. <i>PLOS ONE</i> 10:e0122432.	Unshaded crops	Oil palm	Birds	South America	Perennial
Livingston G, Jha S, Vega A, Gilbert L. 2013. Conservation Value and Permeability of Neotropical Oil Palm Landscapes for Orchid Bees. <i>PLOS ONE</i> 8:e78523.	Unshaded crops	Oil palm	Invertebrates	Central America	Perennial
López-Ricaurte L, Edwards DP, Romero-Rodríguez N, Gilroy JJ. 2017. Impacts of oil palm expansion on avian biodiversity in a Neotropical natural savanna. <i>Biological Conservation</i> 213:225–233.	Unshaded crops	Oil palm	Birds	South America	Perennial
Love K, Kurz DJ, Vaughan IP, Ke A, Evans LJ, Goossens B. 2017. Bearded pig (<i>Sus barbatus</i>) utilisation of a fragmented forest–oil palm landscape in Sabah, Malaysian Borneo. <i>Wildlife Research</i> 44:603.	Unshaded crops	Oil palm	Mammals	Asia	Perennial
Lucey JM, Tawatao N, Senior MJM, Chey VK, Benedick S, Hamer KC, Woodcock P, Newton RJ, Bottrell SH, Hill JK. 2014. Tropical forest fragments contribute to species richness in adjacent oil palm plantations. <i>Biological Conservation</i> 169:268–276.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Luke SH, Fayle TM, Eggleton P, Turner EC, Davies RG. 2014. Functional structure of ant and termite assemblages in old growth forest, logged forest and oil palm plantation in Malaysian Borneo. <i>Biodiversity and Conservation</i> 23:2817–2832.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
MacGregor-Fors I, Blanco-García A, Lindig-Cisneros R. 2010. Bird community shifts related to different forest restoration efforts: A case study from a managed habitat matrix in Mexico. <i>Ecological Engineering</i> 36:1492–1496.	Unshaded crops	Maize	Birds	Central America	Annual
MacGregor-Fors I, González-García F, Hernández-Lara C, Santiago-Alarcon D. 2018. Where are the birds in the matrix? Avian diversity in a Neotropical landscape mosaic. <i>The Wilson Journal of Ornithology</i> 130:81–93.	Shaded crops	Coffee	Birds	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
MacGregor-Fors I, Schondube JE. 2011. Use of Tropical Dry Forests and Agricultural Areas by Neotropical Bird Communities: Birds in Modified Landscapes. <i>Biotropica</i> 43:365–370.	Unshaded crops	Mixed tropical crops	Birds	Central America	Annual
	Unshaded crops	Mixed tropical crops	Birds	Central America	Perennial
Mandal J, Shankar Raman TR. 2016. Shifting agriculture supports more tropical forest birds than oil palm or teak plantations in Mizoram, northeast India. <i>The Condor</i> 118:345–359.	Unshaded crops	Oil palm	Birds	Asia	Perennial
Martin EA, Viano M, Ratsimisetra L, Laloë F, Carrière SM. 2012. Maintenance of bird functional diversity in a traditional agroecosystem of Madagascar. <i>Agriculture, Ecosystems & Environment</i> 149:1–9.	Unshaded crops	Rice	Birds	Africa	Annual
	Unshaded crops	Mixed tropical crops	Birds	Africa	Annual
Mas AH, Dietsch TV. 2004. Linking shade coffee certification to biodiversity conservation: butterflies and birds in Chiapas, Mexico. <i>Ecological Applications</i> 14:642–654.	Shaded crops	Coffee	Invertebrates	Central America	Perennial
Mathieu J, Rossi J-P, Mora P, Lavelle P, Martins PF da S, Rouland C, Grimaldi M. 2005. Recovery of Soil Macrofauna Communities after Forest Clearance in Eastern Amazonia, Brazil. <i>Conservation Biology</i> 19:1598–1605.	Unshaded crops	Rice	Invertebrates	South America	Annual
Mendenhall CD, Frishkoff LO, Santos-Barrera G, Pacheco J, Mesfun E, Quijano FM, Ehrlich PR, Ceballos G, Daily GC, Pringle RM. 2014. Countryside biogeography of Neotropical reptiles and amphibians. <i>Ecology</i> 95:856–870.	Unshaded crops	Coffee	Herpetofauna	Central America	Perennial
Mendes-Oliveira AC, Peres CA, Maués PCR de A, Oliveira GL, Mineiro IGB, de Maria SLS, Lima RCS. 2017. Oil palm monoculture induces drastic erosion of an Amazonian forest mammal fauna. <i>PLOS ONE</i> 12:e0187650.	Unshaded crops	Oil palm	Mammals	South America	Perennial
Méndez-Castro FE, Rao D. 2014. Spider diversity in epiphytes: Can shade coffee plantations promote the conservation of cloud forest assemblages? <i>Biodiversity and Conservation</i> 23:2561–2577.	Shaded crops	Coffee	Invertebrates	Central America	Perennial
Milder JC, DeCLERCK FAJ, Sanfiorenzo A, Sánchez DM, Tobar DE, Zuckerberg B. 2010. Effects of farm and landscape management on bird and butterfly conservation in western Honduras. <i>Ecosphere</i> 1:art2.	Shaded crops	Coffee	Birds	Central America	Perennial
Milheiras SG, Guedes M, Augusto Barbosa Silva F, Aparício P, Mace GM. 2020. Patterns of biodiversity response along a gradient of forest use in Eastern Amazonia, Brazil. <i>PeerJ</i> 8:e8486.	Shaded crops	All other tropical crops	Invertebrates	South America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Moura NG, Lees AC, Andretti CB, Davis BJW, Solar RRC, Aleixo A, Barlow J, Ferreira J, Gardner TA. 2013. Avian biodiversity in multiple-use landscapes of the Brazilian Amazon. <i>Biological Conservation</i> 167:339–348.	Unshaded crops	Mixed tropical crops	Birds	South America	Annual
	Unshaded crops	Mixed tropical crops	Birds	South America	Mixed
Moya-Raygoza G, Cuevas-Guzmán R, Pinedo-Escatel JA, Morales-Arias JG. 2019. Comparison of Leafhopper (Hemiptera: Cicadellidae) Diversity in Maize and Its Wild Ancestor Teosinte, and Plant Diversity in the Teosinte Habitat. <i>Annals of the Entomological Society of America</i> 112:99–106.	Unshaded crops	Maize	Invertebrates	Central America	Annual
Muhamad D, Okubo S, Miyashita T, Takeuchi K. 2013. Effects of habitat type, vegetation structure, and proximity to forests on bird species richness in a forest–agricultural landscape of West Java, Indonesia. <i>Agroforestry Systems</i> 87:1247–1260.	Shaded crops	Mixed tropical crops	Birds	Asia	Perennial
Mulwa RK, Böhning-Gaese K, Schleuning M. 2012. High Bird Species Diversity in Structurally Heterogeneous Farmland in Western Kenya. <i>Biotropica</i> 44:801–809.	Unshaded crops	Sugarcane	Birds	Africa	Annual
	Crops with some vegetation	Mixed tropical crops	Birds	Africa	Unknown
Mulwa RK, Neuschulz EL, Böhning-Gaese K, Schleuning M. 2013. Seasonal fluctuations of resource abundance and avian feeding guilds across forest–farmland boundaries in tropical Africa. <i>Oikos</i> 122:524–532.	Unshaded crops	Mixed tropical crops	Birds	Africa	Annual
	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Annual
Mumme S, Jochum M, Brose U, Haneda NF, Barnes AD. 2015. Functional diversity and stability of litter-invertebrate communities following land-use change in Sumatra, Indonesia. <i>Biological Conservation</i> 191:750–758.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Murillo-Pacheco J, López-Iborra GM, Escobar F, Bonilla-Rojas WF, Verdú JR. 2018. The value of small, natural and man-made wetlands for bird diversity in the east Colombian Piedmont. <i>Aquatic Conservation: Marine and Freshwater Ecosystems</i> 28:87–97.	Unshaded crops	Rice	Birds	South America	Annual
Murrieta-Galindo R, González-Romero A, López-Barrera F, Parra-Olea G. 2013. Coffee agrosystems: an important refuge for amphibians in central Veracruz, Mexico. <i>Agroforestry Systems</i> 87:767–779.	Shaded crops	Coffee	Herpetofauna	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Muvengwi J, Mbiba M, Ndagurwa HGT, Nyamadzawo G, Nhokovedzo P. 2017. Termite diversity along a land use intensification gradient in a semi-arid savanna. <i>Journal of Insect Conservation</i> 21:801–812.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Annual
Naughton-Treves L, Mena JL, Treves A, Alvarez N, Radeloff VC. 2003. Wildlife Survival Beyond Park Boundaries: the Impact of Slash-and-Burn Agriculture and Hunting on Mammals in Tambopata, Peru. <i>Conservation Biology</i> 17:1106–1117.	Unshaded crops	Mixed tropical crops	Mammals	South America	Annual
Ndriantsoa SH, Riemann JC, Raminosoa N, Rödel M-O, Glos JS. 2017. Amphibian Diversity in the Matrix of a Fragmented Landscape Around Ranomafana in Madagascar Depends on Matrix Quality. <i>Tropical Conservation Science</i> 10:194008291668606.	Crops with some vegetation	Banana	Herpetofauna	Africa	Perennial
	Unshaded crops	Rice	Herpetofauna	Africa	Annual
Nicolas V, Barrière P, Papier A, Colyn M. 2009. Shrew species diversity and abundance in Ziama Biosphere Reserve, Guinea: comparison among primary forest, degraded forest and restoration plots. <i>Biodiversity and Conservation</i> 18:2043–2061.	Unshaded crops	Mixed tropical crops	Mammals	Africa	Mixed
Norfolk O, Asale A, Temesgen T, Denu D, Platts PJ, Marchant R, Yewhalaw D. 2017b. Diversity and composition of tropical butterflies along an Afromontane agricultural gradient in the Jimma Highlands, Ethiopia. <i>Biotropica</i> 49:346–354.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Annual
	Shaded crops	Coffee	Invertebrates	Africa	Perennial
Norfolk O, Jung M, Platts PJ, Malaki P, Odeny D, Marchant R. 2017a. Birds in the matrix: the role of agriculture in avian conservation in the Taita Hills, Kenya. <i>African Journal of Ecology</i> 55:530–540.	Shaded crops	Mixed tropical crops	Birds	Africa	Mixed
	Unshaded crops	Mixed tropical crops	Birds	Africa	Annual
Ocampo-Ariza C, Denis K, Njie Motombi F, Bobo KS, Kreft H, Waltert M. 2019. Extinction thresholds and negative responses of Afrotropical ant-following birds to forest cover loss in oil palm and agroforestry landscapes. <i>Basic and Applied Ecology</i> 39:26–37.	Unshaded crops	Oil palm	Birds	Africa	Perennial
	Shaded crops	Mixed tropical crops	Birds	Africa	Mixed
Paoletti A, Darras K, Jayanto H, Grass I, Kusri M, Tschardt T. 2018. Amphibian and reptile communities of upland and riparian sites across Indonesian oil palm, rubber and forest. <i>Global Ecology and Conservation</i> 16:e00492.	Unshaded crops	Oil palm	Herpetofauna	Asia	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Pardo LE, Campbell MJ, Edwards W, Clements GR, Laurance WF. 2018. Terrestrial mammal responses to oil palm dominated landscapes in Colombia. PLOS ONE 13:e0197539.	Unshaded crops	Oil palm	Mammals	South America	Perennial
Perfecto I, Mas A, Dietsch T, Vandermeer J. 2003. Conservation of biodiversity in coffee agroecosystems: a tri-taxa comparison in southern Mexico. Biodiversity and Conservation 12:1239–1252.	Shaded crops	Coffee	Invertebrates	Central America	Perennial
	Shaded crops	Coffee	Birds	Central America	Perennial
Perry J, Lojka B, Quinones Ruiz L, Van Damme P, Houška J, Fernandez Cusimamani E. 2016. How natural Forest Conversion Affects Insect Biodiversity in the Peruvian Amazon: Can Agroforestry Help? Forests 7:82.	Unshaded crops	All other tropical crops	Invertebrates	South America	Annual
	Shaded crops	Cacao	Invertebrates	South America	Perennial
	Shaded crops	All other tropical crops	Invertebrates	South America	Perennial
Peters MK, Fischer G, Schaab G, Kraemer M. 2009. Species compensation maintains abundance and raid rates of African swarm-raiding army ants in rainforest fragments. Biological Conservation 142:668–675.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Mixed
Petit LJ, Petit DR, Christian DG, Powell HDW. 1999. Bird communities of natural and modified habitats in Panama. Ecography 22:292–304.	Shaded crops	Coffee	Birds	Central America	Perennial
Pineda E, Halfiter G. 2004. Species diversity and habitat fragmentation: frogs in a tropical montane landscape in Mexico. Biological Conservation 117:499–508.	Shaded crops	Coffee	Herpetofauna	Central America	Perennial
Potapov AM et al. 2020. Functional losses in ground spider communities due to habitat structure degradation under tropical land-use change. Ecology 101:e02957.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Prabowo WE, Darras K, Clough Y, Toledo-Hernandez M, Arlettaz R, Mulyani YA, Tsharntke T. 2016. Bird Responses to Lowland Rainforest Conversion in Sumatran Smallholder Landscapes, Indonesia. PLOS ONE 11:e0154876.	Unshaded crops	Oil palm	Birds	Asia	Perennial
Pringle CM, Ramírez A. 1998. Use of both benthic and drift sampling techniques to assess tropical stream invertebrate communities along an altitudinal gradient, Costa Rica. Freshwater Biology 39:359–373.	Unshaded crops	Banana	Invertebrates	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Ramamonjisoa N, Sakai M, Ndriantsoa SH, Kakehashi R, Kurabayashi A, Tomaru N, Natuhara Y. 2020. Hotspots of stream tadpole diversity in forest and agricultural landscapes in Ranomafana, Madagascar. <i>Landscape and Ecological Engineering</i> 16:207–221.	Unshaded crops	Mixed tropical crops	Herpetofauna	Africa	Mixed
Raman TRS. 2006. Effects of Habitat Structure and Adjacent Habitats on Birds in Tropical Rainforest Fragments and Shaded Plantations in the Western Ghats, India. <i>Biodiversity and Conservation</i> 15:1577–1607.	Shaded crops	Mixed tropical crops	Birds	Asia	Perennial
Ranganathan J, Chan KMA, Daily GC. 2007. Satellite detection of bird communities in tropical countryside. <i>Ecological Applications</i> 17:1499–1510.	Unshaded crops	Mixed tropical crops	Birds	Central America	Annual
	Shaded crops	Coffee	Birds	Central America	Perennial
Ranganathan J, Daniels RJR, Chandran MDS, Ehrlich PR, Daily GC. 2008. Sustaining biodiversity in ancient tropical countryside. <i>Proceedings of the National Academy of Sciences</i> 105:17852–17854.	Unshaded crops	All other tropical crops	Birds	Asia	Perennial
Ribeiro J, Colli GR, Caldwell JP, Ferreira E, Batista R, Soares A. 2017. Evidence of neotropical anuran community disruption on rice crops: a multidimensional evaluation. <i>Biodiversity and Conservation</i> 26:3363–3383.	Unshaded crops	Rice	Herpetofauna	South America	Annual
Ricketts TH, Daily GC, Ehrlich PR, Fay JP. 2001. Countryside Biogeography of Moths in a Fragmented Landscape: Biodiversity in Native and Agricultural Habitats. <i>Conservation Biology</i> 15:378–388.	Unshaded crops	Coffee	Invertebrates	Central America	Perennial
Roberts DL, Cooper RJ, Petit LJ. 2000. Use of Premontane Moist Forest and Shade Coffee Agroecosystems by Army Ants in Western Panama. <i>Conservation Biology</i> 14:192–199.	Shaded crops	Coffee	Invertebrates	Central America	Perennial
Rocha J, Laps RR, Machado CG, Campiolo S. 2019. The conservation value of cacao agroforestry for bird functional diversity in tropical agricultural landscapes. <i>Ecology and Evolution</i> 9:7903–7913.	Shaded crops	Cacao	Birds	South America	Perennial
Rocha R, Virtanen T, Cabeza M. 2015. Bird Assemblages in a Malagasy Forest-Agricultural Frontier: Effects of Habitat Structure and Forest Cover. <i>Tropical Conservation Science</i> 8:681–710.	Unshaded crops	Mixed tropical crops	Birds	Africa	Mixed
Roth DS, Perfecto I, Rathcke B. 1994. The Effects of Management Systems on Ground-Foraging Ant Diversity in Costa Rica. <i>Ecological Applications</i> 4:423–436.	Unshaded crops	Banana	Invertebrates	Central America	Perennial
	Shaded crops	Cacao	Invertebrates	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Rousseau L, Fonte SJ, Téllez O, van der Hoek R, Lavelle P. 2013. Soil macrofauna as indicators of soil quality and land use impacts in smallholder agroecosystems of western Nicaragua. <i>Ecological Indicators</i> 27:71–82.	Unshaded crops	Mixed tropical crops	Invertebrates	Central America	Annual
	Shaded crops	Mixed tropical crops	Invertebrates	Central America	Annual
Sambhu H, Nankishore A, Turton SM, Northfield TD. 2018. Trade-offs for butterfly alpha and beta diversity in human-modified landscapes and tropical rainforests. <i>Ecology and Evolution</i> 8:12918–12928.	Unshaded crops	Sugarcane	Invertebrates	Oceania	Annual
Sambhu H, Northfield T, Nankishore A, Ansari A, Turton S. 2017. Tropical Rainforest and Human-Modified Landscapes Support Unique Butterfly Communities That Differ in Abundance and Diversity. <i>Environmental Entomology</i> 46:1225–1234.	Unshaded crops	Sugarcane	Invertebrates	South America	Annual
Santos A de C, Sales PCL, Ribeiro DB, Silva PRR. 2020. Habitat conversion affects beta diversity in frugivorous butterfly assemblages. <i>Studies on Neotropical Fauna and Environment</i> :1–13.	Unshaded crops	Sugarcane	Invertebrates	South America	Annual
Schulze CH, Waltert M, Kessler PJA, Pitopang R, Veddeler D, Mühlberg M, Gradstein SR, Leuschner C, Steffan-Dewenter I, Tscharntke T. 2004. Biodiversity indicator groups of tropical land-use systems: comparing plants, birds, and insects. <i>Ecological Applications</i> 14:1321–1333.	Shaded crops	Cacao	Birds	Asia	Perennial
	Unshaded crops	Maize	Birds	Asia	Annual
	Shaded crops	Cacao	Invertebrates	Asia	Perennial
	Unshaded crops	Maize	Invertebrates	Asia	Annual
Scriven SA, Gillespie GR, Laimun S, Goossens B. 2018. Edge effects of oil palm plantations on tropical anuran communities in Borneo. <i>Biological Conservation</i> 220:37–49.	Unshaded crops	Oil palm	Herpetofauna	Asia	Perennial
Seidu I, Danquah E, Ayine Nsor C, Amaning Kwarteng D, Lancaster LT. 2017. Odonata community structure and patterns of land use in the Atewa Range Forest Reserve, Eastern Region (Ghana). <i>International Journal of Odonatology</i> 20:173–189.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Mixed
Sewlal J-AN, Hailey A. 2019. Diversity and species composition of Araneidae, Tetragnathidae and Nephilidae in different levels of disturbed habitats in Trinidad, West Indies. <i>Journal of Natural History</i> 53:1889–1903.	Unshaded crops	Mixed tropical crops	Invertebrates	Central America	Annual
	Shaded crops	Cacao	Invertebrates	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Shahabuddin S, Hidayat P, Manuwoto S, Noerdjito WA, Tscharnke T, Schulze CH. 2010. Diversity and body size of dung beetles attracted to different dung types along a tropical land-use gradient in Sulawesi, Indonesia. <i>Journal of Tropical Ecology</i> 26:53–65.	Shaded crops	Cacao	Invertebrates	Asia	Perennial
Shahabuddin S, Schulze CH, Tscharnke T. 2005. Changes of dung beetle communities from rainforests towards agroforestry systems and annual cultures in Sulawesi (Indonesia). <i>Biodiversity and Conservation</i> 14:863–877.	Unshaded crops	Maize	Invertebrates	Asia	Annual
	Shaded crops	Cacao	Invertebrates	Asia	Perennial
Sidhu S, Raman TRS, Mudappa D. 2015. Prey abundance and leopard diet in a plantation and rainforest landscape, Anamalai Hills, Western Ghats. <i>Current Science</i> 109:323–330.	Unshaded crops	Tea	Mammals	Asia	Perennial
	Shaded crops	Coffee	Mammals	Asia	Perennial
Sodhi NS, Koh LP, Prawiradilaga DM, Tinulele I, Putra DD, Tong Tan TH. 2005. Land use and conservation value for forest birds in Central Sulawesi (Indonesia). <i>Biological Conservation</i> 122:547–558.	Unshaded crops	Mixed tropical crops	Birds	Asia	Mixed
Soh MCK, Sodhi NS, Lim SLH. 2006. High sensitivity of montane bird communities to habitat disturbance in Peninsular Malaysia. <i>Biological Conservation</i> 129:149–166.	Unshaded crops	Tea	Birds	Asia	Perennial
Sreekar R, Srinivasan U, Mammides C, Chen J, Manage Goodale U, Wimalabandara Kotagama S, Sidhu S, Goodale E. 2015. The effect of land-use on the diversity and mass-abundance relationships of understory avian insectivores in Sri Lanka and southern India. <i>Scientific Reports</i> 5:11569.	Crops with some vegetation	Mixed tropical crops	Birds	Asia	Unknown
Srinivas A, Koh LP. 2016. Oil palm expansion drives avifaunal decline in the Pucallpa region of Peruvian Amazonia. <i>Global Ecology and Conservation</i> 7:183–200.	Crops with some vegetation	Oil palm	Birds	South America	Perennial
Strauß L, Faustino de Lima R, Riesbeck F, Rödel M-O. 2018. São Tomé Island Endemic Treefrogs (<i>Hyperolius</i> spp.) and Land-Use Intensification: A Tale of Hope and Caution. <i>Tropical Conservation Science</i> 11:194008291877643.	Unshaded crops	Mixed tropical crops	Herpetofauna	Africa	Mixed
Trainor CR. 2007. Changes in bird species composition on a remote and well-forested Wallacean Island, South-East Asia. <i>Biological Conservation</i> 140:373–385.	Unshaded crops	Mixed tropical crops	Birds	Asia	Mixed
Trimble MJ, van Aarde RJ. 2014. Amphibian and reptile communities and functional groups over a land-use gradient in a coastal tropical forest landscape of high richness and endemism: Herpetofauna over a land-use gradient. <i>Animal Conservation</i> 17:441–453.	Unshaded crops	Sugarcane	Herpetofauna	Africa	Annual

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Turner EC, Foster WA. 2009. The impact of forest conversion to oil palm on arthropod abundance and biomass in Sabah, Malaysia. <i>Journal of Tropical Ecology</i> 25:23–30.	Unshaded crops	Oil palm	Invertebrates	Asia	Perennial
Tylianakis JM, Klein A-M, Lozada T, Tscharntke T. 2006b. Spatial scale of observation affects alpha, beta and gamma diversity of cavity-nesting bees and wasps across a tropical land-use gradient. <i>Journal of Biogeography</i> 33:1295–1304.	Shaded crops	Coffee	Invertebrates	South America	Perennial
	Unshaded crops	Rice	Invertebrates	South America	Annual
Tylianakis JM, Klein A-M, Tscharntke T. 2005. Spatiotemporal variation in the diversity of Hymenoptera across a tropical habitat gradient. <i>Ecology</i> 86:3296–3302.	Unshaded crops	Rice	Invertebrates	South America	Annual
	Shaded crops	Coffee	Invertebrates	South America	Perennial
Tylianakis JM, Tscharntke T, Klein A-M. 2006a. Diversity, ecosystem function, and stability of parasitoid–host interactions across a tropical habitat gradient. <i>Ecology</i> 87:3047–3057.	Shaded crops	Coffee	Invertebrates	South America	Perennial
	Unshaded crops	Rice	Invertebrates	South America	Annual
Umetsu F, Pardini R. 2007. Small mammals in a mosaic of forest remnants and anthropogenic habitats—evaluating matrix quality in an Atlantic forest landscape. <i>Landscape Ecology</i> 22:517–530.	Unshaded crops	Mixed tropical crops	Mammals	South America	Annual
Urrutia-Escobar MX, Armbrrecht I. 2013. Effect of Two Agroecological Management Strategies on Ant (Hymenoptera: Formicidae) Diversity on Coffee Plantations in Southwestern Colombia. <i>Environmental Entomology</i> 42:194–203.	Unshaded crops	Coffee	Invertebrates	South America	Perennial
	Shaded crops	Coffee	Invertebrates	South America	Perennial
van Biervliet O, Wiśniewski K, Daniels J, Vonesh JR. 2009. Effects of Tea Plantations on Stream Invertebrates in a Global Biodiversity Hotspot in Africa: Effect of Tea Plantations on Stream Biodiversity. <i>Biotropica</i> 41:469–475.	Unshaded crops	Tea	Invertebrates	Africa	Perennial
Vasconcelos S, Rodrigues P, Palma L, Mendes LF, Palminha A, Catarino L, Beja P. 2015. Through the eye of a butterfly: Assessing biodiversity impacts of cashew expansion in West Africa. <i>Biological Conservation</i> 191:779–786.	Unshaded crops	All other tropical crops	Invertebrates	Africa	Perennial
Waltert M, Bobo KS, Kaupa S, Montoya ML, Nsanyi MS, Fermon H. 2011. Assessing Conservation Values: Biodiversity and Endemicity in Tropical Land Use Systems. <i>PLOS ONE</i> 6:e16238.	Unshaded crops	Mixed tropical crops	Invertebrates	Africa	Mixed

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
	Shaded crops	Mixed tropical crops	Invertebrates	Africa	Perennial
	Unshaded crops	Mixed tropical crops	Birds	Africa	Mixed
	Shaded crops	Mixed tropical crops	Birds	Africa	Perennial
Waltert M, Bobo KS, Sainge NM, Fermon H, Mühlenberg M. 2005. From forest to farmland: habitat effects on Afrotropical forest bird diversity. <i>Ecological Applications</i> 15:1351–1366.	Unshaded crops	Mixed tropical crops	Birds	Africa	Annual
	Shaded crops	Mixed tropical crops	Birds	Africa	Perennial
Waltert M, Mardiasuti A, Mühlenberg M. 2004. Effects of Land Use on Bird Species Richness in Sulawesi, Indonesia. <i>Conservation Biology</i> 18:1339–1346.	Shaded crops	Cacao	Birds	Asia	Perennial
	Unshaded crops	Maize	Birds	Asia	Annual
Wearn OR, Carbone C, Rowcliffe JM, Bernard H, Ewers RM. 2016. Grain-dependent responses of mammalian diversity to land use and the implications for conservation set-aside. <i>Ecological Applications</i> 26:1409–1420.	Unshaded crops	Oil palm	Mammals	Asia	Perennial
Wearn OR, Rowcliffe JM, Carbone C, Pfeifer M, Bernard H, Ewers RM. 2017. Mammalian species abundance across a gradient of tropical land-use intensity: A hierarchical multi-species modelling approach. <i>Biological Conservation</i> 212:162–171.	Unshaded crops	Oil palm	Mammals	Asia	Perennial
Wilkinson CL, Yeo DCJ, Tan HH, Fikri AH, Ewers RM. 2018b. Land-use change is associated with a significant loss of freshwater fish species and functional richness in Sabah, Malaysia. <i>Biological Conservation</i> 222:164–171.	Unshaded crops	Oil palm	Fish	Asia	Perennial
	Shaded crops	Oil palm	Fish	Asia	Perennial
	Unshaded crops	Oil palm	Fish	Asia	Perennial
	Unshaded crops	Oil palm	Fish	Asia	Perennial
Williams-Guillén K, Perfecto I. 2010. Effects of Agricultural Intensification on the Assemblage of Leaf-Nosed Bats (Phyllostomidae) in a Coffee Landscape in Chiapas, Mexico: Phyllostomid Diversity in Shade Coffee Plantations. <i>Biotropica</i> 42:605–613.	Shaded crops	Coffee	Mammals	Central America	Perennial

Study bibliographic details	Crop shading	Crop category	Taxa	Continent	Perennial or Annual
Williams-Guillén K, Perfecto I. 2011. Ensemble Composition and Activity Levels of Insectivorous Bats in Response to Management Intensification in Coffee Agroforestry Systems. <i>PLOS ONE</i> 6:e16502.	Shaded crops	Coffee	Mammals	Central America	Perennial
Willig MR, Presley SJ, Plante J-L, Bloch CP, Solari S, Pacheco V, Weaver SC. 2019. Guild-level responses of bats to habitat conversion in a lowland Amazonian rainforest: species composition and biodiversity. <i>Journal of Mammalogy</i> 100:223–238.	Unshaded crops	Mixed tropical crops	Mammals	South America	Mixed
Wordley CFR, Sankaran M, Mudappa D, Altringham JD. 2018. Heard but not seen: Comparing bat assemblages and study methods in a mosaic landscape in the Western Ghats of India. <i>Ecology and Evolution</i> 8:3883–3894.	Unshaded crops	Tea	Mammals	Asia	Perennial
	Shaded crops	Tea	Mammals	Asia	Perennial
	Shaded crops	Coffee	Mammals	Asia	Perennial
Zuluaga GJC, Rodewald AD. 2015. Response of mixed-species flocks to habitat alteration and deforestation in the Andes. <i>Biological Conservation</i> 188:72–81.	Shaded crops	Coffee	Birds	South America	Perennial



Table S2. Crops analysed in the meta-analysis per pairwise comparison by region

Crops	Central America	South America	Africa	Asia	Oceania	Total
Allspice	12	-	-	-	-	12
Banana	38	4	2	-	-	44
Cacao	50	45	11	76	-	182
Cacao and coffee	8	-	9	-	-	17
Citrus	8	-	-	-	-	8
Coffee	102	51	31	32	-	216
Maize	9	-	2	23	-	34
Mixed tropical crops	78	65	186	85	-	414
Oil palm	38	50	2	174	-	264
All other tropical crops	28	8	15	4	-	55
Rice	-	32	11	-	-	43
Sugarcane	-	23	8	-	2	33
Tea	-	-	4	38	-	42

Table S3. Effect sizes calculated in the study as displayed in Figure 3. For single crops with fewer than four studies, we grouped these and reported them as ‘all other tropical crops’. When biodiversity values were provided for sites that did not distinguish between multiple different crops, we reported them as ‘mixed tropical crops’

Analysis	Mean Hedges' g	Lower 95% CI	Upper 95% CI	<i>p</i>
Overall	-0.588	-0.671	-0.505	<0.001
Crop (Fig. 3a)				
<i>Allspice</i>	-0.29	-0.848	0.258	0.296
<i>Banana</i>	0.20	-0.258	0.665	0.387
<i>Cacao</i>	-0.43	-0.717	-0.152	0.003
<i>Cacao and coffee</i>	0.05	-0.737	0.841	0.897
<i>Citrus</i>	-0.39	-1.281	0.499	0.390
<i>Coffee</i>	-0.05	-0.198	0.098	0.509
<i>Maize</i>	-2.65	-3.479	-1.820	<0.001
<i>Mixed tropical crops</i>	-0.39	-0.539	-0.247	<0.001
<i>Oil palm</i>	-1.23	-1.429	-1.030	<0.001
<i>All other tropical crops</i>	-0.98	-1.290	-0.675	<0.001
<i>Rice</i>	-0.67	-1.224	-0.118	0.017
<i>Sugarcane</i>	-1.22	-1.610	-0.828	<0.001
<i>Tea</i>	-0.73	-1.155	-0.295	0.001
Shading (Fig. 3b)				
<i>Crops with some vegetation</i>	-0.160	-0.674	0.354	0.542
<i>Shaded crops</i>	-0.168	-0.290	-0.046	0.007
<i>Unshaded crops</i>	-0.876	-0.987	-0.766	<0.001
Crop rotation time (Fig. 3c)				
<i>Annual</i>	-0.927	-1.132	-0.721	<0.001
<i>Mixed</i>	-0.121	-0.378	0.135	0.354
<i>Perennial</i>	-0.605	-0.707	-0.504	<0.001
<i>Unknown</i>	-0.173	-0.525	0.179	0.334
Taxonomic group (Fig. 3d)				
<i>Birds</i>	-0.841	-1.025	-0.656	<0.001
<i>Fish</i>	0.324	-0.551	1.199	0.468
<i>Herpetofauna</i>	-0.756	-1.151	-0.360	<0.001
<i>Invertebrates</i>	-0.626	-0.735	-0.518	<0.001
<i>Mammals</i>	-0.109	-0.252	0.033	0.132

Analysis	Mean Hedges' g	Lower 95% CI	Upper 95% CI	<i>p</i>
Geographic region (Fig. 3e)				
<i>Africa</i>	-0.339	-0.503	-0.174	<0.001
<i>Asia</i>	-1.043	-1.213	-0.873	<0.001
<i>Central America</i>	-0.196	-0.345	-0.046	0.010
<i>Oceania</i>	1.064	-2.950	5.079	0.603
<i>South America</i>	-0.698	-0.862	-0.534	<0.001



895x764mm (72 x 72 DPI)