

# Cyclopropane-Fused *N*-Heterocycles via Aza-Heck-Triggered C(sp<sup>3</sup>)-H Functionalization Cascades

Changcheng Jing, Benjamin T. Jones, Ross J. Adams, and John F. Bower\*



Cite This: <https://doi.org/10.1021/jacs.2c08304>



Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

**ABSTRACT:** Unique examples of aza-Heck-based C(sp<sup>3</sup>)-H functionalization cascades are described. Under Pd(0)-catalyzed conditions, the aza-Heck-type cyclization of *N*-(pentafluorobenzoyloxy)carbamates generates alkyl-Pd(II) intermediates that effect C(sp<sup>3</sup>)-H palladation en route to cyclopropanes. Key factors that control the site selectivity of the cyclopropanation process have been elucidated such that selective access to a wide range of ring- or spiro-fused systems can be achieved.


Cyclopropanes are routinely employed in pharmaceutical design to moderate compound lipophilicity or *N*-centered basicity.<sup>1</sup> Reflecting their relative ease of synthesis, *peripheral* cyclopropane units are featured in many marketed drugs, whereas *core* cyclopropane units are encountered less often (Scheme 1A).<sup>2</sup> An example of the latter is the DPP-4 inhibitor saxagliptin, which possesses a cyclopropane-fused pyrrolidine.<sup>3a</sup> This subunit is derived from pyroglutamic acid via a lengthy Simmons–Smith-based route, making access to

more complex derivatives challenging.<sup>3b</sup> Accordingly, direct and flexible methods that can address these issues are likely to be of interest. A powerful but underdeveloped option involves the intramolecular aza-palladation of an alkene (step a) in advance of C–H palladation-initiated cyclopropanation (step b) (Scheme 1B). Yang and co-workers have demonstrated such processes under oxidative conditions (Scheme 1C).<sup>4</sup> Although conceptually important, specific constraints hamper both steps; for example, step a requires a conformationally biasing and acidifying anilide unit and is not well suited to six-ring cyclizations, whereas step b suffers from limited scope and selectivity.


We and others have demonstrated that the efficiency of alkene aza-palladation-based processes can be enhanced substantially by replacing an external oxidant (e.g., O<sub>2</sub> in Scheme 1C) with a N–O bond.<sup>5</sup> In this approach, N–O oxidative addition unites the key processes of substrate binding and catalyst oxidation, so the catalysis is often more robust. The non-oxidative conditions also mean that highly tunable P-based ligands can be used to moderate the properties of the Pd center. Indeed, aza-Heck-based approaches of this type now allow an expanding range of nonconformationally biased cyclizations and cascades involving sterically and electronically diverse alkenes.<sup>5a–f,i,j</sup> This includes asymmetric aza-Heck cyclizations<sup>5a,d</sup> and cascade reactions involving aryl C(sp<sup>2</sup>)-H palladation.<sup>5a,j</sup> In the present study, we outline aza-Heck-triggered cyclopropanation processes that involve a more challenging C(sp<sup>3</sup>)-H palladation step (Scheme 1D). To the best of our knowledge, these are the first examples of C(sp<sup>3</sup>)-H functionalization cascades that use newer classes of N–O units (i.e., nonoxime ester-based).<sup>5l,m,6</sup> Compared to Scheme 1C, notable features of these new processes include (a) the

## Scheme 1. Introduction


### (A) Pharmaceuticals possessing cyclopropanes




### (B) Cyclopropane-fused heterocycles via alkene aza-palladation:



### (C) Yang's oxidative processes:



### (D) Proposed aza-Heck C(sp<sup>3</sup>)-H functionalization cascades:



Received: August 5, 2022

efficient participation of sterically encumbered alkenes, (b) efficient 5-*exo* cyclizations in the absence of a conformationally biasing and acidifying anilide unit, (c) efficient 6-*exo* cyclizations, and (d) no requirement for the benzylic activation of the target C(sp<sup>3</sup>)-H bond. We also demonstrate that either steric or electronic control can be used to enforce the regioselectivity of the cyclopropanation event, thereby providing selective access to ring- or spiro-fused systems. In broader terms, these studies offer rare examples of C(sp<sup>3</sup>)-H cyclopropanation processes that are triggered by alkene heteropalladation, a sequence that is likely challenging because of the reversibility of the migratory insertion step (vide infra).<sup>4,7,8</sup>

A mechanistic analysis of the processes developed here is outlined in Scheme 2A. The N-O oxidative addition of **1** is expected to provide the aza-Pd(II) intermediate **Int-I**. Prior work has indicated that efficient alkene aza-palladation requires the dissociation of pentafluorobenzoate from **Int-I** to give **Int-I'**.<sup>5b</sup> **Int-I'** undergoes cyclization and carboxylate association to


give the alkyl-Pd(II) intermediate **Int-II**, which can provide palladacyclobutane **Int-III** via concerted metalation deprotonation-type metalation. Reductive elimination from **Int-III** then releases the cyclopropane product **2**. Depending on the nature of R (vide infra), alternative palladacyclobutanes may be accessible. The carboxylic acid released during the cyclopropanation sequence is expected to undergo deprotonation by triethylamine, and the resulting triethylammonium salt will trigger the facile protodecarboxylation of pentafluorobenzoate to release C<sub>6</sub>F<sub>5</sub>H.<sup>9</sup> Previous studies indicated that alkene aza-palladation is reversible under cationic conditions (vide infra),<sup>5j,10</sup> so the success of the process is likely to depend on the efficiency of the carboxylate-mediated C(sp<sup>3</sup>)-H metalation step (**Int-II** to **Int-III**). Here, external carboxylate additives (R'CO<sub>2</sub>M) are likely required because the pentafluorobenzoate released during N-O oxidative addition is highly dissociative and a relatively weak base. Note that the optimal mechanistic scenario requires a cationic species for aza-palladation (**Int-I'**) and a neutral intermediate for C(sp<sup>3</sup>)-H metalation (**Int-II**). As such, optimal conditions require an appropriate trade-off because the complete partitioning of pathways at each stage is likely unattainable.

Proof-of-concept studies focused initially on the cyclization of the *O*-pentafluorobenzoyl system **1a** to target **2a** (Scheme 2B). Under the indicated conditions and in the absence of a carboxylate additive, target **2a** was generated in only a 13% yield (entry 1). The addition of 100 mol% CsOPiv markedly increased the efficiency such that **2a** was formed in a 56% yield (entry 2). During these initial studies, other optimal parameters were established. Most significantly, from a screen of P ligands commonly employed in aza-Heck processes, it was found that **L1** (CgPPh) was by far the most efficient.<sup>5b</sup> Note that P ligands were specifically noted as being incompatible with the method in Scheme 1C.<sup>4</sup> To optimize the process further, a library of approximately 15 derivatives was prepared from CgPH (Scheme 2C and the SI).<sup>11</sup> Selected evaluation results are shown in entries 3–6, with the key finding being that the benzofuryl system **L3** could provide **2a** in an 80% yield over 6 h. At this stage we evaluated the choice of the *O*-based leaving group (entries 7 and 8), which confirmed that the -O<sup>F</sup>Bz system (**1a**) was superior to both -OTs and -OPiv variants **1a'** and **1a''**, respectively. The latter result is consistent with the idea that a cationic aza-palladium intermediate offers an optimal aza-palladation efficiency (**Int-I'** to **Int-II**).


The optimized conditions were proven to be applicable to the synthesis of a wide range of cyclopropane-fused pyrrolidines (Table 1A). For example, a variety of alkyl substituents were tolerated at R<sup>1</sup>, as evidenced by the efficient formation of **2b–f**. Note that the bulky secondary alkyl substituents of **2b** and **2f** did not significantly diminish the reaction efficiency relative to, for example, that of **2c**. The electronically distinct styrenyl system **1g** also participated to provide adduct **2g**, albeit with a more modest efficiency. Substituents can be introduced at C4 and C5 of the targets, as demonstrated by the formation of **2h** and **2i**, respectively. For the former, minimal diastereoselectivity was observed, whereas the latter was generated as a single diastereomer. The hydrogenolytic N-Cbz deprotection of **2i** allowed access to the N-DNs derivative **2i'**. **2i'** was characterized by single-crystal X-ray diffraction, which revealed a *syn*-relationship between the methyl and ethyl substituents.<sup>12</sup> This outcome is consistent with the alkene aza-palladation step being reversible because the productive palladacyclobutane **Int-2i** is formed via

## Scheme 2. Mechanistic Analysis and Optimization of the Cascade Process

### (A) Mechanistic analysis:




### (B) Selected optimization results:



Entry	LG	A	L (B)	Additive (C)	Time (h)	NMR Yield
1	O <sup>F</sup> Bz	5	L1 (20)	none	24	13%
2	O <sup>F</sup> Bz	5	L1 (20)	CsOPiv (100)	24	56%
3	O <sup>F</sup> Bz	5	L2 (20)	CsOPiv (100)	8	68%
4	O <sup>F</sup> Bz	5	L3 (20)	CsOPiv (100)	6	72%
5	O <sup>F</sup> Bz	5	L3 (25)	CsOPiv (100)	6	80% <sup>a</sup>
6	O <sup>F</sup> Bz	5	L4 (25)	CsOPiv (100)	6	58%
7	OTs	5	L3 (25)	CsOPiv (100)	6	43%
8	OPiv	5	L3 (25)	CsOPiv (100)	24	39%


### (C) Ligand synthesis:




<sup>a</sup>Isolated yield. <sup>b</sup>Further optimization results, including the evaluation of other ligands, are given in the SI.

Table 1. Cyclopropane-Fused Pyrrolidines


## (A) Synthesis of ring-fused systems:




## (B) Rationale for the diastereoselectivity observed for 2i:



## (C) Competitive formation of ring- and spiro-fused systems:



## (D) Selective synthesis of spiro-fused systems:




<sup>a</sup>L3 (50 mol %) was used. <sup>b</sup>DNs = 2,4-dinitrophenylsulfonfyl. <sup>c</sup>L3 (30 mol %) was used, and the reaction time was 24 h.

the disfavored cyclization mode (TS-2i) (Table 1B).<sup>13</sup> For system 1j, where R<sup>1</sup> = Me, the expected product 2j was formed in a 55% yield alongside smaller quantities of spiro-fused

cyclopropane *iso*-2j (14% yield) (Table 1C). The formation of the latter is presumably facilitated by the C(sp<sup>3</sup>)–H metalation of the more sterically accessible (versus Table 1A) methyl group of Int-II'.<sup>14</sup> To enforce this selectivity, systems possessing a substitution at the internal allylic position (C3) were investigated (Table 1D). In the reaction, the cyclization of the ethyl-substituted system 1k provided *iso*-2k in a 73% yield, and the corresponding ring-fused cyclopropane was not observed. This process required an increased reaction temperature (160 °C) and an alternate carboxylate additive (KOAc). These modified conditions were also effective for the selective formation of *iso*-2l and the intriguing vicinally dispiro-fused adduct *iso*-2m.<sup>15</sup>

Applying the reaction conditions to, for example, the homologue of 1a did not provide the corresponding cyclopropane-fused piperidine (see the SI), presumably because 6-*exo* aza-palladation is relatively demanding. This limitation can be circumvented by instead using systems that possess a degree of conformational bias (Table 2). Indeed, the cascade

Table 2. Cyclopropane-Fused Tetrahydroisoquinolines



<sup>a</sup>Pd<sub>2</sub>(dba)<sub>3</sub> (5 mol %) was used as the precatalyst. <sup>b</sup>L3 (50 mol %) was used.


cyclization of 1n, which possesses an aromatic linker, was efficient, delivering the cyclopropane-fused tetrahydroisoquinoline 2n in a 76% yield. As mentioned earlier, the process is relatively insensitive to the steric demands of the alkene substituent (R<sup>1</sup>), so the bulky isopropyl group of 2p was well-tolerated. The protocol offers a useful scope with respect to the aromatic component, with both electron-rich (2q, 2r, and 2t) and electron-poor (2s) units participating. Diastereoselective processes are achievable for systems where R<sup>2</sup> ≠ H; this was demonstrated by the highly stereocontrolled cyclization of 1u to 2u, which favored the *syn*-diastereomer (>15:1 dr). For these processes, spiro-fused cyclopropanes (cf. *iso*-2j) were not observed, which likely reflects the inherent preference for C(sp<sup>3</sup>)–H palladation at the benzylic position. For many of the examples in Table 2, the use of Pd<sub>2</sub>(dba)<sub>3</sub> as the precatalyst resulted in purification problems because dba coeluted with

the product during chromatography. This issue was alleviated by instead using  $\text{Pd}_2(\text{p-MeO-dba})_3$ .<sup>16,17</sup>



We have established that the cascade cyclopropanation procedure can be used to generate reactive donor–acceptor cyclopropanes. A preliminary example process involves the conversion **3a** to piperidine **4a** via initial (nonconformationally biased) 5-*exo* aza-palladation (Table 3A). This generates **Int-**

**Table 3. Rearrangement Processes via the Generation of Donor–Acceptor Cyclopropanes**

(A) Preliminary studies and mechanistic considerations:



(B) Optimized conditions and reaction scope:



<sup>a</sup>The major regioisomer is depicted. <sup>b</sup>L3 (25 mol %) was used. <sup>c</sup>The reaction time was 42 h, and the regioisomers were isolated in 6:1 and >15:1 d.r., respectively (see the SI). <sup>d</sup>The substrate was prepared from (*S*)-propylene oxide.

**II''** in which there is a choice of three different  $\text{C}(\text{sp}^3)\text{-H}$  bonds for metalation, leading to either four- or six-membered palladacycles (not depicted).<sup>18</sup> The productive pathway involves metalation at the C3 position en route to donor–acceptor cyclopropane **Int-IV**. This is primed for thermally promoted ring opening to provide piperidine **4a**.<sup>19,20</sup> Accordingly, the rearrangement process transfers the methylene  $\text{CH}_2$  unit of the starting material (**3a**) to C3 of the target. The metalation selectivity at the stage of **Int-II''** contrasts with the processes in Table 1D; this is presumably due to the electronic activation provided by the ketone substituent, which also allows the process to operate at a lower temperature (110 °C versus 140–160 °C). To improve the

efficiency, optimization studies were undertaken, resulting in the conditions shown in Table 3B. The protocol was applicable to a range of systems **3a–g** with different substituents at  $\text{R}^1$ ,  $\text{R}^2$ , or  $\text{R}^3$ . In general, the processes were efficient, and a mixture of alkene regioisomers was obtained in each case. Based on the mechanistic analysis in Table 3A, the C3–C4 regioisomers of **4a–g** result from the isomerization of the initially generated C2–C3 regioisomer under the reaction conditions. Attempts to intercept the donor–acceptor cyclopropane intermediates (**Int-IV**) in cycloaddition processes using either internal (**4d**) or external  $\pi$ -unsaturates (e.g., activated ketones) have so far been unsuccessful.<sup>20</sup>

In summary, we demonstrate the first examples of aza-Heck-triggered  $\text{C}(\text{sp}^3)\text{-H}$  functionalization cascades that lead to ring- or spiro-fused cyclopropanes. To enable these processes, a library of largely novel P ligands was designed and evaluated, from which L3 emerged as the optimal ligand. The resulting methodology provides an attractive approach to the synthesis of diverse heterocycles containing core cyclopropanes. These are medicinally valuable scaffolds that are challenging to access by other means. From a reactivity viewpoint, our observations have elucidated how steric or electronic control can be used to govern the  $\text{C}(\text{sp}^3)\text{-H}$  metalation selectivity. In broader terms, the processes described here are unusual because they exploit alkene heteropalladation to trigger  $\text{C}(\text{sp}^3)\text{-H}$  cyclopropanation.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.2c08304>.

Experimental details, characterization data, and crystallographic data (PDF)

### Accession Codes

CCDC 2189444 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif), or by emailing [data\\_request@ccdc.cam.ac.uk](mailto:data_request@ccdc.cam.ac.uk), or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

## AUTHOR INFORMATION

### Corresponding Author

John F. Bower – Department of Chemistry, University of Liverpool, Liverpool L69 7ZD, United Kingdom; [orcid.org/0000-0002-7551-8221](https://orcid.org/0000-0002-7551-8221); Email: [john.bower@liverpool.ac.uk](mailto:john.bower@liverpool.ac.uk)

### Authors

Changcheng Jing – Department of Chemistry, University of Liverpool, Liverpool L69 7ZD, United Kingdom  
Benjamin T. Jones – School of Chemistry, University of Bristol, Bristol BS8 1TS, United Kingdom  
Ross J. Adams – Department of Chemistry, University of Liverpool, Liverpool L69 7ZD, United Kingdom; [orcid.org/0000-0002-7402-2232](https://orcid.org/0000-0002-7402-2232)

Complete contact information is available at <https://pubs.acs.org/doi/10.1021/jacs.2c08304>

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We thank the European Research Council for financial support via the EU's Horizon 2020 Programme (ERC Grant 639594 CatHet), the Leverhulme Trust (Philip Leverhulme Prize to J.F.B.), the EPSRC (EP/N509619/1) (studentship to B.T.J.), and the University of Liverpool (Regius Chair to J.F.B.). We thank the University of Liverpool's Department of Chemistry X-ray crystallography service for analysis.

## REFERENCES

- (1) (a) Talele, T. T. The "Cyclopropyl Fragment" is a Versatile Player that Frequently Appears in Preclinical/Clinical Drug Molecules. *J. Med. Chem.* **2016**, *59*, 8712. (b) Časar, Z. Synthetic Approaches to Contemporary Drugs that Contain the Cyclopropyl Moiety. *Synthesis* **2020**, *52*, 1315.
- (2) McGrath, N. A.; Brichacek, M.; Njardarson, J. T. A Graphical Journey of Innovative Organic Architectures That Have Improved Our Lives. *J. Chem. Educ.* **2010**, *87*, 1348.
- (3) (a) Tahrani, A. A.; Piya, M. K.; Barnett, A. H. Saxagliptin: a New DPP-4 Inhibitor for the Treatment of Type 2 Diabetes Mellitus. *Adv. Ther.* **2009**, *26*, 249. (b) Dong, J.; Gong, Y.; Liu, J.; Chen, X.; Wen, X.; Sun, H. Synthesis and Biological Evaluation of All Eight Stereoisomers of DPP-IV Inhibitor Saxagliptin. *Bioorg. Med. Chem.* **2014**, *22*, 1383.
- (4) Du, W.; Gu, Q.; Li, Z.; Yang, D. Palladium(II)-Catalyzed Intramolecular Tandem Aminoalkylation via Divergent C(sp<sup>3</sup>)-H Functionalization. *J. Am. Chem. Soc.* **2015**, *137*, 1130.
- (5) For recent processes, see: (a) Bao, X.; Wang, Q.; Zhu, J. Palladium-Catalyzed Enantioselective Narasaka-Heck Reaction/Direct C-H Alkylation of Arenes: Iminoarylation of Alkenes. *Angew. Chem., Int. Ed.* **2017**, *56*, 9577. (b) Hazelden, I. R.; Carmona, R. C.; Langer, T.; Pringle, P. G.; Bower, J. F. Pyrrolidines and Piperidines by Ligand Enabled Aza-Heck Cyclizations and Cascades of N-(Pentafluorobenzoyloxy)carbamates. *Angew. Chem., Int. Ed.* **2018**, *57*, 5124. (c) Hazelden, I. R.; Ma, X.; Langer, T.; Bower, J. F. Diverse N-Heterocyclic Ring Systems via Aza-Heck Cyclizations of N-(Pentafluorobenzoyloxy)sulfonamides. *Angew. Chem., Int. Ed.* **2016**, *55*, 11198. (d) Ma, X.; Hazelden, I. R.; Langer, T.; Munday, R. H.; Bower, J. F. Enantioselective Aza-Heck Cyclizations of N-(Tosyloxy)carbamates: Synthesis of Pyrrolidines and Piperidines. *J. Am. Chem. Soc.* **2019**, *141*, 3356. (e) Shuler, S. A.; Yin, G.; Krause, S. B.; Vesper, C. M.; Watson, D. A. Synthesis of Secondary Unsaturated Lactams via an Aza-Heck Reaction. *J. Am. Chem. Soc.* **2016**, *138*, 13830. (f) Xu, F.; Shuler, S. A.; Watson, D. A. Synthesis of N-H Bearing Imidazolidinones and Dihydroimidazolones Using Aza-Heck Cyclizations. *Angew. Chem., Int. Ed.* **2018**, *57*, 12081. (g) Xu, F.; Korch, K. M.; Watson, D. A. Synthesis of Indolines and Derivatives by Aza-Heck Cyclization. *Angew. Chem., Int. Ed.* **2019**, *58*, 13448. (h) Korch, K. M.; Watson, D. A. Total Synthesis of (±)-Impatien A via Aza-Heck Cyclization. *Org. Lett.* **2021**, *23*, 7285. (i) Gao, R.; Shuler, S. A.; Watson, D. A. Tandem Aza-Heck Suzuki and Carbonylation Reactions of O-Phenyl Hydroxamic Ethers: Complex Lactams via Carboamination. *Chem. Sci.* **2021**, *12*, 8859. (j) Jones, B. T.; García-Cárceles, J.; Caiger, L.; Hazelden, I. R.; Lewis, R. J.; Langer, T.; Bower, J. F. Complex Polyheterocycles and the Stereochemical Reassignment of Pileamartine A via Aza-Heck Triggered Aryl C-H Functionalization Cascades. *J. Am. Chem. Soc.* **2021**, *143*, 15593. (k) Geffers, F. J.; Kurth, F. R.; Jones, P. G.; Werz, D. B. Alkyne Aminopalladation/Heck and Suzuki Cascades: An Approach to Tetrasubstituted Enamines. *Chem.-Eur. J.* **2021**, *27*, 14846–14850. For reviews, see: (l) Race, N. J.; Hazelden, I. R.; Faulkner, A.; Bower, J. F. Recent developments in the use of aza-Heck cyclizations for the synthesis of chiral N-heterocycles. *Chem. Sci.* **2017**, *8*, 5248–5260. (m) Vulovic, B.; Watson, D. A. Heck-like Reactions Involving Heteroatomic Electrophiles. *Eur. J. Org. Chem.* **2017**, *2017*, 4996–5009.
- (6) During the course of this work, oxime ester-based cascades that provide cyclobutanes were reported, see: (a) Wei, W.-X.; Li, Y.; Wen, Y.-T.; Li, M.; Li, X.-S.; Wang, C.-T.; Liu, H.-C.; Xia, Y.; Zhang, B.-S.; Jiao, R.-Q.; Liang, Y.-M. Experimental and Computational Studies of Palladium-Catalyzed Spirocyclization via a Narasaka-Heck/C(sp<sup>3</sup> or sp<sup>2</sup>)-H Activation Cascade Reaction. *J. Am. Chem. Soc.* **2021**, *143*, 7868. For other recent C-H functionalization processes that involve N-O cleavage under Pd catalysis, see: (b) Wei, W.-X.; Kong, X.; Jiao, R.-Q.; Li, X.-S.; Wang, C.-T.; Li, Y.; Liang, Y.-M. Regioselective Synthesis of Spirocyclic Pyrrolines via A Palladium-catalyzed Narasaka-Heck/C-H Activation/[4 + 2] Annulation Cascade Reaction. *Chem. Sci.* **2022**, *13*, 6348. (c) Zhang, B.-S.; Jia, W.-Y.; Gou, X.-Y.; Yang, Y.-H.; Wang, F.; Wang, Y.-M.; Wang, X.-C.; Quan, Z.-J. Synthesis of C8-Aminated Pyrrolo-Phenanthridines or -Indoles via Series C(sp<sup>2</sup> or sp<sup>3</sup>)-H Activation and Fluorescence Study. *Org. Lett.* **2022**, *24*, 2104.
- (7) Selected examples involving alkene carbopalladation, see: (a) Liron, F.; Knochel, P. Reactivity of Stable Neopentyl-Pd Intermediates in the Absence of Nucleophile. *Tetrahedron Lett.* **2007**, *48*, 4943. (b) Kim, H. S.; Gowrisankar, S.; Kim, S. H.; Kim, J. N. Synthesis of 6-Oxacyclopropa[a]indene Derivatives Starting from Baylis-Hillman Adducts via Pd-Mediated C(sp<sup>3</sup>)-H Activation. *Tetrahedron Lett.* **2008**, *49*, 3858. (c) Huang, Q.; Larock, R. C. Synthesis of Cyclopropanes by Pd-Catalyzed Activation of Alkyl C-H Bonds. *Tetrahedron Lett.* **2009**, *50*, 7235. (d) Chung, D. S.; Lee, J. S.; Ryu, H.; Park, J.; Kim, H.; Lee, J. H.; Kim, U. B.; Lee, W. K.; Baik, M.-H.; Lee, S.-g. Palladium-Catalyzed Divergent Cyclopropanation by Regioselective Solvent-Driven C(sp<sup>3</sup>)-H Bond Activation. *Angew. Chem., Int. Ed.* **2018**, *57*, 15460.
- (8) To the best of our knowledge, reference 4 details the only prior examples involving alkene aza-palladation. We are unaware of examples involving alkene oxa-palladation.
- (9) Gierczyk, B.; Wojciechowski, G.; Brzezinski, B.; Grech, E.; Schroeder, G. Study of the Decarboxylation Mechanism of Fluorobenzoic Acids by Strong N-Bases. *J. Phys. Org. Chem.* **2001**, *14*, 691.
- (10) White, P. B.; Stahl, S. S. Reversible Alkene Insertion into the Pd-N Bond of Pd(II)-Sulfonamides and Implications for Catalytic Amidation Reactions. *J. Am. Chem. Soc.* **2011**, *133*, 18594.
- (11) (a) Baber, R. A.; Clarke, M. L.; Heslop, K. M.; Marr, A. C.; Orpen, A. G.; Pringle, P. G.; Ward, A.; Zambrano-Williams, D. E. Phenylphosphatrioxa-adamantanes: Bulky, Robust, Electron-Poor Ligands That Give Very Efficient Rhodium(I) Hydroformylation Catalysts. *Dalton Trans* **2005**, 1079. (b) Le, C. M.; Hou, X.; Sperger, T.; Schoenebeck, F.; Lautens, M. An Exclusively *trans*-Selective Chlorocarbonylation of Alkynes Enabled by a Palladium/Phosphadamtane Catalyst. *Angew. Chem., Int. Ed.* **2015**, *54*, 15897.
- (12) CCDC 2189444 contains the crystallographic information for this manuscript.
- (13) The favored aza-palladation mode is supported by previous observations in processes terminated by (fast) β-hydride elimination (see references 5b and 5c). We favor a mechanism where C(sp<sup>3</sup>)-H palladation (rather than C-C reductive elimination) is the product determining step on the basis of DFT calculations reported in reference 7d and KIE studies reported in reference 4.
- (14) The preference for 2j over *iso*-2j may be due to the minimization of steric interactions with the Cbz unit at the stage of the palladacyclobutane. When the cyclization of 1k to *iso*-2k was performed in the presence of D<sub>2</sub>O (300 mol%), no deuterium incorporation was observed in the product (see the SI); this is consistent with the suggestion that C-H metalation is irreversible.
- (15) For other approaches to spirofused pyrrolidines related to those in Table 1D, see: (a) Melnykov, K. P.; Artemenko, A. N.; Ivanenko, B. O.; Sokolenko, Y. M.; Nosik, P. S.; Ostapchuk, E. N.; Grygorenko, O. O.; Volochnyuk, D. M.; Ryabukhin, S. V. Scalable Synthesis of Biologically Relevant Spirocyclic Pyrrolidines. *ACS Omega* **2019**, *4*, 7498. (b) Lin, K.-W.; Yan, S.; Hsieh, I.-L.; Yan, T.-H. Unusual Amphiphilic Carbenoid Equivalent in Amide Cyclopropanation. *Org. Lett.* **2006**, *8*, 2265.

(16) Fairlamb, I. J. S.; Kapdi, A. R.; Lee, A. F.  $\eta^2$ -dba Complexes of Pd(0): The Substituent Effect in Suzuki–Miyaura Coupling. *Org. Lett.* **2004**, *6*, 4435.

(17) Compounds related to those in Table 2 have been accessed by the intramolecular C(sp<sup>3</sup>)–H arylation of cyclopropanes, see: Pedroni, J.; Saget, T.; Donets, P. A.; Cramer, N. Enantioselective Palladium(0)-Catalyzed Intramolecular Cyclopropane Functionalization: Access to Dihydroquinolones, Dihydroisoquinolones and the BMS-791325 Ring System. *Chem. Sci.* **2015**, *6*, 5164.

(18) Other metalation modes can also be envisaged.

(19) Isomerization of the alkene of **3a** was not observed after 1 h under the conditions shown in Table 3B. At the stage of **Int-II'**, the metalation selectivity may be controlled by the closer proximity of the C3–H bond compared to the C7 methyl group. The full 2D NMR analysis of **4a** is presented in the SI.

(20) For a review encompassing this aspect, see: Sokolova, O. O.; Bower, J. F. Selective Carbon-Carbon Bond Cleavage of Cyclopropylamine Derivatives. *Chem. Rev.* **2021**, *121*, 80.

## Recommended by ACS

### $\alpha,\alpha'$ -C–H Bond Difunctionalization of Unprotected Alicyclic Amines

Daniel A. Valles, Daniel Seidel, *et al.*

JULY 29, 2021  
ORGANIC LETTERS

READ 

### Enabling Reductive C–N Cross-Coupling of Nitroalkanes and Boronic Acids by Steric Design of P(III)/P(V)=O Catalysts

Gen Li, Alexander T. Radosevich, *et al.*

MAY 02, 2022  
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

### Catalytic Synthesis of Cyclopropenium Cations with Rh-Carbynoids

Hang-Fei Tu, Marcos G. Suero, *et al.*

SEPTEMBER 08, 2022  
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

### Complex Polyheterocycles and the Stereochemical Reassignment of Pileamartine A via Aza-Heck Triggered Aryl C–H Functionalization Cascades

Benjamin T. Jones, John F. Bower, *et al.*

SEPTEMBER 21, 2021  
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Get More Suggestions >