

Northwest Atlantic



Fisheries Organization

Serial No. N6822

NAFO SCR Doc. 18/033REV

SCIENTIFIC COUNCIL MEETING – JUNE 2018

The status of redfish (*S. mentella* and *S. fasciatus*) in Divisions 3LN at present and the likelihood its follow up in the near future (under the ongoing Management Strategy or a status quo TAC scenario)

by

A. M. Ávila de Melo ¹, R. Alpoim ¹, D. González Troncoso ², F. González ² and M. Pochtar ³

¹ Instituto Português do Mar e Atmosfera-Departamento do Mar e Recursos Marinhos, IPMA-DMRM,
Rua Alfredo Magalhães Ramalho, nº6, 1495-006, Lisboa, Portugal

² Instituto Español de Oceanografía, Aptdo 1552, E-36280 Vigo (Pontevedra), Spain

³ Knipovich Polar Research Institute of Marine Fisheries and Oceanography (PINRO), 6 Knipovitch street, Murmansk 183763 Russia

Abstract

There are two species of redfish in Divisions 3L and 3N, the deep-sea redfish (*Sebastes mentella*) and the Acadian redfish (*Sebastes fasciatus*) that have been commercially fished and reported collectively as redfish in fishery statistics. Both species, occurring on Div. 3LN and managed as a single stock, don't belong to isolated local populations but, on the contrary, are part of a large Northwest Atlantic complex ranging from the Gulf of Maine to south of Baffin Island.

The ASPIC assessment of this stock is based on the logistic form of a non-equilibrium surplus production model (Schaeffer, 1954; Prager, 1994), adjusted to a standardized catch rate series (Power, 1997) and to all stratified-random bottom trawl surveys conducted in various years and seasons in Div. 3L and Div. 3N from 1978 onwards. Both CPUE and surveys were used with all observations within each series.

The 2018 assessment proceed on the threshold of the new 2014 approach, the main features of the previous input framework were kept, with MSY fixed at 1960-1985 average catch and the rest of the approved 2014 assessment framework updated. Despite its poor performance the 3L Spanish survey has been kept in the analysis due to its high correlation with the autumn 3LN Canadian survey, one of the two backbone series of the assessment.

ASPIC results confirm a stable stock from the 1960's to the first half of the 1980's, sustaining an average yield of 21 000t. Stock declined with a sudden rise of the catch over the late 1980's first half of the 1990's, and started to gradually recover after catches fell to a residual level when the stock collapse. The maximum observed sustainable yield (MSY) of 21 000 t can be a long term sustainable yield if fishing mortality stands at 0.11/year, exploiting a correspondent B_{msy} at 187 000 t.

From assessment results there is a high probability (CL's 80%) that the stock was at the beginning of 2018 above B_{msy} , after crossing 2017 under a fishing mortality most likely below 34% F_{msy} .



From short term stochastic projections there is a high probability (CL's 80%) that catch on 2018 at 14 200 t TAC increasing to 18 100 t on 2019 and 2020 (as predicted in the 2014 Risk-Based Management Strategy for 3LN Redfish) will keep fishing mortality on 2018-2020 below F_{msy} and biomass above B_{msy} till the beginning of 2021.

However, preliminary analysis of the majority of recent data (2016-2017) from ongoing surveys and commercial fisheries suggests that the 3LN redfish management unit has stop growing and may even been declining. No evidence of good recruitments and a likely shrinkage of the length/age population structure are also in line with that hypothesis.

Introduction

There are two species of the genus *Sebastes* with distribution overlapping in several areas of Northwest Atlantic, namely on the Gulf of St. Lawrence, Laurentian Channel, off Newfoundland and south of Labrador Sea: the deep sea redfish (*Sebastes mentella*), with a maximum abundance at depths greater than 350m, and Acadian redfish (*Sebastes fasciatus*), preferring shallower waters of less than 300m (DFO, 2008). They have been commercially fished on the slopes of the Grand Bank, both on Div. 3LN (north-south east) and Div. 3O (south-west).

Due to their external resemblance *S. mentella* and *S. fasciatus* are commonly designated as beaked redfish. Beaked redfish are viviparous with the larvae eclosion occurring right before or after birth, long living and slow growing, with females attaining size of 50% maturity at 30-34cm (Power, 2001). Both species have pelagic and demersal concentrations as well as a long recruitment process to the bottom. Their external characteristics are very similar, making them difficult to distinguish. Therefore, they are reported collectively as "redfish" in the commercial fishery statistics. *S. mentella* and *S. fasciatus* are also treated as a single species in the Grand Bank surveys carried out by Canada, Russia and more recently by EU-Spain.

Either redfish species occurring on Div. 3LN don't belong to isolated local populations but, on the contrary, are part of a large Northwest Atlantic complex ranging from the Gulf of Maine to south of Baffin Island. This complex is centred on the Gulf of St. Lawrence (GSL)- Laurentian Channel - western slope of the Grand Bank system, where the GSL is a main nursery area for *S. fasciatus* and *S. mentella* local populations, due to current patterns that favours larval drift from main regions of larval extrusion towards the gulf.

As regards redfish occurring on the Gran Bank of Newfoundland, differences observed in the "state of the stock" between Div. 3O and Div. 3LN suggests that it would be prudent to keep Div. 3LN as a separate management unit. Being so, beaked redfish in Div. 3LN has been considered by NAFO Scientific Council as a management unit composed by the ensemble *S. mentella* - *S.fasciatus* aggregations overlapping on the north-south east parts of the Grand Bank.

Within this management unit, relative abundance of *S. mentella* - *S.fasciatus* may vary with the income and survival of juveniles from either species, though *S. fasciatus* tend to be more abundant in the south (Div. 3N) while *S. mentella* is more abundant in the northern division (3L). On 2011-2015 most of Canadian spring and autumn surveys found larger redfish concentrations more frequent on Div. 3N (Rideout, 2016), despite the major proportion of the catch being annually taken from Div. 3L. *S. mentella* dominates the commercial catch on either division.

Commercial Fishery

Nominal catches and TAC's

Between 1959 and 1960 reported catches drop from 44 600 to 26 600 t, oscillating over the next 25 years (1960-1985) around an average level of 21 000 t. Catches rise afterwards to a 79 000 t high in 1987 and fall steadily to a 450 t minimum reached in 1996. Catches were kept at a low level (450-3 000 t) until 2009.

The NAFO Fisheries Commission (FC) implemented a moratorium on directed fishing for this stock in 1998. In June 2009 the Scientific Council confirmed the upward trend of the stock as shown by spring and autumn surveys (NAFO, 2009). The Fisheries Commission endorsed the Scientific Council recommendations from the 2011 onwards and catches increased being at 11 800t in 2017, the highest level recorded since 1993 (Table 1, Fig. 1). Since the reopening in 2010 Canada, followed by Russia and EU-Portugal are the main partners of a fishery mostly deployed northwards, in Div. 3L.

The 2011-2016 catches were taken from the NAFO STATLANT 21 data base. Last year's catch (2017) was estimated with the CDAG method and given by the NAFO Joint Commission-Scientific Council Catch Estimation Strategy Advisory Group (COM-SC CESAG, 2018).

The perception of the recent stock status, with biomass at or above B_{msy} and fishing mortality below F_{msy} , justified the adoption by the FC on the 36th Annual Meeting – September 2014 of a Risk-Based Management Strategy (MS) for redfish in Divisions 3LN (Ávila de Melo *et al.*, 2014; FC Working Paper 14/23). This MS was designed to reach 18 100 t of annual catch by 2019-2020 through a stepwise biannual catch increase of constant magnitude. That resulted on a TAC for 2017-2018 of 14 200t.

Description of the fishery

In the early 1980's the former USSR, Cuba and Canada were the primary fleets directing for redfish in Div. 3LN. The rapid expansion of the fishery was due to the entry of EU-Portugal in 1986 and South Korea in 1987, along with various re-flagged fleets. In the early 1990's Russia and the Baltic mid-water trawlers, together with South Korea and Portuguese bottom trawlers, were still responsible for the bulk of fishing effort, concentrated by that time on the "Beothuk Knoll" (divisions 3L-3M-3N border, southwest of the Flemish Cap).

South Korea left the area by the end of 1993 and from 1994 onwards the other fleets reduced their effort substantially on the north and south east of the Grand Bank. The quick decline of catch rates was the main reason for this reduction of redfish fishing effort, and justified its partial shift southwest to Div. 3O. Since 1994 most of the redfish catches in NAFO Divisions 3L and 3N were taken as by-catch of the Greenland halibut fishery pursued from the northern slopes of the Sackville Spur in Div. 3L through Flemish Pass till the canyons of southern Grand Bank in Div. 3N. EU-Portugal and EU-Spain bottom trawl fleets were the main fleets responsible for the 3LN redfish by-catch during the moratorium years.

Since its reopen in 2010 Canada, Russia and EU-Portugal are the main partners of a fishery mainly deployed northwards, on Div. 3L. However, the fishery turn south last year, with Div. 3N having the major share of the catches for the first time in many years, while Canada almost vanish from catch records.

Catch and Effort

On the 1997 assessment (Power, 1997) catch/effort data for Div. 3L and Div. 3N from 1959 to 1995 were analyzed with a multiplicative model (Gavaris, 1980) in order to derive a catch rate series for each division standardized for country-gear-tonnage class, NAFO division, month, and amount of by-catch associated with each observation. Both CPUE series shows much within year variability over time, with no statistically difference between the catch rates for most of the years. That assessment considered that *catch rate indices for Div. 3L and Div. 3N were not reflective of year to year changes in population abundance but they may be indicative of trends over longer periods of time.*

ASPIC assessments recovered the predicted effort series in fishing hours for Div. 3L and Div. 3N from the 1997 multivariate analysis, in order to derive a single annual catch rate for Div. 3LN. For each year of the 1959-1994 interval, this standardized catch rate is given by the ratio between the sum of Div. 3L and Div. 3N STATLANT catch (thousand tons) and the sum of Div. 3L and Div. 3N predicted effort (fishing hours). The catch rate series for Div. 3LN is presented on Table 2 and Fig. 2 (standardized to zero mean and unit standard deviation in the figure). Catch rate for Div. 3LN increased on the first years of the time series, 1959 till 1967,

oscillated around the average on the intermediate years and declined after 1987. On the final years of this CPUE series, 1990-1994, catch rates were stable at a minimum level.

Commercial catch@length

Most of the commercial length sampling data available for the 3LN beaked redfish came, since 1990, from the Portuguese fisheries and have been annually included in the Portuguese research reports on the NAFO SCS Document series (Vargas *et al.*, 2018). Taking into account that the majority of the length sampling was from depths greater than 250m, most of these data should represent *S. mentella* catches. Length data from Spain and Russia were used to estimate the length composition of the commercial catches for those fleets in several years (González *et al.*, 2018; Fomin *et al.*, 2018; Pochtar, *pers comm.* 2018). The 1990-2017 per mille length composition of the Portuguese trawl catch was applied to the rest of the commercial catches (Table 3a). Commercial length weight relationships used to get catch numbers at length were derived from redfish sampling on board of Portuguese vessels fishing on divisions 3L and 3N (Table 3b).

The overall mean length of the 1990-2017 catch (arithmetic mean of the annual mean lengths of the commercial catch) was used to derive length anomalies of the 3LN catch over this period (Table 3a, Fig. 3). The proportion of small redfish (less than 20cm) in the catch is presented as well on the bottom of Table 3a. The purpose of the length anomalies was to detect possible shifts in the length structure of the catch that could reflect changes in the length structure of the exploitable stock.

Above average mean lengths, apparent stable catch at length with no clear trends towards smaller or larger sizes, proportion of small redfish usually below 1% are observed on most of the years of the 1990-2005 interval. However, well below average mean lengths coupled with in excess of 10% of small redfish in the catch occurred afterwards, on most years between 2006 and 2015 (Table 3a, Fig. 3). Under a low exploitation regime such interlinked events should reflect an average level of recruitment over those years well above the low recruitments from the 1990's first half of the 2000's. Average proportion of small redfish in the commercial catch rose from 1.0% (1990-2005) to 13.9% (2006-2015).

However, proportion of small redfish fell to 7.2% in 2016 and again to 1.6% in 2017 while the mean length in the catch gradually increased, approaching the overall 1990-2017 mean and larger sizes are recently the bulk of the catch (Table 3a). An important increase on the numbers of small redfish in the catch can reflect the income of one or more good recruitments but, on the contrary, a noticeable decline on this indicator, as observed on recent years, can signal that year classes coming in the fishery are now below average or even weak. And that exploitable stock is again basically relying on the survival of the year classes already recruited.

Research Surveys

From 1978 till 1990 several stratified-random bottom trawl surveys have been conducted by Canada in various years and seasons in Div. 3L. However only since 1991 Canadian stratified-random surveys covered both Div. 3L and Div. 3N on a regular annual basis: a spring survey (May-Jun.) and an autumn survey (Sep.-Oct. 3N/Nov.-Dec. 3L for most years). No survey was carried out on Div. 3N in spring 2006 and autumn 2014. In the spring of 2017 there were problems with 3L survey coverage and none of the 3L strata in the redfish index were sampled (Rideout and Ings, 2018). As regards Canadian surveys, only Campelen data and Engel data converted into Campelen equivalents are used in this assessment.

Since 1983 Russian bottom trawl surveys in NAFO Div. 3LMNO turn to stratified-random, following the Doubleday stratification for Sub area 3. On 1984 standard tows were set to half hour at 3.5 knots, with a standard gear. On 1992 redfish results of the 1984-1991 stratified-random surveys in Div. 3LN by Russia were revised according to standard methodology (Power and Vaskov, 1992) and since 2008 this "Power revised" 1984-1991 Russian survey series is incorporated in the input framework of the redfish 3LN ASPIC assessment (Ávila de Melo *et al.*, 2008). Between 1992 and 1994 the coverage of NAFO Sub area 3 by the Russian bottom trawl series became irregular and in 1995 was discontinued.

In 1995 EU-Spain started a new stratified-random bottom trawl spring (May-June) survey on NAFO Regulatory Area of Div. 3NO. All strata in the NRA were covered every year following the standard stratification, first till 732m and from 1998 onwards till the 1464 m depth contour. In 2003 the Spanish survey was extended northwards to some strata in Div. 3L, but it was only in 2006 that an adequate prospecting survey was first conducted in Division 3L with over 100 valid hauls (Róman *et al.*, 2018).

Details on the two Canadian survey series, as well as on the Russian series and the two Spanish surveys can be found on previous assessments (Ávila de Melo *et al.*, 2014).

Survey biomass and female spawning biomass

All survey biomass series from stratified-random bottom trawl surveys used in the 2018 ASPIC assessment are presented in Table 4. The 1991-2016 (spring) 2017 (autumn) female SSB Canadian survey indices for Div. 3LN combined are included on Table 4. In order to turn the survey series comparable and facilitate the detection of trends in stock dynamics, the survey biomass series used in the assessment framework and the female SSB survey series were standardized to zero mean and unit standard deviation and so presented on Figure 4a and 4b.

From the late 1970s to the beginning of the 1990s Canadian surveys in Div. 3L and Russian bottom trawl surveys in Div. 3LN suggest that stock size suffered a substantial reduction. Redfish bottom biomass from surveys in Div. 3LN remained well below average level over the 1990's and early 2000's, but since 1997 those indices start to show some dynamics of increase. The inconsistent dynamics turn to clear increases of survey biomass indices on 2007-2015, but, with the exception of the 2016 Canadian 3LN spring, the rest of the ongoing surveys went down on 2016-2017. Still, considering the overall picture given by all available bottom trawl survey series occurring in Div. 3L and Div. 3N from 1978 till 2017, 91% of the observations were above the average of their own series on 1978-1985, only 3% on 1986-2006, and 82% on 2007-2017.

In order to estimate spring and autumn female spawning survey biomass by division, Div. 3L and Div. 3N female maturity at length vectors (Power 2001; Ávila de Melo *et al.*, 2005) were applied to the 1991-2017 female abundances at length of the spring and autumn surveys. Female spawners and stock abundance at length by division were used to calculate SOP female spawning and stock biomass for Div. 3L and Div. 3N, using female and sex combined length weight relationships derived from data collected on board of the Canadian 3LN autumn surveys, 1997-2004 (Power, *pers. comm.*, 2005), of the 3N Spanish survey, 2005, and of the 2006-2017 3LN Spanish survey (González *et al.* 2018; González, *pers. comm.*, 2018). The SOP ratios (SSB/stock biomass) by division were then applied to the respective swept area survey biomasses to give the spring and autumn female SSB in Div. 3L and Div. 3N.

Both 1991-2017 Canadian spring and autumn standardized female SSB survey series for Div. 3LN have trends concurrent to their correspondent biomass series (Fig.4b).

Abundance at length

Spring and autumn survey abundance at length, for Div. 3LN combined, are presented in Table 5a and 5b, respective length anomalies in Fig. 5a and 5b. The overall 1991-2016/2017 mean length for each survey series (arithmetic mean of the annual mean lengths of the survey abundances at length) was used to derive the spring and autumn survey length anomalies for the stock over this period (Table 5a and 5b last line, Fig. 5a and 5b). During the first half of the 1990's, on both surveys, the length anomalies were negative or slightly positive. Mean lengths on most of the years between 1996 and 2007 (spring survey) or 2006 (autumn survey) were above the mean, reflecting a shift on the stock length structure to larger individuals probably justified by a higher survival of the main year classes crossing the stock through this time interval and a regular income of weak year classes. Between 2007-2008 and 2011-2012 mean lengths generally fall and stay below average, just as observed on the commercial catch at length (Fig. 3). This recent pattern on the length structure of both surveys and commercial catch suggests the occurrence of good recruitments by the late 2000's, after a low productivity regime that prevailed for more than 15 years. At the same time larger sizes =>20cm increase their abundance until 2015, both in surveys and commercial catch (Fig. 5c), confirming

that both good recruitments and high survival rates were in place at the time. This picture is not valid anymore.

In fact, on 2016-2017, from Canadian survey data, mean length in the stock increased but the numbers of fish =>20cm declined. This is not only observed in the stock but in commercial catch as well (Fig's 5a, 5b and 5c).

Between 2006-2007 and 2009-2010 the recruitment index (numbers of redfish < 20cm) increased rapidly both in commercial catch and Canadian surveys, reaching by then maximum values. The recruitment index drops fast on the following years and is at low levels since 2014-2015 (Fig. 5d). Nevertheless, unusual high numbers of very small redfish pre recruits (5-12cm) have been observed on recent years (2015-2017) on Canadian spring and autumn surveys.

All these indicators from most recent biomass levels and stock structure, given by the surveys and the fisheries, suggest that the stock is not growing, and has either reach a stable level or is making a downward turn. And they need to be taken into to consideration when analyzing the assessment results next.

ASPIC assessment suite

Brief history and background for the pre fixed *MSY* option

A non-equilibrium surplus production model (ASPIC; Prager, 1994) is used to assess the status of the stock since 2008. The ASPIC operating model is a non-equilibrium implementation of Schaefer's and Pella-Tomlinson models, among others.

Until 2012 the model was adjusted to an array of Canadian, Russian and Spanish surveys series arranged under the formulation adopted on the "*The 2nd Take of the 2008 Assessment of Redfish in NAFO Divisions 3LN,*" (Ávila de Melo and Alpoim, 2010). However, the model was showing an increasing unfitness to survey biomass increases observed by the second half of the 2000's-early 2010's on all ongoing surveys. The approved framework of the 2012 assessment ends up excluding the 3N Spanish survey and several inter annual biomass bumps from the Canadian survey series, either combined (spring) or separate (autumn) (Ávila de Melo *et al*, 2012).

On the next assessment (Ávila de Melo *et al*, 2014) the purpose was to reach an inclusive approach that would incorporate most, if not all, the survey points available for the two divisions with no haircuts, and at the same time delivering a "realistic" output. In other words, resulting on key parameters and biomass and fishing mortality trajectories in line with the perception of stock and fishery dynamics one has from historical survey and commercial time series. To achieve this goal two of the five input frameworks running on the exploratory analysis preceding the 2014 assessment were allowed to run with maximum sustainable yield fixed at a user starting guess of 21 000 t. This *MSY proxy* is the average level of sustained catch for the 1960-1985, when the stock experienced an apparent stability, suggested either by the STATLANT CPUE series or available surveys, and before stock decline in response to a sudden rise of catch level.

From exploratory analysis the better framework to run the 2014 assessment had *MSY* pre fixed at 1960-1985 average catch. This framework also kept negative correlated STATLANT CPUE series and all "outliers" in their respective survey series, with Canadian autumn surveys on Div. 3L and Div. 3N assembled in a single 3LN Canadian autumn series.

Input series

This assessment keeps the selected arrangement of input series considered on 2016 the better framework to run the redfish 3LN ASPIC (Ávila de Melo *et al*, 2016): with *MSY* fixed at 1960-1985 average catch, the suite of survey time series already approved for the 2014 assessment, updated and now including the 3L Spanish survey.

All input series consist of annual observed values and were given equal weight in the analysis. Each Canadian series is referred by its season and division(s), while the Russian and Spanish series are also referred by their country name. The input series of this assessment are

I1 (Statlant CPUE and catch)	Statlant cpue for Div. 3LN, 1959-1994 & catch for Div. 3LN, 1959-2017	
I2 (3LN spring survey)	Canadian spring survey biomass for Div. 3LN, 1991-2005, 2007-2016	
I3 (3LN autumn survey)	Canadian autumn survey biomass for Div. 3LN, 1991-2017	
I4 (3LN Power russian survey)	Russian spring survey biomass for Div. 3LN, 1984-1991 (Power and Vaskov, 1992)	
I5 (3L winter survey)	Canadian winter survey biomass for Div. 3L, 1985-1986 and 1990	
I6 (3L summer survey)	Canadian summer survey biomass for Div. 3L, 1978-1979, 1981, 1984-1985, 1990-1991 and 1993	
I7 (3L autumn survey)	Canadian autumn survey biomass for Div. 3L, 1985-1986, 1990	
I8 _a (3N spring spanish survey _{long})	Spanish survey biomass for Div. 3N, 1995-2017	
I9 (3L summer spanish survey)	Spanish survey biomass for Div. 3L, 2006-2017	

The CPUE series and the short survey series (Russian survey, Canadian summer, autumn and winter surveys on Div. 3L), reflect the stock dynamics from the early 1960's until the first half of the 1990's, while the spring and autumn Canadian surveys reflect the stock dynamics from the 1990's till nowadays. Trends within the two periods differ and overlap of series mostly belonging to different intervals is short. The negative correlations found between "old" and "new" series are expected (and disqualified to halt the ASPIC assessment). Unfitness between observed and estimated STATLANT CPUE series is also expected, as the observed lay over the first half of the assessment interval ending in 1994 while the correspondent estimated STATLANT CPUE series is extended until the last year (2017) in line with the stock biomass estimated by the model.

Basic assumptions on ASPIC fit mode

In this assessment the new ASPIC version 7.03 (Prager, 2015) fit the logistic form of the production model (Schaefer, 1954). Being K the carrying capacity stock biomass, r the intrinsic rate of stock biomass increase, C the catch biomass, MSY and B_{msy} the long term yield and biomass associated with F_{msy} , the model basic assumptions are:

- 1) A logistic population growth over time of the unexploited stock (Schaefer, 1954)

$$dB_t / dt = rB_t - (r/K)B_t^2 \quad (1)$$

- 2) For an exploited stock catch is also incorporated in the population growth

$$dB_t / dt = rB_t - (r/K)B_t^2 - C_t \quad (2)$$

- 3) The biological reference points are

a. $MSY = rK / 4 \quad (3)$

b. $B_{msy} = K / 2 \quad (4)$

c. $F_{msy} = r / 2 \quad (5)$

Starting with user guesses (seeds) for the key parameters and catchability coefficients, ASPIC fit generate iteratively an expected series for each observed series of the input framework. Key parameters of the model are found by a minimization routine that gathers the sums of log squared residuals within each series.

The model assumes that all catchability coefficients are constant over time. Because of the imprecision associated with the estimate of catchability for the various indices, absolute estimates of stock size and fishing mortality are normalized to the stock size and fishing mortality at MSY (B_{msy} and F_{msy} respectively). That is why normalized estimates are used in the trajectories of biomass and fishing mortality. In a production model fishing mortality refers to catch/biomass ratio.

A detailed summary of the ASPIC model (Prager, 1994) is available at the 2003 assessment of redfish in Div. 3M (Ávila de Melo *et al.*, 2003).

Input file settings

ASPIC model requires from the user a set of initial definitions/starting guesses/constraints and data series, all of them included in a single input file. On ASPIC 7.03 input format has changed, but the updated 2018 input file is arranged on version 5 format and then converted to the new format using the utility program ASPIC5to7.

Control parameters are kept from the 2014 assessment and line-by-line details of all input settings can be found on the correspondent SCR Doc. (Ávila de Melo *et al.*, 2014). However, problems were found on the run of ASPIC boot with too many trials replaced due to q 's and B1/K estimates at their bounds. At the start of the 2018 assessment Line 17 of the ASPIC.inp file (version 5 format) stayed as follows:

- STATLANT CPUE, 9.007E-06 (q of STATLANT CPUE for Div. 3M redfish ASPIC assessment, Ávila de Melo *et al.* 2003);
- spring survey on Div. 3LN combined, 1.0000E+00;
- autumn survey on Div. 3LN, 1.0000E+00;
- Russian survey on Div. 3LN combined, 1.0000E+00;
- winter survey on Div. 3L, 0.322d0 (average 1991-2009 3L/3LN spring survey biomass ratio times average 1991-2009 spring 3LN/autumn 3LN survey biomass ratio);
- summer and autumn survey on Div. 3L and Spanish survey on this division 0.275d0 (average 1991-2009 3L/3LN autumn survey biomass ratio);
- Spanish survey on Div. 3N 0.759d0 (average 1991-2009 3N/3LN Canadian autumn survey biomass ratio).

But taking into account the problems found, the user guess catchabilities for each of the 9 data sets were found by a 3 steps proceeding:

1 q user guess = average ratio survey biomass (one division)/survey total biomass (two divisions)
 or 1.00E+00 in the case of q for two divisions combined ## q (starting guesses -- 1 per data series)
 9.01E-06 1.00E+00 1.00E+00 1.00E+00 3.22E-01 2.75E-01 2.75E-01 7.59E-01 2.75E-01

2 q user guess = q max bound of ASPIC.fit run with q user guess 1
 9.01E-04 6.00E+00 6.00E+00 6.00E+00 1.20E+00 1.20E+00 1.20E+00 4.55E+00 1.20E+00

3 q user guess= q estimate of ASPIC.fit run with q user guess 2
1.50E-05 8.85E-01 8.85E-01 3.17E-01 2.49E-01 1.03E+00 2.30E-01 8.75E-01 6.61E-01

From this stage on, the q 's at step 3 were the user guess that started all runs of the 2018 assessment.

The 1959-2010 catches used are the catches adopted by STACFIS for this stock. The 2011-2016 catches were taken from the NAFO STATLANT 21 data base. Last year's catch (2017) was estimated with the CDAG method and given by the NAFO Joint Commission-Scientific Council Catch Estimation Strategy Advisory Group (COM-SC CESAG, 2018).

All data sets have now 59 years' length (1959-2017). Input.a7inp file for the 2018 framework is on Appendix 1.

Assessment results

ASPIC2018 run first on deterministic (FIT) mode. Results are presented on Appendix 2. Relative biomass and fishing mortality fit trajectories are plotted on Fig's 6a and 6b.

In terms of biomass dynamics results showed a good nearness index, crossing twice B_{msy} and presenting good contrast. Besides no correlation between series with a very small number of pair-wise observations, 3L Spanish don't fit with 3LN spring survey and is poorly correlated with 3N Spanish as well. This time the model is unable to fit to this data set. But in turn 3L Spanish has a good correlation with 3LN autumn survey, both declining on 2016-2017, so despite the caveats the survey stayed. From the correlations between series and between model results and respective data sets it is clear at this stage the existing of two conflicting trends, one upwards based on the 3LN spring and another downwards pulled by the 3LN autumn survey, with the help of the 3L Spain set. On the outcome, an upward signal prevails and the 2018 ASPIC fit assessment gives as results a stock biomass still growing away from B_{msy} and a fishing mortality still kept well below F_{msy} . But, as a consequence of these conflicting trends, correlation among input series generally decrease and unfitness of the model to the main surveys increased from last 2016 assessment (Ávila de Melo *et al.*, 2016). Nevertheless the model fit run ended normally with a relatively small number of restarts required for convergence: made possible by the *fixed MSY* approach that allows an assessment with a broader input framework (all surveys and all observations on board).

To investigate whether or not there was statistical evidence of model mis-specification, a Wald-Wolfowitz runs-test (Brandão and Butterworth, 2018) was carried out on the residuals of the fits of the surplus production model to the four abundance indices that cover recent years: 3LSpain, 3NSpain, 3LNautumn and 3LNspring. The respective p-values under the hypothesis of independence of the residuals for each of these series were respectively 0.030, 0.670, 0.313 and 0.369, i.e. only for the 3LSpain series is the hypothesis of independence of residuals rejected at the 5% level, which would in turn indicate model mis-specification. Viewed overall, these results offer weak support only for considering the model fit to the data to be unsatisfactory (Butterworth *pers comm.*, 2018)

The assessment switched afterwards to bootstrap mode (BOT, 1000 trials) to measure variability around parameter point estimates using bootstrap methods to calculate very high and high probability confidence limits, here associated with 80% and 60% CL respectively. Estimates from bootstrap analysis are presented in Appendix 3 (with main results and relative IQ range from ASPIC2016 within brackets for consistency check). Due to the previous approach to the user q 's versus maximum bounds for the nine input data sets (in preparation of the fit run) this year bootstrap run smoothly compared to 2016 with no trials replaced with q at bound and only five trials replaced with $B1/K$ at bound (against 429 and 13 replacements in 2016)

For reasons already explained in previous assessments (Ávila de Melo *et al.*, 2014 and 2016) the best available estimator of central tendency is the point estimate. Being so, on the recent ASPIC version 7.03 (Prager, 2015), both estimated bias in point estimate and estimated relative bias were removed from the bootstrap output summary table (Appendix 3). Prager kept in the revised ASPICbot output the point estimates now associated with the bias-corrected confidence limits, since they are within these limits as well. A description of how these bias corrected confidence intervals are computed can be found in Prager (1994) and in Efron and Gong (1983).

Bootstrap results confirm a stock at the beginning of 2018 with a very high probability to be above B_{msy} and a fishing mortality in 2017 with a very high probability to be well below F_{msy} (Appendix 3). Very similar key parameters (B_{msy} , F_{msy} , $B1/K$) and smooth progression from 2016 to 2018 results, all tied to relative inter-quartile ranges within the same previous order of magnitude (2016 in brackets) highlight the consistence of between the 2018 and the 2016 assessment, the first to run with the same survey data sets.

From ASPIC assessment results the maximum observed sustainable yield (MSY) of 21 000 t can be a long term sustainable yield if fishing mortality stands at a level of 0.112/year. The correspondent B_{msy} for this stock is at the level of 187 000 t.

Catch versus surplus production trajectories are presented on Fig. 7. Between 1960 and 1985 catches form a scattered cloud of points around the surplus production curve. On 1986-1987 catch rise well above surplus production and, though declining continuously since then, was still above equilibrium yield in 1993. Catch has drop well below surplus production on 1995 and from 2010 onwards has been slowly increasing, but always bellow the equilibrium yield roof.

Stock and fishing mortality trajectory within a Precautionary Approach framework

The ASPIC point estimate results were put under the precautionary framework on Fig. 8. During the first 25 years the trajectory shows a stock within $B_{msy} - 1.2 B_{msy}$ exploited in the vicinity of F_{msy} (1960-1985). The stock rapidly declined afterwards to well below B_{msy} when fishing mortality rises to well above F_{msy} (1987-1994). Fishing mortality dropped in 1996 to well below F_{msy} , being kept at a low to very low level ever since.

The NAFO SC Study Group recommendations from the meeting in Lorient in 2004 (NAFO, 2004), as regards Limit Reference Points (LRP's) for stocks evaluated with surplus production models, considered F_{lim} at F_{msy} and F_{target} at $2/3 F_{msy}$. The Study Group also considered that the biomass giving production of 50% MSY was a suitable B_{lim} . With the Schaeffer model used in the present ASPIC assessment this biomass corresponds to (roughly) 30% B_{msy} . The stock was below B_{lim} between 1993 and 2002. Biomass reaches and surpasses B_{msy} ten years after (2010-2011).

ASPIC short term catch projection under the actual management strategy

Background for catch projection

The Risk-Based Management Strategy (MS) for 3LN Redfish adopted by the Fisheries Commission on the 36th Annual Meeting – September 2014 (Ávila de Melo *et al.*, 2014; FC Working Paper 14/23), was designed to reach 18 100 t of annual catch by 2019-2020. It has a Harvest Control Rule (red 3LN HCR) that holds to a stepwise constant catch increase every two years, between 2015 and 2020.

18 100 t was the equilibrium yield given by the 2014 assessment, already carried out under the assumption of an MSY of 21 000 t. Since then the following assessments monitored the impact of the MS on the stock. From their results 3LN redfish has stand over recent years (2016-2018) with biomass sizes above B_{msy} and fishing mortalities well below F_{msy} .

At last short term catch projection should quantify the likelihood of the stock 1) to be exploited below F_{msy} until the end of 2020, assuming that the 2018 TAC will be effectively taken and the 2019-2020 catch will reach the HCR 2019-2020 TAC of 18 100 t and 2) to arrive to the beginning of 2021 still on the safe zone above B_{msy} .

But regardless the good results of the present ASPIC assessment the fact is that of the four ongoing surveys three shown consecutive declines on 2016 and 2017 (3LN autumn survey, 3N spring and 3L Spanish surveys) and the other (3LN spring survey) has no results for redfish on 2017. Furthermore from either commercial or survey the catch there are no signs of the income of good recruitments whereas length spectrum of the stock seem to be shrinking. In short there is no evidence that nowadays the stock is still growing. On top of these signals from observed data the 2018 assessment is giving an equilibrium yield roughly 16% below the next 2019-2020 TAC of 18 100 t.

Those are strong arguments in favour of considering on the short term projections an alternate option to the red 3LN HCR (2019-2020 TAC of 18 100 t), keeping instead the catch level on the next coming years at the 2017-2018 TAC (14 200 t).

ASPIC projection framework

ASPICP, the ASPIC auxiliary program for projections, provided point estimates (with associated bias corrected 80% and 50% confidence limits) of biomass and fishing mortality for the assessment time interval, 1959-2017, extended to the projection years, 2018-2021, with 2018 catch at the present TAC and at the 2019-2020 HCR TAC on the next coming years. ASPICP reads the results from the 1000 trials of the ASPIC_{bot} 2018 assessment stored in a .bio file and project each of these runs three years ahead with the each of the two following 2018-2020 catch options:

1) the red 3LN HCR option	2018: 14 200 t	or the status quo 2017 TAC option	2018: 14 200 t
	2019: 18 100 t		2019: 14 200 t
	2020: 18 100 t		2020: 14 200 t

ASPICP read the specifications for these two short term catch projections from two control files with a .ctl extension. The control file format is presented and explained on Appendix 4 (with the red 3LN HCR option). To run ASPICP the .ctl files for each option were dragged and dropped to the available ASPICP shortcut.

Projection results

The 1959-2021 ASPICP results with the two projection scenarios were stored in .prj files and are presented on Table 6 and Fig. 9a and 9b (HCR option) and Table 7 and Fig. 10a and 10b (status quo 2017 TAC option).

Comparisons of results with the two options are presented in Table 8a and Fig. 11 (for B_{msy} 2017-2021) and Table 8b (for F_{msy} 2017-2021).

Either the HCR predicted catch increase or catch at status quo 2017 TAC on 2019 and 2020 will maintain biomass at the beginning of 2021 above B_{msy} while keeping fishing mortality till 2019 below F_{msy} . However the second option will allow biomass level roughly to be kept at its present level, avoiding the beginning of a marginal decline predicted by the first option that is already suggested by the majority of recent observed data.

Acknowledgements

This assessment is part of a EU research project supported by the European Commission (DG Mare, Program for the Collection of Data in Fisheries Sector), IPMA (EU-Portugal) and IEO (EU-Spain). The authors would like to thank Danny Ings (Science, Oceans and Environment Branch, DFO, St. John's, NL, Canada) for the early submission of the 2016 and 2017 data from the spring and autumn Canadian surveys on Div. 3L and Div. 3N, and Doug Butterworth (Department of Mathematics and Applied Mathematics, University of Cape Town South Africa) for his advice and assistance regarding the testing of the model fit to survey data. We also would like to express our gratitude to Dayna Bell of the NAFO Secretariat (Dartmouth, NS, Canada) for her work on the edition of this research document.

REFERENCES

- Alpoim, R. and Vargas, J., 2004. Length-weight relationships of the Portuguese commercial catches in NAFO, 1998-2003. *NAFO SCR Doc. 04/40* Ser. No N4991, 10pp.
- Ávila de Melo, A.M., Alpoim, R. and Saborido-Rey, F., 2003. An assessment of beaked redfish (*S. mentella* and *S. fasciatus*) in NAFO Div. 3M. *NAFO SCR Doc. 03/45* Ser. No N4863, 72p.
- Ávila de Melo, A.M., Power, D. and Alpoim, R., 2005. An assessment of the status of the redfish resource in NAFO Divisions 3LN. *NAFO SCR Doc. 05/52*, Serial No. N5138, 19 pp.
- Ávila de Melo, A.M. and Alpoim, R., 2010. The 2nd take of 2008 assessment of redfish in NAFO Divisions 3LN: going further on the exploratory analysis of ASPIC formulations. *NAFO SCR Doc. 10/29*, Serial No. N5787, 21 pp.
- Ávila de Melo, A.M., Alpoim, R. and González-Troncoso, D., 2012. An ASPIC Based Assessment of Redfish (*S. mentella* and *S. fasciatus*) in NAFO Divisions 3LN (*can a surplus production model cope with bumpy survey data?*). *NAFO SCR Doc. 12/032*, Serial No. N6059, 63 pp.
- Ávila de Melo, A.M., Brites, N., Alpoim, R. and González-Troncoso, D., 2014. An ASPIC Based Assessment of Redfish (*S. mentella* and *S. fasciatus*) in NAFO Divisions 3LN (*assuming that the highest apparently sustained historical average level of catch is a sound proxy to MSY*). *NAFO SCR Doc. 14/22*, Serial No. N6317, 78 pp.
- Ávila de Melo, A.M., Brites, N., Alpoim, R., González-Troncoso, D., and Fomin, K., 2016. A Revised Update of the 2014 ASPIC Assessment of Redfish (*S. mentella* and *S. fasciatus*) in Divisions 3LN (*how the the stock is coping with the actual Management Strategy and its likely impact on the next coming years*). *NAFO SCR Doc. 16/33*, Serial No. N6576, 69 pp.
- Brandão, A. and Butterworth, D., 2018. Wald-Wolfowitz runs test for independence in time series of residuals for recent 3LN redfish abundance indices. *SC WP 18/016*, 5pp.
- COM-SC CESAG, 2018. Application of the CDAG method to all NAFO managed stocks for 2017. *COM-SC CESAG-WP 18-01 (Rev.2)*, 8pp.
- DFO. 2008. Advice on the stock definition of redfish (*Sebastes fasciatus* and *S. mentella*) in Units 1 and 2. *DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2008/026*.
- Efron, B., Gong, G., 1983. A Leisurely Look at the Bootstrap, the Jackknife, and Cross-Validation. *The American Statistician*, Vol. 37, No. 1, (Feb., 1983), pp. 36-48.
- Fomin, K. and Pochtar, M., 2018. Russian research report for 2017. *NAFO SCS Doc. 18/13*, Serial No. N6824, 13 pp.
- Gavaris, S., 1980. Use of a multiplicative model to estimate catch rate and effort from commercial data. *Canadian Journal of Fisheries and Aquatic Science* 37, p. 2272-2275.
- González-Costas, F., Ramilo, G., Román, E., Lorenzo, J., Gago, A., González-Troncoso, D., Sacau, M., 2018. Spanish Research Report for 2017. *NAFO SCS Doc. 18/07*. Serial Number N6787, 50 pp.
- González-Troncoso, D., Gago A. and Ramilo, L., 2018. Yellowtail flounder, redfish (*Sebastes spp.*) and witch flounder indices from the Spanish Survey conducted in Divisions 3NO of the NAFO Regulatory Area. *NAFO SCR Doc. 18/012*, Serial No. N6796. 34pp
- Mainly, B. F. J., 1997. Randomization, bootstrap and Monte Carlo methods in biology (second edition). *Texts in Statistical Science*, Chapman & Hall, London, 399pp.

- NAFO, 2004. Report of the NAFO Study Group on Limit Reference Points Lorient, France, 15-20 April, 2004. *NAFO SCS Doc.* 04/12, Serial Number N4980, 72 pp.
- NAFO, 2009. Northwest Atlantic Fisheries Organization Scientific Council Reports 2007. Dartmouth, Nova Scotia, Canada, 325 pp.
- Power, D. and Vaskov, A.A., 1992. Abundance and biomass estimates of redfish (*S. mentella*) in Div. 3LN from Russian groundfish surveys from 1984-91. *NAFO SCR Doc.* 92/59. Serial No. N2113. 9 pp.
- Power, D., 1997. Redfish in NAFO Divisions 3LN. *NAFO SCR Doc.* 97/64, Serial No. N2898. 37 pp.
- Power, D., 2001. An assessment of the status of the redfish resource in NAFO Divisions 3LN. *NAFO SCR Doc.* 01/62, Serial No. N4440, 22 pp.
- Prager, M.H., 1994. A suite of extensions to non-equilibrium surplus-production model. *Fish. Bull. U.S.*, 92: 374-389, 29pp.
- Prager, M.H., 2015. User's Guide for ASPIC Suite, version 7: A Stock Production Model Incorporating Covariates, and auxiliary programs. Michael H. Prager — Prager Consulting Portland, Oregon, USA, 33pp.
- Rideout, R.M., 2016. Spatial distribution patterns of NAFO demersal fish stocks based on data from the Canadian multi-species surveys of Divisions 2HJ3KLNO. *NAFO SCR Doc.* 16/20, Serial Number N6563, 45p.
- Rideout, R.M. and Ings, D.W., 2018. Temporal and spatial coverage of Canadian (Newfoundland and Labrador region) Spring and Autumn multi-species RV bottom trawl surveys, with an emphasis on surveys conducted in 2017. *NAFO SCR Doc.* 18/017, Serial Number N6801, 36pp.
- Román, E., D. González-Troncoso and Alvarez, M., 2018. Results for the Atlantic cod, roughhead grenadier, redfish, thorny skate and black dogfish of the Spanish Survey in the NAFO Div. 3L for the period 2003-2017. *NAFO SCR Doc.* 18/018, Serial No. N6802, 68pp.
- Schaefer, M.B., 1954. Some aspects of the dynamics of populations important to management of the commercial marine fisheries. *Bull. Inter-Am. Trop. Tuna Comm.* 1(2): p. 27-56.
- Valentin, A., Power, D., Sévigny, J-M., 2015. Understanding recruitment patterns of historically strong juvenile year classes in redfish (*Sebastes* spp.): The importance of species identity, population structure, and juvenile migration. *Canadian Journal of Fisheries and Aquatic Sciences* 72(5): p. 774-784.
- Vargas, J., Alpoim, R., Santos, E. and Ávila de Melo, A.M., 2018. Portuguese Research Report for 2018. *NAFO SCS Doc.* 18/08, Ser. No N6788, 30 pp.

Table 1. Summary of catch and TAC's of redfish in Div. 3LN estimated from various sources

YEAR	3L	3N	TOTAL	TAC
1959	34107	10478	44585	
1960	10015	16547	26562	
1961	8349	14826	23175	
1962	3425	18009	21439	a
1963	8191	12906	27362	a
1964	3898	4206	10261	a
1965	18772	4694	23466	
1966	6927	10047	16974	
1967	7684	19504	27188	
1968	2378	15265	17660	a
1969	2344	22356	24750	a
1970	1029	13359	14419	a
1971	10043	24310	34370	a
1972	3095	25838	28933	
1973	4709	28588	33297	
1974	11419	10867	22286	28000
1975	3838	14033	17871	20000
1976	15971	4541	20513	20000
1977	13452	3064	16516	16000
1978	6318	5725	12043	16000
1979	5584	8483	14067	18000
1980	4367	11663	16030	25000
1981	9407	14873	24280	25000
1982	7870	13677	21547	25000
1983	8657	11090	19747	25000
1984	2696	12065	14761	25000
1985	3677	16880	20557	25000
1986	27833	14972	42805	25000
1987	30342	40949	79031	25000 b
1988	22317	23049	53266	25000 b
1989	18947	12902	33649	25000 b
1990	15538	9217	29105	25000 b
1991	8892	12723	25815	14000 b
1992	4630	10153	27283	14000 b
1993	5897	9077	21308	14000 bc
1994	379	2274	5741	14000 bc
1995	292	1697	1989	14000
1996	112	339	451	11000
1997	151	479	630	11000
1998	494	405	899	0
1999	518	1318	2318	0 b
2000	657	819	3141	0 bc
2001	653	245	1442	0 b
2002	651	327	1216	0 b
2003	584	751	1334	0
2004	401	236	637	0
2005	581	78	659	0
2006	53	444	496	0
2007	118	1546	1664	0
2008	220	377	597	0
2009	57	994	1051	0
2010	260	3688	4120	3500
2011	2418	1254	3672	6000
2012	2781	1535	4316	6000
2013	4446	1786	6232	6500 d
2014	4245	1450	5695	6500 d
2015	8620	1320	9940	10400 d
2016	6652	1805	8457	10400 d
2017	7790	4026	11816	14200 e
2018				14200

^a Includes catch that could not be identified by division

^b Includes estimates of unreported catches

^c Catch could not be precisely estimate due to discrepancies in figures from available sources: average of the range of the different catch estimates.

^d STATLANT 21A catches as updated on June 2018.

^e Catch estimates with CEDAG method (COM-SC CESAG, 2018)



Table 2. Redfish STATLANT catch and predicted effort for Div. 3L and Div. 3N, 1959-1994 (Power,1997).

	3L		3N		3LN		3LN CPUE annual
	STATLANT Catch	Predicted EFFORT	STATLANT Catch	Predicted EFFORT	STATLANT Catch	Predicted EFFORT	
1959	34107	22604	10478	8659	44585	31263	1.426
1960	10015	5690	16547	10892	26562	16582	1.602
1961	8349	3610	14826	10049	23175	13659	1.697
1962	3425	2049	18009	11090	21434	13139	1.631
1963	8191	3973	12906	8958	21097	12931	1.632
1964	3898	1491	4206	2981	8104	4472	1.812
1965	18772	8190	4694	2551	23466	10741	2.185
1966	6927	4615	10047	4915	16974	9530	1.781
1967	7684	3793	19504	10569	27188	14362	1.893
1968	2378	1446	15265	17684	17643	19130	0.922
1969	2344	1354	22356	17109	24700	18463	1.338
1970	1029	499	13359	10026	14388	10525	1.367
1971	10043	5207	24310	20320	34353	25527	1.346
1972	3095	1877	25838	18982	28933	20859	1.387
1973	4709	2078	28588	18186	33297	20264	1.643
1974	11419	11907	10867	5374	22286	17281	1.290
1975	3838	2443	14033	8265	17871	10708	1.669
1976	15971	11335	4541	4537	20512	15872	1.292
1977	13452	10461	3064	2738	16516	13199	1.251
1978	6318	5961	5725	4925	12043	10886	1.106
1979	5584	3517	8483	6176	14067	9693	1.451
1980	4367	2873	11663	6229	16030	9102	1.761
1981	9407	6020	14873	9216	24280	15236	1.594
1982	7870	4812	13677	8160	21547	12972	1.661
1983	8657	4960	11090	7734	19747	12694	1.556
1984	2696	1804	12065	12263	14761	14067	1.049
1985	3677	2104	16880	16858	20557	18962	1.084
1986	27833	15247	14972	15057	42805	30304	1.413
1987	34212	22369	44819	29517	79031	51886	1.523
1988	26267	19629	26999	24453	53266	44082	1.208
1989	19847	10567	13802	14884	33649	25451	1.322
1990	17713	16774	11392	18513	29105	35287	0.825
1991	8892	12329	12723	20052	21615	32381	0.668
1992	4630	2452	10153	13755	14783	16207	0.912
1993	5897	1576	9077	17116	14974	18692	0.801
1994	379	410	2274	2900	2653	3310	0.802

Table 3a. Length composition (absolute frequencies in '000s) of the 3LN redfish commercial catch and by-catch, 1990-2017.

Length	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017			
10																				0.1			1	2	0.3		23	4			
11														0.03						0.2	3	13	3	5	18	13	60	48			
12														0.03					0.2	10	30	21	14	17	37	202	96				
13		12												1					0.1	23	49	69	30	133	94	198	223				
14		6												1					0.1	33	154	117	52	96	180	358	222				
15		28	28											5					2.6	51	268	224	78	166	212	459	316	11			
16		73	103	9								1	0.3	8					14	362	506	281	95	420	330	566	394	11			
17		199	394	28		2				0.3	1	2	1	21	1	2	34		80	670	768	371	139	460	400	795	492	61			
18		286	1034	412		5		0.01	1		1	1	1	44	2	4	65	0.3	103	747	935	430	201	793	576	1321	767	265			
19		445	2157	1291	5	6	3	1	0.3	2	16	4	4	3	90	6	9	99	43	96	495	1383	766	426	1313	907	2032	962	252		
20		720	3313	2375		16	14	4	2	13	47	6	18	14	151	15	11	182	143	43	357	1737	1253	890	2221	1880	3529	2093	486		
21		1309	3780	2943	235	287	9		11	57	80	10	52	41	218	28	13	300	77	133	428	2284	1942	1564	3518	2891	5416	4042	797		
22		2081	4922	3600	714	683	65	6	17	151	150	26	102	81	269	35	11	347	149	239	475	3084	2545	2235	4913	3865	8176	6959	1371		
23		3212	7340	4358	1141	594	64	17	34	277	128	46	118	101	277	41	16	340	212	303	552	2188	1959	1778	3468	3425	8021	7758	2354		
24		4164	7575	5552	2565	708	99	9	64	296	120	85	114	132	258	54	35	210	170	253	311	1183	1099	1231	2335	2125	4873	5939	3260		
25		5216	6944	4981	5237	944	100	9	98	248	178	195	114	154	261	85	61	147	221	224	268	831	593	992	1605	1510	3485	4980	3872		
26		5560	5981	5145	5115	1297	277	12	118	221	318	364	126	204	309	157	138	111	206	138	271	769	450	746	1719	1118	2558	3693	4295		
27		5410	6197	4579	5433	1404	330	35	144	218	555	546	170	248	324	190	181	99	134	81	193	584	371	670	1156	990	1534	2731	4366		
28		5217	5322	4063	5004	1182	300	75	114	173	712	943	188	289	286	184	201	88	521	32	194	580	462	646	1270	1066	1602	2309	3825		
29		4712	3354	4637	4437	1188	263	76	114	154	673	1003	179	289	245	184	223	62	425	42	140	490	445	620	642	916	1477	1514	2810		
30		4751	4043	3911	3283	1011	310	182	114	120	520	1027	236	294	225	178	176	60	368	44	96	416	434	714	691	1159	1757	1326	2216		
31		4551	2695	3711	2964	912	313	197	154	129	413	564	289	295	204	107	109	35	335	31	64	296	608	691	864	975	1415	772	1631		
32		3943	2478	2187	2313	944	309	98	146	119	434	315	303	276	189	108	91	28	594	37	49	276	674	641	598	862	1207	615	1366		
33		3062	1582	1355	2291	596	226	67	131	110	383	237	298	216	196	95	83	19	316	58	40	242	535	681	517	673	843	412	1582		
34		2737	1179	1569	1527	526	189	30	71	66	268	217	218	132	149	73	71	17	252	83	37	215	223	392	304	438	539	203	1083		
35		2100	928	1604	1059	363	182	35	24	19	141	129	212	83	112	51	63	10	124	62	11	208	170	265	141	345	511	158	885		
36		1681	831	1895	923	202	106	23	19	18	89	60	121	37	62	36	56	5	110	39	13	137	85	157	167	222	315	90	524		
37		1416	580	1571	766	196	160	7	14	11	82	78	82	18	41	17	31	2	4	31	2	70	46	101	70	123	153	43	262		
38		1128	482	1303	807	158	171	5	10	8	51	50	55	11	22	10	15	1	2	12	1	69	54	105	48	122	91	33	132		
39		729	363	1114	489	124	100	11	3	3	37	47	30	3	14	9	8	0.01	23	9	2	32	19	40	16	57	25	21	36		
40		458	292	790	505	69	144	2	4	3	23	23	18	2	7	5	8	0.3	22	1	0.4	17	17	33	5	48	29	5	28		
41		321	188	558	320	49	63	3	1	2	19	12	10	1	2	2	4	0.003	0.4	1	0.5	7	10	20	6	26	33	5	0.01		
42		255	117	420	306	23	1	1	0.1	13	15	7	2	3	1	2	0.003	0.2	0	1	3	5	11	4	10	4	2	0.003			
43		227	68	203	137	15	3	2	2	0.1	3	9	4	2	2	2	6	0.5	3	0.02	0.1	5	7	4	2	4	2				
44		157	83	85	175	7	3	2	1	1	3	1	3	1	2	1	3	0.1	0.05	0.2			2			1	3	0.4			
45		84	33	76	107	1	3	2	0.1		2	1	1	0.1	1	1			1	0.3								1	1		
46		58	8	32	9	3		0.1	0.0	0.2		1	1	2	0.2	1	0.3							0.4				1	0.1		
47		24	9	47	0.2						0.5	0.2		0.04	1	2												1			
48		11	2	8				0.1								0.04															
49		6		1	0.1																										
50																															
51		1	25		2									0.3												1			0.2		
52		2																													
53		1																												0.2	
54		2																												0.01	
no ('000)	66410	74421	66375	47918	13517	3815	910	1411	2422	5457	6020	3076	2929	3999	1681	1632	2295	4454	2199	5901	19825	16289	16274	29702	27598	53616	49232	37792			
weight (tons)	29105	25815	27283	21308	5741	1989	451	630	899	2318	2617	1442	1216	1334	637	659	497	1664	597	1051	3948	3672	4316	6232	5695	10467	8457	11816			
mean weight (g)	438	347	411	445	425	521	496	446	371	425	435	469	415	334	379	404	217	365	271	178	199	225	265	210	206	195	172	313			
mean length	29.3	26.6	28.4	29.6	29.1	31.6	31.2	29.8	27.4	29.9	30.1	30.8	29.5	27.5	29.5	30.1	23.9	29.4	25.4	22.0	23.4	24.4	25.9	24.2	25.0	24.3	24.5	27.8			
length anomalies	1.70	-1.0	0.8	2.0	1.5	4.0	3.6	2.2	-0.2	2.3	2.5	3.2	1.9	-0.1	1.8	2.5	-3.7	1.8	-2.2	-5.6	-4.3	-3.3	-1.7	-3.5	-2.7	-3.4	-3.1	0.3			
%lengths <20cm	1.6%	5.0%	2.6%	0.0%	0.1%	0.2%	0.1%	0.0%	0.1%	0.3%	0.1%	0.2%	0.2%	4.2%	0.5%	0.9%	10.1%	1.0%	13.5%	40.6%	20.7%	14.0%	6.4%	11.5%	10.0%	11.2%	7.2%	1.6%			



Table 3b. length weight relationships from 3LN *Sebastes* sp. Portuguese commercial sampling data used in the computation of 3LN catch at length (Alpoim and Vargas, 2004; Vargas et al, 2005-2018)

<i>Sebastes</i> sp.	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004
a	0.1115	0.1115	0.1115	0.1115	0.1115	0.1115	0.1115	0.1115	0.1115	0.0689	0.0979	0.0769	0.0447	0.0095	0.0208
b	2.4353	2.4353	2.4353	2.4353	2.4353	2.4353	2.4353	2.4353	2.4353	2.5588	2.4602	2.5298	2.6885	3.1279	2.8851

Table 3b (cont.)

<i>Sebastes</i> sp.	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017
a	0.0208	0.0611	0.0207	0.0207	0.0207	0.0214	0.0214	0.0214	0.0360	0.0462	0.0116	0.0290	0.0279
b	2.8851	2.5597	2.8946	2.8946	2.8946	2.8659	2.8659	2.8659	2.6998	2.5880	3.0190	2.7011	2.7896

Table 4 Survey biomass ('000 t) from stratified bottom trawl surveys on Div. 3L and Div.3N included in the 2018 ASPIC framework, survey female SSB from spring and autumn Canadian surveys on Div. 3LN (1991-2017)

	Canadian				Russian	Div. 3L			Spanish	
	Div. 3LN		Div. 3LN		Div. 3LN	Div. 3L	Div. 3L	Div. 3L	Div. 3N	Div. 3L
	I2 _{springcomb}	I2 _{springSSB}	I3 _{autumncomb}	I3 _{autumnSSB}	I4 _{Powercomb}	I5 _{winter}	I6 _{summer}	I7 _{autumn}	I8 _{spring}	I9 _{summer}
1978							311.2			
1979							227.8			
1980										
1981							261.4			
1982										
1983										
1984					215.9		277.7			
1985					94.0	90.2	161.0	98.2		
1986					63.0	36.6		17.1		
1987					70.3					
1988					44.9					
1989					12.3					
1990					8.4	18.2	92.8	20.7		
1991	10.6	1.5	37.9	4.7	18.7		37.6			
1992	10.1	1.8	136.4	15.4						
1993	22.6	4.3	19.2	3.6			20.8			
1994	4.2	0.6	31.8	5.9						
1995	5.9	0.8	90.7	15.9					46.1	
1996	22.8	11.6	16.0	2.6					6.6	
1997	14.9	1.8	70.7	10.7					4.8	
1998	59.4	11.5	112.2	14.5					22.5	
1999	61.5	15.2	72.0	12.6					46.5	
2000	87.8	17.3	100.5	16.6					68.9	
2001	41.6	7.0	132.6	13.8					53.9	
2002	31.0	5.8	50.1	9.4					7.6	
2003	27.7	3.7	71.9	9.6					11.0	
2004	79.6	26.2	49.9	11.4					27.0	
2005	66.5	8.8	58.6	11.2					146.9	
2006			91.9	12.9					87.8	70.1
2007	218.8	39.4	124.8	16.8					87.6	31.4
2008	144.0	23.4	198.5	27.4					68.1	75.6
2009	183.4	20.7	246.7	29.6					735.7	103.7
2010	165.3	21.5	461.5	55.5					359.5	266.8
2011	173.7	22.2	562.3	64.1					418.3	170.6
2012	322.0	45.5	596.0	89.7					265.2	481.5
2013	271.5	48.1	288.8	41.1					429.5	235.2
2014	271.7	38.3							178.1	216.4
2015	480.6	60.1	425.9	64.6					523.5	130.4
2016	654.2		215.2						117.3	98.8
2017			192.0						265.9	56.6

Table 5a. 3LN spring survey abundance at length, 1991-2016 (thousands).

Length	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006 ⁽¹⁾	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016			
4															40										25.0	19.4			
5												62		31				416	46	258			146	182.8		330.6			
6						466		20	16	185	109	170	293	804	108		154	1966	479	137	559	41	695	1349.6	945.9	1246.8			
7						228		40	656	795	1511	472	2057	2400	540		3452	2942	974	562	781	1055	1772	968.4	8495.2	2957.5			
8						149	685	8	3280	378	1302	1073	1682	1236	950		9327	3135	954	936	858	1039	1106	826.4	5173.1	13159.1			
9	849					298	360	39	5877	89	483	1526	1524	2209	2891		2625	3381	371	1361	1073	1596	1791	1038.8	673.4	19021.3			
10	1149					296	251	113	1343	166	240	2518	1197	4107	4892		886	4258	994	2423	1342	1921	1896	1265.7	248.8	6233.8			
11	798	381	122	355		478	730	533	309	403	116	1085	417	2911	7296		1683	5317	1695	2902	2464	2178	2225	1377.0	812.1	2529.4			
12	558	2988	1304	540		806	722	455	430	191	451	1645	1448	1653	8756		2296	2432	3642	2871	1701	2852	1761	1609.7	1127.3	1194.9			
13	2524	7925	2396	500	108	920	540	172	517	412	345	838	1101	1330	9684		1908	1286	16098	2256	2458	2581	2303	1708.2	614.1	468.6			
14	322	5192	5646	536	272	413	1871	561	369	353	1073	517	1278	639	7710		1928	5396	12659	4892	3568	3658	1807	1526.8	332.9	563.0			
15	699	2862	11059	1329	278	716	1859	896	175	2458	1738	766	2609	1235	7437		3631	3841	11260	8481	4481	5998	4327	1149.2	284.6	1692.9			
16	2250	382	13647	1790	966	846	1126	1506	774	2199	1681	1371	3559	1335	7357		5993	15866	75231	14345	8907	9617	2467	1438.3	927.6	1116.0			
17	3865	419	8796	3123	2847	1588	1201	2046	703	2157	3337	2580	6189	2764	8647		14186	45719	197691	26140	17787	52512	3940	4388.2	124.8	2833.3			
18	6226	1111	2719	3084	4285	4356	1860	2121	3455	3525	5257	6444	8643	3668	16472		24586	77478	325440	108928	56811	96670	12330	9410.2	13441.8	2199.3			
19	7749	2480	2474	1403	5014	9476	3280	2849	2988	7017	8267	8161	15473	8995	31506		26943	50553	310284	219289	115709	194730	20116	63952.3	81531.5	2881.0			
20	4522	2574	3839	829	2703	10910	4708	9472	5379	13198	9589	11326	21089	11905	33702		26003	48021	164370	234599	144823	289679	103824	187958.9	388738.0	9036.5			
21	3482	3559	5754	922	1815	12119	6367	24848	16817	22002	14393	13958	23750	16956	33182		43665	49072	92564	178663	221969	398075	210629	323755.1	672239.2	24063.9			
22	5148	1690	5301	783	1335	13844	7008	34265	31067	42769	15551	14932	19290	16584	30967		68143	78864	60965	74436	128066	315995	237958	340215.9	734832.5	57932.1			
23	7253	1732	5708	1181	1257	16629	8191	31121	38232	53557	15590	15882	15120	20423	30644		87375	88837	65881	72484	85379	190758	182210	246336.0	479788.0	90422.7			
24	6187	2721	4756	1498	1359	12502	10669	28376	45394	53956	14839	16034	10814	17004	28561		96975	87288	76912	66508	62237	140353	147932	176272.9	291257.6	85383.6			
25	3366	2865	3398	1748	1004	8318	9469	21275	21482	34350	10166	12606	8036	14657	24305		78847	61337	55777	61001	46547	94274	114790	78066.6	107567.1	75116.0			
26	1963	3250	3701	1564	1600	5649	7757	19512	30227	27946	10041	11224	6889	24397	18438		90996	54230	30388	38296	44947	63492	62515	62684.3	41933.0	91635.4			
27	1426	2411	4478	1057	1693	5106	4047	16075	21654	21918	11330	8887	5102	38936	20027		81118	34946	17043	18645	37756	45182	45089	33607.8	24506.4	94097.4			
28	953	1834	3283	803	1437	4901	2760	12716	15663	13775	10217	7496	3552	43216	15249		36969	28227	14167	18908	32300	32808	40858	24603.8	23672.8	136262.8			
29	1038	1506	2876	731	1154	4264	1871	9632	14331	15612	10385	6419	2778	24426	11907		38023	19445	13076	11302	24988	33669	28454	15455.4	15087.4	155044.6			
30	607	1048	2606	482	721	3323	1797	6120	6698	14650	9523	3741	2701	18145	8832		30266	12314	8659	10701	16753	26246	25858	11810.6	23284.0	143012.3			
31	534	1014	2969	318	474	2231	1354	6513	5732	12804	10450	3588	2176	13713	5769		30137	10571	6011	4704	10141	18307	15530	9541.1	17648.0	156646.4			
32	417	809	3087	244	548	1564	991	6157	4322	10277	8884	2235	2356	9706	3036		21974	7018	4096	4110	8774	14817	14990	7287.5	16681.6	107686.1			
33	369	825	2621	138	264	762	640	5687	3259	6538	5183	1382	1972	3487	2012		9163	7747	3448	2908	4925	5029	9922	4668.4	13284.1	108920.8			
34	399	540	2161	156	144	337	438	3287	2024	5043	3035	996	1009	5391	1617		8158	4329	2327	2565	2999	4685	9065	4325.9	11634.4	71849.5			
35	251	544	1502	109	105	163	160	967	877	3301	990	455	640	2249	832		7223	1860	1609	1804	1662	1757	10028	3834.7	6785.6	75698.1			
36	190	366	880	135	113	105	77	660	534	895	296	227	227	476	592		9422	1361	839	1035	1367	1276	5801	3450.8	4791.6	52144.4			
37	222	216	696	127	151	118	42	402	273	709	378	93	82	877	222		1894	786	312	394	788	958	3558	1381.3	1913.3	50188.8			
38	159	219	669	82	101	28	88	82	102	396	116	43	35	75	112		1945	386	235	197	848	548	1831	769.8	1486.1	21874.4			
39	130	300	726	31	70	55	4	82	67	186	155	59	35	43	86		193	325	90	31	224	897	1406	205.5	657.2	3972.4			
40	118	220	483	46	62	28	0	216	79	183	23		94	23	12		115	189	55	54	71	167	1275	174.5	199.0	8477.5			
41	45	77	371	0	15	15	0	15	51	16		15		4	15		59							119	81	865	10.7	80.7	5888.5
42	88	85	215	9	46	4	0	20	66	47	63	15		15	8		24	53	50					15	1637	1256.2	28.7	2802.6	
43	69	85	83	49	27	35	15	201	0	31	28		15	15			8				76			48	463	552.2	149.0		
44	45	77	189	29	31		31	12	27	31	28						23	60						97	12	11.7	0.0		
45	57	62					15	15	15		31	31	15														417	3050.3	
46		46	51				15	46																			401	0.0	
47		4	20		15		15																					0.0	
48	11	31	31																	15								0.0	
49		31																										0.0	
50																												0.0	
51																												0.0	
52																												0.0	
53																												0.0	
abundance (millions)	66.0	54.5	110.6	26.3	32.0	124.1	83.0	249.1	285.3	374.5	187.2	160.5	175.2	318.1	384.4		868.3	821.3	1576.7	1199.2	1096.3	2055.7	1337.3	1629.9	2993.4	1689.8			
mean length (cm)	21.6	21.6	22.6	21.5	22.7	23.4	23.5	25.1	24.7	25.3	25.2	23.5	22.0	25.7	22.2		25.1	22.9	20.3	21.6	22.6	22.5	24.1	23.1	22.8	29.3			
length anomalies (cm)	-1.5	-1.5	-0.5	-1.6	-0.4	0.3	0.4	2.0	1.6	2.2	2.1	0.4	-1.1	2.6	-0.9		2.0	-0.3	-2.8	-1.5	-0.5	-0.7	1.0	0.0	-0.3	5.9			

(1) Survey data only from Division 3L, no survey on Div. 3N



Table 5b. 3LN autumn survey abundance at length, 1991-2017 (thousands).

Length	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014 ⁽¹⁾	2015	2016	2017								
4																												126.7							
5				15	240	56	86	17		117	445	232	1090	34	0	84	234	31	96	1384	57	263	417			712.0	420.6	1580.9							
6					256	359	330	0	251	481	937	915	2427	85	133	1418	512	641	624	1110	318	1405	4539			2832.0	735.5	15791.8							
7	203				138	88	395	39	50	673	755	873	2185	61	162	1831	2222	2359	318	1405	727	926	2902			35135.9	499.2	18652.0							
8	1298				111	72	386	47	37	602	2114	1614	2714	620	908	466	2924	2745	871	3377	878	2332	4879			86942.0	6969.9	2504.7							
9	1236				241	146	468	252	421	620	3146	1275	2095	1280	2236	829	8313	2359	3452	3788	2878	6976	5697			33864.2	22014.0	1713.6							
10	7263		93	31	292	250	306	214	171	388	4323	1129	2855	1719	1574	1458	8498	4100	9932	4676	2265	9925	5439			5654.4	50159.7	4021.2							
11	22235	371	63	31	213	349	249	203	402	215	2846	2840	1839	1046	3957	1709	7527	5543	5206	6612	1841	8539	8460			2408.8	12040.5	12142.6							
12	62419	62	372		241	106	175	275	786	202	1266	2255	1123	1131	9942	3083	6352	4861	4025	7947	1925	6516	12917			1684.6	7325.0	19385.0							
13	109337	3189	457	335	304	274	366	596	868	320	1056	2072	1488	1436	11090	3970	5871	27297	9473	10315	3250	5463	12133			1763.4	1940.6	21301.3							
14	33876	27936	1775	551	513	1419	728	912	2472	587	445	2545	1451	1015	10309	8256	9046	28768	20311	11133	4187	6377	15089			2478.8	969.5	19948.6							
15	14030	104298	1333	2362	967	722	1104	1768	1548	3635	407	1884	1929	538	8461	13286	21881	23691	17750	8561	8268	5234	28083			3107.3	1493.4	9182.6							
16	7809	113966	3259	3697	1611	919	1405	4159	717	4671	11018	2159	8240	879	6083	20912	40243	116528	35720	12943	14606	3150	12974			3042.9	1133.6	6841.0							
17	7860	106448	5283	12985	9645	825	1848	8155	1144	5480	31421	4694	15193	1984	5713	27177	51164	228751	138765	18474	46427	9778	15860			4207.6	1105.3	5510.0							
18	16191	95896	8707	28684	37932	2227	2095	12225	3185	7035	57695	9082	25813	5468	7248	23009	43358	221311	396982	77810	103647	24081	34943			4920.2	1135.9	3819.8							
19	32214	71577	6425	29295	72192	5062	8438	17373	6536	11926	74228	13661	38672	8222	10928	24342	35091	141084	421539	269160	432556	116751	92467			28085.8	3178.0	7951.4							
20	27189	113846	3906	15292	78316	6479	21672	46005	9069	31680	80538	12568	45262	9790	15982	26793	45870	78263	279787	459453	996936	315398	109915			96232.7	16383.0	18706.1							
21	15810	148628	5306	7701	43397	6621	47562	88726	15347	50184	65575	16481	42849	13134	25645	36447	55971	63995	138841	499979	1138226	664907	283174			285478.4	71853.2	43755.6							
22	7915	153395	6375	5119	27652	6123	52500	124662	23121	66781	130029	20168	39683	19332	23890	49628	61550	65482	67350	303473	587045	653151	454374			521322.4	156745.5	77215.2							
23	6139	89704	6578	6494	20117	6743	44777	92991	29000	60123	118427	23529	39374	16732	29785	71774	84212	89011	53177	261470	300782	501477	309498			455998.7	203673.6	97048.5							
24	8377	28658	5164	5456	10296	4864	31865	56410	26969	52986	85149	25353	31785	15458	20362	67361	81986	80398	65248	260734	126712	314858	193667			350489.1	151520.6	80128.0							
25	8943	14222	3947	6808	12898	4429	24356	30123	29819	50534	64519	21326	21398	13066	15824	34947	57418	66252	46806	165444	97731	203720	122411			221208.4	87513.8	59080.0							
26	6602	13410	4120	8670	8517	4370	21375	23090	27515	40188	39693	19872	18032	10432	12713	32335	39981	49866	39922	120859	82802	152183	64144			115599.5	64421.9	53400.7							
27	4022	14699	4361	7830	17364	2890	21141	20596	25585	21851	33743	16470	17605	9397	10857	19109	26128	48823	34957	95155	49339	135137	32163			95099.7	25728.2	46878.4							
28	3776	8768	4240	8402	17495	2707	14031	18336	24801	17424	20396	10503	13962	12135	12471	11651	19087	37469	24861	72543	35075	76038	23430			62113.2	16855.5	30698.3							
29	2526	4855	3503	7625	16330	2678	8032	13397	16323	16387	14957	7230	7798	13950	12659	10147	13206	21724	24372	38007	30904	67575	16618			44304.7	16201.7	24137.1							
30	2110	3340	2765	6195	12717	2242	6138	7942	11346	12127	11093	5122	4910	12267	9865	7475	7643	18374	14245	26788	35523	46137	14071			31552.2	12742.7	19442.2							
31	1960	3229	1949	4553	16297	3409	4994	6250	7641	10199	9147	5109	3755	9066	7347	9531	6404	11854	10895	15934	17230	29841	8793			35299.7	12981.8	15994.8							
32	1314	2389	1901	2709	10628	2210	4035	5730	6315	7165	5261	4608	3523	6787	5214	7469	4180	6793	7953	14869	11668	28059	8562			5060.8	13155.4	18581.4							
33	1212	3299	1671	1603	7262	1220	2107	3878	5642	5026	4354	3862	3360	4636	4905	4870	3623	6389	6675	9280	4838	18841	5790			8421.2	18683.3	16031.1							
34	1117	1431	1286	916	3447	559	1673	4512	4545	3369	2776	2701	2182	2959	3942	2096	2183	5268	3627	5875	2164	7507	2538			3359.7	15952.4	10590.3							
35	1287	716	1044	610	1966	217	653	2048	3256	1303	1679	1451	1175	1760	2720	1118	1067	2385	2538	1885	1869	4530	2229			1893.2	16453.8	7424.1							
36	1184	595	800	297	1171	118	499	1080	1539	1092	675	560	506	1259	1456	537	416	970	2163	2310	1332	2698	1220			1438.6	7863.2	8629.7							
37	1005	385	460	211	335	64	308	426	339	499	636	325	182	765	1298	444	847	784	1772	1299	817	5530	653			687.7	6419.1	5266.4							
38	1166	401	427	257	998	14	243	247	184	329	282	85	111	392	385	136	275	654	700	1374	138	5691	208			323.4	4493.7	3990.7							
39	787	228	308	274	572	22	176	85	272	227	215	67	115	666	228	55	40	0	300	372	136	1938	257			478.5	2031.0	1221.9							
40	662	93	237	119	75	22	164	17	67	151	180	136		308	60	116	17	391	250	389	0	954	375			62.7	557.5	1376.6							
41	221	124	155	0	20	22	191	40	82	67	81			76	85	61	103			129	208	0	1509			61.6	52.9	553.0							
42	135	77	132	15	24		45			67				232	60			263	505	195	0	630					663.5	218.5							
43	102	31	37	32	32				35	50		4	21							92		45	571				663.5	243.7							
44	128	46	99				42		17	50	4		17															525	274.3	211.6	155.3				
45	46	15	69	15	36		28		17	50	76		17															355	71.4	23.0					
46	24	46			12		14				18	17					16		63	131	46		83				216	125.3	441.6						
47	15	15	15	8			12																						355		9.8				
48												17																		77					
49																				62										77					
50	15																													77					
51																															77		52.9	223.9	
52																																0			
53																																	0		
abundance (millions)		422	1130	89	175	432	71	327	593	288	487	882	245	407	195	297	526	755	1456	1892	2797.7	4205.1	3448.4	1910.9											

Table 6 (HCR option) B/Bmsy and F/Fmsy point estimate and bias corrected trajectories with ASPICprj. 2018, 2018-2021 relative B and F projections under 2014 red 3LN HCR for 2018-2020. (catch 2017 = 14200 t TAC; 2018 and 2019 = 18100 t)

Year	Relative B trajectory and 2018-2021 projection under red 3LN HCR			Relative F trajectory and 2018-2020 projection under red 3LN HCR		
	Point estimate	Approx 80% lower CL	Approx 80% upper CL	Point estimate	Approx 80% lower CL	Approx 80% upper CL
1959	1.395	1.137	2.181	1.604	1.045	1.959
1960	1.257	1.033	1.906	1.021	0.687	1.240
1961	1.221	1.008	1.783	0.910	0.634	1.100
1962	1.204	0.998	1.700	0.850	0.612	1.024
1963	1.198	0.996	1.641	1.106	0.816	1.327
1964	1.160	0.967	1.555	0.412	0.311	0.493
1965	1.213	1.016	1.581	0.928	0.719	1.106
1966	1.196	1.005	1.528	0.671	0.531	0.797
1967	1.212	1.023	1.517	1.085	0.873	1.283
1968	1.175	0.995	1.451	0.711	0.581	0.839
1969	1.189	1.010	1.445	1.001	0.829	1.177
1970	1.166	0.992	1.399	0.581	0.487	0.681
1971	1.197	1.023	1.419	1.412	1.196	1.650
1972	1.123	0.962	1.319	1.252	1.068	1.459
1973	1.079	0.926	1.262	1.517	1.298	1.766
1974	1.013	0.871	1.182	1.051	0.902	1.224
1975	1.006	0.863	1.171	0.839	0.722	0.979
1976	1.023	0.876	1.186	0.954	0.824	1.115
1977	1.026	0.876	1.184	0.758	0.658	0.888
1978	1.049	0.895	1.206	0.534	0.466	0.627
1979	1.097	0.934	1.253	0.601	0.528	0.705
1980	1.132	0.966	1.286	0.667	0.589	0.781
1981	1.156	0.988	1.305	1.009	0.897	1.179
1982	1.136	0.973	1.275	0.905	0.809	1.056
1983	1.132	0.971	1.262	0.829	0.745	0.966
1984	1.136	0.976	1.261	0.610	0.551	0.710
1985	1.167	1.004	1.289	0.839	0.762	0.974
1986	1.166	1.006	1.281	1.845	1.685	2.133
1987	1.048	0.908	1.144	4.275	3.937	4.905
1988	0.736	0.647	0.796	3.987	3.702	4.521
1989	0.548	0.484	0.588	3.217	3.003	3.647
1990	0.452	0.399	0.485	3.387	3.150	3.838
1991	0.370	0.326	0.398	3.727	3.451	4.258
1992	0.293	0.255	0.318	5.397	4.919	6.188
1993	0.195	0.171	0.220	6.776	5.632	7.849
1994	0.112	0.093	0.145	2.523	1.885	3.140
1995	0.105	0.081	0.145	0.854	0.609	1.112
1996	0.118	0.089	0.166	0.166	0.118	0.221
1997	0.142	0.106	0.201	0.192	0.136	0.261
1998	0.171	0.125	0.242	0.229	0.161	0.316
1999	0.205	0.146	0.288	0.502	0.355	0.709
2000	0.236	0.166	0.337	0.593	0.415	0.861
2001	0.269	0.182	0.386	0.235	0.163	0.347
2002	0.317	0.214	0.458	0.168	0.117	0.253
2003	0.375	0.246	0.540	0.156	0.109	0.241
2004	0.440	0.282	0.624	0.063	0.045	0.099
2005	0.518	0.335	0.728	0.056	0.040	0.089
2006	0.605	0.378	0.834	0.036	0.027	0.058
2007	0.701	0.437	0.949	0.106	0.079	0.170
2008	0.797	0.493	1.057	0.033	0.026	0.054
2009	0.904	0.566	1.171	0.052	0.041	0.083
2010	1.010	0.638	1.279	0.186	0.148	0.293
2011	1.100	0.701	1.362	0.153	0.125	0.237
2012	1.190	0.778	1.445	0.167	0.139	0.251
2013	1.273	0.854	1.516	0.227	0.192	0.333
2014	1.341	0.929	1.567	0.197	0.170	0.279
2015	1.407	1.003	1.614	0.350	0.307	0.485
2016	1.443	1.050	1.628	0.275	0.246	0.373
2017	1.486	1.113	1.653	0.376	0.340	0.496
2018	1.507	1.156	1.654	0.448	0.410	0.576
2019	1.514	1.186	1.642	0.572	0.530	0.722
2020	1.501	1.196	1.610	0.577	0.540	0.716
2021	1.489	1.206	1.584			

Table 7 (staus quo TAC option) B/Bmsy and F/Fmsy point estimate and bias corrected trajectories with ASPICprj. 2018 2018-2021 relative B and F projections under status quo catch at 2017 TAC. (catch 2017 = 14200 t TAC; catch 2018 and 2019 = 14200 t)

Year	Relative B trajectory and 2018-2021 projection under catch at 2017 TAC			Relative F trajectory and 2018-2020 projection under catch at 2017 TAC		
	Point estimate	Approx 80% lower CL	Approx 80% upper CL	Point estimate	Approx 80% lower CL	Approx 80% upper CL
1959	1.395	1.137	2.181	1.604	1.045	1.959
1960	1.257	1.033	1.906	1.021	0.687	1.240
1961	1.221	1.008	1.783	0.910	0.634	1.100
1962	1.204	0.998	1.700	0.850	0.612	1.024
1963	1.198	0.996	1.641	1.106	0.816	1.327
1964	1.160	0.967	1.555	0.412	0.311	0.493
1965	1.213	1.016	1.581	0.928	0.719	1.106
1966	1.196	1.005	1.528	0.671	0.531	0.797
1967	1.212	1.023	1.517	1.085	0.873	1.283
1968	1.175	0.995	1.451	0.711	0.581	0.839
1969	1.189	1.010	1.445	1.001	0.829	1.177
1970	1.166	0.992	1.399	0.581	0.487	0.681
1971	1.197	1.023	1.419	1.412	1.196	1.650
1972	1.123	0.962	1.319	1.252	1.068	1.459
1973	1.079	0.926	1.262	1.517	1.298	1.766
1974	1.013	0.871	1.182	1.051	0.902	1.224
1975	1.006	0.863	1.171	0.839	0.722	0.979
1976	1.023	0.876	1.186	0.954	0.824	1.115
1977	1.026	0.876	1.184	0.758	0.658	0.888
1978	1.049	0.895	1.206	0.534	0.466	0.627
1979	1.097	0.934	1.253	0.601	0.528	0.705
1980	1.132	0.966	1.286	0.667	0.589	0.781
1981	1.156	0.988	1.305	1.009	0.897	1.179
1982	1.136	0.973	1.275	0.905	0.809	1.056
1983	1.132	0.971	1.262	0.829	0.745	0.966
1984	1.136	0.976	1.261	0.610	0.551	0.710
1985	1.167	1.004	1.289	0.839	0.762	0.974
1986	1.166	1.006	1.281	1.845	1.685	2.133
1987	1.048	0.908	1.144	4.275	3.937	4.905
1988	0.736	0.647	0.796	3.987	3.702	4.521
1989	0.548	0.484	0.588	3.217	3.003	3.647
1990	0.452	0.399	0.485	3.387	3.150	3.838
1991	0.370	0.326	0.398	3.727	3.451	4.258
1992	0.293	0.255	0.318	5.397	4.919	6.188
1993	0.195	0.171	0.220	6.776	5.632	7.849
1994	0.112	0.093	0.145	2.523	1.885	3.140
1995	0.105	0.081	0.145	0.854	0.609	1.112
1996	0.118	0.089	0.166	0.166	0.118	0.221
1997	0.142	0.106	0.201	0.192	0.136	0.261
1998	0.171	0.125	0.242	0.229	0.161	0.316
1999	0.205	0.146	0.288	0.502	0.355	0.709
2000	0.236	0.166	0.337	0.593	0.415	0.861
2001	0.269	0.182	0.386	0.235	0.163	0.347
2002	0.317	0.214	0.458	0.168	0.117	0.253
2003	0.375	0.246	0.540	0.156	0.109	0.241
2004	0.440	0.282	0.624	0.063	0.045	0.099
2005	0.518	0.335	0.728	0.056	0.040	0.089
2006	0.605	0.378	0.834	0.036	0.027	0.058
2007	0.701	0.437	0.949	0.106	0.079	0.170
2008	0.797	0.493	1.057	0.033	0.026	0.054
2009	0.904	0.566	1.171	0.052	0.041	0.083
2010	1.010	0.638	1.279	0.186	0.148	0.293
2011	1.100	0.701	1.362	0.153	0.125	0.237
2012	1.190	0.778	1.445	0.167	0.139	0.251
2013	1.273	0.854	1.516	0.227	0.192	0.333
2014	1.341	0.929	1.567	0.197	0.170	0.279
2015	1.407	1.003	1.614	0.350	0.307	0.485
2016	1.443	1.050	1.628	0.275	0.246	0.373
2017	1.486	1.113	1.653	0.376	0.340	0.496
2018	1.507	1.156	1.654	0.448	0.410	0.576
2019	1.514	1.186	1.642	0.446	0.413	0.562
2020	1.521	1.214	1.632	0.444	0.416	0.551
2021	1.526	1.238	1.623			

Table 8a. B/Bmsy point estimate and lower bias corrected 80% CL with ASPICP. 2018-2021 relative B projection under red 3LN HCR and status quo 2017 TAC.

Year	Relative B trajectory and 2018-2021 projection under red 3LN HCR		Relative B trajectory and 2018-2021 projection under status quo 2017 TAC	
	Point estimate	Approx 80% lower CL	Point estimate	Approx bias corrected 80% lower CL
2017	1.486	1.113	1.486	1.113
2018	1.507	1.156	1.507	1.156
2019	1.514	1.186	1.514	1.186
2020	1.501	1.196	1.521	1.214
2021	1.489	1.206	1.526	1.238

Table 8b. F/Fmsy point estimate and upper bias corrected 80% CL with ASPICP. 2018-2020 relative F projection under red 3LN HCR and status quo 2017 TAC.

Year	Relative F trajectory and 2018-2020 projection under red 3LN HCR		Relative F trajectory and 2018-2020 projection under status quo 2017 TAC	
	Point estimate	Approx 80% upper CL	Point estimate	Approx bias corrected 80% upper CL
2017	0.376	0.373	0.3759	0.4955
2018	0.448	0.496	0.4475	0.5762
2019	0.572	0.576	0.4456	0.5618
2020	0.577	0.722	0.4438	0.5511




Fig. 1. Redfish catch and TAC's in Div. 3LN, 1959-2017




Fig. 2. Redfish Div. 3LN cpue 1959-1994




Fig. 3. Annual anomalies of the mean length of the commercial catch, 1990-2017.




Fig. 4a. Standardized biomass from bottom trawl surveys included in ASPIC framework, 1978-2017




Fig. 4b. Standardized survey female SSB from spring and autumn Canadian surveys on Div. 3LN, 1991-2017




Fig. 5a. Annual anomalies of the mean length on the spring survey stock, 1991-2016




Fig. 5b. Annual anomalies of the mean length on the autumn survey stock, 1991-2017.




Fig. 5c. Redfish in Div. 3LN =>20cm: numbers in the catch and survey abundance (10⁶)




Fig. 5d. Recruitment index (lengths <20cm) in the survey abundance (10^6)




Fig. 6a. B/B_{msy} from 2018 ASPICfit




Fig. 6b. F/F_{msy} from 2016 ASPICfit




Fig. 7. Catch versus Surplus Production from ASPIC_{fit} 2018




Fig. 8. Stock trajectory under a precautionary framework for ASPIC_{fit} 2018




Fig. 9a. B/Bmsy trajectory and 2018-2021 projection under 2014 red 3LN HCR. Results from ASPICP 2018 (point estimates with approximate 80% bc CL's)




Fig. 9b. F/F_{msy} trajectory and 2018-2020 projection under 2014 red 3LN HCR. results from ASPICP 2018 (point estimates with approximate 80% bc CL's)




Fig. 10a. B/B_{msy} trajectory and 2018-2021 projection under status quo catch at 2017 TAC. Results from ASPICP 2018 (point estimates with approximate 80% bc CL's)




Fig. 10b. F/Fmsy trajectory and 2018-2020 projection under status quo catch at 2017 TAC. Results from ASPICP 2018 (point estimates with approximate 80% bc CL's)




Fig. 11. B/Bmsy 2018-2021 projection under red 3LN HCR versus status quo 2017 TAC. Results from ASPICP 2018 (point estimates with approximate lower 80% bc CL's)

Appendix 1: Input .a7inp file of 2018 framework

ASPIC-V7

File generated by aspic5to7 v.0.62, at 2018-06-08 19:06:56

"3LN redfish"

Program mode (FIT/BOT), verbosity, [if BOT] N bootstraps, [opt] user percentile:

FIT 2

Model shape, conditioning (YLD/EFT), obj. fn. (SSE/LAV/ML/MLP):

LOGISTIC YLD SSE

N years, N series:

59 9

Monte Carlo mode (0/1/2), N trials:

0 20000

Convergence criteria (3 values):

1.00E-08 3.00E-08 1.00E-04

Maximum F, N restarts, [gen. model] N steps/yr:

6.00E+00 18 24

Random seed (large integer):

3941285

Initial guesses and bounds follow:

'B1K', guess, estflag, min, max, ['penalty', penalty], or [priorname, prior params]

B1K 5.00E-01 1 5.00E-02 3.00E+00 penalty 0.00E+00

#B1K 5.00E-01 1 5.00E-02 3.00E+00 prior uniform 5.00E-02 3.00E+00

'MSY', guess, estflag, min, max, [if MAP] priorname, prior params

MSY 2.10E+04 0 5.00E+03 5.00E+04 prior uniform 5.00E+03 5.00E+04

'Fmsy', guess, estflag, min, max, [if MAP] priorname, prior params

Fmsy 8.40E-02 1 4.20E-03 8.40E-01 prior uniform 4.20E-03 8.40E-01

q, guess, estflag, seriesweight, min, max, [if MAP] priorname, prior params

q 1.50E-05 1 1.00E+00 2.50E-07 1.50E-03 prior uniform 2.50E-07 1.50E-03

q 8.85E-01 1 1.00E+00 1.48E-02 5.31E+00 prior uniform 1.48E-02 5.31E+00

q 8.85E-01 1 1.00E+00 1.48E-02 5.31E+00 prior uniform 1.48E-02 5.31E+00

q 3.17E-01 1 1.00E+00 5.29E-03 1.20E+00 prior uniform 5.29E-03 1.20E+00

q 2.49E-01 1 1.00E+00 4.14E-03 1.20E+00 prior uniform 4.14E-03 1.20E+00

q 1.03E+00 1 1.00E+00 1.71E-02 6.17E+00 prior uniform 1.71E-02 6.17E+00

q 2.30E-01 1 1.00E+00 3.84E-03 1.20E+00 prior uniform 3.84E-03 1.20E+00

q 8.75E-01 1 1.00E+00 1.46E-02 5.25E+00 prior uniform 1.46E-02 5.25E+00

q 6.60E-01 1 1.00E+00 1.10E-02 3.96E+00 prior uniform 1.10E-02 3.96E+00

Parameters for GENGRID or GENFIT go here.

DATA

NOTE: Nominal CVs added by aspic5to7.

"Statlant CPUE"

CC

1959	1.4260E+00	4.4585E+04	3.0000E-01
1960	1.6020E+00	2.6562E+04	3.0000E-01
1961	1.6970E+00	2.3175E+04	3.0000E-01
1962	1.6310E+00	2.1439E+04	3.0000E-01
1963	1.6320E+00	2.7362E+04	3.0000E-01
1964	1.8120E+00	1.0261E+04	3.0000E-01
1965	2.1850E+00	2.3466E+04	3.0000E-01
1966	1.7810E+00	1.6974E+04	3.0000E-01
1967	1.8930E+00	2.7188E+04	3.0000E-01
1968	9.2200E-01	1.7660E+04	3.0000E-01
1969	1.3380E+00	2.4750E+04	3.0000E-01
1970	1.3670E+00	1.4419E+04	3.0000E-01
1971	1.3460E+00	3.4370E+04	3.0000E-01
1972	1.3870E+00	2.8933E+04	3.0000E-01
1973	1.6430E+00	3.3297E+04	3.0000E-01
1974	1.2900E+00	2.2286E+04	3.0000E-01
1975	1.6690E+00	1.7871E+04	3.0000E-01
1976	1.2920E+00	2.0513E+04	3.0000E-01
1977	1.2510E+00	1.6516E+04	3.0000E-01
1978	1.1060E+00	1.2043E+04	3.0000E-01
1979	1.4510E+00	1.4067E+04	3.0000E-01
1980	1.7610E+00	1.6030E+04	3.0000E-01
1981	1.5940E+00	2.4280E+04	3.0000E-01
1982	1.6610E+00	2.1547E+04	3.0000E-01
1983	1.5560E+00	1.9747E+04	3.0000E-01
1984	1.0490E+00	1.4761E+04	3.0000E-01
1985	1.0840E+00	2.0557E+04	3.0000E-01
1986	1.4130E+00	4.2805E+04	3.0000E-01
1987	1.5230E+00	7.9031E+04	3.0000E-01
1988	1.2080E+00	5.3266E+04	3.0000E-01
1989	1.3220E+00	3.3649E+04	3.0000E-01
1990	8.2500E-01	2.9105E+04	3.0000E-01
1991	6.6800E-01	2.5815E+04	3.0000E-01
1992	9.1200E-01	2.7283E+04	3.0000E-01
1993	8.0100E-01	2.1308E+04	3.0000E-01
1994	8.0200E-01	5.7410E+03	3.0000E-01
1995	-1.0000E-03	1.9890E+03	3.0000E-01
1996	-1.0000E-03	4.5100E+02	3.0000E-01
1997	-1.0000E-03	6.3000E+02	3.0000E-01
1998	-1.0000E-03	8.9900E+02	3.0000E-01
1999	-1.0000E-03	2.3180E+03	3.0000E-01
2000	-1.0000E-03	3.1410E+03	3.0000E-01
2001	-1.0000E-03	1.4420E+03	3.0000E-01
2002	-1.0000E-03	1.2160E+03	3.0000E-01
2003	-1.0000E-03	1.3340E+03	3.0000E-01
2004	-1.0000E-03	6.3700E+02	3.0000E-01
2005	-1.0000E-03	6.5900E+02	3.0000E-01
2006	-1.0000E-03	4.9600E+02	3.0000E-01
2007	-1.0000E-03	1.6640E+03	3.0000E-01
2008	-1.0000E-03	5.9700E+02	3.0000E-01
2009	-1.0000E-03	1.0510E+03	3.0000E-01
2010	-1.0000E-03	4.1200E+03	3.0000E-01
2011	-1.0000E-03	3.6720E+03	3.0000E-01
2012	-1.0000E-03	4.3160E+03	3.0000E-01
2013	-1.0000E-03	6.2320E+03	3.0000E-01
2014	-1.0000E-03	5.6950E+03	3.0000E-01
2015	-1.0000E-03	1.0467E+04	3.0000E-01
2016	-1.0000E-03	8.4570E+03	3.0000E-01
2017	-1.0000E-03	1.1816E+04	3.0000E-01

"3LN spring survey"

I1

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	-1.0000E-03	3.0000E-01
1979	-1.0000E-03	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	-1.0000E-03	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	-1.0000E-03	3.0000E-01
1985	-1.0000E-03	3.0000E-01
1986	-1.0000E-03	3.0000E-01
1987	-1.0000E-03	3.0000E-01
1988	-1.0000E-03	3.0000E-01
1989	-1.0000E-03	3.0000E-01
1990	-1.0000E-03	3.0000E-01
1991	1.0642E+04	3.0000E-01
1992	1.0066E+04	3.0000E-01
1993	2.2573E+04	3.0000E-01
1994	4.1620E+03	3.0000E-01
1995	5.8560E+03	3.0000E-01
1996	2.2812E+04	3.0000E-01
1997	1.4928E+04	3.0000E-01
1998	5.9402E+04	3.0000E-01
1999	6.1496E+04	3.0000E-01
2000	8.7842E+04	3.0000E-01
2001	4.1573E+04	3.0000E-01
2002	3.0959E+04	3.0000E-01
2003	2.7700E+04	3.0000E-01
2004	7.9631E+04	3.0000E-01
2005	6.6462E+04	3.0000E-01
2006	-1.0000E-03	3.0000E-01
2007	2.1885E+05	3.0000E-01
2008	1.4398E+05	3.0000E-01
2009	1.8338E+05	3.0000E-01
2010	1.6535E+05	3.0000E-01
2011	1.7369E+05	3.0000E-01
2012	3.2198E+05	3.0000E-01
2013	2.7151E+05	3.0000E-01
2014	2.7175E+05	3.0000E-01
2015	4.8056E+05	3.0000E-01
2016	6.5421E+05	3.0000E-01
2017	-1.0000E-03	3.0000E-01

"3LN autumn survey"

I2

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	-1.0000E-03	3.0000E-01
1979	-1.0000E-03	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	-1.0000E-03	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	-1.0000E-03	3.0000E-01
1985	-1.0000E-03	3.0000E-01
1986	-1.0000E-03	3.0000E-01
1987	-1.0000E-03	3.0000E-01
1988	-1.0000E-03	3.0000E-01
1989	-1.0000E-03	3.0000E-01
1990	-1.0000E-03	3.0000E-01
1991	3.7886E+04	3.0000E-01
1992	1.3641E+05	3.0000E-01
1993	1.9233E+04	3.0000E-01
1994	3.1757E+04	3.0000E-01
1995	9.0728E+04	3.0000E-01
1996	1.5968E+04	3.0000E-01
1997	7.0660E+04	3.0000E-01
1998	1.1222E+05	3.0000E-01
1999	7.1986E+04	3.0000E-01
2000	1.0046E+05	3.0000E-01
2001	1.3257E+05	3.0000E-01
2002	5.0123E+04	3.0000E-01
2003	7.1889E+04	3.0000E-01
2004	4.9907E+04	3.0000E-01
2005	5.8561E+04	3.0000E-01
2006	9.1883E+04	3.0000E-01
2007	1.2476E+05	3.0000E-01
2008	1.9849E+05	3.0000E-01
2009	2.4671E+05	3.0000E-01
2010	4.6149E+05	3.0000E-01
2011	5.6228E+05	3.0000E-01
2012	5.9599E+05	3.0000E-01
2013	2.8875E+05	3.0000E-01
2014	-1.0000E-03	3.0000E-01
2015	4.2586E+05	3.0000E-01
2016	2.1518E+05	3.0000E-01
2017	1.9203E+05	3.0000E-01

"3LN Power russian survey"

I1

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	-1.0000E-03	3.0000E-01
1979	-1.0000E-03	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	-1.0000E-03	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	2.1588E+05	3.0000E-01
1985	9.3996E+04	3.0000E-01
1986	6.2975E+04	3.0000E-01
1987	7.0298E+04	3.0000E-01
1988	4.4884E+04	3.0000E-01
1989	1.2268E+04	3.0000E-01
1990	8.3650E+03	3.0000E-01
1991	1.8680E+04	3.0000E-01
1992	-1.0000E-03	3.0000E-01
1993	-1.0000E-03	3.0000E-01
1994	-1.0000E-03	3.0000E-01
1995	-1.0000E-03	3.0000E-01
1996	-1.0000E-03	3.0000E-01
1997	-1.0000E-03	3.0000E-01
1998	-1.0000E-03	3.0000E-01
1999	-1.0000E-03	3.0000E-01
2000	-1.0000E-03	3.0000E-01
2001	-1.0000E-03	3.0000E-01
2002	-1.0000E-03	3.0000E-01
2003	-1.0000E-03	3.0000E-01
2004	-1.0000E-03	3.0000E-01
2005	-1.0000E-03	3.0000E-01
2006	-1.0000E-03	3.0000E-01
2007	-1.0000E-03	3.0000E-01
2008	-1.0000E-03	3.0000E-01
2009	-1.0000E-03	3.0000E-01
2010	-1.0000E-03	3.0000E-01
2011	-1.0000E-03	3.0000E-01
2012	-1.0000E-03	3.0000E-01
2013	-1.0000E-03	3.0000E-01
2014	-1.0000E-03	3.0000E-01
2015	-1.0000E-03	3.0000E-01
2016	-1.0000E-03	3.0000E-01
2017	-1.0000E-03	3.0000E-01

"3L winter survey"

I0

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	-1.0000E-03	3.0000E-01
1979	-1.0000E-03	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	-1.0000E-03	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	-1.0000E-03	3.0000E-01
1985	9.0245E+04	3.0000E-01
1986	3.6568E+04	3.0000E-01
1987	-1.0000E-03	3.0000E-01
1988	-1.0000E-03	3.0000E-01
1989	-1.0000E-03	3.0000E-01
1990	1.8202E+04	3.0000E-01
1991	-1.0000E-03	3.0000E-01
1992	-1.0000E-03	3.0000E-01
1993	-1.0000E-03	3.0000E-01
1994	-1.0000E-03	3.0000E-01
1995	-1.0000E-03	3.0000E-01
1996	-1.0000E-03	3.0000E-01
1997	-1.0000E-03	3.0000E-01
1998	-1.0000E-03	3.0000E-01
1999	-1.0000E-03	3.0000E-01
2000	-1.0000E-03	3.0000E-01
2001	-1.0000E-03	3.0000E-01
2002	-1.0000E-03	3.0000E-01
2003	-1.0000E-03	3.0000E-01
2004	-1.0000E-03	3.0000E-01
2005	-1.0000E-03	3.0000E-01
2006	-1.0000E-03	3.0000E-01
2007	-1.0000E-03	3.0000E-01
2008	-1.0000E-03	3.0000E-01
2009	-1.0000E-03	3.0000E-01
2010	-1.0000E-03	3.0000E-01
2011	-1.0000E-03	3.0000E-01
2012	-1.0000E-03	3.0000E-01
2013	-1.0000E-03	3.0000E-01
2014	-1.0000E-03	3.0000E-01
2015	-1.0000E-03	3.0000E-01
2016	-1.0000E-03	3.0000E-01
2017	-1.0000E-03	3.0000E-01

"3L summer survey"

I1

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	3.1116E+05	3.0000E-01
1979	2.2779E+05	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	2.6138E+05	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	2.7771E+05	3.0000E-01
1985	1.6104E+05	3.0000E-01
1986	-1.0000E-03	3.0000E-01
1987	-1.0000E-03	3.0000E-01
1988	-1.0000E-03	3.0000E-01
1989	-1.0000E-03	3.0000E-01
1990	9.2840E+04	3.0000E-01
1991	3.7572E+04	3.0000E-01
1992	-1.0000E-03	3.0000E-01
1993	2.0838E+04	3.0000E-01
1994	-1.0000E-03	3.0000E-01
1995	-1.0000E-03	3.0000E-01
1996	-1.0000E-03	3.0000E-01
1997	-1.0000E-03	3.0000E-01
1998	-1.0000E-03	3.0000E-01
1999	-1.0000E-03	3.0000E-01
2000	-1.0000E-03	3.0000E-01
2001	-1.0000E-03	3.0000E-01
2002	-1.0000E-03	3.0000E-01
2003	-1.0000E-03	3.0000E-01
2004	-1.0000E-03	3.0000E-01
2005	-1.0000E-03	3.0000E-01
2006	-1.0000E-03	3.0000E-01
2007	-1.0000E-03	3.0000E-01
2008	-1.0000E-03	3.0000E-01
2009	-1.0000E-03	3.0000E-01
2010	-1.0000E-03	3.0000E-01
2011	-1.0000E-03	3.0000E-01
2012	-1.0000E-03	3.0000E-01
2013	-1.0000E-03	3.0000E-01
2014	-1.0000E-03	3.0000E-01
2015	-1.0000E-03	3.0000E-01
2016	-1.0000E-03	3.0000E-01
2017	-1.0000E-03	3.0000E-01

"3L autumn survey"

I2

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	-1.0000E-03	3.0000E-01
1979	-1.0000E-03	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	-1.0000E-03	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	-1.0000E-03	3.0000E-01
1985	9.8233E+04	3.0000E-01
1986	1.7119E+04	3.0000E-01
1987	-1.0000E-03	3.0000E-01
1988	-1.0000E-03	3.0000E-01
1989	-1.0000E-03	3.0000E-01
1990	2.0743E+04	3.0000E-01
1991	-1.0000E-03	3.0000E-01
1992	-1.0000E-03	3.0000E-01
1993	-1.0000E-03	3.0000E-01
1994	-1.0000E-03	3.0000E-01
1995	-1.0000E-03	3.0000E-01
1996	-1.0000E-03	3.0000E-01
1997	-1.0000E-03	3.0000E-01
1998	-1.0000E-03	3.0000E-01
1999	-1.0000E-03	3.0000E-01
2000	-1.0000E-03	3.0000E-01
2001	-1.0000E-03	3.0000E-01
2002	-1.0000E-03	3.0000E-01
2003	-1.0000E-03	3.0000E-01
2004	-1.0000E-03	3.0000E-01
2005	-1.0000E-03	3.0000E-01
2006	-1.0000E-03	3.0000E-01
2007	-1.0000E-03	3.0000E-01
2008	-1.0000E-03	3.0000E-01
2009	-1.0000E-03	3.0000E-01
2010	-1.0000E-03	3.0000E-01
2011	-1.0000E-03	3.0000E-01
2012	-1.0000E-03	3.0000E-01
2013	-1.0000E-03	3.0000E-01
2014	-1.0000E-03	3.0000E-01
2015	-1.0000E-03	3.0000E-01
2016	-1.0000E-03	3.0000E-01
2017	-1.0000E-03	3.0000E-01

"3N spanish survey"

I1

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	-1.0000E-03	3.0000E-01
1979	-1.0000E-03	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	-1.0000E-03	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	-1.0000E-03	3.0000E-01
1985	-1.0000E-03	3.0000E-01
1986	-1.0000E-03	3.0000E-01
1987	-1.0000E-03	3.0000E-01
1988	-1.0000E-03	3.0000E-01
1989	-1.0000E-03	3.0000E-01
1990	-1.0000E-03	3.0000E-01
1991	-1.0000E-03	3.0000E-01
1992	-1.0000E-03	3.0000E-01
1993	-1.0000E-03	3.0000E-01
1994	-1.0000E-03	3.0000E-01
1995	4.6084E+04	3.0000E-01
1996	6.5580E+03	3.0000E-01
1997	4.7530E+03	3.0000E-01
1998	2.2540E+04	3.0000E-01
1999	4.6459E+04	3.0000E-01
2000	6.8928E+04	3.0000E-01
2001	5.3855E+04	3.0000E-01
2002	7.6200E+03	3.0000E-01
2003	1.1031E+04	3.0000E-01
2004	2.7016E+04	3.0000E-01
2005	1.4692E+05	3.0000E-01
2006	8.7830E+04	3.0000E-01
2007	8.7602E+04	3.0000E-01
2008	6.8059E+04	3.0000E-01
2009	7.3574E+05	3.0000E-01
2010	3.5954E+05	3.0000E-01
2011	4.1830E+05	3.0000E-01
2012	2.6524E+05	3.0000E-01
2013	4.2953E+05	3.0000E-01
2014	1.7805E+05	3.0000E-01
2015	5.2346E+05	3.0000E-01
2016	1.1727E+05	3.0000E-01
2017	2.6590E+05	3.0000E-01

"3L spanish survey"

I1

1959	-1.0000E-03	3.0000E-01
1960	-1.0000E-03	3.0000E-01
1961	-1.0000E-03	3.0000E-01
1962	-1.0000E-03	3.0000E-01
1963	-1.0000E-03	3.0000E-01
1964	-1.0000E-03	3.0000E-01
1965	-1.0000E-03	3.0000E-01
1966	-1.0000E-03	3.0000E-01
1967	-1.0000E-03	3.0000E-01
1968	-1.0000E-03	3.0000E-01
1969	-1.0000E-03	3.0000E-01
1970	-1.0000E-03	3.0000E-01
1971	-1.0000E-03	3.0000E-01
1972	-1.0000E-03	3.0000E-01
1973	-1.0000E-03	3.0000E-01
1974	-1.0000E-03	3.0000E-01
1975	-1.0000E-03	3.0000E-01
1976	-1.0000E-03	3.0000E-01
1977	-1.0000E-03	3.0000E-01
1978	-1.0000E-03	3.0000E-01
1979	-1.0000E-03	3.0000E-01
1980	-1.0000E-03	3.0000E-01
1981	-1.0000E-03	3.0000E-01
1982	-1.0000E-03	3.0000E-01
1983	-1.0000E-03	3.0000E-01
1984	-1.0000E-03	3.0000E-01
1985	-1.0000E-03	3.0000E-01
1986	-1.0000E-03	3.0000E-01
1987	-1.0000E-03	3.0000E-01
1988	-1.0000E-03	3.0000E-01
1989	-1.0000E-03	3.0000E-01
1990	-1.0000E-03	3.0000E-01
1991	-1.0000E-03	3.0000E-01
1992	-1.0000E-03	3.0000E-01
1993	-1.0000E-03	3.0000E-01
1994	-1.0000E-03	3.0000E-01
1995	-1.0000E-03	3.0000E-01
1996	-1.0000E-03	3.0000E-01
1997	-1.0000E-03	3.0000E-01
1998	-1.0000E-03	3.0000E-01
1999	-1.0000E-03	3.0000E-01
2000	-1.0000E-03	3.0000E-01
2001	-1.0000E-03	3.0000E-01
2002	-1.0000E-03	3.0000E-01
2003	-1.0000E-03	3.0000E-01
2004	-1.0000E-03	3.0000E-01
2005	-1.0000E-03	3.0000E-01
2006	7.0066E+04	3.0000E-01
2007	3.1410E+04	3.0000E-01
2008	7.5567E+04	3.0000E-01
2009	1.0368E+05	3.0000E-01
2010	2.6675E+05	3.0000E-01
2011	1.7063E+05	3.0000E-01
2012	4.8147E+05	3.0000E-01
2013	2.3516E+05	3.0000E-01
2014	2.1641E+05	3.0000E-01
2015	1.3042E+05	3.0000E-01
2016	9.8807E+04	3.0000E-01
2017	5.6557E+04	3.0000E-01

Appendix 2 ASPIC Fit 2018 results
3LN redfish

Page 1

Friday, 08 Jun 2018 at 19:07:12

ASPIC -- A Surplus-Production Model Including Covariates (BETA Ver. 7.03)

FIT program mode

Author: Michael H. Prager

Prager Consulting

<http://www.mhprager.com>

LOGISTIC model mode

YLD conditioning

SSE optimization

Reference: Prager, M. H. 1994. A suite of extensions to a nonequilibrium
surplus-production model. Fishery Bulletin 92: 374-389.

ASPIC program and user's guide
available gratis at www.mhprager.com

CONTROL PARAMETERS (FROM INPUT FILE) Input file: C:/.../3LN/ASPIC3LN/2018/ASPIC18 fit - Cópia/ASPIC18qmaxbound.a7inp

Operation of ASPIC: Fit logistic (Schaefer) model by direct optimization.

Number of years analyzed: 59 Number of bootstrap trials: 0
Number of data series: 9 Objective function: Least squares
Relative conv. criterion (simplex): 1.000E-08 Monte Carlo search mode, trials: 0 20000
Relative conv. criterion (restart): 3.000E-08 Random number seed: 3941285
Relative conv. criterion (effort): 1.000E-04 Identical convergences required in fitting: 18
Maximum F allowed in fitting: 6.000

PROGRAM STATUS INFORMATION (NON-BOOTSTRAPPED ANALYSIS)

error code 0

Normal convergence

WARNING: Negative correlations detected between some indices. A fundamental assumption of ASPIC is that all indices
represent the abundance of the stock. That assumption should be checked.

Number of restarts required for convergence: 31

CORRELATION AMONG INPUT SERIES EXPRESSED AS CPUE (NUMBER OF PAIRWISE OBSERVATIONS BELOW)

```

-----
1 Statlant CPUE      | 1.000
|                   |
| 36                |
|                   |
2 3LN spring survey | -0.019 1.000
|                   |
| 4 25              |
|                   |
3 3LN autumn survey | 0.700 0.594 1.000
|                   |
| 4 24 26           |
|                   |
4 3LN Power russian survey | 0.108 0.000 0.000 1.000
|                   |
| 8 1 1 8           |
|                   |
5 3L winter survey  | 0.178 0.000 0.000 0.908 1.000
|                   |
| 3 0 0 3 3        |
|                   |
6 3L summer survey  | 0.733 -1.000 1.000 0.964 1.000 1.000
|                   |
| 8 2 2 4 2 8      |
|                   |
7 3L autumn survey  | -0.108 0.000 0.000 0.751 0.959 1.000 1.000
|                   |
| 3 0 0 3 3 2 3    |
|                   |
8 3N spanish survey | 0.000 0.463 0.690 0.000 0.000 0.000 1.000
|                   |
| 0 21 22 0 0 0 0 23
|                   |
9 3L spanish survey | 0.000 -0.036 0.792 0.000 0.000 0.000 0.171 1.000
|                   |
| 0 10 11 0 0 0 0 12 12
-----

```

1 2 3 4 5 6 7 8 9

GOODNESS-OF-FIT AND WEIGHTING (NON-BOOTSTRAPPED ANALYSIS)

Objective function component: source of variance	label and	Weighted SSE	N	Weighted MSE	Weighted weight	Current weight	Inv. var. in CPUE	R-squared
Loss(-1)	Unmatched yield	0.000E+00						
Loss(0)	Penalty on B1 > K	0.000E+00	1	N/A	0.000E+00	N/A		
Loss(1)	Statlant CPUE	7.228E+00	36	2.126E-01	1.000E+00	1.645E+00		-0.306
Loss(2)	3LN spring survey	1.262E+01	25	5.489E-01	1.000E+00	6.370E-01		0.488
Loss(3)	3LN autumn survey	9.811E+00	26	4.088E-01	1.000E+00	8.553E-01		0.558
Loss(4)	3LN Power russian survey	3.446E+00	8	5.743E-01	1.000E+00	6.088E-01		0.265
Loss(5)	3L winter survey	4.345E-01	3	4.345E-01	1.000E+00	8.047E-01		0.417
Loss(6)	3L summer survey	7.658E-01	8	1.276E-01	1.000E+00	2.739E+00		0.751
Loss(7)	3L autumn survey	1.460E+00	3	1.460E+00	1.000E+00	2.395E-01		0.252
Loss(8)	3N spanish survey	1.975E+01	23	9.402E-01	1.000E+00	3.719E-01		0.229
Loss(9)	3L spanish survey	5.146E+00	12	5.146E-01	1.000E+00	6.795E-01		0.002
TOTAL OBJECTIVE FUNCTION, MSE, RMSE:		6.06605699E+01		4.561E-01	6.753E-01			
Estimated contrast index (good=0.5, best=1.0):		0.7014 Mean of B coverage proportions > and < Bmsy					
Estimated nearness index (best=1.0):		1.0000 Proportional closeness of any B to Bmsy					



3LN redfish

Page 2

MODEL PARAMETER ESTIMATES (NON-BOOTSTRAPPED)

Parameter	Estimate	User guess	2nd guess	Min bound	Max bound	Estim?	
B1/K	Starting relative biomass (in 1959)	6.976E-01	5.000E-01	2.874E+00	5.000E-02	3.000E+00	1
MSY	Maximum sustainable yield	2.100E+04	2.100E+04	2.100E+04	2.100E+04	2.100E+04	0
Fmsy	Fishing mortality rate at MSY	1.122E-01	8.400E-02	1.029E-01	4.200E-03	8.400E-01	1
phi	Shape of production curve (Bmsy/K)	0.5000	0.5000	----	----	----	0
q(1)	Statlant CPUE	8.243E-06	1.500E-05	5.320E-06	2.500E-07	1.500E-03	1
q(2)	3LN spring survey	7.814E-01	8.850E-01	4.408E-01	1.480E-02	5.310E+00	1
q(3)	3LN autumn survey	1.299E+00	8.850E-01	4.319E-01	1.480E-02	5.310E+00	1
q(4)	3LN Power russian survey	3.129E-01	3.170E-01	1.408E-01	5.290E-03	1.200E+00	1
q(5)	3L winter survey	2.461E-01	2.490E-01	1.589E-02	4.140E-03	1.200E+00	1
q(6)	3L summer survey	1.009E+00	1.030E+00	3.419E-01	1.710E-02	6.170E+00	1
q(7)	3L autumn survey	2.276E-01	2.300E-01	2.499E-01	3.840E-03	1.200E+00	1
q(8)	3N spanish survey	7.679E-01	8.750E-01	1.097E+00	1.460E-02	5.250E+00	1
q(9)	3L spanish survey	6.024E-01	6.600E-01	1.849E-01	1.100E-02	3.960E+00	1

MANAGEMENT and DERIVED PARAMETER ESTIMATES (NON-BOOTSTRAPPED)

Parameter	Estimate	Logistic formula	General formula
MSY	Maximum sustainable yield	2.100E+04	----
Bmsy	Stock biomass giving MSY	1.871E+05	K/2 $K^n \cdot (1/(1-n))$
K	Carrying capacity	3.743E+05	2*Bmsy Bmsy/phi
n	Exponent in production function	2.0000	----
g	Fletcher's gamma	4.000E+00	---- $[n^{**}(n/(n-1))]/[n-1]$
B./Bmsy	Ratio: B(2018)/Bmsy	1.507E+00	----
F./Fmsy	Ratio: F(2017)/Fmsy	3.759E-01	----
Fmsy/F.	Ratio: Fmsy/F(2017)	2.660E+00	----
Y.(Fmsy)	Approx. yield available at Fmsy in 2018	3.082E+04	MSY*B./Bmsy MSY*B./Bmsy
...	...as proportion of MSY	1.467E+00	----
Ye.	Equilibrium yield available in 2018	1.560E+04	$4*MSY*(B/K-(B/K)**2)$ $g*MSY*(B/K-(B/K)**n)$
...	...as proportion of MSY	7.426E-01	----
----- Fishing effort rate at MSY in units of each CE or CC series -----			
fmsy(1)	Statlant CPUE	1.361E+04	Fmsy/q(1) Fmsy/q(1)



ESTIMATED POPULATION TRAJECTORY (NON-BOOTSTRAPPED)

Year	total	Estimated	Estimated	Observed	Model	Estimated	Ratio of	Ratio of
Obs or ID	F mort	starting	average	total	surplus	F mort	to Fmsy	to Bmsy
		biomass	biomass	yield	yield	production		
1 1959	0.180	2.611E+05	2.476E+05	4.458E+04	4.458E+04	1.877E+04	1.604E+00	1.395E+00
2 1960	0.115	2.353E+05	2.318E+05	2.656E+04	2.656E+04	1.980E+04	1.021E+00	1.257E+00
3 1961	0.102	2.285E+05	2.269E+05	2.317E+04	2.317E+04	2.005E+04	9.101E-01	1.221E+00
4 1962	0.095	2.254E+05	2.247E+05	2.144E+04	2.144E+04	2.015E+04	8.501E-01	1.204E+00
5 1963	0.124	2.241E+05	2.205E+05	2.736E+04	2.736E+04	2.033E+04	1.106E+00	1.198E+00
6 1964	0.046	2.171E+05	2.221E+05	1.026E+04	1.026E+04	2.026E+04	4.116E-01	1.160E+00
7 1965	0.104	2.271E+05	2.254E+05	2.347E+04	2.347E+04	2.012E+04	9.279E-01	1.213E+00
8 1966	0.075	2.237E+05	2.253E+05	1.697E+04	1.697E+04	2.012E+04	6.713E-01	1.196E+00
9 1967	0.122	2.269E+05	2.233E+05	2.719E+04	2.719E+04	2.021E+04	1.085E+00	1.212E+00
10 1968	0.080	2.199E+05	2.213E+05	1.766E+04	1.766E+04	2.030E+04	7.113E-01	1.175E+00
11 1969	0.112	2.225E+05	2.203E+05	2.475E+04	2.475E+04	2.034E+04	1.001E+00	1.189E+00
12 1970	0.065	2.181E+05	2.211E+05	1.442E+04	1.442E+04	2.031E+04	5.811E-01	1.166E+00
13 1971	0.159	2.240E+05	2.168E+05	3.437E+04	3.437E+04	2.046E+04	1.412E+00	1.197E+00
14 1972	0.140	2.101E+05	2.059E+05	2.893E+04	2.893E+04	2.078E+04	1.252E+00	1.123E+00
15 1973	0.170	2.020E+05	1.956E+05	3.330E+04	3.330E+04	2.095E+04	1.517E+00	1.079E+00
16 1974	0.118	1.896E+05	1.890E+05	2.229E+04	2.229E+04	2.100E+04	1.051E+00	1.013E+00
17 1975	0.094	1.883E+05	1.899E+05	1.787E+04	1.787E+04	2.099E+04	8.385E-01	1.006E+00
18 1976	0.107	1.915E+05	1.917E+05	2.051E+04	2.051E+04	2.099E+04	9.536E-01	1.023E+00
19 1977	0.085	1.919E+05	1.942E+05	1.652E+04	1.652E+04	2.097E+04	7.579E-01	1.026E+00
20 1978	0.060	1.964E+05	2.009E+05	1.204E+04	1.204E+04	2.088E+04	5.343E-01	1.049E+00
21 1979	0.067	2.052E+05	2.086E+05	1.407E+04	1.407E+04	2.072E+04	6.009E-01	1.097E+00
22 1980	0.075	2.119E+05	2.142E+05	1.603E+04	1.603E+04	2.056E+04	6.669E-01	1.132E+00
23 1981	0.113	2.164E+05	2.145E+05	2.428E+04	2.428E+04	2.055E+04	1.009E+00	1.156E+00
24 1982	0.102	2.127E+05	2.122E+05	2.155E+04	2.155E+04	2.062E+04	9.048E-01	1.136E+00
25 1983	0.093	2.118E+05	2.122E+05	1.975E+04	1.975E+04	2.062E+04	8.293E-01	1.132E+00
26 1984	0.068	2.126E+05	2.156E+05	1.476E+04	1.476E+04	2.051E+04	6.102E-01	1.136E+00
27 1985	0.094	2.184E+05	2.183E+05	2.056E+04	2.056E+04	2.042E+04	8.391E-01	1.167E+00
28 1986	0.207	2.182E+05	2.068E+05	4.280E+04	4.280E+04	2.074E+04	1.845E+00	1.166E+00
29 1987	0.480	1.962E+05	1.647E+05	7.903E+04	7.903E+04	2.053E+04	4.275E+00	1.048E+00
30 1988	0.447	1.377E+05	1.191E+05	5.327E+04	5.327E+04	1.816E+04	3.987E+00	7.357E-01
31 1989	0.361	1.026E+05	9.322E+04	3.365E+04	3.365E+04	1.569E+04	3.217E+00	5.481E-01
32 1990	0.380	8.462E+04	7.658E+04	2.910E+04	2.910E+04	1.366E+04	3.387E+00	4.522E-01
33 1991	0.418	6.917E+04	6.172E+04	2.582E+04	2.582E+04	1.156E+04	3.727E+00	3.696E-01
34 1992	0.606	5.491E+04	4.505E+04	2.728E+04	2.728E+04	8.876E+03	5.397E+00	2.934E-01
35 1993	0.760	3.651E+04	2.802E+04	2.131E+04	2.131E+04	5.806E+03	6.776E+00	1.951E-01
36 1994	0.283	2.101E+04	2.028E+04	5.741E+03	5.741E+03	4.304E+03	2.523E+00	1.122E-01
37 1995	0.096	1.957E+04	2.075E+04	1.989E+03	1.989E+03	4.399E+03	8.541E-01	1.046E-01
38 1996	0.019	2.198E+04	2.423E+04	4.510E+02	4.510E+02	5.085E+03	1.659E-01	1.175E-01
39 1997	0.022	2.661E+04	2.925E+04	6.300E+02	6.300E+02	6.050E+03	1.920E-01	1.422E-01
40 1998	0.026	3.203E+04	3.507E+04	8.990E+02	8.990E+02	7.131E+03	2.285E-01	1.712E-01
41 1999	0.056	3.826E+04	4.116E+04	2.318E+03	2.318E+03	8.220E+03	5.019E-01	2.045E-01
42 2000	0.067	4.417E+04	4.717E+04	3.141E+03	3.141E+03	9.250E+03	5.934E-01	2.360E-01
43 2001	0.026	5.028E+04	5.469E+04	1.442E+03	1.442E+03	1.048E+04	2.349E-01	2.687E-01
44 2002	0.019	5.931E+04	6.458E+04	1.216E+03	1.216E+03	1.199E+04	1.678E-01	3.169E-01
45 2003	0.018	7.008E+04	7.610E+04	1.334E+03	1.334E+03	1.360E+04	1.562E-01	3.745E-01
46 2004	0.007	8.235E+04	8.953E+04	6.370E+02	6.370E+02	1.528E+04	6.340E-02	4.400E-01
47 2005	0.006	9.699E+04	1.050E+05	6.590E+02	6.590E+02	1.694E+04	5.593E-02	5.183E-01
48 2006	0.004	1.133E+05	1.221E+05	4.960E+02	4.960E+02	1.845E+04	3.619E-02	6.053E-01
49 2007	0.012	1.312E+05	1.402E+05	1.664E+03	1.664E+03	1.966E+04	1.058E-01	7.012E-01
50 2008	0.004	1.492E+05	1.591E+05	5.970E+02	5.970E+02	2.051E+04	3.343E-02	7.974E-01
51 2009	0.006	1.691E+05	1.791E+05	1.051E+03	1.051E+03	2.094E+04	5.230E-02	9.038E-01
52 2010	0.021	1.890E+05	1.975E+05	4.120E+03	4.120E+03	2.092E+04	1.859E-01	1.010E+00
53 2011	0.017	2.058E+05	2.143E+05	3.672E+03	3.672E+03	2.054E+04	1.527E-01	1.100E+00
54 2012	0.019	2.227E+05	2.306E+05	4.316E+03	4.316E+03	1.986E+04	1.668E-01	1.190E+00
55 2013	0.025	2.382E+05	2.447E+05	6.232E+03	6.232E+03	1.900E+04	2.269E-01	1.273E+00
56 2014	0.022	2.510E+05	2.573E+05	5.695E+03	5.695E+03	1.804E+04	1.972E-01	1.341E+00
57 2015	0.039	2.634E+05	2.668E+05	1.047E+04	1.047E+04	1.719E+04	3.496E-01	1.407E+00
58 2016	0.031	2.701E+05	2.742E+05	8.457E+03	8.457E+03	1.645E+04	2.749E-01	1.443E+00
59 2017	0.042	2.781E+05	2.801E+05	1.182E+04	1.182E+04	1.581E+04	3.759E-01	1.486E+00
60 2018		2.821E+05				1.507E+00		

RESULTS FOR DATA SERIES # 1 (NON-BOOTSTRAPPED)

Statlant CPUE

Data type CC: CPUE-catch series


Series weight: 1.000

Obs	Year	Observed CPUE	Estimated CPUE	Estim F	Observed yield	Model yield	Resid log scale	in weight	Statist
1	1959	1.426E+00	2.041E+00	0.1800	4.458E+04	4.458E+04	0.35875	1.000E+00	
2	1960	1.602E+00	1.911E+00	0.1146	2.656E+04	2.656E+04	0.17628	1.000E+00	
3	1961	1.697E+00	1.871E+00	0.1021	2.317E+04	2.317E+04	0.09736	1.000E+00	
4	1962	1.631E+00	1.853E+00	0.0954	2.144E+04	2.144E+04	0.12737	1.000E+00	
5	1963	1.632E+00	1.818E+00	0.1241	2.736E+04	2.736E+04	0.10771	1.000E+00	
6	1964	1.812E+00	1.831E+00	0.0462	1.026E+04	1.026E+04	0.01055	1.000E+00	
7	1965	2.185E+00	1.858E+00	0.1041	2.347E+04	2.347E+04	-0.16227	1.000E+00	
8	1966	1.781E+00	1.858E+00	0.0753	1.697E+04	1.697E+04	0.04207	1.000E+00	
9	1967	1.893E+00	1.841E+00	0.1218	2.719E+04	2.719E+04	-0.02801	1.000E+00	
10	1968	9.220E-01	1.824E+00	0.0798	1.766E+04	1.766E+04	0.68216	1.000E+00	
11	1969	1.338E+00	1.816E+00	0.1124	2.475E+04	2.475E+04	0.30540	1.000E+00	
12	1970	1.367E+00	1.823E+00	0.0652	1.442E+04	1.442E+04	0.28779	1.000E+00	
13	1971	1.346E+00	1.788E+00	0.1585	3.437E+04	3.437E+04	0.28369	1.000E+00	
14	1972	1.387E+00	1.698E+00	0.1405	2.893E+04	2.893E+04	0.20204	1.000E+00	
15	1973	1.643E+00	1.612E+00	0.1702	3.330E+04	3.330E+04	-0.01877	1.000E+00	
16	1974	1.290E+00	1.558E+00	0.1179	2.229E+04	2.229E+04	0.18855	1.000E+00	
17	1975	1.669E+00	1.566E+00	0.0941	1.787E+04	1.787E+04	-0.06398	1.000E+00	
18	1976	1.292E+00	1.580E+00	0.1070	2.051E+04	2.051E+04	0.20137	1.000E+00	
19	1977	1.251E+00	1.601E+00	0.0851	1.652E+04	1.652E+04	0.24654	1.000E+00	
20	1978	1.106E+00	1.656E+00	0.0600	1.204E+04	1.204E+04	0.40350	1.000E+00	
21	1979	1.451E+00	1.720E+00	0.0674	1.407E+04	1.407E+04	0.16982	1.000E+00	
22	1980	1.761E+00	1.766E+00	0.0748	1.603E+04	1.603E+04	0.00259	1.000E+00	
23	1981	1.594E+00	1.768E+00	0.1132	2.428E+04	2.428E+04	0.10369	1.000E+00	
24	1982	1.661E+00	1.749E+00	0.1015	2.155E+04	2.155E+04	0.05178	1.000E+00	
25	1983	1.556E+00	1.749E+00	0.0931	1.975E+04	1.975E+04	0.11706	1.000E+00	
26	1984	1.049E+00	1.777E+00	0.0685	1.476E+04	1.476E+04	0.52703	1.000E+00	
27	1985	1.084E+00	1.800E+00	0.0942	2.056E+04	2.056E+04	0.50691	1.000E+00	
28	1986	1.413E+00	1.705E+00	0.2070	4.280E+04	4.280E+04	0.18763	1.000E+00	
29	1987	1.523E+00	1.358E+00	0.4798	7.903E+04	7.903E+04	-0.11475	1.000E+00	
30	1988	1.208E+00	9.814E-01	0.4474	5.327E+04	5.327E+04	-0.20772	1.000E+00	
31	1989	1.322E+00	7.685E-01	0.3610	3.365E+04	3.365E+04	-0.54250	1.000E+00	
32	1990	8.250E-01	6.312E-01	0.3801	2.910E+04	2.910E+04	-0.26770	1.000E+00	
33	1991	6.680E-01	5.088E-01	0.4182	2.582E+04	2.582E+04	-0.27220	1.000E+00	
34	1992	9.120E-01	3.713E-01	0.6057	2.728E+04	2.728E+04	-0.89858	1.000E+00	
35	1993	8.010E-01	2.310E-01	0.7603	2.131E+04	2.131E+04	-1.24340	1.000E+00	
36	1994	8.020E-01	1.672E-01	0.2831	5.741E+03	5.741E+03	-1.56820	1.000E+00	
37	1995	*	1.711E-01	0.0958	1.989E+03	1.989E+03	0.00000	1.000E+00	
38	1996	*	1.997E-01	0.0186	4.510E+02	4.510E+02	0.00000	1.000E+00	
39	1997	*	2.411E-01	0.0215	6.300E+02	6.300E+02	0.00000	1.000E+00	
40	1998	*	2.891E-01	0.0256	8.990E+02	8.990E+02	0.00000	1.000E+00	
41	1999	*	3.393E-01	0.0563	2.318E+03	2.318E+03	0.00000	1.000E+00	
42	2000	*	3.888E-01	0.0666	3.141E+03	3.141E+03	0.00000	1.000E+00	
43	2001	*	4.509E-01	0.0264	1.442E+03	1.442E+03	0.00000	1.000E+00	
44	2002	*	5.324E-01	0.0188	1.216E+03	1.216E+03	0.00000	1.000E+00	
45	2003	*	6.273E-01	0.0175	1.334E+03	1.334E+03	0.00000	1.000E+00	
46	2004	*	7.380E-01	0.0071	6.370E+02	6.370E+02	0.00000	1.000E+00	
47	2005	*	8.656E-01	0.0063	6.590E+02	6.590E+02	0.00000	1.000E+00	
48	2006	*	1.007E+00	0.0041	4.960E+02	4.960E+02	0.00000	1.000E+00	
49	2007	*	1.155E+00	0.0119	1.664E+03	1.664E+03	0.00000	1.000E+00	
50	2008	*	1.312E+00	0.0038	5.970E+02	5.970E+02	0.00000	1.000E+00	
51	2009	*	1.476E+00	0.0059	1.051E+03	1.051E+03	0.00000	1.000E+00	
52	2010	*	1.628E+00	0.0209	4.120E+03	4.120E+03	0.00000	1.000E+00	
53	2011	*	1.767E+00	0.0171	3.672E+03	3.672E+03	0.00000	1.000E+00	
54	2012	*	1.901E+00	0.0187	4.316E+03	4.316E+03	0.00000	1.000E+00	
55	2013	*	2.017E+00	0.0255	6.232E+03	6.232E+03	0.00000	1.000E+00	
56	2014	*	2.121E+00	0.0221	5.695E+03	5.695E+03	0.00000	1.000E+00	
57	2015	*	2.199E+00	0.0392	1.047E+04	1.047E+04	0.00000	1.000E+00	
58	2016	*	2.260E+00	0.0308	8.457E+03	8.457E+03	0.00000	1.000E+00	
59	2017	*	2.309E+00	0.0422	1.182E+04	1.182E+04	0.00000	1.000E+00	

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 1



RESULTS FOR DATA SERIES # 2 (NON-BOOTSTRAPPED)

3LN spring survey

Data type I1: Abundance index (annual average)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid log index	weight	Statistic
1	1959	0.000E+00	0.000E+00	-- *	1.935E+05	0.00000	1.000E+00		
2	1960	0.000E+00	0.000E+00	-- *	1.811E+05	0.00000	1.000E+00		
3	1961	0.000E+00	0.000E+00	-- *	1.773E+05	0.00000	1.000E+00		
4	1962	0.000E+00	0.000E+00	-- *	1.756E+05	0.00000	1.000E+00		
5	1963	0.000E+00	0.000E+00	-- *	1.723E+05	0.00000	1.000E+00		
6	1964	0.000E+00	0.000E+00	-- *	1.736E+05	0.00000	1.000E+00		
7	1965	0.000E+00	0.000E+00	-- *	1.761E+05	0.00000	1.000E+00		
8	1966	0.000E+00	0.000E+00	-- *	1.761E+05	0.00000	1.000E+00		
9	1967	0.000E+00	0.000E+00	-- *	1.745E+05	0.00000	1.000E+00		
10	1968	0.000E+00	0.000E+00	-- *	1.729E+05	0.00000	1.000E+00		
11	1969	0.000E+00	0.000E+00	-- *	1.721E+05	0.00000	1.000E+00		
12	1970	0.000E+00	0.000E+00	-- *	1.728E+05	0.00000	1.000E+00		
13	1971	0.000E+00	0.000E+00	-- *	1.694E+05	0.00000	1.000E+00		
14	1972	0.000E+00	0.000E+00	-- *	1.609E+05	0.00000	1.000E+00		
15	1973	0.000E+00	0.000E+00	-- *	1.528E+05	0.00000	1.000E+00		
16	1974	0.000E+00	0.000E+00	-- *	1.476E+05	0.00000	1.000E+00		
17	1975	0.000E+00	0.000E+00	-- *	1.484E+05	0.00000	1.000E+00		
18	1976	0.000E+00	0.000E+00	-- *	1.498E+05	0.00000	1.000E+00		
19	1977	0.000E+00	0.000E+00	-- *	1.517E+05	0.00000	1.000E+00		
20	1978	0.000E+00	0.000E+00	-- *	1.569E+05	0.00000	1.000E+00		
21	1979	0.000E+00	0.000E+00	-- *	1.630E+05	0.00000	1.000E+00		
22	1980	0.000E+00	0.000E+00	-- *	1.674E+05	0.00000	1.000E+00		
23	1981	0.000E+00	0.000E+00	-- *	1.676E+05	0.00000	1.000E+00		
24	1982	0.000E+00	0.000E+00	-- *	1.658E+05	0.00000	1.000E+00		
25	1983	0.000E+00	0.000E+00	-- *	1.658E+05	0.00000	1.000E+00		
26	1984	0.000E+00	0.000E+00	-- *	1.684E+05	0.00000	1.000E+00		
27	1985	0.000E+00	0.000E+00	-- *	1.706E+05	0.00000	1.000E+00		
28	1986	0.000E+00	0.000E+00	-- *	1.616E+05	0.00000	1.000E+00		
29	1987	0.000E+00	0.000E+00	-- *	1.287E+05	0.00000	1.000E+00		
30	1988	0.000E+00	0.000E+00	-- *	9.303E+04	0.00000	1.000E+00		
31	1989	0.000E+00	0.000E+00	-- *	7.284E+04	0.00000	1.000E+00		
32	1990	0.000E+00	0.000E+00	-- *	5.983E+04	0.00000	1.000E+00		
33	1991	1.000E+00	1.000E+00	--	1.064E+04	4.823E+04	-1.51115	1.000E+00	
34	1992	1.000E+00	1.000E+00	--	1.007E+04	3.520E+04	-1.25177	1.000E+00	
35	1993	1.000E+00	1.000E+00	--	2.257E+04	2.190E+04	0.03042	1.000E+00	
36	1994	1.000E+00	1.000E+00	--	4.162E+03	1.584E+04	-1.33678	1.000E+00	
37	1995	1.000E+00	1.000E+00	--	5.856E+03	1.622E+04	-1.01852	1.000E+00	
38	1996	1.000E+00	1.000E+00	--	2.281E+04	1.893E+04	0.18651	1.000E+00	
39	1997	1.000E+00	1.000E+00	--	1.493E+04	2.285E+04	-0.42581	1.000E+00	
40	1998	1.000E+00	1.000E+00	--	5.940E+04	2.740E+04	0.77379	1.000E+00	
41	1999	1.000E+00	1.000E+00	--	6.150E+04	3.216E+04	0.64831	1.000E+00	
42	2000	1.000E+00	1.000E+00	--	8.784E+04	3.686E+04	0.86853	1.000E+00	
43	2001	1.000E+00	1.000E+00	--	4.157E+04	4.273E+04	-0.02756	1.000E+00	
44	2002	1.000E+00	1.000E+00	--	3.096E+04	5.046E+04	-0.48853	1.000E+00	
45	2003	1.000E+00	1.000E+00	--	2.770E+04	5.946E+04	-0.76384	1.000E+00	
46	2004	1.000E+00	1.000E+00	--	7.963E+04	6.996E+04	0.12953	1.000E+00	
47	2005	1.000E+00	1.000E+00	--	6.646E+04	8.204E+04	-0.21063	1.000E+00	
48	2006	0.000E+00	0.000E+00	-- *	9.543E+04	0.00000	1.000E+00		
49	2007	1.000E+00	1.000E+00	--	2.188E+05	1.095E+05	0.69237	1.000E+00	
50	2008	1.000E+00	1.000E+00	--	1.440E+05	1.243E+05	0.14670	1.000E+00	
51	2009	1.000E+00	1.000E+00	--	1.834E+05	1.399E+05	0.27051	1.000E+00	
52	2010	1.000E+00	1.000E+00	--	1.653E+05	1.543E+05	0.06921	1.000E+00	
53	2011	1.000E+00	1.000E+00	--	1.737E+05	1.675E+05	0.03649	1.000E+00	
54	2012	1.000E+00	1.000E+00	--	3.220E+05	1.801E+05	0.58072	1.000E+00	
55	2013	1.000E+00	1.000E+00	--	2.715E+05	1.912E+05	0.35060	1.000E+00	
56	2014	1.000E+00	1.000E+00	--	2.717E+05	2.010E+05	0.30141	1.000E+00	
57	2015	1.000E+00	1.000E+00	--	4.806E+05	2.085E+05	0.83521	1.000E+00	
58	2016	1.000E+00	1.000E+00	--	6.542E+05	2.142E+05	1.11641	1.000E+00	
59	2017	0.000E+00	0.000E+00	-- *	2.189E+05	0.00000	1.000E+00		

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 2



RESULTS FOR DATA SERIES # 3 (NON-BOOTSTRAPPED)

3LN autumn survey

Data type I2: Abundance index (end of year)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid log index	weight	Statistic
1	1959	0.000E+00	0.000E+00	-- *	3.057E+05	0.00000	1.000E+00		
2	1960	0.000E+00	0.000E+00	-- *	2.969E+05	0.00000	1.000E+00		
3	1961	0.000E+00	0.000E+00	-- *	2.928E+05	0.00000	1.000E+00		
4	1962	0.000E+00	0.000E+00	-- *	2.911E+05	0.00000	1.000E+00		
5	1963	0.000E+00	0.000E+00	-- *	2.820E+05	0.00000	1.000E+00		
6	1964	0.000E+00	0.000E+00	-- *	2.950E+05	0.00000	1.000E+00		
7	1965	0.000E+00	0.000E+00	-- *	2.907E+05	0.00000	1.000E+00		
8	1966	0.000E+00	0.000E+00	-- *	2.948E+05	0.00000	1.000E+00		
9	1967	0.000E+00	0.000E+00	-- *	2.857E+05	0.00000	1.000E+00		
10	1968	0.000E+00	0.000E+00	-- *	2.891E+05	0.00000	1.000E+00		
11	1969	0.000E+00	0.000E+00	-- *	2.834E+05	0.00000	1.000E+00		
12	1970	0.000E+00	0.000E+00	-- *	2.910E+05	0.00000	1.000E+00		
13	1971	0.000E+00	0.000E+00	-- *	2.730E+05	0.00000	1.000E+00		
14	1972	0.000E+00	0.000E+00	-- *	2.624E+05	0.00000	1.000E+00		
15	1973	0.000E+00	0.000E+00	-- *	2.463E+05	0.00000	1.000E+00		
16	1974	0.000E+00	0.000E+00	-- *	2.447E+05	0.00000	1.000E+00		
17	1975	0.000E+00	0.000E+00	-- *	2.487E+05	0.00000	1.000E+00		
18	1976	0.000E+00	0.000E+00	-- *	2.493E+05	0.00000	1.000E+00		
19	1977	0.000E+00	0.000E+00	-- *	2.551E+05	0.00000	1.000E+00		
20	1978	0.000E+00	0.000E+00	-- *	2.666E+05	0.00000	1.000E+00		
21	1979	0.000E+00	0.000E+00	-- *	2.753E+05	0.00000	1.000E+00		
22	1980	0.000E+00	0.000E+00	-- *	2.811E+05	0.00000	1.000E+00		
23	1981	0.000E+00	0.000E+00	-- *	2.763E+05	0.00000	1.000E+00		
24	1982	0.000E+00	0.000E+00	-- *	2.751E+05	0.00000	1.000E+00		
25	1983	0.000E+00	0.000E+00	-- *	2.762E+05	0.00000	1.000E+00		
26	1984	0.000E+00	0.000E+00	-- *	2.837E+05	0.00000	1.000E+00		
27	1985	0.000E+00	0.000E+00	-- *	2.835E+05	0.00000	1.000E+00		
28	1986	0.000E+00	0.000E+00	-- *	2.549E+05	0.00000	1.000E+00		
29	1987	0.000E+00	0.000E+00	-- *	1.789E+05	0.00000	1.000E+00		
30	1988	0.000E+00	0.000E+00	-- *	1.333E+05	0.00000	1.000E+00		
31	1989	0.000E+00	0.000E+00	-- *	1.099E+05	0.00000	1.000E+00		
32	1990	0.000E+00	0.000E+00	-- *	8.986E+04	0.00000	1.000E+00		
33	1991	1.000E+00	1.000E+00	--	3.789E+04	7.134E+04	-0.63291	1.000E+00	
34	1992	1.000E+00	1.000E+00	--	1.364E+05	4.743E+04	1.05643	1.000E+00	
35	1993	1.000E+00	1.000E+00	--	1.923E+04	2.729E+04	-0.34989	1.000E+00	
36	1994	1.000E+00	1.000E+00	--	3.176E+04	2.542E+04	0.22244	1.000E+00	
37	1995	1.000E+00	1.000E+00	--	9.073E+04	2.855E+04	1.15605	1.000E+00	
38	1996	1.000E+00	1.000E+00	--	1.597E+04	3.457E+04	-0.77252	1.000E+00	
39	1997	1.000E+00	1.000E+00	--	7.066E+04	4.162E+04	0.52941	1.000E+00	
40	1998	1.000E+00	1.000E+00	--	1.122E+05	4.971E+04	0.81422	1.000E+00	
41	1999	1.000E+00	1.000E+00	--	7.199E+04	5.738E+04	0.22680	1.000E+00	
42	2000	1.000E+00	1.000E+00	--	1.005E+05	6.532E+04	0.43053	1.000E+00	
43	2001	1.000E+00	1.000E+00	--	1.326E+05	7.705E+04	0.54261	1.000E+00	
44	2002	1.000E+00	1.000E+00	--	5.012E+04	9.105E+04	-0.59690	1.000E+00	
45	2003	1.000E+00	1.000E+00	--	7.189E+04	1.070E+05	-0.39753	1.000E+00	
46	2004	1.000E+00	1.000E+00	--	4.991E+04	1.260E+05	-0.92612	1.000E+00	
47	2005	1.000E+00	1.000E+00	--	5.856E+04	1.472E+05	-0.92141	1.000E+00	
48	2006	1.000E+00	1.000E+00	--	9.188E+04	1.705E+05	-0.61810	1.000E+00	
49	2007	1.000E+00	1.000E+00	--	1.248E+05	1.939E+05	-0.44074	1.000E+00	
50	2008	1.000E+00	1.000E+00	--	1.985E+05	2.197E+05	-0.10165	1.000E+00	
51	2009	1.000E+00	1.000E+00	--	2.467E+05	2.456E+05	0.00464	1.000E+00	
52	2010	1.000E+00	1.000E+00	--	4.615E+05	2.674E+05	0.54573	1.000E+00	
53	2011	1.000E+00	1.000E+00	--	5.623E+05	2.893E+05	0.66449	1.000E+00	
54	2012	1.000E+00	1.000E+00	--	5.960E+05	3.095E+05	0.65525	1.000E+00	
55	2013	1.000E+00	1.000E+00	--	2.888E+05	3.261E+05	-0.12163	1.000E+00	
56	2014	0.000E+00	0.000E+00	-- *	3.421E+05	0.00000	1.000E+00		
57	2015	1.000E+00	1.000E+00	--	4.259E+05	3.509E+05	0.19368	1.000E+00	
58	2016	1.000E+00	1.000E+00	--	2.152E+05	3.613E+05	-0.51814	1.000E+00	
59	2017	1.000E+00	1.000E+00	--	1.920E+05	3.665E+05	-0.64623	1.000E+00	

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 3



RESULTS FOR DATA SERIES # 4 (NON-BOOTSTRAPPED)

3LN Power russian survey

Data type I1: Abundance index (annual average)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid log index	weight	Statistic
1	1959	0.000E+00	0.000E+00	-- *	7.748E+04	0.00000	1.000E+00		
2	1960	0.000E+00	0.000E+00	-- *	7.252E+04	0.00000	1.000E+00		
3	1961	0.000E+00	0.000E+00	-- *	7.099E+04	0.00000	1.000E+00		
4	1962	0.000E+00	0.000E+00	-- *	7.031E+04	0.00000	1.000E+00		
5	1963	0.000E+00	0.000E+00	-- *	6.898E+04	0.00000	1.000E+00		
6	1964	0.000E+00	0.000E+00	-- *	6.950E+04	0.00000	1.000E+00		
7	1965	0.000E+00	0.000E+00	-- *	7.051E+04	0.00000	1.000E+00		
8	1966	0.000E+00	0.000E+00	-- *	7.050E+04	0.00000	1.000E+00		
9	1967	0.000E+00	0.000E+00	-- *	6.986E+04	0.00000	1.000E+00		
10	1968	0.000E+00	0.000E+00	-- *	6.922E+04	0.00000	1.000E+00		
11	1969	0.000E+00	0.000E+00	-- *	6.892E+04	0.00000	1.000E+00		
12	1970	0.000E+00	0.000E+00	-- *	6.918E+04	0.00000	1.000E+00		
13	1971	0.000E+00	0.000E+00	-- *	6.784E+04	0.00000	1.000E+00		
14	1972	0.000E+00	0.000E+00	-- *	6.443E+04	0.00000	1.000E+00		
15	1973	0.000E+00	0.000E+00	-- *	6.120E+04	0.00000	1.000E+00		
16	1974	0.000E+00	0.000E+00	-- *	5.912E+04	0.00000	1.000E+00		
17	1975	0.000E+00	0.000E+00	-- *	5.942E+04	0.00000	1.000E+00		
18	1976	0.000E+00	0.000E+00	-- *	5.997E+04	0.00000	1.000E+00		
19	1977	0.000E+00	0.000E+00	-- *	6.076E+04	0.00000	1.000E+00		
20	1978	0.000E+00	0.000E+00	-- *	6.284E+04	0.00000	1.000E+00		
21	1979	0.000E+00	0.000E+00	-- *	6.526E+04	0.00000	1.000E+00		
22	1980	0.000E+00	0.000E+00	-- *	6.701E+04	0.00000	1.000E+00		
23	1981	0.000E+00	0.000E+00	-- *	6.711E+04	0.00000	1.000E+00		
24	1982	0.000E+00	0.000E+00	-- *	6.639E+04	0.00000	1.000E+00		
25	1983	0.000E+00	0.000E+00	-- *	6.639E+04	0.00000	1.000E+00		
26	1984	1.000E+00	1.000E+00	--	2.159E+05	6.744E+04	1.16349	1.000E+00	
27	1985	1.000E+00	1.000E+00	--	9.400E+04	6.830E+04	0.31932	1.000E+00	
28	1986	1.000E+00	1.000E+00	--	6.298E+04	6.470E+04	-0.02697	1.000E+00	
29	1987	1.000E+00	1.000E+00	--	7.030E+04	5.154E+04	0.31045	1.000E+00	
30	1988	1.000E+00	1.000E+00	--	4.488E+04	3.725E+04	0.18647	1.000E+00	
31	1989	1.000E+00	1.000E+00	--	1.227E+04	2.917E+04	-0.86601	1.000E+00	
32	1990	1.000E+00	1.000E+00	--	8.365E+03	2.396E+04	-1.05224	1.000E+00	
33	1991	1.000E+00	1.000E+00	--	1.868E+04	1.931E+04	-0.03324	1.000E+00	
34	1992	0.000E+00	0.000E+00	-- *	1.409E+04	0.00000	1.000E+00		
35	1993	0.000E+00	0.000E+00	-- *	8.768E+03	0.00000	1.000E+00		
36	1994	0.000E+00	0.000E+00	-- *	6.344E+03	0.00000	1.000E+00		
37	1995	0.000E+00	0.000E+00	-- *	6.493E+03	0.00000	1.000E+00		
38	1996	0.000E+00	0.000E+00	-- *	7.580E+03	0.00000	1.000E+00		
39	1997	0.000E+00	0.000E+00	-- *	9.150E+03	0.00000	1.000E+00		
40	1998	0.000E+00	0.000E+00	-- *	1.097E+04	0.00000	1.000E+00		
41	1999	0.000E+00	0.000E+00	-- *	1.288E+04	0.00000	1.000E+00		
42	2000	0.000E+00	0.000E+00	-- *	1.476E+04	0.00000	1.000E+00		
43	2001	0.000E+00	0.000E+00	-- *	1.711E+04	0.00000	1.000E+00		
44	2002	0.000E+00	0.000E+00	-- *	2.021E+04	0.00000	1.000E+00		
45	2003	0.000E+00	0.000E+00	-- *	2.381E+04	0.00000	1.000E+00		
46	2004	0.000E+00	0.000E+00	-- *	2.801E+04	0.00000	1.000E+00		
47	2005	0.000E+00	0.000E+00	-- *	3.285E+04	0.00000	1.000E+00		
48	2006	0.000E+00	0.000E+00	-- *	3.821E+04	0.00000	1.000E+00		
49	2007	0.000E+00	0.000E+00	-- *	4.385E+04	0.00000	1.000E+00		
50	2008	0.000E+00	0.000E+00	-- *	4.978E+04	0.00000	1.000E+00		
51	2009	0.000E+00	0.000E+00	-- *	5.602E+04	0.00000	1.000E+00		
52	2010	0.000E+00	0.000E+00	-- *	6.178E+04	0.00000	1.000E+00		
53	2011	0.000E+00	0.000E+00	-- *	6.706E+04	0.00000	1.000E+00		
54	2012	0.000E+00	0.000E+00	-- *	7.213E+04	0.00000	1.000E+00		
55	2013	0.000E+00	0.000E+00	-- *	7.656E+04	0.00000	1.000E+00		
56	2014	0.000E+00	0.000E+00	-- *	8.050E+04	0.00000	1.000E+00		
57	2015	0.000E+00	0.000E+00	-- *	8.347E+04	0.00000	1.000E+00		
58	2016	0.000E+00	0.000E+00	-- *	8.578E+04	0.00000	1.000E+00		
59	2017	0.000E+00	0.000E+00	-- *	8.764E+04	0.00000	1.000E+00		

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 4



RESULTS FOR DATA SERIES # 5 (NON-BOOTSTRAPPED)

3L winter survey

Data type 10: Abundance index (start of year)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid log index	weight	Statistic
1	1959	0.000E+00	0.000E+00	-- *	6.426E+04	0.00000	1.000E+00		
2	1960	0.000E+00	0.000E+00	-- *	5.791E+04	0.00000	1.000E+00		
3	1961	0.000E+00	0.000E+00	-- *	5.624E+04	0.00000	1.000E+00		
4	1962	0.000E+00	0.000E+00	-- *	5.547E+04	0.00000	1.000E+00		
5	1963	0.000E+00	0.000E+00	-- *	5.516E+04	0.00000	1.000E+00		
6	1964	0.000E+00	0.000E+00	-- *	5.343E+04	0.00000	1.000E+00		
7	1965	0.000E+00	0.000E+00	-- *	5.589E+04	0.00000	1.000E+00		
8	1966	0.000E+00	0.000E+00	-- *	5.507E+04	0.00000	1.000E+00		
9	1967	0.000E+00	0.000E+00	-- *	5.584E+04	0.00000	1.000E+00		
10	1968	0.000E+00	0.000E+00	-- *	5.412E+04	0.00000	1.000E+00		
11	1969	0.000E+00	0.000E+00	-- *	5.477E+04	0.00000	1.000E+00		
12	1970	0.000E+00	0.000E+00	-- *	5.369E+04	0.00000	1.000E+00		
13	1971	0.000E+00	0.000E+00	-- *	5.514E+04	0.00000	1.000E+00		
14	1972	0.000E+00	0.000E+00	-- *	5.171E+04	0.00000	1.000E+00		
15	1973	0.000E+00	0.000E+00	-- *	4.971E+04	0.00000	1.000E+00		
16	1974	0.000E+00	0.000E+00	-- *	4.667E+04	0.00000	1.000E+00		
17	1975	0.000E+00	0.000E+00	-- *	4.635E+04	0.00000	1.000E+00		
18	1976	0.000E+00	0.000E+00	-- *	4.712E+04	0.00000	1.000E+00		
19	1977	0.000E+00	0.000E+00	-- *	4.724E+04	0.00000	1.000E+00		
20	1978	0.000E+00	0.000E+00	-- *	4.833E+04	0.00000	1.000E+00		
21	1979	0.000E+00	0.000E+00	-- *	5.051E+04	0.00000	1.000E+00		
22	1980	0.000E+00	0.000E+00	-- *	5.215E+04	0.00000	1.000E+00		
23	1981	0.000E+00	0.000E+00	-- *	5.326E+04	0.00000	1.000E+00		
24	1982	0.000E+00	0.000E+00	-- *	5.234E+04	0.00000	1.000E+00		
25	1983	0.000E+00	0.000E+00	-- *	5.212E+04	0.00000	1.000E+00		
26	1984	0.000E+00	0.000E+00	-- *	5.233E+04	0.00000	1.000E+00		
27	1985	1.000E+00	1.000E+00	--	9.024E+04	5.375E+04	0.51820	1.000E+00	
28	1986	1.000E+00	1.000E+00	--	3.657E+04	5.371E+04	-0.38451	1.000E+00	
29	1987	0.000E+00	0.000E+00	-- *	4.829E+04	0.00000	1.000E+00		
30	1988	0.000E+00	0.000E+00	-- *	3.389E+04	0.00000	1.000E+00		
31	1989	0.000E+00	0.000E+00	-- *	2.525E+04	0.00000	1.000E+00		
32	1990	1.000E+00	1.000E+00	--	1.820E+04	2.083E+04	-0.13470	1.000E+00	
33	1991	0.000E+00	0.000E+00	-- *	1.702E+04	0.00000	1.000E+00		
34	1992	0.000E+00	0.000E+00	-- *	1.352E+04	0.00000	1.000E+00		
35	1993	0.000E+00	0.000E+00	-- *	8.985E+03	0.00000	1.000E+00		
36	1994	0.000E+00	0.000E+00	-- *	5.170E+03	0.00000	1.000E+00		
37	1995	0.000E+00	0.000E+00	-- *	4.816E+03	0.00000	1.000E+00		
38	1996	0.000E+00	0.000E+00	-- *	5.410E+03	0.00000	1.000E+00		
39	1997	0.000E+00	0.000E+00	-- *	6.550E+03	0.00000	1.000E+00		
40	1998	0.000E+00	0.000E+00	-- *	7.884E+03	0.00000	1.000E+00		
41	1999	0.000E+00	0.000E+00	-- *	9.418E+03	0.00000	1.000E+00		
42	2000	0.000E+00	0.000E+00	-- *	1.087E+04	0.00000	1.000E+00		
43	2001	0.000E+00	0.000E+00	-- *	1.237E+04	0.00000	1.000E+00		
44	2002	0.000E+00	0.000E+00	-- *	1.460E+04	0.00000	1.000E+00		
45	2003	0.000E+00	0.000E+00	-- *	1.725E+04	0.00000	1.000E+00		
46	2004	0.000E+00	0.000E+00	-- *	2.027E+04	0.00000	1.000E+00		
47	2005	0.000E+00	0.000E+00	-- *	2.387E+04	0.00000	1.000E+00		
48	2006	0.000E+00	0.000E+00	-- *	2.788E+04	0.00000	1.000E+00		
49	2007	0.000E+00	0.000E+00	-- *	3.230E+04	0.00000	1.000E+00		
50	2008	0.000E+00	0.000E+00	-- *	3.673E+04	0.00000	1.000E+00		
51	2009	0.000E+00	0.000E+00	-- *	4.163E+04	0.00000	1.000E+00		
52	2010	0.000E+00	0.000E+00	-- *	4.652E+04	0.00000	1.000E+00		
53	2011	0.000E+00	0.000E+00	-- *	5.066E+04	0.00000	1.000E+00		
54	2012	0.000E+00	0.000E+00	-- *	5.481E+04	0.00000	1.000E+00		
55	2013	0.000E+00	0.000E+00	-- *	5.864E+04	0.00000	1.000E+00		
56	2014	0.000E+00	0.000E+00	-- *	6.178E+04	0.00000	1.000E+00		
57	2015	0.000E+00	0.000E+00	-- *	6.482E+04	0.00000	1.000E+00		
58	2016	0.000E+00	0.000E+00	-- *	6.647E+04	0.00000	1.000E+00		
59	2017	0.000E+00	0.000E+00	-- *	6.844E+04	0.00000	1.000E+00		

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 5



RESULTS FOR DATA SERIES # 6 (NON-BOOTSTRAPPED)

3L summer survey

Data type 11: Abundance index (annual average)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid log index	weight	Statistic
1	1959	0.000E+00	0.000E+00	-- *	2.499E+05	0.00000	1.000E+00		
2	1960	0.000E+00	0.000E+00	-- *	2.339E+05	0.00000	1.000E+00		
3	1961	0.000E+00	0.000E+00	-- *	2.290E+05	0.00000	1.000E+00		
4	1962	0.000E+00	0.000E+00	-- *	2.268E+05	0.00000	1.000E+00		
5	1963	0.000E+00	0.000E+00	-- *	2.225E+05	0.00000	1.000E+00		
6	1964	0.000E+00	0.000E+00	-- *	2.242E+05	0.00000	1.000E+00		
7	1965	0.000E+00	0.000E+00	-- *	2.274E+05	0.00000	1.000E+00		
8	1966	0.000E+00	0.000E+00	-- *	2.274E+05	0.00000	1.000E+00		
9	1967	0.000E+00	0.000E+00	-- *	2.253E+05	0.00000	1.000E+00		
10	1968	0.000E+00	0.000E+00	-- *	2.233E+05	0.00000	1.000E+00		
11	1969	0.000E+00	0.000E+00	-- *	2.223E+05	0.00000	1.000E+00		
12	1970	0.000E+00	0.000E+00	-- *	2.231E+05	0.00000	1.000E+00		
13	1971	0.000E+00	0.000E+00	-- *	2.188E+05	0.00000	1.000E+00		
14	1972	0.000E+00	0.000E+00	-- *	2.078E+05	0.00000	1.000E+00		
15	1973	0.000E+00	0.000E+00	-- *	1.974E+05	0.00000	1.000E+00		
16	1974	0.000E+00	0.000E+00	-- *	1.907E+05	0.00000	1.000E+00		
17	1975	0.000E+00	0.000E+00	-- *	1.916E+05	0.00000	1.000E+00		
18	1976	0.000E+00	0.000E+00	-- *	1.934E+05	0.00000	1.000E+00		
19	1977	0.000E+00	0.000E+00	-- *	1.960E+05	0.00000	1.000E+00		
20	1978	1.000E+00	1.000E+00	--	3.112E+05	2.027E+05	0.42863	1.000E+00	
21	1979	1.000E+00	1.000E+00	--	2.278E+05	2.105E+05	0.07893	1.000E+00	
22	1980	0.000E+00	0.000E+00	-- *	2.161E+05	0.00000	1.000E+00		
23	1981	1.000E+00	1.000E+00	--	2.614E+05	2.165E+05	0.18861	1.000E+00	
24	1982	0.000E+00	0.000E+00	-- *	2.141E+05	0.00000	1.000E+00		
25	1983	0.000E+00	0.000E+00	-- *	2.141E+05	0.00000	1.000E+00		
26	1984	1.000E+00	1.000E+00	--	2.777E+05	2.175E+05	0.24429	1.000E+00	
27	1985	1.000E+00	1.000E+00	--	1.610E+05	2.203E+05	-0.31334	1.000E+00	
28	1986	0.000E+00	0.000E+00	-- *	2.087E+05	0.00000	1.000E+00		
29	1987	0.000E+00	0.000E+00	-- *	1.662E+05	0.00000	1.000E+00		
30	1988	0.000E+00	0.000E+00	-- *	1.201E+05	0.00000	1.000E+00		
31	1989	0.000E+00	0.000E+00	-- *	9.407E+04	0.00000	1.000E+00		
32	1990	1.000E+00	1.000E+00	--	9.284E+04	7.727E+04	0.18352	1.000E+00	
33	1991	1.000E+00	1.000E+00	--	3.757E+04	6.229E+04	-0.50550	1.000E+00	
34	1992	0.000E+00	0.000E+00	-- *	4.546E+04	0.00000	1.000E+00		
35	1993	1.000E+00	1.000E+00	--	2.084E+04	2.828E+04	-0.30535	1.000E+00	
36	1994	0.000E+00	0.000E+00	-- *	2.046E+04	0.00000	1.000E+00		
37	1995	0.000E+00	0.000E+00	-- *	2.094E+04	0.00000	1.000E+00		
38	1996	0.000E+00	0.000E+00	-- *	2.445E+04	0.00000	1.000E+00		
39	1997	0.000E+00	0.000E+00	-- *	2.951E+04	0.00000	1.000E+00		
40	1998	0.000E+00	0.000E+00	-- *	3.539E+04	0.00000	1.000E+00		
41	1999	0.000E+00	0.000E+00	-- *	4.153E+04	0.00000	1.000E+00		
42	2000	0.000E+00	0.000E+00	-- *	4.760E+04	0.00000	1.000E+00		
43	2001	0.000E+00	0.000E+00	-- *	5.519E+04	0.00000	1.000E+00		
44	2002	0.000E+00	0.000E+00	-- *	6.517E+04	0.00000	1.000E+00		
45	2003	0.000E+00	0.000E+00	-- *	7.679E+04	0.00000	1.000E+00		
46	2004	0.000E+00	0.000E+00	-- *	9.035E+04	0.00000	1.000E+00		
47	2005	0.000E+00	0.000E+00	-- *	1.060E+05	0.00000	1.000E+00		
48	2006	0.000E+00	0.000E+00	-- *	1.232E+05	0.00000	1.000E+00		
49	2007	0.000E+00	0.000E+00	-- *	1.414E+05	0.00000	1.000E+00		
50	2008	0.000E+00	0.000E+00	-- *	1.606E+05	0.00000	1.000E+00		
51	2009	0.000E+00	0.000E+00	-- *	1.807E+05	0.00000	1.000E+00		
52	2010	0.000E+00	0.000E+00	-- *	1.993E+05	0.00000	1.000E+00		
53	2011	0.000E+00	0.000E+00	-- *	2.163E+05	0.00000	1.000E+00		
54	2012	0.000E+00	0.000E+00	-- *	2.327E+05	0.00000	1.000E+00		
55	2013	0.000E+00	0.000E+00	-- *	2.470E+05	0.00000	1.000E+00		
56	2014	0.000E+00	0.000E+00	-- *	2.596E+05	0.00000	1.000E+00		
57	2015	0.000E+00	0.000E+00	-- *	2.692E+05	0.00000	1.000E+00		
58	2016	0.000E+00	0.000E+00	-- *	2.767E+05	0.00000	1.000E+00		
59	2017	0.000E+00	0.000E+00	-- *	2.827E+05	0.00000	1.000E+00		

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 6



RESULTS FOR DATA SERIES # 7 (NON-BOOTSTRAPPED)

3L autumn survey

Data type I2: Abundance index (end of year)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid log index	weight	Statistic
1	1959	0.000E+00	0.000E+00	-- *	5.354E+04	0.00000	1.000E+00		
2	1960	0.000E+00	0.000E+00	-- *	5.201E+04	0.00000	1.000E+00		
3	1961	0.000E+00	0.000E+00	-- *	5.129E+04	0.00000	1.000E+00		
4	1962	0.000E+00	0.000E+00	-- *	5.100E+04	0.00000	1.000E+00		
5	1963	0.000E+00	0.000E+00	-- *	4.940E+04	0.00000	1.000E+00		
6	1964	0.000E+00	0.000E+00	-- *	5.168E+04	0.00000	1.000E+00		
7	1965	0.000E+00	0.000E+00	-- *	5.092E+04	0.00000	1.000E+00		
8	1966	0.000E+00	0.000E+00	-- *	5.163E+04	0.00000	1.000E+00		
9	1967	0.000E+00	0.000E+00	-- *	5.005E+04	0.00000	1.000E+00		
10	1968	0.000E+00	0.000E+00	-- *	5.065E+04	0.00000	1.000E+00		
11	1969	0.000E+00	0.000E+00	-- *	4.964E+04	0.00000	1.000E+00		
12	1970	0.000E+00	0.000E+00	-- *	5.098E+04	0.00000	1.000E+00		
13	1971	0.000E+00	0.000E+00	-- *	4.782E+04	0.00000	1.000E+00		
14	1972	0.000E+00	0.000E+00	-- *	4.596E+04	0.00000	1.000E+00		
15	1973	0.000E+00	0.000E+00	-- *	4.315E+04	0.00000	1.000E+00		
16	1974	0.000E+00	0.000E+00	-- *	4.286E+04	0.00000	1.000E+00		
17	1975	0.000E+00	0.000E+00	-- *	4.357E+04	0.00000	1.000E+00		
18	1976	0.000E+00	0.000E+00	-- *	4.368E+04	0.00000	1.000E+00		
19	1977	0.000E+00	0.000E+00	-- *	4.469E+04	0.00000	1.000E+00		
20	1978	0.000E+00	0.000E+00	-- *	4.670E+04	0.00000	1.000E+00		
21	1979	0.000E+00	0.000E+00	-- *	4.822E+04	0.00000	1.000E+00		
22	1980	0.000E+00	0.000E+00	-- *	4.925E+04	0.00000	1.000E+00		
23	1981	0.000E+00	0.000E+00	-- *	4.840E+04	0.00000	1.000E+00		
24	1982	0.000E+00	0.000E+00	-- *	4.819E+04	0.00000	1.000E+00		
25	1983	0.000E+00	0.000E+00	-- *	4.839E+04	0.00000	1.000E+00		
26	1984	0.000E+00	0.000E+00	-- *	4.970E+04	0.00000	1.000E+00		
27	1985	1.000E+00	1.000E+00	--	9.823E+04	4.967E+04	0.68199	1.000E+00	
28	1986	1.000E+00	1.000E+00	--	1.712E+04	4.465E+04	-0.95859	1.000E+00	
29	1987	0.000E+00	0.000E+00	-- *	3.133E+04	0.00000	1.000E+00		
30	1988	0.000E+00	0.000E+00	-- *	2.334E+04	0.00000	1.000E+00		
31	1989	0.000E+00	0.000E+00	-- *	1.926E+04	0.00000	1.000E+00		
32	1990	1.000E+00	1.000E+00	--	2.074E+04	1.574E+04	0.27588	1.000E+00	
33	1991	0.000E+00	0.000E+00	-- *	1.250E+04	0.00000	1.000E+00		
34	1992	0.000E+00	0.000E+00	-- *	8.308E+03	0.00000	1.000E+00		
35	1993	0.000E+00	0.000E+00	-- *	4.780E+03	0.00000	1.000E+00		
36	1994	0.000E+00	0.000E+00	-- *	4.454E+03	0.00000	1.000E+00		
37	1995	0.000E+00	0.000E+00	-- *	5.002E+03	0.00000	1.000E+00		
38	1996	0.000E+00	0.000E+00	-- *	6.057E+03	0.00000	1.000E+00		
39	1997	0.000E+00	0.000E+00	-- *	7.290E+03	0.00000	1.000E+00		
40	1998	0.000E+00	0.000E+00	-- *	8.708E+03	0.00000	1.000E+00		
41	1999	0.000E+00	0.000E+00	-- *	1.005E+04	0.00000	1.000E+00		
42	2000	0.000E+00	0.000E+00	-- *	1.144E+04	0.00000	1.000E+00		
43	2001	0.000E+00	0.000E+00	-- *	1.350E+04	0.00000	1.000E+00		
44	2002	0.000E+00	0.000E+00	-- *	1.595E+04	0.00000	1.000E+00		
45	2003	0.000E+00	0.000E+00	-- *	1.874E+04	0.00000	1.000E+00		
46	2004	0.000E+00	0.000E+00	-- *	2.207E+04	0.00000	1.000E+00		
47	2005	0.000E+00	0.000E+00	-- *	2.578E+04	0.00000	1.000E+00		
48	2006	0.000E+00	0.000E+00	-- *	2.986E+04	0.00000	1.000E+00		
49	2007	0.000E+00	0.000E+00	-- *	3.396E+04	0.00000	1.000E+00		
50	2008	0.000E+00	0.000E+00	-- *	3.849E+04	0.00000	1.000E+00		
51	2009	0.000E+00	0.000E+00	-- *	4.302E+04	0.00000	1.000E+00		
52	2010	0.000E+00	0.000E+00	-- *	4.684E+04	0.00000	1.000E+00		
53	2011	0.000E+00	0.000E+00	-- *	5.068E+04	0.00000	1.000E+00		
54	2012	0.000E+00	0.000E+00	-- *	5.422E+04	0.00000	1.000E+00		
55	2013	0.000E+00	0.000E+00	-- *	5.712E+04	0.00000	1.000E+00		
56	2014	0.000E+00	0.000E+00	-- *	5.993E+04	0.00000	1.000E+00		
57	2015	0.000E+00	0.000E+00	-- *	6.146E+04	0.00000	1.000E+00		
58	2016	0.000E+00	0.000E+00	-- *	6.328E+04	0.00000	1.000E+00		
59	2017	0.000E+00	0.000E+00	-- *	6.419E+04	0.00000	1.000E+00		

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 7



RESULTS FOR DATA SERIES # 8 (NON-BOOTSTRAPPED)

3N spanish survey

Data type I1: Abundance index (annual average)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid in log index	Statist weight
1	1959	0.000E+00	0.000E+00	-- *	1.902E+05	0.00000	1.000E+00	
2	1960	0.000E+00	0.000E+00	-- *	1.780E+05	0.00000	1.000E+00	
3	1961	0.000E+00	0.000E+00	-- *	1.742E+05	0.00000	1.000E+00	
4	1962	0.000E+00	0.000E+00	-- *	1.726E+05	0.00000	1.000E+00	
5	1963	0.000E+00	0.000E+00	-- *	1.693E+05	0.00000	1.000E+00	
6	1964	0.000E+00	0.000E+00	-- *	1.706E+05	0.00000	1.000E+00	
7	1965	0.000E+00	0.000E+00	-- *	1.730E+05	0.00000	1.000E+00	
8	1966	0.000E+00	0.000E+00	-- *	1.730E+05	0.00000	1.000E+00	
9	1967	0.000E+00	0.000E+00	-- *	1.715E+05	0.00000	1.000E+00	
10	1968	0.000E+00	0.000E+00	-- *	1.699E+05	0.00000	1.000E+00	
11	1969	0.000E+00	0.000E+00	-- *	1.692E+05	0.00000	1.000E+00	
12	1970	0.000E+00	0.000E+00	-- *	1.698E+05	0.00000	1.000E+00	
13	1971	0.000E+00	0.000E+00	-- *	1.665E+05	0.00000	1.000E+00	
14	1972	0.000E+00	0.000E+00	-- *	1.581E+05	0.00000	1.000E+00	
15	1973	0.000E+00	0.000E+00	-- *	1.502E+05	0.00000	1.000E+00	
16	1974	0.000E+00	0.000E+00	-- *	1.451E+05	0.00000	1.000E+00	
17	1975	0.000E+00	0.000E+00	-- *	1.458E+05	0.00000	1.000E+00	
18	1976	0.000E+00	0.000E+00	-- *	1.472E+05	0.00000	1.000E+00	
19	1977	0.000E+00	0.000E+00	-- *	1.491E+05	0.00000	1.000E+00	
20	1978	0.000E+00	0.000E+00	-- *	1.542E+05	0.00000	1.000E+00	
21	1979	0.000E+00	0.000E+00	-- *	1.602E+05	0.00000	1.000E+00	
22	1980	0.000E+00	0.000E+00	-- *	1.645E+05	0.00000	1.000E+00	
23	1981	0.000E+00	0.000E+00	-- *	1.647E+05	0.00000	1.000E+00	
24	1982	0.000E+00	0.000E+00	-- *	1.629E+05	0.00000	1.000E+00	
25	1983	0.000E+00	0.000E+00	-- *	1.629E+05	0.00000	1.000E+00	
26	1984	0.000E+00	0.000E+00	-- *	1.655E+05	0.00000	1.000E+00	
27	1985	0.000E+00	0.000E+00	-- *	1.676E+05	0.00000	1.000E+00	
28	1986	0.000E+00	0.000E+00	-- *	1.588E+05	0.00000	1.000E+00	
29	1987	0.000E+00	0.000E+00	-- *	1.265E+05	0.00000	1.000E+00	
30	1988	0.000E+00	0.000E+00	-- *	9.142E+04	0.00000	1.000E+00	
31	1989	0.000E+00	0.000E+00	-- *	7.158E+04	0.00000	1.000E+00	
32	1990	0.000E+00	0.000E+00	-- *	5.880E+04	0.00000	1.000E+00	
33	1991	0.000E+00	0.000E+00	-- *	4.740E+04	0.00000	1.000E+00	
34	1992	0.000E+00	0.000E+00	-- *	3.459E+04	0.00000	1.000E+00	
35	1993	0.000E+00	0.000E+00	-- *	2.152E+04	0.00000	1.000E+00	
36	1994	0.000E+00	0.000E+00	-- *	1.557E+04	0.00000	1.000E+00	
37	1995	1.000E+00	1.000E+00	--	4.608E+04	1.594E+04	1.06189 1.000E+00	
38	1996	1.000E+00	1.000E+00	--	6.558E+03	1.860E+04	-1.04268 1.000E+00	
39	1997	1.000E+00	1.000E+00	--	4.753E+03	2.246E+04	-1.55286 1.000E+00	
40	1998	1.000E+00	1.000E+00	--	2.254E+04	2.693E+04	-0.17783 1.000E+00	
41	1999	1.000E+00	1.000E+00	--	4.646E+04	3.160E+04	0.38532 1.000E+00	
42	2000	1.000E+00	1.000E+00	--	6.893E+04	3.622E+04	0.64347 1.000E+00	
43	2001	1.000E+00	1.000E+00	--	5.386E+04	4.200E+04	0.24870 1.000E+00	
44	2002	1.000E+00	1.000E+00	--	7.620E+03	4.959E+04	-1.87301 1.000E+00	
45	2003	1.000E+00	1.000E+00	--	1.103E+04	5.843E+04	-1.66715 1.000E+00	
46	2004	1.000E+00	1.000E+00	--	2.702E+04	6.875E+04	-0.93403 1.000E+00	
47	2005	1.000E+00	1.000E+00	--	1.469E+05	8.063E+04	0.60004 1.000E+00	
48	2006	1.000E+00	1.000E+00	--	8.783E+04	9.378E+04	-0.06559 1.000E+00	
49	2007	1.000E+00	1.000E+00	--	8.760E+04	1.076E+05	-0.20580 1.000E+00	
50	2008	1.000E+00	1.000E+00	--	6.806E+04	1.222E+05	-0.58518 1.000E+00	
51	2009	1.000E+00	1.000E+00	--	7.357E+05	1.375E+05	1.67724 1.000E+00	
52	2010	1.000E+00	1.000E+00	--	3.595E+05	1.516E+05	0.86338 1.000E+00	
53	2011	1.000E+00	1.000E+00	--	4.183E+05	1.646E+05	0.93283 1.000E+00	
54	2012	1.000E+00	1.000E+00	--	2.652E+05	1.770E+05	0.40428 1.000E+00	
55	2013	1.000E+00	1.000E+00	--	4.295E+05	1.879E+05	0.82671 1.000E+00	
56	2014	1.000E+00	1.000E+00	--	1.780E+05	1.976E+05	-0.10400 1.000E+00	
57	2015	1.000E+00	1.000E+00	--	5.235E+05	2.049E+05	0.93813 1.000E+00	
58	2016	1.000E+00	1.000E+00	--	1.173E+05	2.105E+05	-0.58512 1.000E+00	
59	2017	1.000E+00	1.000E+00	--	2.659E+05	2.151E+05	0.21202 1.000E+00	

* Asterisk indicates missing value(s).



UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 8



RESULTS FOR DATA SERIES # 9 (NON-BOOTSTRAPPED)

3L spanish survey

Data type I1: Abundance index (annual average)


Series weight: 1.000

Obs	Year	Observed effort	Estimated effort	Estim F	Observed index	Model index	Resid log index	weight	Statistic
1	1959	0.000E+00	0.000E+00	-- *	1.492E+05	0.00000	1.000E+00		
2	1960	0.000E+00	0.000E+00	-- *	1.396E+05	0.00000	1.000E+00		
3	1961	0.000E+00	0.000E+00	-- *	1.367E+05	0.00000	1.000E+00		
4	1962	0.000E+00	0.000E+00	-- *	1.354E+05	0.00000	1.000E+00		
5	1963	0.000E+00	0.000E+00	-- *	1.328E+05	0.00000	1.000E+00		
6	1964	0.000E+00	0.000E+00	-- *	1.338E+05	0.00000	1.000E+00		
7	1965	0.000E+00	0.000E+00	-- *	1.358E+05	0.00000	1.000E+00		
8	1966	0.000E+00	0.000E+00	-- *	1.357E+05	0.00000	1.000E+00		
9	1967	0.000E+00	0.000E+00	-- *	1.345E+05	0.00000	1.000E+00		
10	1968	0.000E+00	0.000E+00	-- *	1.333E+05	0.00000	1.000E+00		
11	1969	0.000E+00	0.000E+00	-- *	1.327E+05	0.00000	1.000E+00		
12	1970	0.000E+00	0.000E+00	-- *	1.332E+05	0.00000	1.000E+00		
13	1971	0.000E+00	0.000E+00	-- *	1.306E+05	0.00000	1.000E+00		
14	1972	0.000E+00	0.000E+00	-- *	1.240E+05	0.00000	1.000E+00		
15	1973	0.000E+00	0.000E+00	-- *	1.178E+05	0.00000	1.000E+00		
16	1974	0.000E+00	0.000E+00	-- *	1.138E+05	0.00000	1.000E+00		
17	1975	0.000E+00	0.000E+00	-- *	1.144E+05	0.00000	1.000E+00		
18	1976	0.000E+00	0.000E+00	-- *	1.155E+05	0.00000	1.000E+00		
19	1977	0.000E+00	0.000E+00	-- *	1.170E+05	0.00000	1.000E+00		
20	1978	0.000E+00	0.000E+00	-- *	1.210E+05	0.00000	1.000E+00		
21	1979	0.000E+00	0.000E+00	-- *	1.257E+05	0.00000	1.000E+00		
22	1980	0.000E+00	0.000E+00	-- *	1.290E+05	0.00000	1.000E+00		
23	1981	0.000E+00	0.000E+00	-- *	1.292E+05	0.00000	1.000E+00		
24	1982	0.000E+00	0.000E+00	-- *	1.278E+05	0.00000	1.000E+00		
25	1983	0.000E+00	0.000E+00	-- *	1.278E+05	0.00000	1.000E+00		
26	1984	0.000E+00	0.000E+00	-- *	1.298E+05	0.00000	1.000E+00		
27	1985	0.000E+00	0.000E+00	-- *	1.315E+05	0.00000	1.000E+00		
28	1986	0.000E+00	0.000E+00	-- *	1.246E+05	0.00000	1.000E+00		
29	1987	0.000E+00	0.000E+00	-- *	9.923E+04	0.00000	1.000E+00		
30	1988	0.000E+00	0.000E+00	-- *	7.172E+04	0.00000	1.000E+00		
31	1989	0.000E+00	0.000E+00	-- *	5.616E+04	0.00000	1.000E+00		
32	1990	0.000E+00	0.000E+00	-- *	4.613E+04	0.00000	1.000E+00		
33	1991	0.000E+00	0.000E+00	-- *	3.718E+04	0.00000	1.000E+00		
34	1992	0.000E+00	0.000E+00	-- *	2.713E+04	0.00000	1.000E+00		
35	1993	0.000E+00	0.000E+00	-- *	1.688E+04	0.00000	1.000E+00		
36	1994	0.000E+00	0.000E+00	-- *	1.221E+04	0.00000	1.000E+00		
37	1995	0.000E+00	0.000E+00	-- *	1.250E+04	0.00000	1.000E+00		
38	1996	0.000E+00	0.000E+00	-- *	1.459E+04	0.00000	1.000E+00		
39	1997	0.000E+00	0.000E+00	-- *	1.762E+04	0.00000	1.000E+00		
40	1998	0.000E+00	0.000E+00	-- *	2.112E+04	0.00000	1.000E+00		
41	1999	0.000E+00	0.000E+00	-- *	2.479E+04	0.00000	1.000E+00		
42	2000	0.000E+00	0.000E+00	-- *	2.841E+04	0.00000	1.000E+00		
43	2001	0.000E+00	0.000E+00	-- *	3.295E+04	0.00000	1.000E+00		
44	2002	0.000E+00	0.000E+00	-- *	3.890E+04	0.00000	1.000E+00		
45	2003	0.000E+00	0.000E+00	-- *	4.584E+04	0.00000	1.000E+00		
46	2004	0.000E+00	0.000E+00	-- *	5.393E+04	0.00000	1.000E+00		
47	2005	0.000E+00	0.000E+00	-- *	6.325E+04	0.00000	1.000E+00		
48	2006	1.000E+00	1.000E+00	--	7.007E+04	7.357E+04	-0.04881	1.000E+00	
49	2007	1.000E+00	1.000E+00	--	3.141E+04	8.443E+04	-0.98874	1.000E+00	
50	2008	1.000E+00	1.000E+00	--	7.557E+04	9.585E+04	-0.23780	1.000E+00	
51	2009	1.000E+00	1.000E+00	--	1.037E+05	1.079E+05	-0.03959	1.000E+00	
52	2010	1.000E+00	1.000E+00	--	2.667E+05	1.190E+05	0.80761	1.000E+00	
53	2011	1.000E+00	1.000E+00	--	1.706E+05	1.291E+05	0.27886	1.000E+00	
54	2012	1.000E+00	1.000E+00	--	4.815E+05	1.389E+05	1.24323	1.000E+00	
55	2013	1.000E+00	1.000E+00	--	2.352E+05	1.474E+05	0.46702	1.000E+00	
56	2014	1.000E+00	1.000E+00	--	2.164E+05	1.550E+05	0.33385	1.000E+00	
57	2015	1.000E+00	1.000E+00	--	1.304E+05	1.607E+05	-0.20883	1.000E+00	
58	2016	1.000E+00	1.000E+00	--	9.881E+04	1.652E+05	-0.51369	1.000E+00	
59	2017	1.000E+00	1.000E+00	--	5.656E+04	1.687E+05	-1.09311	1.000E+00	



* Asterisk indicates missing value(s).







UNWEIGHTED LOG RESIDUAL PLOT FOR DATA SERIES # 9





3LN redfish











3LN redfish

Page 26



Time Plot of Estimated F/Fmsy and B/Bmsy (dashed line = 1.0)



Elapsed time: 0 hours, 0 minutes, 5.538 seconds.



Appendix 3 ASPIC BOT 2018 results (some 2016 key parameters and diagnostics in brackets)

3LN redfish

Page 1

Friday, 08 Jun 2018 at 19:20:52

ASPIC -- A Surplus-Production Model Including Covariates (BETA Ver. 7.03)

BOT program mode

Author: Michael H. Prager

LOGISTIC model mode

Prager Consulting

YLD conditioning

<http://www.mhprager.com>

SSE optimization

Reference: Prager, M. H. 1994. A suite of extensions to a nonequilibrium surplus-production model. Fishery Bulletin 92: 374-389. ASPIC program and user's guide available gratis at www.mhprager.com

CONTROL PARAMETERS (FROM INPUT FILE) Input file: C:/...LN/2018/ASPIC18 bot 60CL 14200 prj/ASPIC18qmaxboundbot.a7inp

Operation of ASPIC: Fit logistic (Schaefer) model by direct optimization with bootstrap.

Number of years analyzed: 59 Number of bootstrap trials: 1000
 Number of data series: 9 Objective function: Least squares
 Relative conv. criterion (simplex): 1.000E-08 Monte Carlo search mode, trials: 0 20000
 Relative conv. criterion (restart): 3.000E-08 Random number seed: 3941285
 Relative conv. criterion (effort): 1.000E-04 Identical convergences required in fitting: 18
 Maximum F allowed in fitting: 6.000

3LN redfish

Page13

ESTIMATES FROM BOOTSTRAP ANALYSIS

(Notation X. means terminal estimate of X)

Param name	Point estimate	Bias-corrected approximate confidence limits			Inter-quartile		Relative range	2016 IQ range
		80% lower	80% upper	60% lower	60% upper			
B1/K	6.976E-01 (6.874E-01)	5.683E-01	1.090E+00	5.984E-01	8.817E-01	2.186E-01	0.313	(0.256)
MSY	2.100E+04	NA	NA	NA	NA	NA	NA	
Fmsy	1.122E-01 (1.116E-01)	9.290E-02	1.264E-01	9.896E-02	1.220E-01	1.952E-02	0.174	(0.164)
q(1)	8.243E-06	7.220E-06	9.836E-06	7.660E-06	9.374E-06	1.327E-06	0.161	(0.166)
q(2)	7.814E-01	5.713E-01	1.026E+00	6.237E-01	9.188E-01	2.313E-01	0.296	(0.334)
q(3)	1.299E+00	9.505E-01	1.705E+00	1.054E+00	1.525E+00	3.836E-01	0.295	(0.344)
q(4)	3.129E-01	2.259E-01	4.118E-01	2.533E-01	3.802E-01	1.001E-01	0.320	(0.336)
q(5)	2.461E-01	1.356E-01	3.790E-01	1.639E-01	3.197E-01	1.241E-01	0.504	(0.509)
q(6)	1.009E+00	7.136E-01	1.367E+00	8.120E-01	1.235E+00	3.282E-01	0.325	(0.181)
q(7)	2.276E-01	1.246E-01	3.494E-01	1.527E-01	2.993E-01	1.160E-01	0.510	(0.511)
q(8)	7.679E-01	5.298E-01	9.987E-01	5.912E-01	8.937E-01	2.352E-01	0.306	(0.340)
q(9)	6.024E-01	4.270E-01	8.056E-01	4.787E-01	7.201E-01	1.926E-01	0.320	(0.373)
Ye(2018)	1.560E+04	1.204E+04	2.033E+04	1.289E+04	1.906E+04	4.907E+03	0.315	
Ye(2016)	(1.782E+04)					(0.263)		
Y.(Fmsy)	3.082E+04 (2.858E+04)	2.409E+04	3.343E+04	2.691E+04	3.284E+04	4.555E+03	0.148	(0.218)
Bmsy	1.871E+05 (1.882E+05)	1.662E+05	2.261E+05	1.722E+05	2.122E+05	3.394E+04	0.181	(0.172)



Param name	Bias-corrected approximate confidence limits				Inter- quartile		Relative range	2016 IQ range
	Point estimate	80% lower	80% upper	60% lower	60% upper			
fmsy(1)	1.361E+04	1.084E+04	1.678E+04	1.170E+04	1.565E+04	3.136E+03	0.230	
fmsy(2)	1.436E-01	1.038E-01	2.001E-01	1.192E-01	1.811E-01	4.992E-02	0.348	
fmsy(3)	8.638E-02	6.058E-02	1.171E-01	7.039E-02	1.072E-01	2.897E-02	0.335	
fmsy(4)	3.587E-01	2.631E-01	5.131E-01	2.935E-01	4.560E-01	1.284E-01	0.358	
fmsy(5)	4.559E-01	2.869E-01	8.263E-01	3.394E-01	6.764E-01	2.692E-01	0.590	
fmsy(6)	1.112E-01	8.056E-02	1.568E-01	9.066E-02	1.405E-01	4.023E-02	0.362	
fmsy(7)	4.931E-01	3.192E-01	9.083E-01	3.813E-01	7.511E-01	2.922E-01	0.593	
fmsy(8)	1.461E-01	1.076E-01	2.207E-01	1.222E-01	1.940E-01	5.739E-02	0.393	
fmsy(9)	1.863E-01	1.276E-01	2.646E-01	1.508E-01	2.380E-01	7.101E-02	0.381	
B./Bmsy (1.389E+00)	1.507E+00 (1.389E+00)	1.156E+00	1.654E+00	1.300E+00	1.622E+00 (0.237)	2.433E-01	0.161	
F./Fmsy (3.640E-01)	3.759E-01 (3.640E-01)	3.403E-01	4.955E-01	3.477E-01	4.385E-01 (0.268)	6.753E-02	0.180	
Ye./MSY	7.426E-01	5.735E-01	9.682E-01	6.137E-01	9.078E-01	2.337E-01	0.315	
q2/q1	9.479E+04	6.381E+04	1.204E+05	7.098E+04	1.076E+05	2.906E+04	0.307	
q3/q1	1.576E+05	1.087E+05	2.005E+05	1.193E+05	1.788E+05	4.642E+04	0.295	
q4/q1	3.795E+04	2.658E+04	5.265E+04	3.026E+04	4.727E+04	1.346E+04	0.355	
q5/q1	2.986E+04	1.556E+04	4.613E+04	1.934E+04	3.949E+04	1.630E+04	0.546	
q6/q1	1.224E+05	8.333E+04	1.682E+05	9.380E+04	1.496E+05	4.494E+04	0.367	
q7/q1	2.761E+04	1.487E+04	4.311E+04	1.839E+04	3.676E+04	1.501E+04	0.544	
q8/q1	9.315E+04	6.323E+04	1.205E+05	6.891E+04	1.076E+05	3.023E+04	0.324	
q9/q1	7.307E+04	4.864E+04	9.582E+04	5.467E+04	8.382E+04	2.329E+04	0.319	

INFORMATION FOR REPAST (Prager, Porch, Shertzer, & Caddy. 2003. NAJFM 23: 349-361)

Unitless limit reference point in F (Fmsy/F.): 2.660
 CV of above (from bootstrap distribution): 0.148

NOTES ON BOOTSTRAPPED ESTIMATES:

- Bootstrap results were computed from 1000 trials.
- Results are conditional on parameter bounds in the input file.
- If many trials were replaced, consider relaxing bounds and re-running.
- All bootstrapped intervals are approximate. The statistical literature recommends using at least 1000 trials for accurate 95% intervals. The default 80% intervals used by ASPIC should require fewer trials for equivalent accuracy. Using at least 500 trials is recommended.

Trials replaced for lack of convergence: 0 Trials replaced for MSY at bound or MSY >= K: 0
 Trials replaced for q at bound: 0 (429) Trials replaced for B1/K at bound: 10 (13)
 Trials replaced for Fmsy at bound: 0
 Residual inflation factor: 1.0405 (1.0427)

Elapsed time: 0 hours, 15 minutes, 4.104 seconds.

ASPICP-V4

"2018-2020 ASPIC18qmaxboundbot.bio prj under red3LN HCR"

"ASPIC18qmaxboundbot.bio"

0 "CV on MSY during projections"

BC 1 "bias corrected and smooth CI's"

0 "no years skipped from the start in plots"

1 1 1 "AGRAPH open to plot results; write a .prb file; write a R friendly version of the .prj file"

123456789 "random number seed only used when user CV of MSY is non zero"

14200 YABS "yield in same units as assessment"

18100 YABS

18100 YABS

%% END

