

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Insecta Mundi

Center for Systematic Entomology, Gainesville,  
Florida

---

5-2022

## First record of *Ancognatha erythrodera* (Blanchard, 1846) (Coleoptera: Scarabaeidae: Dynastinae: Cyclocephalini) in the altiplano of Chile

José Mondaca

Servicio Agrícola y Ganadero (SAG), jose.mondaca@sag.gob.cl

Sergio Rothmann

Servicio Agrícola y Ganadero (SAG), sergio.rothmann@sag.gob.cl

Follow this and additional works at: <https://digitalcommons.unl.edu/insectamundi>



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

---

Mondaca, José and Rothmann, Sergio, "First record of *Ancognatha erythrodera* (Blanchard, 1846)  
(Coleoptera: Scarabaeidae: Dynastinae: Cyclocephalini) in the altiplano of Chile" (2022). *Insecta Mundi*.  
1434.

<https://digitalcommons.unl.edu/insectamundi/1434>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

A journal of world insect systematics

# INSECTA MUNDI

---

---

0936

First record of *Ancognatha erythrodera* (Blanchard, 1846)  
(Coleoptera: Scarabaeidae: Dynastinae: Cyclocephalini)  
in the altiplano of Chile

José Mondaca

Servicio Agrícola y Ganadero (SAG)  
Camino La Pólvora Km 12,5  
Valparaíso, Chile

Sergio Rothmann

Servicio Agrícola y Ganadero (SAG)  
Laboratorio de Entomología Complejo Lo Aguirre  
Santiago, Chile

Date of issue: May 27, 2022

Center for Systematic Entomology, Inc., Gainesville, FL

**Mondaca J, Rothmann S. 2022.** First record of *Ancognatha erythrodera* (Blanchard, 1846) (Coleoptera: Scarabaeidae: Dynastinae: Cyclocephalini) in the altiplano of Chile. *Insecta Mundi* 0936: 1–4.

Published on May 27, 2022 by  
**Center for Systematic Entomology, Inc.**  
P.O. Box 141874  
Gainesville, FL 32614-1874 USA  
<http://centerforsystematicentomology.org/>

**INSECTA MUNDI** is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

*Insecta Mundi* is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

**Chief Editor:** David Plotkin, [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Assistant Editor:** Paul E. Skelley, [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Layout Editor:** Robert G. Forsyth

**Editorial Board:** Davide Dal Pos, Oliver Keller, M. J. Paulsen

**Founding Editors:** Ross H. Arnett, Jr., J. H. Frank, Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff

**Review Editors:** Listed on the *Insecta Mundi* webpage

**Printed copies (ISSN 0749-6737) annually deposited in libraries**

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

The Natural History Museum, London, UK

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

**Electronic copies (Online ISSN 1942-1354) in PDF format**

Archived digitally by Portico

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

**Copyright held by the author(s).** This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

# First record of *Ancognatha erythrodera* (Blanchard, 1846) (Coleoptera: Scarabaeidae: Dynastinae: Cyclocephalini) in the altiplano of Chile

José Mondaca

Servicio Agrícola y Ganadero (SAG)  
Camino La Pólvora Km 12,5  
Valparaíso, Chile  
jose.mondaca@sag.gob.cl

Sergio Rothmann

Servicio Agrícola y Ganadero (SAG)  
Laboratorio de Entomología Complejo Lo Aguirre  
Santiago, Chile  
sergio.rothmann@sag.gob.cl

**Abstract.** The presence of *Ancognatha erythrodera* (Blanchard, 1846) (Coleoptera: Scarabaeidae) is confirmed for the first time in Chile based on male and female specimens collected in the extreme north of the country. This is the second species of *Ancognatha* Erichson, 1847 recorded in Chile. Morphological characters, illustrations of male genitalia, male and female habitus photographs of this species, and additional records in Argentina and Bolivia are provided. A map with the collection sites and montane habitats photograph in Chile are included.

**Key words.** Andes, dynastine, species diversity, Visviri.

**ZooBank registration.** urn:lsid:zoobank.org:pub:BF6C7863-E859-4772-B51F-CCC2BEDF69C9

## Introduction

The Dynastinae (Coleoptera: Scarabaeidae) that occur in Chile were comprehensively revised by Ratcliffe et al. (2021), who recorded nine species in six genera. One of these genera, *Ancognatha* Erichson, 1847, includes 23 valid species found from the southwestern United States (Arizona and New Mexico) to northern Argentina, Chile, and Bolivia (Mondaca 2016; Moore et al. 2018a; Paucar-Cabrera and Ratcliffe 2018; Ratcliffe et al. 2021).

In this note, we report the first record of *Ancognatha erythrodera* (Blanchard, 1846) in Chile, based on specimens collected in the altiplanic locality of Visviri, Región de Arica y Parinacota. Previously, *Ancognatha erythrodera* was recorded for the Andes of southern Peru, Bolivia, and northwestern Argentina (Moore et al. 2018b). This new record corresponds to the natural distribution of this species on both sides of the Andes mountain range, which is not surprising due to the similarity of habitats in neighboring localities in Peru, Bolivia, and Argentina.

## Materials and Methods

Six specimens of *A. erythrodera* were studied from Chile and deposited at the entomological collection of the Servicio Agrícola y Ganadero, located in Arica city, Chile (SAGC). The other material examined was borrowed from the collections of the first author (JMEC) and of Víctor Manuel Diéguez's private collection, Santiago, Chile (VMDC).

The male genitalia of one specimen was extracted by relaxing the specimen in hot water, cleansing in a hot solution of KOH at 90°C for 10 minutes, and then gluing on a cardboard point for photographing. Habitus and male genitalia photographs were taken with a Canon DCM510 camera. Geographic coordinates of the collecting sites were recorded using Google Earth Pro (Fig. 5).


## Results

### *Ancognatha erythrodera* (Blanchard, 1846)

(Fig. 1–4)

**Material examined.** 6 specimens (1 male, 5 females) at SAGC from: CHILE, Provincia de Parinacota, Visviri, 28-II-2018 (4 females), 17-VII-2018 (1 male), col. C. Avila; Visviri, 14-II-2022, col. M. Arias (1 female). 5 specimens were collected at an ultraviolet light trap, and 1 specimen at a funnel trap.

**Other material examined.** *Ancognatha erythrodera*: Frontera tripartita Perú, Chile, Bolivia, 3-IV-2015, col. J. Mondaca E. (1 JMEC). ARGENTINA, Jujuy, Iturbe 3.412 m, 7-II-2009, col. V.M. Diéguez y G. Arriagada (1




**Figures 1–4.** *Ancognatha erythrodera* (Blanchard, 1846). 1) Male, dorsal habitus. 2) Female, dorsal habitus. Scale bar: 1.0 mm. 3–4) Parameres. 3) Lateral view. 4) Caudal view. Scale bar: 0.5 mm.



VMDC). BOLIVIA, Departamento Oruro, Caracollo 3790 m, 3–4-II-2008, col. M. Ferrú (2 JMEC, 3 VMDC); Challapata, 1–2-II-2008, col. M. Ferrú (5 VMDC). BOLIVIA, Departamento La Paz, Patacamaya, 4-II-2008, col. M. Ferrú (1 VMDC); Viscachani, 19-III-1995, col. D. Jackson (2 VMDC); Tiahuanaco, 9-II-2008, col. M. Ferrú (1 JMEC).

**Diagnosis** ( $n = 9$ ). The species most similar to *Ancognatha erythrodera* is *A. aymara* Mondaca, but these species have different body color patterns; males and females of *A. aymara* are unicolored, castaneous or reddish-brown, and are smaller (body length 13–19 mm). *Ancognatha erythrodera* is larger (17–27 mm) and dorsally bicolored in both sexes with the pronotum reddish-brown to brown and the elytra dark-brown to black. The pronotum of *A. erythrodera* generally has a longitudinal dark line down the middle that may be either continuous or interrupted; this dark line is entirely absent in *A. aymara*. The head, scutellum, and legs are dark brown in *A. erythrodera* (Fig. 2) and not reddish brown as in *A. aymara*. The head and male genitalia (Fig. 3–4) are similar in both species.

**Distribution and habitat.** This species lives at high elevations (3,300–4,080 m) in the altiplano of Argentina (Jujuy, Tucumán), Bolivia (La Paz, Oruro), Peru (Arequipa, Puno) (Moore et al. 2018b), and northern Chile (Parinacota province) (**new record**) (Fig. 5d). This distribution corresponds to the biogeographic province of the Puna of the Andean Region proposed by Morrone (2015). *Ancognatha erythrodera* has been collected in a semi-arid steppe environment dominated by low and resinous scrubs of *Adesmia* sp. (Fabaceae), *Parastrephia* sp. (Asteraceae), and coarse grasses as *Stipa* sp. and *Poa* sp. (Poaceae) (Fig. 5b, c). The adults are nocturnal and often attracted to lights at night and emerge during the rainy season (highland winter) that occurs between the months



**Figure 5.** Distribution of *Ancognatha erythrodera*. **a–c**) Montane habitats of *A. erythrodera* in northern Chile. **d**) Collecting sites of *Ancognatha erythrodera* in the altiplano of Chile (blue circles). **e**) Location of the Región de Arica y Parinacota in northern Chile.

of December and April. In Chile, adults were collected at an elevation of 4,080 m. The immature stages of this species are not described; the larvae feed on roots of coarse grasses known as “paja brava” and have been cited as a pest of potato in Bolivia (Mellini and Verenini 1986).

**Remarks.** This contribution increases the number of *Ancognatha* species found in Chile to two. The new finding of *A. erythrodera* extends the distributional range to the west by 17 km from the Tripartite milestone (the border between Peru, Chile, and Bolivia; Fig. 5a) and 4.8 km to the east from the Bolivian locality of Charaña (Fig. 5d). *Ancognatha erythrodera* is found at high elevations (3,300–4,080 m) on both sides of the Andes mountain range of Peru, Chile, Bolivia, and Argentina, while *Ancognatha aymara* Mondaca, 2016 is a similar species that lives at slightly lower elevations (3,000–3,600 m) in the Andes of northern Chile and northwestern Argentina (Mondaca 2016, 2020).

## Acknowledgments

We acknowledge the collaboration of Víctor Manuel Diéguez and SAGC, who provided specimens for study. We also thank Marcelo Guerrero for the photographs illustrating this paper. We are grateful to Andrew Smith and Brett Ratcliffe for reviewing this paper.

## Literature Cited

- Mellini E, Verenini M. 1986.** Note di biologia sugli insetti delle aree semiaride dell'Altopiano Boliviano all'inizio della stagione estiva. Bollettino dell'Istituto di Entomologia della Università degli Studi di Bologna 40: 185–214.
- Mondaca J. 2016.** A new, high-elevation species of the genus *Ancognatha* Erichson (Coleoptera: Scarabaeidae: Dynastinae) from Chile. The Coleopterists Bulletin 70: 59–64.
- Mondaca J. 2020.** First records of *Ancognatha aymara* Mondaca, 2016 (Coleoptera: Scarabaeidae: Dynastinae: Cyclocephalini) in Argentina. Insecta Mundi 776: 1–4.
- Moore MR, Cave RD, Branham MA. 2018a.** Synopsis of the cyclocephaline scarab beetles (Coleoptera, Scarabaeidae, Dynastinae). ZooKeys 745: 1–99.
- Moore MR, Cave RD, Branham MA. 2018b.** Annotated catalog and bibliography of the cyclocephaline scarab beetles (Coleoptera, Scarabaeidae, Dynastinae, Cyclocephalini). ZooKeys 745: 101–378.
- Morrone JJ. 2015.** Biogeographical regionalisation of the Andean region. Zootaxa 3936(2): 207–236.
- Paucar-Cabrera A, Ratcliffe BC 2018.** The *Ancognatha* Erichson (Coleoptera: Scarabaeidae: Dynastinae: Cyclocephalini) of Ecuador, with description of a new species. The Coleopterists Bulletin 72(4): 665–687.
- Ratcliffe BC, Cave RD, Mondaca J. 2021.** The dynastine scarab beetles (Coleoptera: Scarabaeidae: Dynastinae) of Chile. The Coleopterists Bulletin 75(2): 279–309.

Received March 22, 2022; accepted May 6, 2022.

Review editor David Plotkin.