

living planet symposium BONN 23-27 May 2022

TAKING THE PULSE OF OUR PLANET FROM SPACE

SMOS sea ice thickness - a review and way forward

Lars Kaleschke | Xiangshan Tian-Kunze | Stefan Hendricks Alfred-Wegener-Institut Helmholtz-Zentrum für Polar und Meeresforschung | Germany Robert Ricker | Norwegian Research Centre | Norway Raffaele Crapolicchio | Serco Italia SpA - for European Space Agency (ESA) | Italy Nairobi 3-4

(อมุรคด

ESA UNCLASSIFIED – For ESA Official Use Only


→ THE EUROPEAN SPACE AGENCY

27.5.2022

14:00


Low-frequency microwave radiometry for sea ice






How to measure sea-ice thickness from space





Parameterization: statistical thickness distribution






Tian-Kunze, X., Kaleschke, L., Maaß, N., Mäkynen, M., Serra, N., Drusch, M., and Krumpen, T.: SMOS-derived thin sea ice thickness: algorithm baseline, product specifications and initial verification, The Cryosphere, 8, 997-1018, 2014.

Level 4 product: SMOS and CryoSat2 combined



- Taking advantage of the complementary thickness retrievals, derived from the CS2 altimeter and the SMOS radiometer
- Weekly Arctic-wide sea-ice thickness fields with reduced relative uncertainties


Ricker, R., et al (2017): A weekly Arctic sea-ice thickness data record from merged CryoSat-2 and SMOS satellite data




12th Arctic season finished 13th Antarctic season started



Available SMOS L3 sea ice thickness products version v3.3 based on L1C v724 for the northern (top) and southern (bottom) hemisphere winter periods.






- SMOS L3 sea ice algorithm v3.3 based on L1C v724
- SMOS & CS2 L4 algorithm v204 based on CryoSat2 baseline E

SMOS & CryoSat2 Arctic Sea Ice Volume Time Series






 \rightarrow LPS22 presentation of Stefan Hendricks

SMOS L3 sea ice thickness extent at the end of the season





 12 yrs SMOS data at the end of the freezing season

15

km²

million


- SMOS sea ice thickness extent >0.025m agrees with NSIDC sea ice index based on SSMIS
- Upward trend over the past 4 years
- Long-term trend shows declining Arctic sea ice
- Negative trend is consistent for ice up to 1m

SMOS sea ice data evolution and quality control



Substantial gain in coverage due to improved RFI flags in L1C data version v724 compared to previous L1C version v620.


RFI-related data gaps filled e.g. in the Atlantic.



L. Kaleschke and X. Tian-Kunze, "SMOS Sea Ice Thickness Data Product Quality Control by Comparison with the Regional Sea Ice Extent," 2021 IEEE International Geoscience and Remote Sensing Symposium IGARSS, 2021, pp. 1110-1113, doi: 10.1109/IGARSS47720.2021.9553630.


Antarctic thin sea ice thickness from SMOS 2010-2020





Tian-Kunze, X., Kaleschke, L. (2021): SMOS-derived sea ice thickness in the Antarctic from 2010 to 2020. https://doi.org/10.1594/PANGAEA.934732

Validation with upward-looking sonar (ULS) < 1m



Overview of validation data. The background shows the SMOS sea-ice thickness averaged from April to October over the years 2010-2020.

Time periodULS $\mu \pm \sigma$ (cm)SMOS $\mu \pm \sigma$ (cm)NRMD(cm)RMSD (cm)Apr. - Oct.46±3249±38210.94+314

Same numbers for v3.2 (L1C v620) and v3.3. (L1C v724).

Summary and way forward



- Complementary sensor synergy has been proven successful: ESA's SMOS low-frequency radiometry together with CryoSat-2 altimetry can be used to measure the sea ice thickness with reduced uncertainty and better data coverage
- SMOS L3 sea ice thickness data v3.3 based on L1C v724 data
 →Sea ice thickness extent
- SMOS and CryoSat2 L4 sea ice thickness data v204 based on baseline E
- →Sea ice volume
- SMOS L3 product for Antarctic sea ice initially validated and operational soon
- Field campaigns such as MOSAiC are necessary for process studies, parameterizations and validation to further improve EO methods
- Ongoing work in ESA-funded project SMOS Expert Support Laboratories (ESL) for further algorithm evolution and improvement of data quality
 - Evolution of L3 & L4 sea ice algorithms
 - Continuing MOSAiC data analysis
 - New metrics for enhanced quality monitoring
 - Extended triple collocation (ETC) for validation
 - More synergy: increasing sampling & grid resolution by including Sentinel-3



Selected recent references



Brogioni, M., Andrews, M. J., Urbini, S., Jezek, K. C., Johnson, J. T., Leduc-Leballeur, M., Macelloni, G., Ackley, S. F., Bringer, A., Brucker, L., Demir, O., Fontanelli, G., Yardim, C., Kaleschke, L., Montomoli, F., Tsang, L., Becagli, S., and Frezzotti, M.: Ice Sheet and Sea Ice Ultrawideband Microwave Airborne eXperiment (ISSIUMAX) in Antarctica: first results from Terra Nova Bay, The Cryosphere Discuss. [preprint], https://doi.org/10.5194/tc-2022-59, in review, 2022.

O. Demir *et al.*, "Studies of Sea-Ice Thickness and Salinity Retrieval Using 0.5–2 GHz Microwave Radiometry," in *IEEE Transactions on Geoscience and Remote Sensing*, vol. 60, pp. 1-12, 2022, Art no. 4304412, doi: 10.1109/TGRS.2022.3168646.

Nicolaus, M, Perovich, DK, Spreen, G, Granskog, MA, Albedyll, LV, Angelopoulos, M, Anhaus, P, Arndt, S, Belter, HJ, Bessonov, V, Birnbaum, G, Brauchle, J, Calmer, R, Cardellach, E, Cheng, B, Clemens-Sewall, D, Dadic, R, Damm, E, de Boer, G, Demir, O, Dethloff, K, Divine, DV, Fong, AA, Fons, S, Frey, MM, Fuchs, N, Gabarró, C, Gerland, S, Goessling, HF, Gradinger, R, Haapala, J, Haas, C, Hamilton, J, Hannula, H-R, Hendricks, S, Herber, A, Heuzé, C, Hoppmann, M, Høyland, KV, Huntemann, M, Hutchings, JK, Hwang, B, Itkin, P, Jacobi, H-W, Jaggi, M, Jutila, A, Kaleschke, L, Katlein, C, Kolabutin, N, Krampe, D, Kristensen, SS, Krumpen, T, Kurtz, N, Lampert, A, Lange, BA, Lei, R, Light, B, Linhardt, F, Liston, GE, Loose, B, Macfarlane, AR, Mahmud, M, Matero, IO, Maus, S, Morgenstern, A, Naderpour, R, Nandan, V, Niubom, A, Oggier, M, Oppelt, N, Pätzold, F, Perron, C, Petrovsky, T, Pirazzini, R, Polashenski, C, Rabe, B, Raphael, IA, Regnery, J, Rex, M, Ricker, R, Riemann-Campe, K, Rinke, A, Rohde, J, Salganik, E, Scharien, RK, Schiller, M, Schneebeli, M, Semmling, M, Shimanchuk, E, Shupe, MD, Smith, MM, Smolyanitsky, V, Sokolov, V, Stanton, T, Stroeve, J, Thielke, L, Timofeeva, A, Tonboe, RT, Tavri, A, Tsamados, M, Wagner, DN, Watkins, D, Webster, M, Wendisch, M. 2022. Overview of the MOSAiC expedition: Snow and sea ice. Elementa: Science of the Anthropocene 10(1). DOI: https://doi.org/10.1525/elementa.2021.000046

Luo, H., Yang, Q., Mu, L., Tian-Kunze, X., Nerger, L., Mazloff, M., . . . Chen, D. (2021). DASSO: A data assimilation system for the Southern Ocean that utilizes both sea-ice concentration and thickness observations. Journal of Glaciology, 67(266), 1235-1240. doi:10.1017/jog.2021.57

O. Demir et al., "Measurements of 540–1740 MHz Brightness Temperatures of Sea Ice During the Winter of the MOSAiC Campaign," in IEEE Transactions on Geoscience and Remote Sensing, vol. 60, pp. 1-11, 2022, Art no. 5302011, doi: 10.1109/TGRS.2021.3105360.

Krumpen, T., von Albedyll, L., Goessling, H. F., Hendricks, S., Juhls, B., Spreen, G., Willmes, S., Belter, H. J., Dethloff, K., Haas, C., Kaleschke, L., Katlein, C., Tian-Kunze, X., Ricker, R., Rostosky, P., Rückert, J., Singha, S., and Sokolova, J.: MOSAiC drift expedition from October 2019 to July 2020: sea ice conditions from space and comparison with previous years, The Cryosphere, 15, 3897–3920, https://doi.org/10.5194/tc-15-3897-2021, 2021.

J. T. Johnson et al., "Microwave Radiometry at Frequencies From 500 to 1400 MHz: An Emerging Technology for Earth Observations," in IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, vol. 14, pp. 4894-4914, 2021, doi: 10.1109/JSTARS.2021.3073286.

→ THE EUROPEAN SPACE AGENCY

13