

Elevational range extension of the marsupial frog, *Gastrotheca marsupiata* (Duméril & Bibron, 1841) (Anura, Hemiphractidae), from southern Peru

Emma Steigerwald^{1,2}, Anton Sorokin³, F. Peter Condori^{4,5}, Y. Jared Guevara^{5,6},
Gumercindo Crispin⁷, Juan C. Chaparro^{4,5}

1 Department of Environmental Science, Policy, and Management, University of California, Berkeley, CA, USA • emma.c.steigerwald@gmail.com
 <https://orcid.org/0000-0003-1027-1421>

2 Museum of Vertebrate Zoology, University of California, Berkeley, CA, USA

3 Department of Biology, East Carolina University, Greenville, NC, USA • antonsrkn@gmail.com  <https://orcid.org/0000-0001-8596-7269>

4 Museo de Biodiversidad del Perú, Cusco, Perú • cfrankpeter@gmail.com  <https://orcid.org/0000-0002-8095-6349> • jchapparouza@yahoo.com
 <https://orcid.org/0000-0001-9121-8600>

5 Museo de Historia Natural de la Universidad Nacional de San Antonio Abad del Cusco, Portal Belén, Cusco, Perú

6 Universidad Nacional de San Antonio Abad del Cusco, Cusco, Perú • jaredyonatan@gmail.com

7 Independent researcher, Pucarami, Cusco, Perú

* Corresponding author

Abstract

We extend by 300 m the known upper elevational range of the Marsupial frog, *Gastrotheca marsupiata* (Duméril & Bibron, 1841), to 4,660 m a.s.l. This record makes *G. marsupiata* the highest occurring frog in its genus, which is already characterized by high-elevation distributions. We suggest that this record may represent a case of climate-induced range shifting and discuss this record in the context of our still limited understanding of how amphibian distributions are being affected by climate change.

Keywords

Amphibian, Andes, Cusco, *Gastrotheca*, Vilcanota mountains

Academic editor: Javier Ernesto Cortés Suárez | Received 8 September 2020 | Accepted 2 January 2021 | Published 1 February 2021

Citation: Steigerwald E, Sorokin A, Condori FP, Guevara YJ, Crispin G, Chaparro JC (2021) Elevational range extension of the marsupial frog, *Gastrotheca marsupiata* (Duméril & Bibron, 1841) (Anura, Hemiphractidae), from southern Peru. Check List 17 (1): 145–150. <https://doi.org/10.15560/17.1.145>

Introduction

The genus *Gastrotheca* (Fitzinger, 1843), known as marsupial frogs, comprises 75 formally described species (Amphibiaweb 2020; Frost 2020). Although some *Gastrotheca* occur in Central America and eastern South

America, their diversity peaks in the Andes (Frost 2020). *Gastrotheca marsupiata* (Duméril & Bibron, 1841), the type species of *Gastrotheca*, is distributed along the Cordillera of the Andes within the Amazonian drainage

systems (Frost 2020). Its type locality is in the Peruvian department of Cusco (Guibé 1950) and its range is one of the largest in the genus, stretching from central Peru to southern Bolivia (Duellman and Trueb 2015; Frost 2020; IUCN 2020). *Gastrotheca marsupiata* occurs in montane cloud forests and treeless, puna habitat characterized by bunchgrasses (Duellman and Trueb 2015) and can persist despite agricultural conversion and forest degradation (IUCN 2020).

Like other marsupial frog species, female *G. marsupiata* have a dorsal pouch where the male places the eggs following fertilization (Dumeril and Bibron 1841; Duellman and Trueb 2015). Fertilization and transfer usually happen away from water, and the eggs hatch in the pouch (Del Pino 1989). In some *Gastrotheca* species, the tadpoles remain in the pouch until metamorphosis, but *G. marsupiata* deposit their tadpoles into stagnant or slow-flowing water to live as free-swimming tadpoles prior to metamorphosis (Duellman and Maness 1980; Sinsch and Joermann 1989). Observations of this and closely related species suggest that the tadpoles remain in the free-swimming stage between 1.5 months (Hoogmoed 1967) and nearly a year (Del Pino and Escobar 1981). Therefore, stable water resources are an important habitat requirement for this species.

Gastrotheca marsupiata, reported as ranging between 2,500 and 4,360 m a.s.l. (IUCN 2020), is one of only seven *Gastrotheca* species ranging above 4,000 m a.s.l., along with *G. dissimilis* (Vellard, 1957), *G. excubitor* (Duellman & Fritts, 1972), *G. pacchamama* (Duellman, 1987), *G. pseustes* (Duellman & Hillis, 1987), *G. griswoldi* (Shreve, 1941), and *G. peruana* (Boulenger, 1900) (Duellman 2013; Duellman and Trueb 2015; Vargas 2015; Frost 2020; IUCN 2020). Of those, only *G. peruana* and *G. pacchamama* have been recorded up to 4,600 m a.s.l., making them the highest-elevation members of the genus until the present record. Here, we report a 300-m elevational range extension for *G. marsupiata* into Osjollo Pass in the core of the Cordillera Vilcanota. This brings *G. marsupiata* to a maximal elevation of 4,660 m a.s.l. and makes it the highest-occurring species in a genus already characterized by high elevation ranges.

The Cordillera Vilcanota, which extends approximately 80 km east–west and 40 km north–south across part of the departments of Cusco and Puno, has the largest Peruvian glacier and the second-highest density of glaciers in Peru (Morales-Arno and Hastenrath 1999). As Osjollo Pass deglaciated following the Little Ice Age maximum (ca. 1850–1900), three anurans expanded their elevational range into this newly ice-free habitat: *Telmatobius marmoratus* (Duméril & Bibron, 1841), *Pleurodema marmoratum* (Duméril & Bibron, 1840), and *Rhinella spinulosa* (Wiegmann, 1834). The former two species, in so doing, have become the highest-living amphibian species worldwide (Seimon et al. 2007). *Gastrotheca marsupiata* is sympatric with these other three amphibians lower in their range, and may similarly be expanding upslope with the shifting climatic conditions.

Methods

On 21 March 2018, our team found tadpoles and metamorphosing froglets in a small pond on the northern side of Osjollo Pass in the Cordillera Vilcanota, Peru (Fig. 1). The tadpoles and froglets were identified as *Gastrotheca marsupiata*, as other regionally occurring *Gastrotheca* species deposit froglets rather than tadpoles from their pouches (Duellman and Fritts 1972) and have a lower maximum elevational range than *G. marsupiata* (IUCN 2020). One froglet was photographed with a Nikon D7200 and Tokina 100mm macro lens to later confirm field species identification using the descriptions of Duellman and Fritts (1972) (Fig. 2), though no voucher was taken.

On 8 May 2019, the field team returned to search for an adult individual by diurnal rock flipping. One specimen was collected, photographed as previously, euthanized by an anesthetic overdose of 20% benzocaine hydrochloride gel applied topically to the ventrum, and preserved in 70% ethanol. The species was identified using the descriptions of Duellman and Fritts (1972). The voucher specimen was deposited at the Museo de Biodiversidad del Perú (MUBI), Cusco, Peru.

Coordinates were taken using a handheld Garmin GPSMAP 64s. The datum used for recording geographical coordinates was WGS84. Collections were made under a permit issued by SERFOR (no. AUT-IFS-2019-014) and an animal care and use protocol approved by University of California, Berkeley's ACUC (no. AUP-2018-12-11648).

Results

Gastrotheca marsupiata (Duméril & Bibron, 1841)

Figure 2

New record. PERU • 1 subadult ♂; Cusco Department, Quispicanchis Province, Cordillera Vilcanota; 13° 70.00'S, 071°13.87'W; 4,660 m a.s.l.; 8 May 2019; E. Steigerwald, Y. J. Guevara, and G. Crispin leg.; under a stone in moraine that was thinly colonized by grasses, ca. 5m from a stream; MUBI 16429.

Identification. Identification was made using information about the geographic distribution of *Gastrotheca* species as well as key traits. The observation of free-swimming tadpoles and metamorphosing froglets differentiates this species from other *Gastrotheca* species in the region, which carry their young past the tadpole stage (Duellman and Fritts 1972). Additionally, other *Gastrotheca* species in the region are found at lower elevations, well below the elevational limit of *G. marsupiata* (IUCN 2020). Further identifying characteristics of the adult frogs were the acutely rounded snout, round canthus, dark canthal and postorbital striping on the head, lighter labial stripe, pronounced supratympanic fold over a vertically ovoid tympanum, smooth tympanic annulus, and slightly concave loreal region. Diagnostic features




Figure 1. Distribution map of *Gastrotheca marsupiata*. The white lines represent national borders, and the black outline represents the species distribution (IUCN 2020). The small, hatched box marks the location of the inset map, and the black star on the inset map represents the location of our new record.

observed on the limbs were the lack of webbing between the fingers on the hands, toe pads that are slightly wider than they are long, and pronounced large round subarticular tubercles on the hands. A slightly granular dorsal skin with dark green markings on the dorsal side, transverse bars or spots on the dorsal portion of limbs, and mottling on the lateral portions of the body all confirm that individuals observed were *G. marsupiata*.

Remarks. This population represents a 300 m increase in the maximum elevation known for this species to 4,660 m a.s.l., the highest record of any *Gastrotheca* species. In March 2018, more than 20 tadpoles were encountered. The following year, in May 2019, seven adult frogs were found and one was collected. The frogs were found on the northern side of Osjollo Pass in rocky puna (grassland) habitat situated above a series of tarns and bofedales (peatlands), fed by precipitation and glacial meltwater. Vegetation at the site is composed of a mixture of native cushion plants and grasses alongside non-native grasses introduced to feed cattle, which are present in low numbers at the site. The tadpoles were found in a small stagnant pond. The presence of both

tadpoles and adult frogs over two consecutive years suggests an established breeding population. The weather at the site is typical of the high Andes, with rapid temperature swings and inclement weather such as hard frosts, snow, hail, and rain, including during the season when the frogs were observed. Temperature loggers at a site 2.5 km southwards into Osjollo Pass recorded an annual terrestrial high of 25 °C and low of −4.5 °C, and an annual aquatic high of 27.5 °C and low of −3 °C.

Discussion

We present new records for *Gastrotheca marsupiata* at 4,660 m a.s.l. in the Cordillera Vilcanota. This represents a 300 m vertical elevational extension of their known range, making *Gastrotheca marsupiata* the highest occurring species in its genus. Previous regular amphibian surveys of the Vilcanota that have been conducted since 2003 did not find *G. marsupiata* on transects along Osjollo Pass from 4,860–6,100 m a.s.l. However, these surveys did not include the site where this species was found (T. Seimon pers. comm. 7 July 2020; K. Reider pers. comm. 19 July 2020), so we cannot know how long




Figure 2. Tadpole and voucher individual of *Gastrotheca marsupiata*. **A.** Tadpole. **B–D.** Voucher individual: **(B)** dorsal, **(C)** ventral, and **(D)** lateral views.

G. marsupiata has occupied this habitat. Additionally, unlike the habitat recently colonized by *Telmatobius marmoratus*, *Pleurodema marmoratum*, and *Rhinella spinulosa* at the southern end and core of Osjollo Pass (Seimon et al. 2007), this site was not glaciated in 1931 aerial photographs (Shippee–Johnson Collection, Department of Library Services, American Museum Natural History; date inferred from Shippee 1933) and was probably not glaciated during the Little Ice Age maximum (1850–1900) given abundant lichen cover on nearby rocks. However, if this record does represent an upslope range extension into mountain pass habitat, it may be attributable to local changes in climatic conditions. Regardless, what *G. marsupiata* certainly has in common with these other local anuran species is that their occupancy of this extreme high elevation habitat will most likely be short-lived due to continued changing climatic conditions, as the supply of glacial meltwater feeding local streams and ponds ebbs.

The evidence for climate-driven range shifts in amphibians is still scarce. This is particularly true for poleward shifts, whereas limited evidence for climate-induced amphibian elevational range shifts has accumulated (Li et al. 2013). Over the last decades, amphibian elevational range shifts have been documented on the Tsaratanana Massif in Madagascar (Raxworthy et al. 2008), in Mount Gede Pangrango National Park in Indonesia (Kusriani et al. 2017), in the Ecuadorian Andes (Bustamante et al. 2005), in the Appalachian Mountains of the United States (Moskwick 2014), and in South Africa (Botts et al. 2015). In all of these cases, the authors suggest that the range shifts documented may have been driven by changes in the regional climate, and the authors of the Indonesian and Madagascan studies were able to rule out land use change entirely as a contributing factor since these studies were conducted within stably protected areas. Our new record for *G. marsupiata* represents a possible further case of contemporary climate-induced elevational range shifting, but we emphasize the critical role of extensive monitoring in being able to distinguish where, when, and how natural communities are truly changing over time.

Batrachochytrium dendrobatidis (Bd), the fungal pathogen that has caused declines and even extinctions of diverse amphibian species globally (Skerratt et al. 2007), has been detected from skin swabs of various Peruvian *Gastrotheca* species, including *G. excubitor* (Catenazzi et al. 2011), which is sympatric with *G. marsupiata* in portions of the Cordillera Vilcanota. However, a study examining the response of Andean frogs to Bd found that different *Gastrotheca* species have vastly different reactions and survival rates (Catenazzi et al. 2017), so it is difficult to predict whether *G. marsupiata* is susceptible prior to trial infection experiments. Although Bd has not yet been recorded in *G. marsupiata*, the species has experienced declines in neighboring Bolivia, even vanishing from parts of its former range (De La Riva and Reichle 2014). As the other amphibian species co-inhabiting Osjollo Pass have tested positive for Bd (Seimon

et al. 2017), the susceptibility of *G. marsupiata* to Bd is another factor that may determine how long this species will persist in these extreme reaches of its elevational range.

Acknowledgements

Fieldwork was funded by the Sophie Danforth Conservation Biology Fund from the Roger Williams Park Zoo & Rhode Island Zoological Society, Karl Koford Grant from the Museum of Vertebrate Zoology, Reed Scholarship from the Northern California Association of Phi Beta Kappa, and the Fellowship of Graduate Student Travel from the Society for Integrative and Comparative Biology. Thanks to Felipe and Severino Crispin for their generous loan of some important field equipment. Thanks to Kelsey Reider and Tracie and Anton Seimon for information on their previous amphibian surveys in the Cordillera Vilcanota. We appreciate the helpful feedback from Javier Ernesto Cortés Suárez and two anonymous reviewers. ES was supported by the NSF GRFP.

Authors' Contributions

ES, AS, FPC, YJG, and GC conducted biological fieldwork. GC translated Quechua. JCC assisted with permit acquisition and species deposition in the Museo de Biodiversidad del Perú. ES and AS wrote and FPC, YJG, and JCC edited the manuscript.

References

- AmphibiaWeb (2020) <https://amphibiaweb.org>. Accessed on: 2020–08–03.
- Botts E, Erasmus B, Alexander G (2015) Observed range dynamics of South African amphibians under conditions of global change. *Austral Ecology* 40 (3): 309–317. <https://doi.org/10.1111/aec.12215>
- Bustamante M, Ron S, Coloma L (2005) Cambios en la diversidad en siete comunidades de anuros en los Andes de Ecuador. *Biotropica: The Journal of Biology and Conservation* 37 (2): 180–189. <https://doi.org/10.1111/j.1744-7429.2005.00025.x>
- Catenazzi A, Lehr E, Rodríguez LO, Vredenburg VT (2011) *Batrachochytrium dendrobatidis* and the collapse of montane anuran communities in the upper Manu National Park, southeastern Peru. *Conservation Biology* 25: 382–391. <https://doi.org/10.1590/S1676-06032013000400024>
- Catenazzi A, Swei A, Finkle J, Foreyt E, Wyman L, Vredenburg VT (2017) Epizootic to enzootic transition of a fungal disease in tropical Andean frogs: are surviving species still susceptible? *PLoS ONE* 12 (10): e0186478. <https://doi.org/10.1371/journal.pone.0186478>
- De la Riva I, Reichle S (2014) Diversity and conservation of the amphibians of Bolivia. *Herpetological Monographs* 28 (1): 46–65. <https://doi.org/10.1655/herpmonographs-d-13-00009>
- Del Pino E (1989) Marsupial frogs. *Scientific American* 260 (5): 110–119.
- Del Pino E, Escobar B (1981) Embryonic stages of *Gastrotheca riobambae* (Fowler) during maternal incubation and comparison of development with that of other egg-brooding hylid frogs. *Journal of Morphology* 167 (3): 277–295. <http://doi.org/10.1002/jmor.1051670303>
- Duellman W (2013) An elusive new species of Marsupial Frog (Anura: Hemiphraactidae: *Gastrotheca*) from the Andes of northern Peru.

- Phyllomedusa 12 (1): 3–11. <https://doi.org/10.11606/issn.2316-9079.v12i1p3-11>
- Duellman W, Fritts T (1972) A taxonomic review of the southern Andean marsupial frogs (Hylidae: *Gastrotheca*). Occasional papers of the Museum of Natural History, the University of Kansas 9: 1–37.
- Duellman W, Maness S (1980) The reproductive behavior of some hylid marsupial frogs. *Journal of Herpetology* 14 (3): 213. <http://doi.org/10.2307/1563542>
- Duellman W, Trueb L (2015) Marsupial frogs, *Gastrotheca* and allied genera. Johns Hopkins University Press, Baltimore, USA, 427 pp.
- Dumeril A, Bibron G (1841) *Erpétologie générale ou histoire naturelle complète des reptiles*, Vol. 8. Roret, Paris, France, 824 pp.
- Frost D R (2020) Amphibian species of the world: an online reference. American Museum of Natural History, New York, USA. <https://amphibiansoftheworld.amnh.org/index.php>. Accessed on: 2020-8-13. <https://doi.org/10.5531/db.vz.0001>
- Guibé J (1950) Catalogue des types d'amphibiens du Muséum national d'Histoire naturelle. Imprimerie Nationale, Paris, France, 71 pp.
- Hoogmoed M (1967) Mating and early development of *Gastrotheca marsupiata* (Dumeril and Bibron) in captivity (Hylidae, Anura, Amphibia). *British Journal of Herpetology* 4: 1–7.
- IUCN (International Union for the Conservation of Nature) (2020) The IUCN Red List of threatened species. Version 2020-1. <https://www.iucnredlist.org>. Accessed on: 2020-8-3.
- Kusrini M, Lubis M, Enderwin W, Yazid M, Darmawan B, Ul-Hasanah A, Sholihata N, Tajallia A, Lestaria V, Utama H, Nasira D, Ardiansyaha D, Rachmadia R (2017) Elevation range shift after 40 years: the amphibians of Mount Gede Pangrango National Park revisited. *Biological Conservation* 206: 75–84. <https://doi.org/10.1016/j.biocon.2016.12.018>
- Li Y, Cohen J, Rohr J (2013) Review and synthesis of the effects of climate change on amphibians. *Integrative Zoology* 8 (2): 145–61. <https://doi.org/10.1111/1749-4877.12001>
- Morales-Arnao B, Hastenrath S, Williams R Jr., Ferrigno J (1999) *Glaciers of South America: glaciers of Peru*. Professional Paper 1386-I-4. U.S. Geological Survey, U.S. Department of the Interior. <https://pubs.usgs.gov/prof/p1386i/peru/orient.html>. Accessed on 2020-8-3. <https://doi.org/10.3133/pp1386I>
- Moskwick M (2014) Recent elevational range expansions in plethodontid salamanders (Amphibia: Plethodontidae) in the southern Appalachian Mountains. *Journal of Biogeography* 41 (10): 1957–1966. <https://doi.org/10.1111/jbi.12352>
- Raxworthy C, Pearson R, Rabibisoa N, Rakotondrazafy A, Ramamanjato J, Raselimanana A, Wu S, Nussbaum R, Stone D (2008) Extinction vulnerability of tropical montane endemism from warming and upslope displacement: a preliminary appraisal for the highest massif in Madagascar. *Global Change Biology* 14 (8): 1703–1720. <https://doi.org/10.1111/j.1365-2486.2008.01596.x>
- Seimon T, Seimon A, Daszak P, Halloy S, Schloegel L, Aguilar C, Sowell P, Hyatt A, Konecky B, Simmons J (2007) Upward range extension of Andean anurans and chytridiomycosis to extreme elevations in response to tropical deglaciation. *Global Change Biology* 13 (1): 288–299. <https://doi.org/10.1111/j.1365-2486.2006.01278.x>
- Shippie R (1933) Air adventures in Peru. *National Geographic Magazine* 64 (1): 81–120.
- Sinsch U, Joermann G (1989) Vocalization and mating behavior of the marsupial frog, *Gastrotheca marsupiata* (Dumeril and Bibron, 1841) (Amphibia: Hylidae). *Copeia* 1989 (3): 755. <http://doi.org/10.2307/1445510>
- Skerratt L, Berger L, Speare R, Cashins S, McDonald K, Phillott A, Hines H, Kenyon N (2007) Spread of Chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth* 4: 125–134. <https://doi.org/10.1007/s10393-007-0093-5>
- Vargas V (2015) *Guía de identificación de anfibios y reptiles*. PERU LNG, Lima, Peru, 111 pp.