

<https://helda.helsinki.fi>

---

## Enhanced activity of hyperthermostable *Pyrococcus horikoshii* endoglucanase in superbase ionic liquids

Hebal, Hakim

2022-08

---

Hebal , H , Hamalainen , J , Makkonen , L , King , A W T , Kilpelainen , I , Bankar , S , Boucherba , N & Turunen , O 2022 , ' Enhanced activity of hyperthermostable *Pyrococcus horikoshii* endoglucanase in superbase ionic liquids ' , *Biotechnology Letters* , vol. 44 , pp. 961-974 . <https://doi.org/10.1007/s10529-022-03268-5>

---

<http://hdl.handle.net/10138/346991>

<https://doi.org/10.1007/s10529-022-03268-5>

---

cc\_by

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*


*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



# Enhanced activity of hyperthermostable *Pyrococcus horikoshii* endoglucanase in superbase ionic liquids

Hakim Hebal · Joonas Hämäläinen · Laura Makkonen · Alistair W. T. King ·  
Ilkka Kilpeläinen · Sandip Bankar · Nawel Boucherba · Ossi Turunen 

Received: 29 December 2021 / Accepted: 31 May 2022 / Published online: 28 June 2022  
© The Author(s) 2022

## Abstract

**Objectives** Ionic liquids (ILs) that dissolve biomass are harmful to the enzymes that degrade lignocellulose. Enzyme hyperthermostability promotes a tolerance to ILs. Therefore, the limits of hyperthermophilic *Pyrococcus horikoshii* endoglucanase (PhEG) to tolerate 11 superbase ILs were explored.

**Results** PhEG was found to be most tolerant to 1-ethyl-3-methylimidazolium acetate ([EMIM]OAc) in soluble 1% carboxymethylcellulose (CMC) and insoluble 1% Avicel substrates. At 35% concentration, this IL caused an increase in enzyme activity (up to 1.5-fold) with CMC. Several ILs were more enzyme inhibiting with insoluble Avicel than with soluble CMC.  $K_m$  increased greatly in the presence of ILs, indicating significant competitive inhibition. Increased hydrophobicity of the IL cation or anion

was associated with the strongest enzyme inhibition and activation. Surprisingly, PhEG activity was increased 2.0–2.5-fold by several ILs in 4% substrate. Cations exerted the main role in competitive inhibition of the enzyme as revealed by their greater binding energy to the active site.

**Conclusions** These results reveal new ways to design a beneficial combination of ILs and enzymes for the hydrolysis of lignocellulose, and the strong potential of PhEG in industrial, high substrate concentrations in aqueous IL solutions.

**Keywords** Hyperthermostable endoglucanase · Enzyme kinetics · Enzyme inhibition · Ionic liquids · Biomass engineering

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s10529-022-03268-5>.

H. Hebal · N. Boucherba  
Laboratoire de Microbiologie Appliquée (LMA),  
Faculté des Sciences de La Nature et de La Vie (FSNV),  
Université de Bejaia, Bejaia, Algeria

H. Hebal  
Faculty of Exact Sciences and Sciences of Nature and Life,  
Department of Biology, Mohamed Khider University  
of Biskra, Biskra, Algeria

J. Hämäläinen  
St1 Oy, Firdonkatu 2, PL 68, 00521 Helsinki, Finland

L. Makkonen · S. Bankar  
Department of Bioproducts and Biosystems, School  
of Chemical Engineering, Aalto University, 00076 Aalto,  
Finland

A. W. T. King · I. Kilpeläinen  
Department of Chemistry, University of Helsinki,  
00014 Helsinki, Finland

O. Turunen (✉)  
School of Forest Sciences, University of Eastern Finland,  
80101 Joensuu, Finland  
e-mail: [ossi.turunen@uef.fi](mailto:ossi.turunen@uef.fi)

## Introduction

As the main component of biomass, lignocellulose is comprised of cellulose, hemicellulose and lignin (Mckendry 2002). Cellulose is the most abundant polysaccharide on the planet and is, therefore, considered a potentially important source of carbohydrate fibres, and monosaccharides, which can be then used as a substrate for microbial fermentations to produce fuels and value-added chemicals (Chacón et al. 2021; Puligundla et al. 2021). Cellulose is a linear polymer of  $\beta$ -1,4 linked-D-glucose units with varying degrees of polymerisation, and is hydrolysed by a consortium of enzymes to glucose: endo-1,4- $\beta$ -glucanase (EC3.2.1.4), cellobiohydrolase (EC 3.2.1.91) and  $\beta$ -glucosidase (EC 3.2.1.21). Endo-1,4- $\beta$ -glucanase acts by randomly hydrolysing cellulose chains (Barbosa et al. 2020). However, cellulose is recalcitrant to enzymatic hydrolysis requiring therefore physico-chemical pretreatment to decrease its crystallinity and degree of polymerisation, in one hand, and improving of enzyme efficiency, in the other hand (Alvira et al. 2010; Raulo et al. 2021; Contreras et al. 2020).

Ionic liquids (ILs) are salts that exist in liquid form below 100 °C and have the capacity to dissolve lignocellulose for improved hydrolysis by enzymes. They are potentially low polluting and display high thermal and chemical stability, high polarity and very low toxicity (Patel et al. 2014). Dissolved cellulose can be precipitated away from the IL solution using an antisolvent. However, traces of ILs (in the range of 10–15%) may remain in the substrate after precipitation (Engel et al. 2010), and most commercial cellulases are inhibited or deactivated by even low amounts of ILs (Wahlström and Suurnäkki 2015; Nemestóthy et al. 2017; Hebal et al. 2021). Therefore, the discovery or development of new enzymes with greater tolerance towards ILs is a prerequisite for the development of lignocellulose processes that use ILs.

Typically, the highly IL-tolerant endoglucanases and  $\beta$ -glucosidases originate from hyperthermophilic microbes (Hebal et al. 2021), which suggests a link between enzyme thermostability and IL tolerance (Ferdjani et al. 2011; Ilmberger et al. 2012). A hyperthermophilic GH5 endoglucanase (PhEG) with the capability of hydrolysing crystalline cellulose from the hyperthermophilic archaeon *Pyrococcus horikoshii* has been reported previously (Ando et al. 2002). The extreme thermostability associated with this enzyme

(Kang and Ishikawa 2007; Hämäläinen et al. 2016) is probably related to its greater tolerance to ILs. The enzyme was found to retain 90% of its specific activity at 20% 1-ethyl-3-methylimidazolium acetate ([EMIM]OAc) concentration at 80 °C, and 79% of activity was recovered after 15 h of incubation at 15% [EMIM]OAc at 80 °C (Datta et al. 2010). Furthermore, PhEG has been found to be the most IL-tolerant enzyme across a range of studied cellulases in the presence of a 40% concentration of seven cellulose-dissolving or swelling ILs (Rahikainen et al. 2018).

The objective of the current study is to study the effect of diluted aqueous solutions of a wide range of superbase ILs on the functioning of the hyperthermostable PhEG. Definition of a superbase includes *a neutral organic base that is more basic than sodium hydroxide* (Ishikawa 2009). Detailed study of the hyperthermostable enzymes that resist protein denaturation by ILs is expected to better reveal how the enzyme activity itself is affected by ILs. Acetate-propionate ILs show rapid dissolution of cellulose and are potentially distillable (King et al. 2011; Parviainen et al. 2013), while guaiacolate ILs do not dissolve biomass (Hebal et al. 2020). In this study, special attention was paid to enzyme performance on a soluble cellulose substrate (carboxymethyl cellulose, CMC) and an insoluble cellulose substrate (Avicel).

## Materials and methods

### Sources of enzymes and ionic liquids

*Pyrococcus horikoshii* endoglucanase (PhEG) (family GH5) was expressed from *Escherichia coli* and partly purified, as described in Hämäläinen et al. (2016). Ionic liquids (Table 1) were prepared, as described in Parviainen et al. (2013, 2014) and Hebal et al. (2020). [EMIM]OAc was purchased from BASF (Ludwigshafen, Germany, purity  $\geq 95\%$ ). The superbase-derived ILs were prepared by combining several superbases ([DBNH]<sup>+</sup>, [DBUH]<sup>+</sup>, [mDBN]<sup>+</sup> and [TMGH]<sup>+</sup>) with organic acids, such as acetate, propionate and guaiacolate (Table 1).

### Enzyme assays

Enzyme activity on soluble cellulose (Na-CMC; Sigma-Aldrich, St. Louis, Montana) and crystalline

**Table 1** The tested biomass-dissolving ionic liquids

Ionic liquid		Molecular weight (g/mol)
[DBNH]OAc	1,5-Diazabicyclo[4.3.0]non-5-enium acetate	125.18 + 59.04 = 184.22
[DBNH]CO <sub>2</sub> Et	1,5-Diazabicyclo[4.3.0]non-5-enium propionate	125.18 + 73.07 = 198.25
[DBNH]guaiacolate	1,5-Diazabicyclo[4.3.0]non-5-enium 2-Hydroxy-3-methoxybenzoate	125.18 + 124.13 = 249.31
[DBUH]OAc	1,8-Diazabicyclo[5.4.0]undec-7-enium acetate	153.23 + 59.04 = 212.27
[DBUH]CO <sub>2</sub> Et	1,8-Diazabicyclo[5.4.0]undec-7-enium propionate	153.23 + 73.07 = 226.3
[DBUH]guaiacolate	1,8-Diazabicyclo[5.4.0]undec-7-enium 2-Hydroxy-3-methoxybenzoate	153.23 + 124.13 = 277.36
[EMIM]OAc	1-Ethyl-3-methylimidazolium acetate	111.17 + 59.04 = 170.21
[mDBN]Me <sub>2</sub> PO <sub>4</sub>	Methyl-1,5-diazabicyclo[4.3.0]non-5-enium dimethyl phosphate	139.22 + 126.05 = 265.27
[TMGH]OAc	1,1,3,3-Tetramethylguanidinium acetate	116.18 + 59.04 = 175.22
[TMGH]CO <sub>2</sub> Et	1,1,3,3-Tetramethylguanidinium propionate	116.18 + 73.07 = 189.25
[TMGH]guaiacolate	1,1,3,3-Tetramethylguanidinium 2-Hydroxy-3-methoxybenzoate	116.18 + 124.13 = 240.31

cellulose (Avicel PH-101; Sigma-Aldrich) were measured with the 3,5-dinitrosalicylic acid (DNS) method (Miller 1959), as described in Hebal et al. (2020). The endoglucanase acted on 1% or 4% (m/v) substrates in 50 mM citrate–phosphate buffer at pH 6 for 30 min at 70 °C, in the absence or presence of 5–35% (v/v) IL solutions. Reciprocal shaking at 200 rpm was used in the enzyme reaction with Avicel and the mixture was centrifuged before the absorbance was measured. The specific activity of the enzymes used in the assays was 14 U/mg with CMC as substrate. One unit (U) of enzyme activity was defined as the amount of enzyme required to liberate 1 μmol of product per min. Since ILs induce an increase in the absorbance, calibration graphs in the presence of ILs were constructed to correct the observed absorbance values (Supplementary Fig. 1).

#### Kinetic experiments

Kinetic parameters were determined in standard assay conditions at 70 °C (pH 6) with and without 5% (v/v) IL. Initial velocities were determined using 4, 6, 10 and 20 mg/ml CMC at different incubation times (10, 20, 30 and 40 min). Initial velocity was determined from the slope of the graph (Supplementary Fig. 2 and Supplementary Fig. 3). All experiments were performed three times with triplicates. Kinetic values were calculated by the hyperbolic regression analysis function (without weighting) in the Hyper 32 program.

#### Molecular docking

The IL cations and anions were docked by Swiss-Dock (Grosdidier et al. 2011) to the active site of PhEG (PDB code 2ZUM; Kim and Ishikawa 2009), as described in Hebal et al. (2020).


## Results and discussion

The use of ILs in combination with enzymes profoundly affects the kinetic behaviour and stability of enzymes. Therefore, in this study, the kinetic behaviour of an hyperthermostable endoglucanase was studied in aqueous IL solutions. The goal was to obtain a better picture of the competitive inhibition caused by ILs with regard to the variation of IL cations and the anions, and the nature of the substrate. Although several glycoside hydrolase enzymes (including PhEG) have been studied earlier, our study revealed new behaviour associated with a hyperthermostable endoglucanase in a range of ILs.

#### Activity of PhEG in 5–35% ionic liquids

The activity of PhEG was studied in 5–35% acetate-propionate ILs (Fig. 1), and in 5% guaiacolate-based ILs (Fig. 2), with 1% CMC as a substrate. In acetate-propionate ILs (Fig. 1), [EMIM]OAc was the only IL that favoured PhEG activity at higher IL concentrations. An increasing concentration of [EMIM]OAc


**Fig. 1** Activity of *Pyrococcus horikoshii* endoglucanase (PhEG) in 5–35% ionic liquids with 1% carboxymethylcellulose (CMC) as substrate. Activity was measured at 70 °C and pH 6. 100% activity denotes the activity without IL



up to 35% increased enzyme activity. Other ILs at 25–35% concentrations greatly decreased enzyme activity. Only [EMIM]OAc, [TMGH]OAc and [TMGH]CO<sub>2</sub>Et showed activity at 35% concentration, although the latter two reduced the activity. At 15% concentration, [mDBN]Me<sub>2</sub>PO<sub>4</sub> also showed clear elevation of PhEG activity. Altogether, [DBNH]OAc, [DBNH]CO<sub>2</sub>Et and [mDBN]Me<sub>2</sub>PO<sub>4</sub> were well tolerated by PhEG at 15% IL concentration, but activity was rapidly lost at 25% concentration. No consistent trend of inhibition was observed at 25% IL concentration when the cation or the anion were varied (Fig. 1). These superbase ILs exerted a clear inhibition on the hyperthermostable *Thermoplasma flexuosa* GH10 xylanase TFXYN10A at this concentration (Hebal et al., 2020). At 35% ILs, PhEG seemed to be most inhibited by hydrophobic cations, i.e. ([DBNH]<sup>+</sup>, [DBUH]<sup>+</sup> and [mDBN]<sup>+</sup>), and was less inhibited or even stimulated by the less hydrophobic cations ([EMIM]<sup>+</sup>, [TMGH]<sup>+</sup>). A similar finding was observed when TFXYN10A was assayed

with superbase ILs (Hebal et al. 2020). However, in general, TFXYN10A seems to be more inhibited by superbase ILs than PhEG at the same range of IL concentrations. In addition, PhEG was most inhibited by the [DBUH]<sup>+</sup> cation, whereas TFXYN10A was more inhibited by [DBNH]<sup>+</sup> than [DBUH]<sup>+</sup> cation. Wahlstrom et al. (2013) studied the hydrolysis of microcrystalline cellulose (MCC) by *Trichoderma reesei* endoglucanase Cel5A in the presence of 20% (w/w) superbase ILs. In line with our results, the authors found that the greatest MCC hydrolysis yield was obtained with [EMIM]AOc, followed by [TMGH]AOc, and that the activity was strongly inhibited by [DBNH]OAc and [DBNH]CO<sub>2</sub>Et.

When ILs were used at 5% concentration, a slight increase in enzyme activity was observed in most acetate-propionate ILs but especially in [TMGH]OAc, [DBUH]CO<sub>2</sub>Et and [DBNH]OAc (Fig. 1). Of these, [TMGH]OAc induced the greatest increase in activity. Enzyme activity was enhanced with guaiacolate IL when the cations were [DBUH]<sup>+</sup> and [TMGH]<sup>+</sup>



**Fig. 2** Activity of *Pyrococcus horikoshii* endoglucanase (PhEG) in 5% guaiacolate ionic liquids with 1% carboxymethylcellulose (CMC) as substrate. Activity was measured at 70 °C and pH 6. 100% activity denotes the activity without IL

but not with [DBNH]<sup>+</sup> (Fig. 2). In contrast, 5% guaiacolate ILs inhibited hyperthermostable GH10 and GH11 xylanases (Hebal et al. 2020). The unconjugated cations displayed the following degrees of declining basicity: [mDBN]<sup>+</sup> > [EMIM]<sup>+</sup> > [DBUH]<sup>+</sup> > [DBNH]<sup>+</sup> > [TMGH]<sup>+</sup> (Parviainen et al. 2013). The more basic cations, [mDBN]<sup>+</sup> and [EMIM]<sup>+</sup> are probably more prone to interact with water molecules than with protein and, therefore, exert a less activating effect on the enzyme. No consistent trend in enzyme activation was observed for the anions.

Our results for PhEG activity in [EMIM]OAc are, to some degree, different from those reported by Datta et al. (2010), since [EMIM]OAc was observed to activate the enzyme in our study. However, experimental conditions were different in these two studies. In Datta et al. (2010), a temperature of 80 °C was used in the enzyme reaction compared to 70 °C in our

study. Moreover, the use of bovine serum albumin (BSA) as a stabiliser and pH adjustment after IL addition was not reported by Datta et al. (2010) (initial pH was 6.4 in Datta et al. (2010); adjusted pH was 6 in our study). Other studies have shown that ILs can decrease the temperature optimum of enzyme activity (Yu et al. 2016; Anbarasan et al. 2017) and modify the pH of the reaction mixture, while stabilizing agents may also increase the tolerance of enzymes to ILs (Zhao 2010). It is probable that our experimental conditions better supported the functioning of the enzymes in ILs. These differences in the results suggest that reaction condition engineering is important in improving enzyme tolerance to ILs. The influence of reaction conditions on enzyme tolerance to ILs was also seen with TfXYN10A, since [EMIM]OAc stabilised the enzyme at 80 °C but not at 60–70 °C (Anbarasan et al. 2017).

In particular, the stimulation of PhEG at 35% [EMIM]OAc, could contribute to the simplification of the overall IL-pretreatment-hydrolysis process by allowing the enzymatic saccharification of cellulose by eliminating the need to recover the regenerated cellulose (Kamiya et al. 2008). In this strategy, after the IL pretreatment of lignocellulose, a buffer solution is added to decrease the IL concentration to <50%, followed by addition of the IL tolerant enzymes. This would overcome the challenges caused by viscosity of the solution and the excessive washing required to remove the ILs and should, therefore, reduce the process costs. However, the one pot process with ILs seems not to be possible with existing commercial cellulases, since it has been shown that [EMIM]OAc concentrations >10% compromise the performance of these enzymes (Auxenfans et al. 2017; Husson et al. 2018).


PhEG was slightly activated in the presence of 40% [TMGH]n-PrCOO (Rahikainen et al. 2018), and enzyme activation by ILs has also been observed in other hyperthermophilic endoglucanases (Hebal et al. 2021). This phenomenon has been proposed to reflect the need for salt to achieve maximal enzyme activity (Ajloo et al. 2013; Gladden et al. 2014). However, PhEG activity was not stimulated with NaCl or KCl (0.5–4 M) in this study (Supplementary Fig. 4). On the contrary, the activity of a halophilic endoglucanase that was enhanced by NaCl was not stimulated by ILs (Ben Hmad et al. 2017). This suggests a different mechanism of enzyme activation by ILs and that

other types of IL-protein interactions are important, in addition to electrostatic effects. In this respect, a major property of ILs is the hydrophobic portion of the cations.

Protein denaturation has been shown to be absent or minimal for several hyperthermostable glycoside hydrolases in dilute IL solutions (Hebal et al. 2021). The unfolding temperature of PhEG only decreased by 10.5 °C in 20% [EMIM]OAc, from 102.3 to 91.8 °C (Datta et al. 2010). Only minimal structural changes were observed in this enzyme when simulated in the presence of 15% and 50% [EMIM]OAc at 80 °C (Jaeger et al. 2015). These results suggest that the enzyme is not very prone to unfolding at the assay temperature of 70 °C used in our study, and it is likely that the ILs affect the enzyme by inhibiting activity rather than by unfolding the protein. In several studies, loss of hyperthermostable enzyme activity in dilute IL solutions was found to be caused by competitive inhibition of IL molecules, especially the cation, during the binding of the substrate to the active site (Hebal et al. 2021).

#### Activity of PhEG in 4% substrate


To clarify the role of competitive inhibition, the effect of ILs on PhEG activity was compared in 1% and 4% substrate concentrations in the presence of 15% acetate-propionate ILs, [mDBN]Me<sub>2</sub>PO<sub>4</sub> (Fig. 3) and 5% guaiacolate ILs (Fig. 4). In competitive inhibition, the greater substrate concentration counteracts inhibition by outcompeting the inhibitor. With PhEG, increasing the substrate concentration resulted not only in full recovery of activity but also in the activation of PhEG with all acetate-propionate ILs and [mDBN]Me<sub>2</sub>PO<sub>4</sub> (Fig. 3). With guaiacolate ILs, an increase in substrate concentration further stimulated enzyme activity, with [TMGH]guaiacolate exerting the most stimulating effect (Fig. 4). The elevated xylan concentration also counteracted the competitive inhibition of TfXYN10A by superbase ILs (Hebal et al. 2020) and of TasXyn10A by 25% [EMIM]OAc (Chawachart et al. 2014). Increased activity in the presence of elevated substrate concentrations suggests that competitive inhibition was counteracted by the substrate. However, the noteworthy increase observed in PhEG activity in 4% substrate with ILs may even indicate that IL molecules in high substrate concentrations stimulate the reaction interaction between



**Fig. 3** Activity of *Pyrococcus horikoshii* endoglucanase (PhEG) in 1% and 4% substrates in the presence of 15% ionic liquids. Activity was measured at 70 °C and pH 6. 100% activity denotes the activity in each substrate concentration without IL

the substrate and the enzyme. These results reveal that the IL molecules may both activate and inhibit the enzymes, and that the balance between these two opposing behaviours depends at least partly on the conditions.

Unlike PhEG under the same conditions, TfXYN10A was not activated or was only slightly activated by superbase ILs in 4% substrate (Hebal et al. 2020). This difference suggests that superbase ILs exert a differing effect on PhEG compared to TfXYN10A. No difference between acetate and propionate was observed when activating PhEG in 4% substrate when they were conjugated to more hydrophobic cations ([DBNH]<sup>+</sup> and [DBUH]<sup>+</sup>). Propionate and guaiacolate showed greater activation than acetate when they were conjugated to the less hydrophobic cation [TMGH]<sup>+</sup> (Figs. 3 and 4). These results suggest that increased hydrophobicity of the IL ions is among the main factors that contribute to the activation of PhEG, and that activation by the




**Fig. 4** Activity of *Pyrococcus horikoshii* endoglucanase (PhEG) in 1% and 4% substrates in the presence of 5% guaiacolate ionic liquids. Activity was measured at 70 °C and pH 6. 100% activity denotes the activity in each substrate concentration without IL

anion depends on the degree of hydrophobicity of the cation. With TfXYN10A, the acetate anion caused minor activation in the 4% substrate, while the propionate and the guaiacolate ILs did not cause any significant activation (Hebal et al. 2020). With [DBUH]OAc and [DBUH]CO<sub>2</sub>Et, the relative PhEG activity was doubled with 4% CMC when compared to the level without ILs (Fig. 3). Similarly, the activity more than doubled with [TMGH]guaiacolate and [DBUH]guaiacolate (Fig. 4). With such enhanced activity, the use of these types of ILs would be advantageous in industrial, high substrate conditions.

#### Comparison of PhEG activity in CMC and avicel

Since PhEG can hydrolyse both soluble cellulose (CMC) and insoluble microcrystalline cellulose




**Fig. 5** Comparison between activities of *Pyrococcus horikoshii* endoglucanase (PhEG) on 1% carboxymethylcellulose (CMC) and 1% Avicel in the presence of 15% ionic liquids. Activity was measured at 70 °C and pH 6. 100% activity denotes the activity without IL

(Avicel), we tested how ILs affect enzyme activity with these substrates. Comparison of the two substrates revealed two distinct activity patterns. In [DBNH]CO<sub>2</sub>Et, [DBUH]OAc, [DBUH]CO<sub>2</sub>Et and [EMIM]OAc, enzyme activity was at the same level for both substrates (Fig. 5). These ILs have a greater cellulose solubilising capability compared to the remaining ILs (Parviainen et al. 2013, 2014), which suggests that at least partial Avicel solubilisation or swelling could have occurred, leading to similar relative enzyme activity as observed with soluble CMC. Rahikainen et al. (2018) found that the most endoglucanase-compatible ILs had the greatest potential for pulp fibre swelling. Although [DBNH]OAc has an enhanced cellulose-dissolving capability, only 30% activity was observed on Avicel. This IL formed a viscous solution with Avicel, which could be the reason for the low activity. This observation indicates that some modification in the state of Avicel occurs in these ILs. For the ILs that are less dissolving, i.e. ([mDBN]Me<sub>2</sub>PO<sub>4</sub>, [TMGH]OAc and



[TMGH]CO<sub>2</sub>Et), or those that do not dissolve cellulose (i.e. guaiacolate ILs) (Parviainen et al. 2013, 2014; Hebal et al. 2020), the activity on Avicel was lower than the activity on CMC (Figs. 5 and 6). These results directly show the influence of substrate solubility on enzyme tolerance to ILs. Chawachart et al. (2014) reported that the xylanase TasXyn10A was more strongly inhibited with [EMIM]OAc when short molecule pNP-xylose was used as a substrate instead of the longer xylan chains. Work by Hu et al. (2016) showed that inhibition of cellulase by ILs was substrate-dependant. However, the studied cellulase was a mixture of endoglucanases, exoglucanases and  $\beta$ -glucosidases, and the IL may differentially affect each enzyme specific for each substrate.


In this study, [EMIM]OAc permitted full activity (even slightly enhanced activity) of PhEG in both substrates at 15% concentration at 70 °C (Fig. 5). In



**Fig. 6** Comparison between activities of *Pyrococcus horikoshii* endoglucanase (PhEG) on 1% carboxymethylcellulose (CMC) and 1% Avicel in the presence of 5% guaiacolate ionic liquids. Activity was measured at 70 °C and pH 6. 100% activity denotes the activity without IL

contrast, Datta et al. (2010) observed a 10% decrease in the sugar yield when PhEG hydrolysed pretreated Avicel in the presence of 15% [EMIM]OAc at 80 °C. This difference may be due to the differences in the reaction conditions, as discussed above. In our study, a clear decrease in enzyme activity was observed with Avicel but not with CMC when [EMIM]OAc was increased to 35% concentration (Fig. 7). This decrease in activity suggests that the extent of Avicel solubilisation at 35% concentration was not sufficient to completely overcome the greater competitive inhibition at the more elevated IL concentration.

The tolerance of PhEG to 35% [EMIM]OAc in hydrolysing crystalline cellulose may have practical importance. It has been shown that relatively low [EMIM]OAc concentrations (25–50% w/v) in water may be effective in pretreating biomass (Fu and Mazza 2011). Therefore, it could be possible to perform with IL tolerant enzymes a simple one pot reaction procedure combining simultaneous aqueous IL and enzyme treatments.



**Fig. 7** Comparison between activities of *Pyrococcus horikoshii* endoglucanase (PhEG) on 1% carboxymethylcellulose (CMC) and 1% Avicel in the presence of 15–35% [EMIM]OAc. Activity was measured at 70 °C and pH 6. 100% activity denotes the activity without IL

## Effect of ionic liquids on the kinetic parameters of PhEG

To further clarify the mechanism by which IL molecules affect enzyme activity, the Michaelis–Menten kinetic parameters  $K_m$  (Michaelis constant) and  $V_{max}$  (maximum reaction velocity) values of the enzymatic reaction were determined in the presence of 5% ILs (Table 2).  $V_{max}$  was measured as a relative value regarding the value without ILs, which gives a comparative result for the effect of ILs.

In this study, all ILs in 5% concentration increased  $K_m$ , which is compatible with the premise that ILs cause competitive inhibition by binding transiently to the active site (Hebal et al. 2021). Among the cations, [DBUH]<sup>+</sup> conjugated to all anions and [TMGH]<sup>+</sup> conjugated to guaiacolate, caused the greatest increase in  $K_m$  (Table 2). Moreover,  $V_{max}$  also increased, but the increase was lower than that observed for  $K_m$ , which indicates that ILs had a greater inhibiting than activating effect on the enzyme in these conditions.

There was an approximate linear correlation between the increase in  $V_{max}$  in the presence of ILs and the increase in enzyme activity with 4% substrate in the presence of ILs (Supplementary Fig. 5). This suggests that  $V_{max}$  reflects the ability of the enzyme to be stimulated when competitive inhibition is relieved. Similarly, enzyme activity in 25% ILs correlates with the relative catalytic efficiency (relative  $V_{max}/K_m$ ) in the presence of ILs (Supplementary Fig. 6), which indicates that the theoretical  $K_m$  and  $V_{max}$  values in the presence of ILs successfully capture the effects of ILs on enzyme activity. An increase in  $K_m$  tends

to lead to an increase in  $V_{max}$ , which was clearly seen in the effect of ILs on PhEG activity (Supplementary Fig. 7).

The increase in both  $K_m$  and  $V_{max}$  values or  $k_{cat}$  (catalytic rate constant) has also been observed for a hyperthermostable  $\beta$ -glucosidase (Kudou et al. 2014) and other enzymes (Daneshjoo et al. 2011; da Silva and de Castro 2018) in ILs. The increase in  $V_{max}$  partly mitigates the effect of competitive inhibition when weaker interaction with the enzyme may improve the product release. In addition, if the ILs reshape the active site interactions, this could affect the activity levels, even positively, in the presence of ILs. Yu et al. (2013) concluded that the increase in both  $V_{max}$  and  $K_m$  values for laccase in aqueous solutions of tetramethylammonium trifluoromethanesulfonate ([TMA]TfO) was associated with a more compact protein structure (resulting from alteration of enzyme surroundings by IL). They suggested that this compact structure hinder the approaching of the substrate to the active site, but also enhances the catalytic rate. A simulation study of GH11 xylanase in an aqueous [EMIM]OAc solution suggested that the strong binding of cations in the enzyme active site may potentially reduce or enhance activity by inducing new geometries or electronic structures that affect the transition states (Jaeger and Pfaendtner 2013). Another simulation study of hyperthermostable xylanase TmXYN10B in an aqueous [EMIM]OAc solution revealed conformational changes to the active site induced by the [EMIM]<sup>+</sup> cations (Manna and Ghosh 2021). It is possible that the strong increase in enzyme activity by ILs in elevated substrate

**Table 2** Effect of 5% ionic liquids on *Pyrococcus horikoshii* endoglucanase (PhEG) kinetic parameters. Activity was measured at 70 °C and pH 6

Ionic liquid	Relative $V_{max}$	$K_m$ (mg/ml)	Relative $V_{max}/K_m$
None	1.000 ± 0.002	1.007 ± 0.306	1.064 ± 0.357
[DBNH]OAc	1.495 ± 0.098	3.273 ± 1.136	0.484 ± 0.119
[DBNH]CO <sub>2</sub> Et	1.72 ± 0.085	5.580 ± 0.225	0.308 ± 0.003
[DBNH]guaiacolate	1.418 ± 0.175	4.233 ± 0.92	0.34 ± 0.031
[DBUH]OAc	2.498 ± 0.152	13.466 ± 2.3	0.187 ± 0.02
[DBUH]CO <sub>2</sub> Et	2.697 ± 0.166	7.698 ± 0.768	0.282 ± 0.015
[DBUH]guaiacolate	2.464 ± 0.123	8.474 ± 0.125	0.3 ± 0.016
[EMIM]OAc	1.200 ± 0.104	1.668 ± 0.215	0.724 ± 0.053
[mDBN]Me <sub>2</sub> PO <sub>4</sub>	1.551 ± 0.069	1.955 ± 0.405	0.811 ± 0.13
[TMGH]OAc	1.879 ± 0.104	4.570 ± 0.269	0.413 ± 0.036
[TMGH]CO <sub>2</sub> Et	1.647 ± 0.069	3.760 ± 1.598	0.488 ± 0.177
[TMGH]guaiacolate	3.071 ± 0.342	11.92 ± 0.444	0.258 ± 0.016

concentrations is mediated both by IL interactions with the substrate and the enzyme.

In contrast to PhEG, which experienced even a strong increase in  $V_{\max}$  by 5% ILs, the  $V_{\max}$  value associated with TfXYN10A remained the same, increased only slightly with two ILs or decreased at 5% superbase ILs (Hebal et al., 2020). However, an increase in  $V_{\max}$  was seen for TfXYN10A at elevated concentrations of [EMIM]OAc and [DBNH]OAc (15 and 35%) (Anbarasan et al. 2017). In addition, the  $V_{\max}$  values of other hyperthermostable xylanases, which show less tolerance to [EMIM]OAc (in contrast to PhEG), were not affected or decreased in the presence of this IL (Li et al. 2013; Chawachart et al. 2014; Yu et al. 2016). These results indicate that activation of PhEG, as reflected by the increase in  $V_{\max}$ , contributes to a general greater tolerance to superbase ILs of this enzyme. Moreover, PhEG has been found to be partly tolerant to 40% concentration of another set of ILs, i.e. ([DMIM]DMP, [BMIM]DBP, [Chol]AcO, [BMIM]DMP, [TMGH]n-PrCOO, and [EMIM]DMP), and retained full activity (107%) in [TMGH]n-PrCOO (Rahikainen et al. 2018). However, the effects of insoluble Avicel and elevated substrate concentration were not studied by Rahikainen et al. (2018).

The [DBUH]<sup>+</sup> cation systematically triggered the greatest increase in kinetic parameters with all anions, which indicates the strong effect of this cation on enzyme functioning. In line with this finding, almost no differences were observed in the kinetic parameters between the different anions when conjugated to [DBUH]<sup>+</sup>. This indicates that the studied anions, in general, have a lesser effect on enzyme activity than cations. Anions showed the strongest effect when the larger guaiacolate anion was conjugated to [TMGH]<sup>+</sup>, which is also an indication of the cooperative effect of these two ions in influencing PhEG activity.

[DBUH]<sup>+</sup> differs from the other cations by its greater hydrophobicity (the amount of hydrophobic C and H atoms: [DBUH]<sup>+</sup> 25; [mDBN]<sup>+</sup> 23; [DBNH]<sup>+</sup> 19; [EMIM]<sup>+</sup> 17 and [TMGH]<sup>+</sup> 17), which suggests a role for hydrophobic interactions in the IL interaction with the active site and in activation of the enzyme. Furthermore, PhEG showed lowered activity in the presence of 15% [Chol]OAc (Rahikainen et al. 2018). This IL has lower hydrophobicity compared to the acetate-based ILs used in our study. Differences between [HMIM]<sup>+</sup> and [BMIM]<sup>+</sup> ILs in


concentrations < 40% in the activation of lipase have been attributed to the difference in the length of the alkyl chain (hydrophobicity) attached to the cation (Daneshjoo et al. 2011). [HMIM]Cl has been reported to increase  $K_m$  and  $V_{\max}$  values more than [BMIM]Cl in IL concentrations < 25%, and enzyme activation has been suggested to result from structural changes induced by ILs in the protein (Daneshjoo et al. 2011).

In a simulation study by Jaeger et al. (2015), PhEG exhibited greater stability at 80 °C in 50% [EMIM]OAc concentration compared to stability in pure water, and the presence of IL resulted in the formation of non-native salt bridges in PhEG. Dilute solutions of [EMIM]OAc also induced stabilisation of hyperthermostable  $\beta$ -glucosidase and xylanases (Kudou et al. 2014; Yu et al. 2016; Anbarasan et al. 2017). The stabilisation of a three  $\alpha$ -helix bundle in [BMIM]Cl was found to be due to electrostatic and hydrophobic interactions of the [BMIM]<sup>+</sup> cation with residues on the protein surface, and such interactions could remove the surrounding water molecules, reduce the hydrogen bonding from water to protein, and thus stabilise the backbone hydrogen bonds (Shao, 2013). Although ILs seem to unfold the proteins (Zhao 2016; Smiatek 2017), these results indicate that the effect of ILs on the activity of enzymes may also include stabilisation of the enzyme structure by interactions at the active site and other parts of the protein. When the enzyme is hyperthermostable, the stabilising effect of ILs could become dominating.

#### Binding of ionic liquid molecules to the active site

Molecular docking with [DBNH]<sup>+</sup>, [DBUH]<sup>+</sup>, [EMIM]<sup>+</sup>, [mDBN]<sup>+</sup> and [TMGH]<sup>+</sup> was performed to correlate the overall binding behaviour among the observed, about 250 binding poses to the enzyme activity in ILs (Supplementary Table 1). The IL cations are trapped at the active site canyon of the enzyme in the same areas where the substrate binds. The binding areas for the [EMIM]<sup>+</sup> cation and the cellotetraose substrate are shown in Fig. 8. The figure indicates how the IL cations are trapped onto the active site canyon of the enzyme in the same areas where the substrate binds. The tetrameric substrate shows the key binding areas for the substrate, which typically is much longer than four sugar units. Therefore, the actual substrate that replaces ionic liquid cations from the active site may bind to the enzyme

**Fig. 8** The potential [EMIM]<sup>+</sup> cation binding sites (a), and the cello-tetraose substrate molecule (b) shown on the surface of *Pyrococcus horikoshii* endoglucanase (PhEG). Cellotetraose is from 3QHM. Ligands are shown in cyan



in a larger area than where the IL cations are detected to bind with the observed binding energies (see Supplementary Table 1). Basically, only a few cations can bind concurrently to the active site, and they largely fill it (Supplementary Fig. 9). Other sites for binding of the cation are also observed outside the enzyme active site (Fig. 8). The location and interaction pattern of the highest energy binder of the [EMIM]<sup>+</sup> cation in a pocket above the catalytic residues of PhEG are shown in Supplementary Fig. 10. The positive charge of [EMIM]<sup>+</sup> is located close to the catalytic glutamate side chains of PhEG, as was reported earlier for this enzyme (Jaeger et al. 2015), and for other endoglucanases and xylanases (Hebal et al. 2021). The differences in the potential cation binding areas in the active site among the five IL cations are shown in Supplementary Fig. 11. The [EMIM]<sup>+</sup> cation shows the smallest potential binding area, which reflects its lower inhibition.

In our study, docking with IL anions (acetate, propionate, butyrate and guaiacolate) indicates that as expected the anion with a larger hydrophobic group is better able to fill the PhEG active site canyon (Supplementary Fig. 12), as has also been shown for GH10 xylanase TfXYN10A (Hebal et al. 2020). The difference is that, in PhEG, the anions bind at sites all over the active site, whereas there is a separate anion binding site at the end of the active site cleft in TfXYN10A (Hebal et al. 2020). The greater competitive inhibition of PhEG may be explained by the stronger binding of cations to PhEG (−8.06 to −8.40 kcal/mol)

compared to TfXYN10A (−6.4 to −7.0 kcal/mol) (Hebal et al. 2020). Moreover, in PhEG, the cations show a greater binding energy compared to the anions (−6.4 to −7.1), thereby indicating that cation binding to the active site could be stronger than the binding of anions. In contrast, similar binding energies were observed for the same cations (−6.4 to −7.0 kcal/mol) and anions (−6.48 to −7.65 kcal/mol) in TfXYN10A xylanase (Hebal et al. 2020).

The ring structures of inhibiting cations ([DBUH]<sup>+</sup>, [DBUH]<sup>+</sup> and [mDBN]<sup>+</sup> (see Supplementary Fig. 8) are likely to bind to the same active site positions as the sugar ring structures of the substrate (Chawachart et al. 2014; Summers et al. 2020). Reasons why the [EMIM]<sup>+</sup> cation is more enzyme friendly compared to the other studied IL molecules could be because it lacks a double ring structure, it is a basic cation, and is not too hydrophobic (see Supplementary Fig. 8). Our docking results indicate that binding of several IL molecules simultaneously onto the active site contributes to the overall effect on enzyme activity, and that there are significant differences in the binding profile between the various IL molecules and between the different enzymes.

## Conclusions

Although the different studies suggest a quite similar general inhibition mechanism for the different enzymes, there exists great variation in the effects

of ILs between the enzymes. Superbase ILs can either inhibit or activate PhEG, depending on conditions and the nature of the substrate. The activation of PhEG, in larger extent than TfXYN10A, seems to be the main difference explaining the greater tolerance of this endoglucanase to these ILs. Furthermore, the greater tolerance displayed by PhEG to many ILs at elevated substrate concentrations indicates that it could be successfully used in industrial, high substrate conditions with a range of ILs. Differences in the binding of ILs to the active sites of PhEG and TfXYN10A appears to determine the differing effect of ILs. These findings could assist in the planning of rational approaches to the design of new enzyme friendly ILs and to the engineering of new IL-tolerant enzymes. The success achieved to date in improving the tolerance of cellulases to ILs in some of the rational approaches is promising (Lee et al. 2017; Pramanik et al. 2021; Summers et al. 2020). Important aspects to consider in planning are that enzymes differ in their response to various ILs and that different substrates may behave differently in IL solutions.

**Acknowledgements** We thank Johanna Aura for technical assistance and Arno Parviainen for assistance in the manufacture of the ionic liquids. We acknowledge the funding from CLIC Innovation (ACel Programme) and Business Finland.

**Supplementary information** Calibration graphs for glucose (Supplementary Fig. 1); time-dependent hydrolysis graphs for PhEG (Supplementary Fig. 2); graphs for the calculation of kinetic parameters in ILs (Supplementary Fig. 3); effect of NaCl and KCl on the activity of PhEG (Supplementary Fig. 4); correlation between relative enzyme activity with 4% substrate in the presence of ILs and relative  $V_{\max}$  (Supplementary Fig. 5); correlation between relative enzyme activity in 25% IL and relative  $V_{\max}/K_m$  in the presence of ILs (Supplementary Fig. 6); correlation between  $K_m$  and  $V_{\max}$  in the presence of ILs (Supplementary Fig. 7); molecular docking of ionic liquids cations and anions to PhEG by SwissDock (Supplementary Figs. 8–12); molecular docking results and PhEG activities in ILs (Supplementary Table 1).

**Author contributions** HH: performed all experimental work, analysed and interpreted the data, and wrote the manuscript; OT: supervised the work, performed modelling, analysed and interpreted the data, wrote and revised the manuscript; JH: produced the enzyme; LM: performed modelling; AWTK and IK: synthesised the ionic liquids; SB, NB: research administration. All authors approved the final manuscript.

**Funding** Open access funding provided by University of Eastern Finland (UEF) including Kuopio University Hospital.

This work was supported by CLIC Innovation (ACel Programme) and Business Finland.

## Declarations

**Competing interests** The authors declare that they have no competing interests.

**Ethical approval and consent to participate** This article does not contain any studies performed by any of the authors on human participants or animals.

**Consent to publish** Not applicable.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Ajlou D, Sangian M, Ghadamgahi M, Evini M, Saboury AA (2013) Effect of two imidazolium derivatives of ionic liquids on the structure and activity of adenosine deaminase. *Int J Biol Macromol* 55:47–61. <https://doi.org/10.1016/j.ijbiomac.2012.12.042>
- Alvira P, Tomas-Pejo E, Ballesteros M, Negro MJ (2010) Pre-treatment technologies for an efficient bioethanol production process based on enzymatic hydrolysis: a review. *Bioresour Technol* 101:4851–4861. <https://doi.org/10.1016/j.biortech.2009.11.093>
- Anbarasan S, Wahlström R, Hummel M, Ojamo H, Sixta H, Turunen O (2017) High stability and low competitive inhibition of thermophilic *Thermopolyspora flexuosa* GH10 xylanase in biomass-dissolving ionic liquids. *Appl Microbiol Biotechnol* 101:1487–1498. <https://doi.org/10.1007/s00253-016-7922-9>
- Ando S, Ishida H, Kosugi Y, Ishikawa K (2002) Hyperthermostable endoglucanase from *Pyrococcus horikoshii*. *Appl Environ Microbiol* 68:430–433. <https://doi.org/10.1128/AEM.68.1.430-433.2002>
- Auxenfans T, Husson E, Sarazin C (2017) Simultaneous pre-treatment and enzymatic saccharification of (ligno) celluloses in aqueous-ionic liquid media: a compromise. *Biochem Eng J* 117:77–86. <https://doi.org/10.1016/j.bej.2016.10.004>

- Barbosa FC, Silvello MA, Goldbeck R (2020) Cellulase and oxidative enzymes: new approaches, challenges and perspectives on cellulose degradation for bioethanol production. *Biotechnol Lett* 42:875–884. <https://doi.org/10.1007/s10529-020-02875-4>
- Ben Hmad I, Boudabbous M, Belghith H, Gargouri A (2017) A novel ionic liquid-stable halophilic endoglucanase from *Stachybotrys microspore*. *Process Biochem* 54:59–66. <https://doi.org/10.1016/j.procbio.2017.01.007>
- Chacón MG, Ibenegbu C, Leak DJ (2021) Simultaneous saccharification and lactic acid fermentation of the cellulosic fraction of municipal solid waste using *Bacillus smithii*. *Biotechnol Lett* 43:667–675. <https://doi.org/10.1007/s10529-020-03049-y>
- Chawachart N, Anbarasan S, Turunen S, Li H, Khanongnuch C, Hummel M, Sixta H, Granstrom T, Lumyong S, Turunen O (2014) Thermal behaviour and tolerance to ionic liquid [emim]OAc in GH10 xylanase from *Thermoascus aurantiacus* SL16W. *Extremophiles* 18:1023–1034. <https://doi.org/10.1007/s00792-014-0679-0>
- Contreras F, Pramanik S, Rozhkova AM, Zorov IN, Korotkova O, Sinitsyn AP, Schwaneberg U, Davari MD (2020) Engineering robust cellulases for tailored lignocellulosic degradation cocktails. *Int J Mol Sci* 21:1589. <https://doi.org/10.3390/ijms21051589>
- da Silva VG, de Castro RJS (2018) Biocatalytic action of proteases in ionic liquids: Improvements on their enzymatic activity, thermal stability and kinetic parameters. *Int J Biol Macromol* 114:124–129. <https://doi.org/10.1016/j.ijbiomac.2018.03.084>
- Daneshjoo S, Akbari N, Sepahi AA, Ranjbar B, Khavarinejad RA, Khajeh K (2011) Imidazolium chloride-based ionic liquid-assisted improvement of lipase activity in organic solvents. *Eng Life Sci* 11:259–263. <https://doi.org/10.1002/elsc.201000154>
- Datta S, Holmes B, Park JI, Chen Z, Dibble DC, Hadi M, Blanch HW, Simmons BA, Sapra R (2010) Ionic liquid tolerant hyperthermophilic cellulases for biomass pretreatment and hydrolysis. *Green Chem* 12:338–345. <https://doi.org/10.1039/b916564a>
- Engel P, Mladenov R, Wulfhorst H, Jäger G, Spiess AC (2010) Point by point analysis: how ionic liquid affects the enzymatic hydrolysis of native and modified cellulose. *Green Chem* 12:1959–1966. <https://doi.org/10.1039/c0gc00135j>
- Ferdjani S, Ionita M, Roy B, Dion M, Djeghaba Z, Rabiller C, Tellier C (2011) Correlation between thermostability and stability of glycosidases in ionic liquid. *Biotechnol Lett* 33:1215–1219. <https://doi.org/10.1007/s10529-011-0560-5>
- Fu D, Mazza G (2011) Aqueous ionic liquid pretreatment of straw. *Bioresour Technol* 102:7008–7011. <https://doi.org/10.1016/j.biortech.2011.04.049>
- Gladden JM, Park JI, Bergmann J, Reyes-Ortiz V et al (2014) Discovery and characterization of ionic liquid-tolerant thermophilic cellulases from a switchgrass-adapted microbial community. *Biotechnol Biofuels* 7:15. <https://doi.org/10.1186/1754-6834-7-15>
- Grosdidier A, Zoete V, Michielin O (2011) SwissDock, a protein-small molecule docking web service based on EADock DSS. *Nucleic Acid Res* 39:W270–W277. <https://doi.org/10.1093/nar/gkr366>
- Hämäläinen J, Granström T, Mollerup F, Wang Y, Xiong H, Turunen O (2016) Effect of enzymatic high temperature prehydrolysis on the subsequent cellulose hydrolysis of steam-pretreated spruce in high solids concentration. *J Chem Technol Biotechnol* 91:1844–1852. <https://doi.org/10.1002/jctb.4777>
- Hebal H, Boucherba N, Binay B, Turunen O (2021) Activity and stability of hyperthermostable cellulases and xylanases in ionic liquids. *Biocatal Biotransform* 39:242–259. <https://doi.org/10.1080/10242422.2021.1882430>
- Hebal H, Parviainen A, Anbarasan S, Li H, Makkonen L, Bankar S, King AWT, Kilpeläinen I, Benallaoua S, Turunen O (2020) Inhibition of hyperthermostable xylanases by superbase ionic liquids. *Process Biochem* 95:148–156. <https://doi.org/10.1016/j.procbio.2020.03.022>
- Hu D, Xiao L, Li L, Zhong C, Ju X, Yan L, Wu T, Qing M, Hu Z (2016) Effects of ionic liquid 1-ethyl-3-methylimidazolium diethylphosphate on cellulase produced by *Paenibacillus* sp. LLZ1. *ACS Sustain Chem Eng* 4:4922–4926. <https://doi.org/10.1021/acssuschemeng.6b01229>
- Husson E, Auxenfans T, Herbaut M, Baralle M, Lambertyn V, Rakotoarivonina H, Remond C, Sarazin C (2018) Sequential and simultaneous strategies for biorefining of wheat straw using room temperature ionic liquids, xylanases and cellulases. *Bioresour Technol* 251:280–287. <https://doi.org/10.1016/j.biortech.2017.12.047>
- Ilmberger N, Meske D, Juergensen J, Schulte M, Barthen P, Rabausch U, Angelov A, Mientus M, Liebl W, Schmitz RA, Streit WR (2012) Metagenomic cellulases highly tolerant towards the presence of ionic liquids-linking thermostability and halotolerance. *Appl Microbiol Biotechnol* 95:135–146. <https://doi.org/10.1007/s00253-011-3732-2>
- Ishikawa T (2009) Superbases for organic synthesis: guanidines, amidines, phosphazenes and related organocatalysts. Wiley, Chichester
- Jaeger VW, Pfaendtner J (2013) Structure, dynamics, and activity of xylanase solvated in binary mixtures of ionic liquid and water. *ACS Chem Biol* 8:1179–1186. <https://doi.org/10.1021/cb3006837>
- Jaeger VW, Burney P, Pfaendtner J (2015) Comparison of three ionic liquid-tolerant cellulases by molecular dynamics. *Biophys J* 108:880–892. <https://doi.org/10.1016/j.bpj.2014.12.043>
- Kamiya N, Matsushita Y, Hanaki M, Nakashima K, Narita M, Goto M, Takahashi H (2008) Enzymatic in situ saccharification of cellulose in aqueous-ionic liquid media. *Biotechnol Lett* 30:1037–1040. <https://doi.org/10.1007/s10529-008-9638-0>
- Kang HJ, Ishikawa K (2007) Analysis of active center in hyperthermophilic cellulase from *Pyrococcus horikoshii*. *J Microbiol Biotechnol* 17:1249–1253
- Kim HW, Ishikawa K (2009) Structure of hyperthermophilic endocellulase from *Pyrococcus horikoshii*. *Proteins* 78:496–500. <https://doi.org/10.1002/prot.22602>
- King AWT, Asikkala J, Mutikainen I, Järvi P, Kilpeläinen I (2011) Distillable acid-base conjugate ionic liquids for cellulose dissolution and processing. *Angew Chem Int Ed* 50:6301–6305. <https://doi.org/10.1002/anie.201100274>
- Kudou M, Kubota Y, Nakashima N, Okazaki F, Nakashima K, Ogino C, Kondo A (2014) Improvement of enzymatic

- activity of  $\beta$ -glucosidase from *Thermotoga maritima* by 1-butyl-3-methylimidazolium acetate. *J Mol Catal B Enzym* 104:17–22. <https://doi.org/10.1016/j.molcatb.2014.02.013>
- Lee JM, Moon SY, Kim YR, Kim KW, Lee BJ, Kong IS (2017) Improvement of thermostability and halostability of b-1,3-1,4-glucanase by substituting hydrophobic residue for Lys48. *Int J Biol Macromol* 94:594–602. <https://doi.org/10.1016/j.ijbiomac.2016.10.043>
- Li H, Kankaanpää A, Xiong H, Hummel M, Sixta H, Ojamo H, Turunen O (2013) Thermostabilization of extremophilic *Dictyoglomus thermophilum* GH11 xylanase by an N-terminal disulfide bridge and the effect of ionic liquid [EMIM]OAc on the enzymatic performance. *Enzyme Microb Technol* 53:414–419. <https://doi.org/10.1016/j.enzmictec.2013.09.004>
- Manna B, Ghosh A (2021) Understanding the conformational change and inhibition of hyperthermophilic GH10 xylanase in ionic liquid. *J Mol Liq* 332:115875. <https://doi.org/10.1016/j.molliq.2021.115875>
- Mckendry P (2002) Energy production from biomass (part 1): overview of biomass. *Bioresour Technol* 83:37–46. [https://doi.org/10.1016/S0960-8524\(01\)00118-3](https://doi.org/10.1016/S0960-8524(01)00118-3)
- Miller GL (1959) Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem* 31:426–428. <https://doi.org/10.1021/ac60147a030>
- Nemestóthy N, Megyeri G, Bakonyi P, Lakatos P, Koók L, Polakovic M, Gubicza L, Bélafi-Bakó K (2017) Enzyme kinetics approach to assess biocatalyst inhibition and deactivation caused by [bmim][Cl] ionic liquid during cellulose hydrolysis. *Bioresour Technol* 229:190–195. <https://doi.org/10.1016/j.biortech.2017.01.004>
- Parviainen A, King AWT, Mutikainen I, Hummel M, Selg C, Hauru LKJ, Sixta H, Kilpelainen I (2013) Predicting cellulose solvating capabilities of acid-base conjugate ionic liquids. *Chemsuschem* 6:2161–2169. <https://doi.org/10.1002/cssc.201300143>
- Parviainen H, Parviainen A, Virtanen T, Kilpelainen I, Ahvenainen P, Serimaa R, Gronqvist S, Maloney T, Maunu SL (2014) Dissolution enthalpies of cellulose in ionic liquids. *Carbohydr Polym* 113:67–76. <https://doi.org/10.1016/j.carbpol.2014.07.001>
- Patel R, Kumari M, Khan AB (2014) Recent advances in the applications of ionic liquids in protein stability and activity: a review. *Appl Biochem Biotechnol* 172:3701–3720. <https://doi.org/10.1007/s12010-014-0813-6>
- Pramanik S, Semenova MVM, Rozhkova A, Zorov IN, Korotkova O, Sinitsyn AP, Davari MD (2021) An engineered cellobiohydrolase I for sustainable degradation of lignocellulosic biomass. *Biotechnol Bioeng* 118:4014–4027. <https://doi.org/10.1002/bit.27877>
- Puligundla P, Mok C (2021) Valorization of sugar beet pulp through biotechnological approaches: recent developments. *Biotechnol Lett* 43:1253–1263. <https://doi.org/10.1007/s10529-021-03146-6>
- Rahikainen J, Anbarasan S, Wahlström R, Parviainen A, King AWT, Puranen T, Kruus K, Kilpelainen I, Turunen O, Suurnakki A (2018) Screening of glycoside hydrolases and ionic liquids for fibre modification. *J Chem Technol Biotechnol* 93:818–826. <https://doi.org/10.1002/jctb.5435>
- Raulo R, Heuson E, Froidevaux R, Phalip V (2021) Combining analytical approaches for better lignocellulosic biomass degradation: a way of improving fungal enzymatic cocktails? *Biotechnol Lett* 43:2283–2298. <https://doi.org/10.1007/s10529-021-03201-2>
- Shao Q (2013) On the influence of hydrated imidazolium-based ionic liquid on protein structure stability: a molecular dynamics simulation study. *J Chem Phys* 139:115102. <https://doi.org/10.1063/1.4821588>
- Smiatek J (2017) Aqueous ionic liquids and their influence on protein structures: an overview on recent theoretical and experimental insights. *J Phys Condens Matter* 29:233001. <https://doi.org/10.1088/1361-648X/aa6c9d>
- Summers S, Kraft C, Alamdari S, Pfaendtner J, Kaar JL (2020) Enhanced activity and stability of *Acidothermus cellulolyticus* endoglucanase I in ionic liquids via engineering active site residues and non-native disulfide bridges. *ACS Sustain Chem Eng* 8:11299–11307. <https://doi.org/10.1021/acssuschemeng.0c03242>
- Wahlström RM, Suurnakki A (2015) Enzymatic hydrolysis of lignocellulosic polysaccharides in the presence of ionic liquids. *Green Chem* 17:694–714. <https://doi.org/10.1039/c4gc01649a>
- Wahlström R, King A, Parviainen A, Kruus K, Suurnakki A (2013) Cellulose hydrolysis with thermo- and alkali-tolerant cellulases in cellulose-dissolving superbase ionic liquids. *RSC Adv* 3:20001–20009. <https://doi.org/10.1039/b000000x>
- Yu X, Zou F, Li Y, Lu L, Huang X, Qu Y (2013) Effect of three trifluoromethanesulfonate ionic liquids on the activity, stability and conformation of laccase. *Int J Biol Macromol* 56:62–68. <https://doi.org/10.1016/j.ijbiomac.2013.02.005>
- Yu T, Anbarasan S, Wang Y, Telli K, Aslan AS, Su Z, Zhou Y, Zhang L, Iivonen P, Havukainen S, Mentunen T, Hummel M, Sixta H, Binay B, Turunen O, Xiong H (2016) Hyperthermostable *Thermotoga maritima* xylanase XYN10B shows high activity at high temperatures in the presence of biomass-dissolving hydrophilic ionic liquids. *Extremophiles* 20:515–524. <https://doi.org/10.1007/s00792-016-0841-y>
- Zhao H (2010) Methods for stabilizing and activating enzymes in ionic liquids - a review. *J Chem Technol Biotechnol* 85:891–907. <https://doi.org/10.1002/jctb.2375>
- Zhao H (2016) Protein stabilization and enzyme activation in ionic liquids: specific ion effects. *J Chem Technol Biotechnol* 91:25–50. <https://doi.org/10.1002/jctb.4837>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.