

<https://helda.helsinki.fi>

Energy Intake and Severity of Dementia Are Both Associated with Health-Related Quality of Life among Older Long-Term Care Residents

KS, Salminen

Multidisciplinary Digital Publishing Institute
2019-09-20

KS, S.; MH, S.; H, K.; HM, R.; KH, P. Energy Intake and Severity of Dementia Are Both Associated with Health-Related Quality of Life among Older Long-Term Care Residents. *Nutrients* 2019, 11, 2261.

<http://hdl.handle.net/10138/348569>

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Article

Energy Intake and Severity of Dementia Are Both Associated with Health-Related Quality of Life among Older Long-Term Care Residents

Salminen KS ^{1,2,*} , Suominen MH ¹, Kautiainen H ¹, Roitto HM ^{1,3} and Pitkala KH ^{1,4}

¹ University of Helsinki, Department of General Practice and Primary Health Care, POB 20, 00014 Helsinki, Finland; merja.suominen@gery.fi (S.M.H.); hannu.kautiainen@medcare.fi (K.H.); hanna-maria.roitto@hel.fi (R.H.M.); kaisu.pitkala@helsinki.fi (P.K.H.)

² Vantaa Social Welfare and Health Care, POB 1600, 01030 Vantaa, Finland

³ Department of Social Services and Health Care, Geriatric Clinic, Helsinki Hospital, POB 6600, 00099 Helsinki, Finland

⁴ Helsinki University Hospital, Unit of Primary Health Care, 00014 Helsinki, Finland

* Correspondence: karoliina.salminen@helsinki.fi; Tel.: +35-840-540-0675

Received: 17 August 2019; Accepted: 18 September 2019; Published: 20 September 2019



Abstract: Our aim was to investigate how energy intake modifies the association of the stage of dementia with health related quality of life (HRQoL) among institutionalized older people. A cross-sectional sample of 538 older long-term care residents with dementia in Helsinki, Finland were assessed with HRQoL (15D), energy intake (from one to two days), and the stage of dementia by the clinical dementia rating (CDR) scale. The energy intakes were standardized by z-scores to include both men and women in the same analyses. Severity of dementia was associated with HRQoL (15D index in CDR 0.5–1: 0.65 (0.11), CDR 2: 0.60 (0.10), CDR 3: 0.52 (0.10)). When the three groups of dementia severity were divided according to their energy intake quartiles, there was an association between the HRQoL and the stage of dementia ($p < 0.001$) and energy intake ($p = 0.013$); however, no interaction was observed ($p = 0.30$). While partial correlation analysis showed that energy intake correlated with HRQoL among residents with very mild/mild or moderate dementia, this was not observed among those with severe dementia. In moderate dementia, the dimensions of mobility and usual activities correlated significantly with higher energy intake. Both energy intake and severity of dementia are associated with HRQoL.

Keywords: energy intake; health-related quality of life; stage of dementia; long-term care

1. Introduction

When measured by the Mini-Nutritional Assessment (MNA), the prevalence of malnutrition in nursing homes in Europe ranges from 14–21% and from 21–40% in long-term care [1]. However, in an earlier study on a heterogeneous sample of older people whose nutritional status was normal according to the MNA, only approximately 40% of participants reached the amount of energy and 25% reached the amount of protein recommended by the Nordic Nutrition Recommendations [2,3]. Engelheart et al. observed that there was no correlation between nutritional status and energy or protein intake among older people living at home or in a nursing home [4]. Thus, assessing energy and protein intake might be a more sensitive assessment than a mere MNA screening tool to detect those at risk of developing malnutrition [2,4,5].

The energy and protein intake in institutionalized older people varies between studies depending on culture, dementia stage, age, and gender distribution. A Dutch study showed that only 18% of residents from a nursing home rehabilitation ward received adequate energy and protein intake [6].

In this Dutch study ($n = 79$), protein intake was considered sufficient if the resident reached ≥ 1.2 g/bodyweight, and energy intake was considered sufficient if the resident reached $\geq 85\%$ of their energy need, as calculated by Harris and Benedict + 30%. In a Swedish study ($n = 128$), 16% of the residents living in a nursing home were malnourished according to the MNA, and received a mean energy intake of 1724 kcal/day and mean protein intake of 63 g/day [4]. A study on Finnish residents ($n = 374$) from assisted living facilities revealed a mean energy intake of 1762 kcal/day and a protein intake of 64 g/day; 17% of residents were malnourished according to MNA [7]. In a recently published Spanish study among nursing home residents ($n = 249$), the mean energy intake was 1624 kcal/day, protein was 60 g/day, and 17% of the residents were malnourished according to MNA short-form [5]. Previous studies on long-term care have shown that especially residents with severe dementia and disabilities are at risk for malnutrition [2,8]. Low energy intake has been associated with sarcopenia [5,9] and frailty [10].

Health-related quality of life (HRQoL) should be considered a goal of good aging [11]. HRQoL is a multidimensional concept that includes an individual's perceived physical and mental health. Quality of life (QoL) is a broader concept; in addition to health and functioning, it includes psychological, social, and environmental dimensions [12].

In some studies, QoL decreased over follow-up time, while in others, QoL was unchanged or even increased among institutionalized persons with dementia [13–17]. These studies differ from each other in follow-up time, methods used to evaluate QoL, and settings. This may explain the differences between studies. Thus, there are contradictory findings concerning how HRQoL is associated with the stage of dementia among institutionalized older residents.

Malnutrition is associated with poor HRQoL among non-institutionalized older people [11,18–22]. Physical performance and functioning declines with malnutrition, and this decline may also impair HRQoL [21]. Wei et al. observed that nutritional deterioration is associated with poor QoL during follow-up among community-living older people [23]. In a previous study, increased energy intake is associated with better HRQoL among non-institutionalized older people and home-care patients [11,24]. Studies investigating the association between energy intake and HRQoL in different stages of dementia are scarce.

Nutrition and food intake are important aspects of QoL among residents in nursing home settings. Nutritional status is an important aspect of HRQoL and psychological well-being [22,25,26]. However, less is known about how nutrient intake is associated with HRQoL. To our knowledge, there are no studies that have explored how energy and food intake modify HRQoL in various stages of dementia. The aim of this study was to determine how dementia severity is associated with HRQoL and how nutrient intake modifies this association.

2. Material and Methods

2.1. Study Sample

This was a cross-sectional study designed to assess various aspects of nutrition among older residents in institutional settings in Helsinki, Finland. The study aimed to obtain a comprehensive picture of nutritional status, nutrient intake, nutritional care, and related factors in institutional settings. The participants were recruited from three nursing homes and 14 assisted living facilities in Helsinki. Although the level of care is basically the same in nursing homes and in assisted living facilities, the residents are more mobile in the latter. In assisted living facilities, residents have their own room and furnishings, while in nursing homes, residents have shared rooms. For residents with dementia, there are group homes in assisted living facilities. Continuous assistance is available and a registered nurse is in charge of the ward.

The inclusion criteria for the present study were: aged ≥ 65 years; native Finnish or Swedish speaker (all participants were Caucasian); living permanently in institutional care; sufficient information on demographic factors; and assessed with a clinical dementia rating (CDR), a one to two day food record, and HRQoL according to the 15D instrument. Altogether 538 residents met these criteria and were included.

The local ethics committee of the Helsinki University Hospital approved the study (No. HUS/2042/2016). Informed consent was acquired from all participants or from their closest proxies in case of moderate-severe dementia (Mini-Mental State Examination (MMSE) < 20 points).

2.2. Measurements

Trained nurses collected the data. The resident's demographic information, diagnoses, and current use of medications were retrieved from medical records. The Charlson Comorbidity Index (CCI) was used to evaluate comorbidities. The CCI includes both the number and severity of a resident's medical conditions. A higher score indicates a greater burden of comorbidities [27].

Evaluation of the stage of the dementia was performed according to the CDR [28]. Participants were divided into categories of having very mild/mild dementia (CDR 0.5–1), moderate dementia (CDR 2), and severe dementia (CDR 3) according to this measurement. Cognition was assessed by the MMSE [29]. The resident's dependence in physical functioning was assessed by the CDR "personal care" question (1 = totally independent; 2 = needs prompting; 3 = requires assistance in dressing, personal hygiene, and keeping of personal belongings; 4 = requires much help with personal care, repeatedly incontinent). CDR "personal care" ≥ 2 was defined as dependence in physical functioning.

The MNA was used to evaluate nutritional status. The MNA is a validated nutrition screening and assessment tool [30]. The MNA has a maximum score of 30 points; <17 points indicates malnutrition, 17–23.5 indicates risk of malnutrition, and >23.5 indicates good nutrition status [30,31]. In addition, each resident was weighed and body mass index (BMI) was calculated as weight divided by height squared (kg/m^2).

Energy and protein intake was determined by a one to two day food record collected by ward nurses. The food records were analyzed by using AivoDiet dietary software (version 2.2.0.0, Aivo Oy, Turku, Finland). The AivoDiet software includes the Fineli Food Composition database Release 16 (2013). The database includes recipes for typical Finnish mixed dishes that are usually served in long-term care. The instruction was to record all the foods and beverages consumed by the resident. Nurses estimated portion sizes by household measures. For prepacked products, the exact brand and product name was given. A suitable recipe was used during data entry.

Nutrition care was assessed with several questions addressed to ward nurses. The consumption of snacks was determined as follows: "Does the resident eat snacks between meals?" (yes/no). The amount of offered food eaten by the resident was evaluated with the following question: "How much does the resident on average eat from the main meal?" with the five options "eats only a little", "eats less than half", "eats half their meal", "eats most of their meal", or "eats all or nearly all of their meal" available as responses. The responses "eats only a little" and "eats less than half" were dichotomized as eats less than half and the responses "eats half their meal", "eats most of their meal", or "eats all or nearly all of their meal" were considered as eats more than half. Use of nutritional supplements was surveyed (yes/no). The nurses were also asked whether the participant ate normal (normal or soft) food or liquid/pureed food (yes/no). The presence of symptoms such as constipation and chewing problems were inquired with yes/no questions.

The 15D instrument was used to evaluate the resident's HRQoL. This instrument is a validated, generic measure for HRQoL [32]. The dimensions of 15D are mobility, vision, hearing, breathing, sleeping, eating, speech, excretion/elimination, usual activities, mental function, discomfort and symptoms, depression, distress, vitality, and sexual activity. The 15D can be completed during a discussion with the resident but also completed by a proxy who knows the patient well. The 15D combines the advantages of a 15-dimension profile and a single-index measure [32]. A score of 0 indicates the poorest HRQoL and 1 indicates perfect HRQoL. The 15D index is reliable, sensitive, and responsive to change [32]. If the subject was unable to respond due to poor cognition, the nurse most familiar with the resident was interviewed about the 15D.

2.3. Statistical Analysis

The statistical significance for the unadjusted hypothesis of linearity across categories (quartiles) of standardized energy intake and characteristics of the study participants were evaluated using the Cochran-Armitage test for trend, analysis of variance (ANOVA), and logistic (ordinal) models with an appropriate contrast. The relationship of the HRQoL (15D) at different cognition levels as the function of the standardized energy intake per day was evaluated using the analysis of variance with an appropriate contrast. A bootstrap method was used when the theoretical distribution of the test statistics was unknown or in case of violation of assumptions (e.g., non-normality). Partial correlations were calculated using 15D dimensions and standardized energy intake adjusted for age, Charlson score, and BMI. The normality of variables was evaluated using the Shapiro–Wilk W test. The Stata 15.1 (StataCorp LP; College Station, Texas, TX, USA) statistical package was used for the analysis.

3. Results

The study population consisted of 538 older long-term care residents of whom 78% were female. Table 1 shows the characteristics of the study population according to CDR. Of the residents, 28% had very mild or mild dementia (CDR 0.5–1), 38% had moderate dementia (CDR 2), and 34% had severe dementia (CDR 3) (Table 1).

Table 1. Characteristics of the study population according to Clinical Dementia Rating [28].

Population Characteristics	CDR 0.5–1 N = 150	CDR 2 N = 206	CDR 3 N = 182	p-Value ¹
<i>Background characteristics</i>				
Education (≤ 8 years), n (%)	70 (50)	87 (49)	82 (53)	0.61
Age (years), mean (SD ²)	85 (7)	84 (8)	83 (7)	0.002
Female (%)	116 (77)	166 (81)	137 (75)	0.60
Dependence in ADLs ³ , n (%)	132 (88)	193 (94)	178 (99)	<0.001
Charlson ⁴ , mean (SD)	2.3 (1.2)	2.1 (1.3)	1.9 (1.1)	<0.001
MMSE ⁵ , mean (SD)	14.8 (5.5)	11.6 (5.7)	7.6 (6.2)	<0.001
<i>Nutrition status</i>				
Mean weight, kg (SD)	69 (15)	68 (15)	63 (11)	<0.001
BMI ⁶ , mean (SD)	26.1 (4.8)	26.0 (5.3)	23.8 (4.2)	<0.001
MNA ⁷ , n (%)				<0.001
Malnourished (<17 points)	12 (9)	30 (17)	46 (28)	
At risk of malnutrition (17–23.5 points)	91 (68)	124 (69)	109 (67)	
Normal nutritional status (>23.5 points)	30 (23)	26 (14)	8 (5)	
<i>Nutrition intake</i>				
Energy total kcal, mean (SD)	1684 (434)	1674 (419)	1648 (383)	0.41
Energy total kcal, mean (SD)				
Women	1611 (432)	1629 (415)	1599 (384)	0.79
Men	1936 (344)	1860 (391)	1797 (341)	0.093
Protein g/kg BW/d, mean (SD)	0.91 (0.32)	0.89 (0.30)	0.91 (0.31)	0.97

Table 1. Cont.

Population Characteristics	CDR 0.5–1 N = 150	CDR 2 N = 206	CDR 3 N = 182	p-Value ¹
<i>Nutrition care</i>				
Eats snacks between meals, n (%)	116 (79)	163 (80)	142 (81)	0.70
Receives oral nutritional supplements, n (%)	24 (16)	45 (22)	54 (30)	0.003
Eats normal food, n (%)	119 (80)	125 (61)	68 (38)	<0.001
Eats less than half of the food portion, n (%)	35 (23)	44 (22)	26 (15)	0.046
<i>Symptoms related to eating and digestion</i>				
Chewing problems, n (%)	23 (16)	56 (30)	78 (45)	<0.001
Constipation, n (%)	44 (32)	54 (29)	45 (28)	0.45
<i>HRQoL</i>				
15D index ⁸ , mean (SD)	0.65 (0.11)	0.60 (0.10)	0.52 (0.10)	<0.001

¹ p-value for linearity was evaluated by using the Cochran-Armitage test for trend and analysis of variance with an appropriate contrast; ² SD = Standard deviation; ³ ADL = Activities of daily living measured by Clinical Dementia Rating (CDR) scale “personal care” score ≥ 2 [28]; ⁴ Charlson comorbidity index [27]; ⁵ MMSE = Mini-Mental State Examination [29]; ⁶ BMI = body mass index (kg/m²); ⁷ MNA = Mini Nutritional Assessment [30]; ⁸ [32]. Italic blackening in the table makes the table clearer.

The residents with severe dementia were younger, more often exhibited dependence in activities of daily living (ADL) functioning, and had lower Charlson comorbidity index and lower MMSE scores than residents with very mild/mild or moderate dementia.

The residents with severe dementia more often had a lower BMI and malnutrition than those with moderate to mild dementia. There were no significant differences in energy intake or protein intake (g/kg) between the CDR groups. Approximately 80% of the residents received snacks between meals in all CDR groups. In addition, residents with severe dementia more often received nutritional supplements and liquid/pureed food. There was no difference between groups in the proportions of those eating snacks. However, residents with severe dementia ate on average less during their meals and more often had chewing problems.

The mean (SD) 15D index was 0.65 (0.11) among residents with very mild/mild dementia, 0.60 (0.10) among residents with moderate dementia, and 0.52 (0.10) among residents with severe dementia. The difference was significant ($p < 0.001$).

Figure 1 panel A presents energy intake per day in males and females, and panel B shows the standardized energy intake per day (z-scores). Figure 2 shows the correlation between standardized energy quartiles and HRQoL in different stages of dementia. Although both stage of dementia ($p < 0.001$) and energy intake ($p = 0.013$) were significantly associated with HRQoL (Figure 2), the interaction was not significant ($p = 0.30$).

Figure 3 presents partial correlation analysis between standardized energy intake per day and dimensions of 15D according to CDR groups. Among residents with very mild/mild dementia, none of the dimensions of 15D were associated with standardized energy intake. In contrast, the total 15D index was significantly associated with higher standardized energy intake (Figure 3). In residents with moderate dementia, the dimensions of mobility and usual activities and the 15D index were significantly associated with standardized energy intake. Among residents with severe dementia, no significant associations emerged between the different dimensions or the 15D index and standardized energy intake.




Figure 1. Distribution with normal curve overlay of energy intake/day in women and men (A) and standardized (z-score) energy intake/day (B). Energy intakes were standardized by z-scores to include both men and women in the same analyses.




Figure 2. Relationship between the standardized energy intake per day at various stages of dementia according to the Clinical Dementia Rating (CDR) [28] with health-related quality of life (15D) [32].




Figure 3. Partial correlations of standardized energy intake per day with health-related quality of life according to 15D [32] at various stages of dementia according to the Clinical Dementia Rating (CDR) [28]. The correlations were adjusted for age, Charlson Comorbidity Index [27], and body mass index. * $p < 0.05$, ** $p < 0.01$.

4. Discussion

We observed that the stage of dementia according to CDR and standardized energy intake were both associated with HRQoL. The interaction between the stage of dementia and standardized energy intake was not significant. The partial correlation between the 15D index and standardized energy intake was significant in very mild to moderate dementia but not in severe dementia.

The strength of this study is the large sample size concerning food records. The assessment tools are well-validated and widely used internationally. The food records were collected by trained nurses. Food records with direct observation have been considered more appropriate than food frequency questionnaires in institutional settings [33]. We included consecutive long-term care residents with mild to severe stages of dementia. We assessed the resident's HRQoL with the 15D tool that can be completed by proxy and may thus also be used among residents with severe dementia. One limitation is that this study is cross sectional and causality cannot be determined between cognitive impairment and HRQoL or standardized energy intake and HRQoL. In addition, we did not explore the contents and servings of the foods, which may be even more important than energy and protein intake with regards to HRQoL.

In our study, residents with severe dementia had poorer HRQoL than residents with moderate or very mild/mild dementia. Our findings are consistent with some previous studies [34–36] but not with all [37]. In addition, few studies have included people with severe dementia. For example, Abrahamson et al. excluded nursing home residents with the most severe dementia [38]. This large American study suggested that residents with more impaired cognitive impairment reported lower QoL in several domains, including activities, individuality, privacy, and meaningful relationships. Comparisons across studies may not be possible due to the various instruments used to assess HRQoL and stage of dementia, and the differing study populations (community samples or from various long-term care settings). However, it seems that HRQoL, at least its functional dimensions, is associated with the stage of dementia.

In our study, standardized energy intake was associated with better HRQoL. This finding is consistent with previous studies among older home-care patients and non-institutionalized older people [11,24]. In addition to energy intake, malnutrition stage is associated with poor HRQoL in institutionalized and community-living older people [22,25,26,39]. These results suggest that adequate energy intake and good nutritional status may lead to better HRQoL in institutionalized older residents.

We did not observe a significant interaction between HRQoL, standardized energy intake, and dementia stage. At very mild/mild and moderate stages of dementia, a higher standardized energy intake was associated with higher HRQoL. However, at the severe stage there was no such relationship. To our knowledge, there are no previous studies that have explored how energy intake modifies the association between the stage of dementia and HRQoL. According to our findings, adequate energy intake has the potential to modify the association between HRQoL and the stage of dementia, at least in mild-moderate stages.

There was a significant partial correlation between standardized energy intake and HRQoL of residents with very mild/mild or moderate dementia, but not with residents with severe dementia. In moderate dementia, the 15D dimensions of mobility and usual activities correlated significantly with higher energy intake. Both mobility and usual activities are part of ADL functions. Thus, energy intake may modify HRQoL through ADL functions. In previous studies among long-term care residents, increased energy intake has maintained ADL functions [40,41]. ADL functions are an important part of HRQoL and may explain the association between standardized energy intake and HRQoL. ADL functions tend to decrease as dementia progresses [42]. There was no correlation between standardized energy intake and HRQoL among residents with severe dementia. Among residents with severe dementia, the ability to perform ADL functions is already greatly compromised regardless of energy intake. This may explain why higher energy intake does not modify HRQoL among these residents.

We did not observe differences between the CDR groups in energy intake, although there was a larger proportion of malnutrition among those with severe dementia than among those with mild-moderate dementia. In addition, there was no difference in eating snacks between the various stages of dementia. However, residents with severe dementia were assessed to eat on average less of their food portion than those at the very mild-moderate stages. The nurses more often administered oral nutritional supplements to those with severe dementia than to those at the very mild-moderate stages. Furthermore, residents at the severe stage of dementia more often received modified (liquid/pureed food) food to maintain their energy intake than those at the very mild-moderate stages. Thus, several means to maintain energy intake were used in these long-term care settings.

5. Conclusions

This study provides evidence that both standardized energy intake and stage of dementia are significantly associated with HRQoL in institutionalized older residents. Our study emphasizes that adequate energy intake may maintain HRQoL especially among residents with mild to moderate dementia, among whom food seems to modify HRQoL through mobility and ADL functions.

Author Contributions: Conceptualization, S.K.S., S.M.H., K.H., R.H.M., and P.K.H.; methodology, S.K.S., K.H., and P.K.H.; formal analysis, S.K.S., K.H., and P.K.H.; investigation, S.K.S. and R.H.M.; data curation, S.M.H. and P.K.H.; writing (original draft preparation), S.K.S. and P.K.H.; writing (review and editing), S.K.S., S.M.H., K.H., R.H.M., and P.K.H.; visualization, S.K.S., K.H., and P.K.H.; supervision, S.M.H. and P.K.H.; project administration, S.M.H. and P.K.H.; funding acquisition, S.M.H. and P.K.H.

Funding: This study was supported by the Päivikki and Sakari Sohlberg Foundation and the Helsinki University Hospital. The sponsor did not have any role in study design, analysis or interpretation of data, writing the report, or decision to submit for publication. The authors are independent researchers not associated with the funding agencies.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Cereda, E.; Pedrolli, C.; Klersy, C.; Bonardi, C.; Quarleri, L.; Cappello, S.; Turri, A.; Rondanelli, M.; Caccialanza, R. Nutritional status in older persons according to healthcare setting: A systematic review and meta-analysis of prevalence data using MNA((R)). *Clin. Nutr.* **2016**, *35*, 1282–1290. [[CrossRef](#)] [[PubMed](#)]
2. Jyvakorpi, S.K.; Pitkala, K.H.; Puranen, T.M.; Bjorkman, M.P.; Kautiainen, H.; Strandberg, T.E.; Soini, H.H.; Suominen, M.H. High proportions of older people with normal nutritional status have poor protein intake and low diet quality. *Arch. Gerontol. Geriatr.* **2016**, *67*, 40–45. [[CrossRef](#)] [[PubMed](#)]
3. Recommendations, N.N. *Integrating Nutrition and Physical Activity 2012*; Nordic Council of Ministers: Copenhagen, Denmark, 2014.
4. Engelheart, S.; Akner, G. Dietary intake of energy, nutrients and water in elderly people living at home or in nursing home. *J. Nutr. Health Aging* **2015**, *19*, 265–272. [[CrossRef](#)] [[PubMed](#)]
5. Rodriguez-Rejon, A.I.; Ruiz-Lopez, M.D.; Artacho, R. Dietary Intake and Associated Factors in Long-Term Care Homes in Southeast Spain. *Nutrients* **2019**, *11*, 266. [[CrossRef](#)] [[PubMed](#)]
6. Van Zwiene-Pot, J.I.; Visser, M.; Kruizenga, H.M. Predictors for achieving adequate protein and energy intake in nursing home rehabilitation patients. *Aging Clin. Exp. Res.* **2018**, *30*, 799–809. [[CrossRef](#)] [[PubMed](#)]
7. Jyvakorpi, S.K.; Pitkala, K.H.; Puranen, T.M.; Bjorkman, M.P.; Kautiainen, H.; Strandberg, T.E.; Soini, H.; Suominen, M.H. Low protein and micronutrient intakes in heterogeneous older population samples. *Arch. Gerontol. Geriatr.* **2015**, *61*, 464–471. [[CrossRef](#)]
8. Batchelor-Murphy, M.; Kennerly, S.M.; Horn, S.D.; Barrett, R.; Bergstrom, N.; Boss, L.; Yap, T.L. Impact of Cognition and Handfeeding Assistance on Nutritional Intake for Nursing Home Residents. *J. Nutr. Gerontol. Geriatr.* **2019**, *38*, 1–15. [[CrossRef](#)]
9. Ter Borg, S.; de Groot, L.C.; Mijnders, D.M.; de Vries, J.H.; Verlaan, S.; Meijboom, S.; Luiking, Y.C.; Schols, J.M. Differences in Nutrient Intake and Biochemical Nutrient Status Between Sarcopenic and Nonsarcopenic Older Adults—Results From the Maastricht Sarcopenia Study. *J. Am. Med. Dir. Assoc.* **2016**, *17*, 393–401. [[CrossRef](#)]

10. Schoufour, J.D.; Franco, O.H.; Kieffe-de Jong, J.C.; Trajanoska, K.; Stricker, B.; Brusselle, G.; Rivadeneira, F.; Lahousse, L.; Voortman, T. The association between dietary protein intake, energy intake and physical frailty: Results from the Rotterdam Study. *Br. J. Nutr.* **2019**, *121*, 393–401. [[CrossRef](#)]
11. Jimenez-Redondo, S.; Beltran de Miguel, B.; Gavidia Banegas, J.; Guzman Mercedes, L.; Gomez-Pavon, J.; Cuadrado Vives, C. Influence of nutritional status on health-related quality of life of non-institutionalized older people. *J. Nutr. Health Aging* **2014**, *18*, 359–364. [[CrossRef](#)]
12. Perales, J.; Cosco, T.D.; Stephan, B.C.M.; Haro, J.M.; Brayne, C. Health-related quality-of-life instruments for Alzheimer’s disease and mixed dementia. *Int. Psychogeriatr.* **2013**, *25*, 691–706. [[CrossRef](#)] [[PubMed](#)]
13. Naylor, M.D.; Hirschman, K.B.; Hanlon, A.L.; Abbott, K.M.; Bowles, K.H.; Foust, J.; Shah, S.; Zubritsky, C. Factors Associated With Changes in Perceived Quality of Life Among Elderly Recipients of Long-Term Services and Supports. *J. Am. Med Dir. Assoc.* **2016**, *17*, 44–52. [[CrossRef](#)] [[PubMed](#)]
14. Lyketsos, C.G.; Gonzales-Salvador, T.; Chin, J.J.; Baker, A.; Black, B.; Rabins, P. A follow-up study of change in quality of life among persons with dementia residing in a long-term care facility. *Int. J. Geriatr. Psychiatry* **2003**, *18*, 275–281. [[CrossRef](#)] [[PubMed](#)]
15. Hoe, J.; Hancock, G.; Livingston, G.; Woods, B.; Challis, D.; Orrell, M. Changes in the quality of life of people with dementia living in care homes. *Alzheimer Dis. Assoc. Disord.* **2009**, *23*, 285–290. [[CrossRef](#)] [[PubMed](#)]
16. Funaki, Y.; Kaneko, F.; Okamura, H. Study on factors associated with changes in quality of life of demented elderly persons in group homes. *Scand. J. Occup. Ther.* **2005**, *12*, 4–9. [[CrossRef](#)] [[PubMed](#)]
17. Oudman, E.; Veurink, B. Quality of life in nursing home residents with advanced dementia: A 2-year follow-up. *Psychogeriatr. Off. J. Jpn. Psychogeriatr. Soc.* **2014**, *14*, 235–240. [[CrossRef](#)] [[PubMed](#)]
18. Borowiak, E.; Kostka, T. Predictors of quality of life in older people living at home and in institutions. *Aging Clin. Exp. Res.* **2004**, *16*, 212–220. [[CrossRef](#)] [[PubMed](#)]
19. Jimenez-Redondo, S.; Beltran de Miguel, B.; Gomez-Pavon, J.; Cuadrado Vives, C. Non-institutionalized nonagenarians health-related quality of life and nutritional status: Is there a link between them? *Nutr. Hosp.* **2014**, *30*, 602–608. [[PubMed](#)]
20. Hernandez-Galiot, A.; Goni, I. Quality of life and risk of malnutrition in a home-dwelling population over 75 years old. *Nutrition* **2017**, *35*, 81–86. [[CrossRef](#)] [[PubMed](#)]
21. Rasheed, S.; Woods, R.T. Malnutrition and quality of life in older people: A systematic review and meta-analysis. *Ageing Res. Rev.* **2013**, *12*, 561–566. [[CrossRef](#)] [[PubMed](#)]
22. Kvamme, J.M.; Olsen, J.A.; Florholmen, J.; Jacobsen, B.K. Risk of malnutrition and health-related quality of life in community-living elderly men and women: The Tromso study. *Qual. Life Res.* **2011**, *20*, 575–582. [[CrossRef](#)] [[PubMed](#)]
23. Wei, K.; Nyunt, M.S.Z.; Gao, Q.; Wee, S.L.; Ng, T.P. Long-term changes in nutritional status are associated with functional and mortality outcomes among community-living older adults. *Nutrition* **2019**, *66*, 180–186. [[CrossRef](#)] [[PubMed](#)]
24. Adıgüzel, E.; Acar-Tek, N. Nutrition-related parameters predict the health-related quality of life in home care patients. *Exp. Gerontol.* **2019**, *120*, 15–20. [[CrossRef](#)] [[PubMed](#)]
25. Keller, H.H.; Ostbye, T.; Goy, R. Nutritional risk predicts quality of life in elderly community-living Canadians. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2004**, *59*, 68–74. [[CrossRef](#)] [[PubMed](#)]
26. Salminen, K.S.; Suominen, M.H.; Soini, H.; Kautiainen, H.; Savikko, N.; Saarela, R.K.T.; Muurinen, S.; Pitkala, K.H. Associations between Nutritional Status and Health-Related Quality of Life among Long-Term Care Residents in Helsinki. *J. Nutr. Health Aging* **2019**, *23*, 474–478. [[CrossRef](#)] [[PubMed](#)]
27. Charlson, M.E.; Pompei, P.; Ales, K.L.; MacKenzie, C.R. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *J. Chronic Dis.* **1987**, *40*, 373–383. [[CrossRef](#)]
28. Hughes, C.P.; Berg, L.; Danziger, W.L.; Coben, L.A.; Martin, R.L. A new clinical scale for the staging of dementia. *Br. J. Psychiatry* **1982**, *140*, 566–572. [[CrossRef](#)] [[PubMed](#)]
29. Folstein, M.F.; Folstein, S.E.; McHugh, P.R. “Mini-mental state”. A practical method for grading the cognitive state of patients for the clinician. *J. Psychiatr. Res.* **1975**, *12*, 189–198. [[CrossRef](#)]
30. Guigoz, Y. The Mini Nutritional Assessment (MNA) review of the literature—What does it tell us? *J. Nutr. Health Aging* **2006**, *10*, 7.
31. Vellas, B.; Guigoz, Y.; Garry, P.J.; Nourhashemi, F.; Bennahum, D.; Lauque, S.; Albaredo, J.-L. The mini nutritional assessment (MNA) and its use in grading the nutritional state of elderly patients. *Nutrition* **1999**, *15*, 116–122. [[CrossRef](#)]

32. Sintonen, H. The 15D instrument of health-related quality of life: Properties and applications. *Ann. Med.* **2001**, *33*, 328–336. [[CrossRef](#)] [[PubMed](#)]
33. Thompson, F.E.; Byers, T. Dietary Assessment Resource Manual. *J. Nutr.* **1994**, *124*, 2245s–2317s. [[PubMed](#)]
34. Almomani, F.M.; McDowd, J.M.; Bani-Issa, W.; Almomani, M. Health-related quality of life and physical, mental, and cognitive disabilities among nursing home residents in Jordan. *Qual. Life Res.* **2014**, *23*, 155–165. [[CrossRef](#)] [[PubMed](#)]
35. Pan, C.-W.; Wang, X.; Ma, Q.; Sun, H.-P.; Xu, Y.; Wang, P. Cognitive dysfunction and health-related quality of life among older Chinese. *Sci. Rep.* **2015**, *5*, 17301. [[CrossRef](#)] [[PubMed](#)]
36. Kazazi, L.; Foroughan, M.; Nejati, V.; Shati, M. Association between age associated cognitive decline and health related quality of life among Iranian older individuals. *Electron. Physician* **2018**, *10*, 6663–6671. [[CrossRef](#)] [[PubMed](#)]
37. Banerjee, S.; Samsi, K.; Petrie, C.D.; Alvir, J.; Treglia, M.; Schwam, E.M.; del Valle, M. What do we know about quality of life in dementia? A review of the emerging evidence on the predictive and explanatory value of disease specific measures of health related quality of life in people with dementia. *Int. J. Geriatr. Psychiatry* **2009**, *24*, 15–24. [[CrossRef](#)] [[PubMed](#)]
38. Abrahamson, K.; Clark, D.; Perkins, A.; Arling, G. Does cognitive impairment influence quality of life among nursing home residents? *Gerontologist* **2012**, *52*, 632–640. [[CrossRef](#)]
39. Johansson, L.; Sidenvall, B.; Malmberg, B.; Christensson, L. Who will become malnourished? A prospective study of factors associated with malnutrition in older persons living at home. *J. Nutr. Health Aging* **2009**, *13*, 855–861. [[CrossRef](#)]
40. Odlund Olin, A.; Armyr, I.; Soop, M.; Jerstrom, S.; Classon, I.; Cederholm, T.; Ljungren, G.; Ljungqvist, O. Energy-dense meals improve energy intake in elderly residents in a nursing home. *Clin. Nutr.* **2003**, *22*, 125–131. [[CrossRef](#)]
41. Nijs, K.A.; de Graaf, C.; Kok, F.J.; van Staveren, W.A. Effect of family style mealtimes on quality of life, physical performance, and body weight of nursing home residents: Cluster randomised controlled trial. *BMJ* **2006**, *332*, 1180–1184. [[CrossRef](#)]
42. Sun, M.; Mainland, B.J.; Ornstein, T.J.; Mallya, S.; Fiocco, A.J.; Sin, G.L.; Shulman, K.I.; Herrmann, N. The association between cognitive fluctuations and activities of daily living and quality of life among institutionalized patients with dementia. *Int. J. Geriatr. Psychiatry* **2018**, *33*, e285. [[CrossRef](#)] [[PubMed](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).