

<https://helda.helsinki.fi>

Erratum: Saastamoinen, M.; Särkijärvi, S.; Valtonen, E. The Effect of Diet Composition on the Digestibility and Fecal Excretion of Phosphorus in Horses: A Potential Risk of P Leaching? *Animals* 2020, 10, 140

Saastamoinen, Markku

Multidisciplinary Digital Publishing Institute

2020-02-12

Saastamoinen, M.; Särkijärvi, S.; Valtonen, E. Erratum: Saastamoinen, M.; Särkijärvi, S.; Valtonen, E. The Effect of Diet Composition on the Digestibility and Fecal Excretion of Phosphorus in Horses: A Potential Risk of P Leaching? *Animals* 2020, 10, 140. *Animals* 2020, 10, 285.

<http://hdl.handle.net/10138/348660>

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Erratum

Erratum: Saastamoinen, M.; Särkijärvi, S.; Valtonen, E. The Effect of Diet Composition on the Digestibility and Fecal Excretion of Phosphorus in Horses: A Potential Risk of P Leaching? *Animals* 2020, 10, 140

Markku Saastamoinen ^{1,*}, Susanna Särkijärvi ¹ and Elisa Valtonen ²

¹ Natural Resources Institute Finland, FI-00790 Helsinki, Finland; susanna.sarkijarvi@luke.fi

² Department of Animal Science, University of Helsinki, FI-00790 Helsinki, Finland;
elisa.mj.valtonen@gmail.com

* Correspondence: markku.saastamoinen@luke.fi

Published: 12 February 2020



The authors wish to make the following corrections to their paper [1]:

In Table 3, the rows showing “Intake P”, “Excretion P” and “Digestibility P” have typing errors and are now corrected (see the corrected version of Table 3 below).

Table 3. Daily intake (g), fecal excretion (g), digestibility (%) and retention (g) of phosphorus (P).

| Diet/Forage | A | B | C | D | E | F | Pooled SEM | Statistical Significance (<i>p</i> -Values) | | | | |
|-----------------|------|---------|------|------|------|------|------------|--|--------------|---------------|--------|-------------|
| | Hay | Haylage | Hay | Hay | Hay | Hay | | Haylage vs. Others | Hay vs. ConS | Oats vs. Comp | ConL | ConT × ConL |
| ConL | 0 | 0 | O20 | O35 | C20 | C35 | | | | | | |
| Intake P | 20.6 | 22.1 | 22.8 | 24.8 | 20.9 | 24.1 | 0.21 | 0.036 | <0.001 | <0.001 | <0.001 | 0.027 |
| Excretion P | 20.0 | 20.2 | 21.5 | 22.1 | 19.9 | 21.5 | 0.42 | 0.125 | 0.021 | 0.025 | 0.033 | 0.251 |
| Digestibility P | 2.7 | 8.0 | 5.6 | 11.1 | 4.9 | 10.6 | 1.98 | 0.652 | 0.037 | 0.761 | 0.024 | 0.974 |
| Retention | 0.6 | 1.9 | 1.0 | 2.8 | 0.9 | 1.9 | 0.45 | 0.354 | 0.075 | 0.25 | 0.014 | 0.379 |

O = oats; C = complete feed; ConL = concentrate level (20 or 35% of oats O or complete feed C); ConS = concentrate supplementation; ConT = concentrate type (oats/complete feed); Comp = complete feed.

Conflicts of Interest: The authors declare no conflict of interest.

Reference

1. Saastamoinen, M.; Särkijärvi, S.; Valtonen, E. The Effect of Diet Composition on the Digestibility and Fecal Excretion of Phosphorus in Horses: A Potential Risk of P Leaching? *Animals* **2020**, *10*, 140. [[CrossRef](#)] [[PubMed](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).