

## Daftar Jurnal Ilmiah Hasil Akreditasi DIKTI

NO.	JUDUL JURNAL	ISSN	PENERBIT	ALAMAT	BIDANG KEILMUAN	PERINGKAT	NOMOR SK	MASA BERLAKU
1	AL-BAYAN	1411-5743	Fakultas Dakwah Institut Agama Islam Negeri Ar-Raniry	Kampus IAIN Darussalam. Banda Aceh 23111. Telp.: (0651) 52548	Agama	C	39/Dikti/Kep/2004	10-14 s.d. 11-07
2	KHAZANAH	1412-372X	Program Pascasarjana Universitas Islam Negeri Sunan Gunung Djati	Jln. A.H. Nasution No.105, Cibiru, Ujung Berung. Bandung 40614. Telp./Fax.: (022) 7800249	Agama	B	39/Dikti/Kep/2004	10-14 s.d. 11-07
3	ALHADHARAH	1412-9515	Fakultas Dakwah Institut Agama Islam Negeri Antasari	Jln. A. Yani Km.4,5. Banjarmasin 70235. Telp.: (0511) 250771. E-mail: <a href="mailto:alhadharah@mailcity.com">alhadharah@mailcity.com</a>	Agama	C	56/Dikti/Kep/2005	12-05 s.d. 05-08
4	TAJDID	0854-9850	Institut Agama Islam Darussalam	Jln. K.H. Ahmad Fadlil, Kampus IAD Pondok Pesantren Darussalam. P.O. Box 2. Ciamis 46271	Agama	B	55a/Dikti/Kep/2006	10-06 s.d. 10-09
5	JOURNAL ANALYTICA ISLAMICA	1411-4380	Program Pascasarjana Institut Agama Islam Negeri Sumatera Utara	Jln. Pembangunan Komplek Pondok Surya, Helvetia Timur. Medan 29124. Telp.: (061) 8465290. Fax.: (061) 615683. E-mail: <a href="mailto:ppsiain@indosat.net.id">ppsiain@indosat.net.id</a>	Agama	B	56/Dikti/Kep/2005	12-05 s.d. 05-08
6	NURANI	1693-8437	Fakultas Syari'ah Institut Agama Islam Negeri Raden Fatah	Jln. Prof. K.H. Zainal Abidin Fikri Km.3,5. Palembang 30126. Telp.: (0711) 362427. Fax.: (0711) 356209. E-mail: <a href="mailto:ikhi2003@yahoo.com">ikhi2003@yahoo.com</a>	Agama	C	56/Dikti/Kep/2005	12-05 s.d. 05-08
7	TA'DIB	1410-6973	Fakultas Tarbiyah Institut Agama Islam Negeri Raden Fatah	Jln. Jend. Sudirman Km.3,5. Palembang 30126. Telp.: (0711) 362502, 354668	Agama	C	56/Dikti/Kep/2005	12-05 s.d. 05-08
8	JURNAL THEOLOGIA	0853-3857	Fakultas Ushuluddin Institut Agama Islam Negeri Walisongo Semarang	Jln. Prof. Dr. Hamka Km.2, Ngaliyan. Semarang 50185	Agama	B	48/Dikti/Kep/2006	06-06 s.d. 06-09
9	AL QALAM	1410-3222	Sekolah Tinggi Agama Islam Negeri Sultan Maulana Hasanuddin Banten	Jln. Jend. Sudirman No.30. Serang 42118. Telp.: (0254) 200323. Fax.: (0254) 200022	Agama	B	108/Dikti/Kep/2007	08-07 s.d. 08-10
			Pusat Penelitian dan					

135	HAYATI (ENGLISH EDITION)	1978-3019	Perhimpunan Biologi Indonesia joint with Departemen Biologi, Fakultas Matematika dan Ilmu Pengetahuan Alam Institut Pertanian Bogor	Kampus IPB Darmaga. Bogor 16680. Telp./Fax.: (0251) 622833. E-mail: <a href="mailto:hayati@ipb.ac.id">hayati@ipb.ac.id</a> . Website: <a href="http://www.ipb.ac.id/gallery/journal/hayati">http://www.ipb.ac.id/gallery/journal/hayati</a>	MIPA	B	55a/Dikti/Kep/2006	10-06 s.d. 10-09
136	MAKARA SERI SAINS	1693-6671	Direktorat Riset dan Pengabdian Masyarakat Universitas Indonesia	Gd.A Lt.2, Kampus UI Pondok Cina. Depok 16424. Telp.: (021) 7270020, 7270021, 7270022, 7270023. Fax.: (021) 7270017, 7863460, 7863447. E-mail: <a href="mailto:pusadmui@makara.cso.ui.ac.id">pusadmui@makara.cso.ui.ac.id</a> . Website: <a href="http://www.http://www.research-ui.org">http://www.http://www.research-ui.org</a>	MIPA	B	39/Dikti/Kep/2004	10-14 s.d. 11-07
137	JURNAL BAHAN ALAM INDONESIA	1412-2855	Perhimpunan Peneliti Bahan Obat Alami	Kantor Sekretariat: KPPOT Puslitbang Farmasi Balitangkes Depkes R.I., Jln. Percetakan Negara No.29. Jakarta 10560. Telp.: (021) 42871807. Fax.: (021) 4243933, 4216059. E-mail: <a href="mailto:kppot@indo.net.id">kppot@indo.net.id</a>	MIPA	B	48/Dikti/Kep/2006	06-06 s.d. 06-09
138	JURNAL MIKROBIOLOGI INDONESIA	0853-358X	Perhimpunan Mikrobiologi Indonesia	Jln. Jend. Sudirman No.51. Jakarta Pusat 12930. Telp.: (021) 5703306 Pes.449. Fax.: (021) 5719060. E-mail: <a href="mailto:microbiol.indonesia@yahoo.co.id">microbiol.indonesia@yahoo.co.id</a> . Wbsite: <a href="http://www.permi.or.id">http://www.permi.or.id</a>	MIPA	A	39/Dikti/Kep/2004	10-14 s.d. 11-07
139	MATEMATIKA DAN KOMPUTER	0216-4728	Sekolah Tinggi Informatika dan Komputer Gunadarma	Jln. Salemba Raya No.31	MIPA	C	39/Dikti/Kep/2004	10-14 s.d. 11-07
140	MIPA	0854-8269	Fakultas Keguruan Ilmu Eksakta Institut Keguruan dan Ilmu Pendidikan Malang	Jln. Surabaya No.6	MIPA	C	39/Dikti/Kep/2004	10-14 s.d. 11-07
141	BERKALA PENELITIAN HAYATI	0852-6834	Perhimpunan Biologi Indonesia Cabang Surabaya	Kantor Sekretariat: Jur. Biologi FMIPA-Unair, Kampus C Unair, Jln. Mulyorejo. Surabaya 60115. Telp.: (031) 5936501. Fax.: (031) 5936502	MIPA	B	56/Dikti/Kep/2005	12-05 s.d. 05-08
142	ALCHEMY	1412-4092	Jurusan Kimia, Fakultas Matematika dan Ilmu Pengetahuan Alam Universitas Sebelas Maret	Jln. Ir. Sutami No.36-A., Ketingan. Surakarta 57126. Telp.: (0271) 663375, 646994 Pes.387. Fax.: (0271) 663375	MIPA	C	56/Dikti/Kep/2005	12-05 s.d. 05-08
			Fakultas Geografi					

*Juntas*

# HAYATI

## Journal of Biosciences

Vol. 14 No. 3 September 2007

Accredited by Indonesian Directorate General of Higher Education No. 55a/DIKTI/Kep/2006

### MANAGING EDITOR

Bambang Suryobroto (~2008)

### CHIEF EDITOR

Rika Raffiudin (~2008)

### EDITORIAL BOARD

Alex Hartana (~2008), Antonius Suwanto (~2012), Aris Tri Wahyudi (~2007), Darjono (~2012),  
Dedy Duryadi Solihin (~2010), Ence Darmo Jaya Supena (~2012), Erin P. Riley (~2008),  
Ikin Mansjoer (~2008), Iman Rusmana (2011), Koji Nakamura (~2012),  
Lisdar Idwan Sudirman (~2008), Miftahudin (~2012), Sri Sudarmiyati Tjitrosoedirdjo (~2009),  
Tedja Imas (~2007), Tri Heru Widarto (~2010), Tukirin Partomihardjo (~2008),  
and Utut Widyastuti (~2007)

### TECHNICAL EDITORS

Berry Juliandi (2012), Dyah Perwitasari (~2007), Juliarni (~2009),  
Kanthi Arum Widayati (~2007), and Nina Ratna Djuita (~2007)

### BUSINESS MANAGER

Utut Widyastuti (~2007)

### PUBLISHER

Indonesian Society for Biology and  
Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University

### EDITORIAL ADDRESS

HAYATI Journal of Biosciences  
Department of Biology, Faculty of Mathematics and Natural Sciences  
Bogor Agricultural University, Darmaga Campus, Bogor 16680  
Phone/Fax: +62-251-421258; E-mail: hayati\_j\_biosci@cbn.net.id  
<http://www.ipb.ac.id/gallery/journal/hayati>

HAYATI Journal of Biosciences publishes articles, short communications, and reviews in bioscientific fields such as development, biotechnology, biodiversity and environmental issue. The articles cover wide range of organisms including virus, microbes, fungi, plants, animals, and human. Published four times a year: March, June, September, and December.

### SUBSCRIPTION RATES - not including shipping and handling

Subscriber	One year	
	Overseas (USD)	Indonesian (Rp)
Personal	10	75.000,00
Institution/library	20	150.000,00

## SHORT COMMUNICATION

# Isolation and Characterization of Silaffin that Catalyze Biosilica Formation from Marine Diatom *Chaetoceros gracilis*

AGNES IMELDA MANURUNG<sup>1</sup>, ALBERTA RIKA PRATIWI<sup>2</sup>, DAHRUL SYAH<sup>1</sup>,  
MAGGY THENAWIDJAJA SUHARTONO<sup>1\*</sup>

<sup>1</sup>Department of Food Science and Technology, Bogor Agricultural University, Darmaga Campus, Bogor 16680, Indonesia

<sup>2</sup>Department of Food Technology, Soegijapranata Chatholic University, Jalan Pawiyatan Luhur IV/1,  
Bendan Duwur, Semarang 50234, Indonesia

Received September 8, 2006/Accepted August 1, 2007

The method of making silica in industries requires extreme conditions. The finding of proteins involved in the formation of biosilica from diatoms, has opened up an alternative way of production. *Chaetoceros gracilis* is one of the diatoms, which is potential in producing silaffin protein. This study aimed to isolate and to characterize the protein. We also analyzed the protein activity toward *tetraethoxyorthosilicate* (TEOS) substrate in *in vitro* reaction. Diatom biomass was harvested and further kept in 2% SDS/100 mM EDTA solution. Protein isolation was conducted by dissolving the silica and separating the protein by soaking in 2 M HF/8 M NH<sub>4</sub>F. Protein concentration was analyzed using Bradford method and the molecular weight was estimated through SDS-PAGE. Protein activity was observed by reacting it with TEOS substrate to form silica polymer and measured by colorimetric molybdate assay. Protein concentration was 1.20 mg/ml and appeared filamentous. The apparent molecular weights consisted of 12, 23, 42, 44 kDa. These protein was able to polymerize the silica at room temperature within 10 min. As much as 85.65 μmol TEOS was polymerized per 1.4 x 10<sup>6</sup> silaffin protein per min. SEM analysis showed the formation of spherical, aggregate biosilica.

Key words: *Chaetoceros gracilis*, silaffin protein, biosilica, polymerization

The application of silica-based material is widely distributed. Silica and silicates are extensively used in cosmetic, paint, food industry, catalyst, semiconductor, and biosensor for various analysis (Perry 2003). Chemical synthesis of silica in industries requires extreme temperatures, pressure, pH, and dangerous chemical compound. In contrast, biological synthesis of silica in nature proceeds (such as occurring in Diatom) at ambient temperatures, pressure, and neutral pH. Diatoms are one of organism that produces nanostructure silica as the component of the cell wall. This organism is unicellular photosynthetic eukaryotes within the class *Bacillariophyceae* whose peculiarity amongst other microalgae is the siliceous cell wall.

Diatom silica nanostructure is precisely controlled by the cell, which involves protein as the biocatalyst. The nature of this organic molecule was clarified through the characterization of diatom biosilica-associated peptides (silaffin) which accelerate silica formation from a silicic acid solution *in vitro* (Kröger *et al.* 1999). Finding of silaffin protein that catalyzes the silica condensation is important for the design of technological process for silica production under mild and friendly environment and economically favorable.

Indonesia is a maritim country with high diversity in its marine diatoms. Diatom *Chaetoceros gracilis* is species that

can be a potential source of silaffin like-proteins, which probably shows unique characteristic in their capability of catalyzing biosilification. This research was aimed to isolate and to characterize the silaffin protein as well as to analyze its activity of toward TEOS substrate.

**Culture Condition.** The axenic culture of *C. gracilis* diatom was provided by Mariculture laboratory of Research Centre for Oceanography- Indonesian Institute of Science (LIPI). The culture was grown in modified f/2 medium. The f/2 medium was containing mayor nutrient (0.99 mM NaNO<sub>3</sub>, 0.07 mM NaH<sub>2</sub>PO<sub>4</sub>·2H<sub>2</sub>O, 5.28 μM Na<sub>2</sub>SiO<sub>3</sub>·9H<sub>2</sub>O), minor nutrient (5.36 μM FeCl<sub>3</sub>·6H<sub>2</sub>O and 26.86 μM Na<sub>2</sub>EDTA), vitamins (0.59 μM vitamin B1, 0.001 μM vitamin B12, 0.004 μM biotin) also trace metal (0.781 μM CuSO<sub>4</sub>·5H<sub>2</sub>O, 2.12 μM ZnSO<sub>4</sub>·7H<sub>2</sub>O, 0.521 μM NaMoO<sub>4</sub>·2H<sub>2</sub>O, 0.005 μM (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>·4H<sub>2</sub>O, 18.19 μM MnCl<sub>2</sub>·4H<sub>2</sub>O, 0.61 μM CoCl<sub>2</sub>·6H<sub>2</sub>O). The medium was adjusted to pH 8. The culture was continuous bubbled with sterile air and maintained at 25 °C as well as under constant light at 4000 lux.

**Biomass Collection and Siliceous Cell Wall (*Frustule*) Extraction.** Cell biomasses were harvested by centrifugation at 10,000 x g, 4 °C for 15 min. These biomasses were stored frozen at -20 °C until use.

Extraction of *frustule* was performed as described by Kröger *et al.* (2002). The cell biomasses were boiled twice in 2% SDS/ 100 mM EDTA to remove intracellular component and

\*Corresponding author. Phone: +62-251-330559,  
Fax: +62-251-621724, E-mail: mthenawidjaja@yahoo.com



membrane. *Frustule* were pelleted by centrifugation at 2,800 xg, extracted with acetone twice, washed extensively with distilled water, and dried at room temperature until further analysis.

**Isolation and Characteristic of Silaffin Protein.** Isolation of *C. gracilis* silaffin protein was performed as described by Shimizu *et al.* (1998). Silaffin protein is tightly associated with diatom biosilica and can only be solubilized by complete removal of the silica. The *frustule* was dissolved in 50 ml 2 M HF (hydrogen fluoride)/8 M  $\text{NH}_4\text{F}$  (ammonium fluoride) (pH 5). After incubation for 30 min at 0 °C, HF/ $\text{NH}_4\text{F}$  was evaporated. This extract was dialyzed (2 kDa-cut off) once against distilled water at 4 °C for 4h and repeated 9 times. The dialysate was centrifuged (10,000 xg, 20 min) at 4 °C. The precipitate protein was dissolved with distilled water and stored at 4 °C until further analysis. The morphology of protein was observed by microscopic analysis. Concentrations of protein were measured by Bradford methods (dye-binding method) with protein Bovine Serum Albumin as standard (Dunn 1989). The protein was counted by using Haemacytometer.

Estimation of protein molecular weight were conducted using SDS-PAGE as described by Laemmli (1970). Protein (10-15  $\mu\text{l}$ ) was dissolved in sample buffer solution (0.3 ml of 1 M Tris-HCl pH 6.8, 2.5 ml of 50% glycerol, 1.0 ml of 10% SDS, 0.25 ml of 2- $\beta$ -mercaptoetanol, 0.5 ml of 1% bromophenol blue and 0.45 ml aquadest). These suspension was boiled at 95 °C for 5 min and was run on 10% polyacrilamide gel for 100 Volt for 1.5 h. Low Molecular Weight protein as marker contained phosphorylase b (97.0 kDa), albumin (66.0 kDa), ovalbumin (45.0 kDa), carbonic anhydrase (30.0 kDa), trypsin inhibitor (20.1 kDa),  $\alpha$ -lactalbumin (14.4 kDa) (Amersham Bioscience, USA). Silver staining was used following SDS-PAGE.

**Silaffin Protein Activity in the *In Vitro* Reaction.** Protein solution was reacted in 50 ml of 50 mM  $\text{CH}_3\text{COONa}/\text{CH}_3\text{COOH}$  pH 5.5 and incubated at 18 °C for 30 min. Assay of protein activity was performed using monosilicic acid as a substrate. A stock of substrate solution was prepared by hydrolyzing 10 ml of 1 M TEOS in 100 ml of 1 mM HCl for 10 min at 25 °C. Immediately after incubation, aliquots of the silicic acid solution were withdrawn and added to the reaction mixtures to give a final concentration of 100 mM silicic acid. Subsequently, the mixtures of reaction were incubated with different incubation time (10, 30 min, 1, 12, and 24 h) at 18 °C by shaker incubator (100 rpm).

The reaction mixture were centrifuged and immediately processed either for quantification of silica or silica structure analysis. Silica quantification was conducted by Colorimetric molybdate assay (Strickland & Parsons 1972). The precipitate silica was washed 3 times with ethanol to remove free TEOS (that was not polymerized), and recentrifuged. Subsequently, the precipitated silica was incubated in 1 ml of 1 M NaOH at 85-95 °C for 48 h. The samples of 0.2 ml was reacted with 0.4 ml of 0.8% ammonium molybdate and 12 ml of 36% HCl solution for 10 min, mixed with 0.6 ml reduction reagent (2 ml of metol sulfite containing 1.2%  $\text{Na}_2\text{SO}_3$  and 2% p-metylamino phenol sulfate, 1.2 ml of 50%  $\text{H}_2\text{SO}_4$ , 1.2 ml of 10% saturated oxalate acid, 1.6 ml water distilled), for 3 h. The absorbance was read

at  $\lambda$  810 nm. Different silicon concentrations (TEOS in DMSO solution) between 20-240  $\mu\text{mol}/\text{ml}$  were used as standard.

The silica polymer structure was observed by Scanning Electron Microscope (SEM). Sample was centrifuged at 16,000 xg for 2 min and resuspended in 5  $\mu\text{l}$  aquadest, mounted onto a 200-mesh copper grid (diameter 3.05 mm; Agar Scientific, Stansted, Essex, U.K.), washed and air dried. Photograph with JEOL JSM-5310LV SEM was prepared before photographing with JEOL JFC-1200 Fine Coater film.

**Biomass and Frustule.** Biomass cell of 19.3 g produced frustules as much as 0.06 g (Figure 1). The silica *frustule* is composed of the silaffin protein. Protein was tied very strongly with the silica. Therefore, we needed to dissolve the silica to obtain the silaffin molecule.

**Characteristic of Silaffin Protein.** After extraction using HF/ $\text{NH}_4\text{F}$  solvent the macromolecule protein was released. The microscopic analysis of protein showed filamentous appearance filament (Figure 2). As much as  $1.4 \times 10^6$  protein filaments per ml of the solution was found. We found the concentration of silaffin protein at 1.2 mg/ml by using Bradford methods.

SDS-PAGE analysis of these proteins revealed four distinct protein bands. The molecular weight of each protein was 44.65, 42.18, 23.88, and 12.06 kDa (Figure 3).

**Activity of Silaffin Protein.** Results of the silaffin protein reaction with the TEOS substrate showed that this substrate was polymerized up to 856.58  $\mu\text{mol}/\text{ml}$  with incubation time of ten min. Therefore, the TEOS was polymerized at 8.56  $\mu\text{mol}/\text{ml}$  per min and at  $1.5 \times 10^{11}$  molecules per protein filament (Table 1).

Analysis using SEM (3500 x) showed clearly the formation of the silica polymer with round (spherical) shape. In addition, the spherical silica structures also showed aggregate formation (Figure 4).

The silica component in the diatom *frustules* (silica *frustules*) could read up to approximately 90% of the dry cell (Round *et al.* 1990). However, this concentration depends on the diatom species. Each diatom species possesses specific characteristics of each wall that was marked by the content and the type of silica structure. The process of frustules formation is not yet known clearly but is suspected to involve silica diffusion. The process of this diffusion, known as sintering, is affected by several factors for example pH and temperatures, which can be varied if the diatom is cultivated in different conditions (Parkinson & Gordon 1999).

The protein isolated from diatom *C. gracilis* in this study was found at 1.2 mg/ml. This protein is strongly attached onto the silica cell wall of the diatom. Therefore, the success in isolating the protein is determined by how far the silica could be dissolved. Frigeri *et al.* (2006) stated that EDTA plays a role in eliminating  $\text{Ca}^{2+}$  from diatom cell wall and SDS disrupt the membrane to expose the silica containing silaffin protein. Addition of HF/ $\text{NH}_4\text{F}$ , further dissolves the silica and reveals the silaffin.

Biosilica protein (silaffin protein) in this study appeared filamentous with length in range at 0.036-0.096 mm. As much as  $1.4 \times 10^6$  protein filament was obtained per ml solution. However, silicatein protein isolated from sponge *Tetya aurantia* that catalyzed silica polymerization showed needle



Figure 1. Silica frustule of Diatom *C. gracilis*.




Figure 2. SEM micrograph of silaffin protein with filamentous shape from diatom *C. gracilis*.




Figure 3. Apparent of molecular weight of protein fraction from diatom *C. gracilis* analysis by SDS-PAGE (a), LMW marker (b).

or stick shape (Shimizu *et al.* 1998; Krasko *et al.* 2000). Polymerization of silica that was formed attached to the protein. Both silaffin and silicatein protein were used as template for silica polymerization (Cha *et al.* 1999; Kröger *et al.* 2002).

Estimation of the protein molecular weight based on analysis of SDS PAGE with 10% polyacrilamide showed four bands of 44, 42, 23, and 12 kDa. Poulsen *et al.* (2003) reported that on the biosilica protein from diatom *Cylindrotheca fuciformis* found 3 peptides with molecular weight of 6.5, 10, and 40 kDa known as natSil-1A, natSil-1B, and natSil-2 respectively. Based on the weight similarity, the protein with molecular weight of 42 kDa from diatom *C. gracilis* was probably natSil-2 and the 12 kDa protein was probably similar

Table 1. Bioactivity of silaffin protein from *C. gracilis* with TEOS substrate

Incubation time (minute)	Silica attached to protein		
	μmole/ml	μmole/ml per minute	Molecule per protein filament
10	856.58	85.66	$1.5 \times 10^{11}$
30	992.10	33.07	$5.8 \times 10^{10}$
60	1094.75	18.25	$3.2 \times 10^{10}$
720	1200.00	1.67	$2.9 \times 10^9$
1440	1003.95	0.70	$1.2 \times 10^9$

Number of protein filament adds in each reaction are  $1.4 \times 10^6$  filament/ml




Figure 4. The spherical morphology of silica precipitated forming aggregate after in vitro reaction between silaffin and TEOS substrate.

to natSil-1B. In this study natSil-1A was not detected, possibly this peptide was present at very low concentration. This result is in agreement with the study of biosilica protein from diatom *Thalassiosira pseudonana*, which revealed four peptides. Three peptides have molecular weight of 35 kDa and two peptides have molecular weight of 19 kDa and both of them were isomers (Poulsen & Kröger 2004). The cell wall silica shows specific structure that is controlled by a specific protein biosilica in each diatom cell. According to Sumper and Kröger (2004), the silaffin diverse chemical structure found in various diatoms implied no homology in their protein sequence the proteins found in different kind diatom evidently have no similarity or homology in their amino acid. In nature, each kind of diatom is formed by unique silica structure, which is genetically programmed. Reaction of the protein catalyst from different diatom will be produce precipitation of silica with different structures.

The result of *in vitro* reaction between biosilica protein from *C. gracilis* and TEOS substrate in this study proved that the biosilica protein (silaffin) had capability to catalyze the silica polymerization in a very short time (10 mins) at room temperature and pH 5.5. This is clearly advantageous compare to chemical production of silica in industry with temperature of more of hundreds to thousands °C which also required long period of time (Harsono 2002). The formation of silica precipitation in short time is thus promising for production of environment-friendly silica biofabrications. Table 1 showed the highest activity of the protein, which polymerized TEOS at 856.58 μmol/ml per 1.2 mg protein filament within ten minutes or 85.658 μmol/ml per min. This result was higher than the polymerization capacity of silicatein (biosilica protein) found in sponge. Research in biosilica protein from sponge ST1

(Binuangen) and ST3 (Nias Island) reported polymerization of TEOS as much as 22 and 42  $\mu\text{mol/ml}$  per min (Nurjanah 2005). This showed that the sponge silicatein polymerized the silica slower than the diatom silaffin.

Observation of the silica polymerization by SEM showed that silaffin diatom *C. gracilis* could polymerize the silica. The polymerized silica on the silaffin was found as aggregate structure with spherical shape. This study result was in agreement with silica precipitation from diatom *T. pseudonana* produced spherical silica. However, *T. pseudonana* produced various spherical forms like silica with the diameter of 230 nm, silica porous sheet with the diameter of 20-200 nm, silica plates of densely packed and silica sphere polydisperse with the diameter of 0.9-42  $\mu\text{m}$  (Poulsen & Kröger 2004).

Cell morphology of *T. pseudonana* and *C. gracilis* were centric, however they produce different type of silica form in the *in vitro* reaction. *C. fuciformis* with pinnate morphology was reported to produce spherical silica precipitate (Kröger *et al.* 1999). Figure 4 shows the formation of silica aggregate by the protein catalyst (silaffin protein) isolated from diatom *C. gracilis*. The protein with silica polymers appeared different from the protein before the reaction (Figures 4 & 2).

According to Sumper and Kröger (2004), formation of silica from silicate acid by silaffin protein can be divided into three stages i.e. polymerization of silicate acid through the condensation of siloxane, forming dimer, trimer, and further into cyclic oligomers. Oligosilicate acids tend to polymerize silica by forming the maximal association of (Si-O-Si). In this stage, polysilicate acid was formed as the core of further formation of biosilica. In the second stage, the core of the polysilicate acid grows forming spherical particle with further polymerization. The monomer and oligomer of silicate acid were fused into these particles. During the last stage, nanosphere silica form network of three dimensions from the particle's cross chains of the associated siloxan.

Silica that was produced by protein silaffin shows diameter of 100-1000 nm and appear as round in shape (Kröger *et al.* 2002). According to Poulsen *et al.* (2003), the size of silica particle is determined by the silaffin concentrations. The increasing silaffin concentration will increase the size of the silica particle as well. However, the amount of silica that is precipitated will be decreased and this biosilica layer will be thinner. In this case, the polymerized silica will be more easily broken during SEM analysis. According to Kröger *et al.* (1999), silaffin could carry out polymerization and precipitation of silica spontaneously with diameter of the particle being ~50 nm when peptides Sil 1A, IB, and 2 are available. However, if only silaffins 1A exist, the size of the silica particle formed will be approximately 700 nm. Moreover, it is known that type of silicatein fraction such as Sil 1A, Sil 1B, and Sil 2 affected silica diameter. To understand further about the activity of

the silaffin fractions, it is necessary to conduct the polymerization reaction in the presence of various composite fractions of isolated silaffin.

## ACKNOWLEDGEMENT

This research is part of the project "Biofabrication of nanosilica structure from Indonesian marine sponge and diatom", supported from Basic Research Project-DIKTI-2005/2006 to Dahrul Syah (Department of Food Science and Technology, Bogor Agricultural University).

## REFERENCES

- Cha JN *et al.* 1999. Silicatein filaments and subunits from a marine sponge direct the polymerization of silica and silicones *in vitro*. *Proc Nat Acad Sci USA* 96:361-365.
- Dunn MJ. 1989. Determination of total protein concentration. In: Harris EL, Angal S (eds). *Protein purification methods*. Oxford IRL.
- Frigeri LG, Radabaugh TR, Haynes PA, Hildebrand M. 2006. Identification of proteins from diatom cell wall fraction of the *Thalassiosira pseudonana*: insight into silica structure formation. *Mol Cell Proteomic* 5:182-193.
- Harsono H. 2002. Pembuatan silika amorf dari limbah sekam padi. *J Ilmu Dasar* 3:98-103.
- Krasko A *et al.* 2000. Expression of silicatein and collagen genes in the marine sponge *Suberites domuncula* is controlled by silicate and myotrophin. *Eur J Biochem* 267:4878-4887.
- Kröger N, Deutzmann R, Sumper M. 1999. Polycationic peptides from biosilica that direct silica nanosphere formation. *Science* 286:1129-1132.
- Kröger N, Lorenz S, Brunner E, Sumper M. 2002. Self-assembly of highly phosphorylated silaffins and their function in biosilica morphogenesis. *Science* 298:584-586.
- Laemmli UK. 1970. Cleavage of structural proteins during the assembly of the head of the bacteriophage T4. *Nature* 227:680-685.
- Nurjanah S. 2005. Eksplorasi protein dari sponge asal perairan pulau Nias dan Binuangen yang mengkatalisis polimerisasi silika [Thesis]. Bogor: Bogor Agricultural Univ.
- Parkinson J, Gordon R. 1999. Beyond micromachining: the potential of diatom. *Trends Biotechnol* 17:190-195.
- Perry CC. 2003. Silification: The processes by which organism capture and mineralize silica. *J Mineral Geochem* 54:291-327.
- Poulsen N, Kröger N. 2004. Silica morphogenesis by alternative processing of silaffin in the diatom *Thalassiosira pseudonana*. *J Biol Chem* 279:42993-42999.
- Poulsen N, Sumper M, Kröger N. 2003. Biosilica formation in diatoms: characterization of native silaffin-2 and its role in silica morphogenesis. *Proc Nat Acad Sci* 100:12075-12080.
- Round FE, Crawford RM, Mann DG. 1990. *The Diatom*. USA: Cambridge Univ.
- Shimizu K, Cha J, Stucky GD, Morse DE. 1998. Silicatein alpha: cathepsin L-like protein in sponge biosilica. *Proc Nat Acad Sci* 95:6234-6238.
- Strickland JDH, Parsons T. 1972. *A Practical Handbook of Seawater Analysis*. 2nd ed. Ottawa: Fisheries Research Board of Canada.
- Sumper M, Kröger N. 2004. Silica Formation in Diatoms: the function of long-chain polyamines and silaffins. *J Mater Chem* 14:2059-2065.