



# LUND UNIVERSITY

## Aerosol formation and emissions from realistic compartment fires

Malmborg, Vilhelm; Lovén, Karin; Dobric, Julia; Hagvall, Lina; Wierzbicka, Aneta; Strandberg, Bo; Hedmer, Maria; Pagels, Joakim

2022

*Document Version:*  
Other version

[Link to publication](#)

*Citation for published version (APA):*

Malmborg, V., Lovén, K., Dobric, J., Hagvall, L., Wierzbicka, A., Strandberg, B., Hedmer, M., & Pagels, J. (2022). *Aerosol formation and emissions from realistic compartment fires*. 10-10. Abstract from International Aerosol Conference 2022, Aten, Greece.

*Total number of authors:*  
8

*Creative Commons License:*  
Unspecified

### General rights

Unless other specific re-use rights are stated the following general rights apply:

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117  
221 00 Lund  
+46 46-222 00 00

# Aerosol formation and emissions from realistic compartment fires

V. Malmberg<sup>1</sup>, K. Lovén<sup>2</sup>, J. Dobric<sup>1</sup>, L. Hagvall<sup>2</sup>, <sup>1</sup>A. Wierzbicka, B. Strandberg<sup>2</sup>, M. Hedmer<sup>2</sup>, J. Pagels<sup>1</sup>

<sup>1</sup>Ergonomics and Aerosol Technology, LTH, Lund University, Box 118, SE-22100, Lund, Sweden

<sup>2</sup>Occupational and Environmental Medicine, Lund University, Box 117, SE-22100, Lund, Sweden

Keywords: Fire Emissions, Fire Aerosols, Firefighter Occupational Exposure

Presenting author email: Vilhelm.malmberg@design.lth.se

Firefighters are occupationally exposed to a large number of toxic compounds (IARC 2010). The occupational exposure of firefighters has been classified as potentially carcinogenic (class 2B, IARC; (Straif K. et al. 2007)). Poorly quantified emission factors and low understanding of when various aerosol emissions are likely to form during a fire event (initiation, combustion, extinguishing) inhibit efforts to reduce exposure by interventions to the firefighting strategy.

The study was designed to evaluate firefighters' exposure to air pollutants and to allow identification of how aerosol emissions respond to burning conditions and interventions of the firefighting. The study was conducted at the MSB firefighter training facility in Revinge outside Lund, Sweden. Eight small (5x3x2 m) sheds were built to imitate small compartment environments: apartment, bedroom, workshop, etc. These sheds were ignited under realistic fire scenarios (e.g., accident, arson) and later used for training new fire investigators (forensic police). Firefighter students and teachers monitored and extinguished the fires in similar procedures to real fire events. A supervisor monitored the combustion conditions, allowing or restricting fresh-air flow into the fire by opening or closing of the main door.

Fire emissions were extracted from the fire through a 10 m (Ø 6 mm) stainless steel pipe, diluted ~1:50 with HEPA and active charcoal filtered air. The diluted emissions were monitored with a battery of aerosol monitoring instruments. Instrumentation included an aerosol mass spectrometer (Aerodyne SP-AMS, Billerica USA), an aethalometer (AE33, Magee Sci. USA), a differential mobility spectrometer (DMS500, Cambustion, UK), CO<sub>2</sub> monitor (LI-COR, USA), and a NO/NO<sub>2</sub> monitor (2B Tech, USA). Complementary background measurements were positioned downwind or sidewind of the fires. With this equipment we collected data with the aim to resolve relationships between combustion conditions and pollution formation during different phases of a fire response.

The results showed that total particle mass (PM1) emissions correlated with CO<sub>2</sub> emissions and thus fire intensity. The emissions were speciated according to equivalent black carbon (eBC), organic aerosol (OA) and polycyclic aromatic hydrocarbon (PAH) derived from AMS data. When speciated, different particle emissions were found to depend on activities of the firefighting and the supervisor responsible for allowing or restricting fresh air into the combustion

environment. Most evidently, we found that restricting the access to O<sub>2</sub> by closing the door resulted in a sharp increase of OA and even more pronounced, PAH. PAH increased by several orders of magnitude, suggesting that PAH exposure-risks may increase drastically when fires become under-ventilated. Extinguishing the fire with water quickly decreased all particle emissions. The results described are illustrated in Figure 1.

Further analysis involves additional off-line analyses, derivation of emission factors, time-resolved speciated emission analysis and evaluation of relationships between emissions, burning conditions and firefighting strategies.




Figure 1. Time-resolved emissions after 1:65 dilution of net carbon dioxide  $\Delta\text{CO}_2$ , equivalent black carbon (eBC), AMS organic aerosol (OA), AMS polycyclic aromatic hydrocarbon (PAH) and total particle mass (PM1).

This work was supported by the Swedish Research Council for Health, Working Life and Welfare grant 2019-00775\_Forte. We acknowledge the Swedish Contingency Agency (MSB) and the Swedish Police Authority for their important support to this study.

IARC. 2010. *Painting, Firefighting and Shiftwork*. Lyon, France: IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 98. WHO, International Agency for Research on Cancer.

Straif K., et al. 2007. *The Lancet Oncology* 8(12):1065-6. doi: 10.1016/S1470-2045(07)70373-X.