

PAPER • OPEN ACCESS

## Reduction and reconstruction of multisymplectic Lie systems

To cite this article: Javier de Lucas *et al* 2022 *J. Phys. A: Math. Theor.* **55** 295204

View the [article online](#) for updates and enhancements.

You may also like

- [Numerical methods for Hamiltonian PDEs](#)  
Thomas J Bridges and Sebastian Reich
- [Multisymplectic geometry and multisymplectic Preissmann scheme for the KdV equation](#)  
Ping Fu Zhao and Meng Zhao Qin
- [Multisymplectic Geometry for the Seismic Wave Equation](#)  
Chen Jing-Bo



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# Reduction and reconstruction of multisymplectic Lie systems

Javier de Lucas<sup>1,\*</sup> , Xavier Gràcia<sup>2</sup> , Xavier Rivas<sup>3</sup> ,  
Narciso Román-Roy<sup>3</sup>  and Silvia Vilarino<sup>4</sup> 

<sup>1</sup> Department of Mathematical Methods in Physics, University of Warsaw, Warszawa, Poland

<sup>2</sup> Department of Mathematics, Universitat Politècnica de Catalunya, Barcelona, Spain

<sup>3</sup> Escuela Superior de Ingeniería y Tecnología, Universidad Internacional de La Rioja, Logroño, Spain

<sup>4</sup> Centro Universitario de la Defensa de Zaragoza & I.U.M.A., Zaragoza, Spain

E-mail: [javier.de.lucas@fuw.edu.pl](mailto:javier.de.lucas@fuw.edu.pl)

Received 22 March 2022, revised 23 May 2022

Accepted for publication 14 June 2022

Published 4 July 2022



CrossMark

## Abstract

A Lie system is a non-autonomous system of first-order ordinary differential equations describing the integral curves of a non-autonomous vector field taking values in a finite-dimensional real Lie algebra of vector fields, a so-called Vessiot–Guldberg Lie algebra. In this work, multisymplectic forms are applied to the study of the reduction of Lie systems through their Lie symmetries. By using a momentum map, we perform a reduction and reconstruction procedure of multisymplectic Lie systems, which allows us to solve the original problem by analysing several simpler multisymplectic Lie systems. Conversely, we study how reduced multisymplectic Lie systems allow us to retrieve the form of the multisymplectic Lie system that gave rise to them. Our results are illustrated with examples from physics, mathematics, and control theory.

Keywords: Lie system, multisymplectic manifold, multisymplectic reduction and reconstruction, Vessiot–Guldberg Lie algebra, Lie group, time-dependent harmonic oscillator, energy–momentum method

(Some figures may appear in colour only in the online journal)

\* Author to whom any correspondence should be addressed.



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

## 1. Introduction

A *Lie system* is a system of first-order ordinary differential equations whose general solution can be expressed as an autonomous function depending on a generic finite family of particular solutions and a set of constants. The autonomous function is called a *superposition rule* of the Lie system. Standard examples of Lie systems are most types of Riccati equations [32] and non-autonomous linear systems of ordinary differential equations [21, 61].

The Lie–Scheffers theorem states that a Lie system amounts to a  $t$ -dependent vector field taking values in a finite-dimensional real Lie algebra of vector fields, a so-called *Vessiot–Guldberg Lie algebra* [21]. The latter property gave rise to a number of methods for determining superposition rules [19, 21, 61]. Meanwhile, the Lie–Scheffers theorem also showed that being a Lie system is the exception rather than the rule [21]. Despite this, Lie systems admit numerous relevant physical and mathematical applications, as witnessed by the many works on the topic [2, 15, 18, 29, 34, 38, 58, 59, 61].

Recently, a lot of attention has been paid to Lie systems admitting a Vessiot–Guldberg Lie algebra of Hamiltonian vector fields and/or Lie symmetries relative to several types of geometric structures: Poisson [15, 18, 22], symplectic [3, 4, 15, 22, 29], Dirac [15, 17],  $k$ -symplectic [49], multisymplectic [33], Jacobi [36], Riemann [37], and others [15, 46]. Surprisingly, this led to finding much more applications of Lie systems than in the literature dealing with mere Lie systems [5, 15, 46]. Such structures allow for the construction of superposition rules, constants of motion, and other properties of Lie systems in an algebraic manner without relying on the solving of complicated systems of partial or ordinary differential equations [18, 19, 21, 61]. Geometric structures also explain the geometric meaning of superposition rules [4] and gave rise to the study of more general differential equations [6] as well as physical and mathematical problems [15, 46].

This work pioneers the analysis of the reduction of Lie systems with compatible geometric structures [48], in particular with multisymplectic structures [33]. The reduction of multisymplectic Lie systems allows us to simplify the system by reducing its number of variables while ensuring the existence of a compatible multisymplectic structure for the reduced system, which is again a multisymplectic Lie system. This may be used, for instance, to obtain superposition rules for the reduced multisymplectic Lie systems, which in turn can be derived through the multisymplectic structure [33]. It should be noted that our paper presents for the first time the use of a type of reduction procedure for Lie systems (cf [21, 48]).

More particularly, by proposing a certain type of reduction process of multisymplectic structures, which is interesting on its own due to the very wide interest in any kind of multisymplectic reduction process (see [27, 60, 61] and references therein), we perform a reduction process of the so-called multisymplectic Lie systems introduced in [33]. More specifically, our reduction process is focused on Lie systems of locally automorphic type, i.e. they can be considered locally as Lie systems on Lie groups of a very particular but relevant type [33]. Finally, we also describe a reconstruction procedure to recover the initial  $t$ -dependent multisymplectic Lie system from several of its multisymplectic reductions. It is worth noting that our techniques are illustrated by the study of relevant physical and mathematical systems that appear in the study of quantum mechanical systems, control systems, and other topics.

The structure of the paper goes as follows. Section 2 presents the basic theory of Lie systems, including the Lie–Scheffers theorem, and some results on automorphic Lie systems. It also offers a review on multisymplectic structures and the notion of multisymplectic Lie system [33] is presented. The last part of this section is devoted to a survey of the notions of unimodular Lie algebras and unimodular Lie groups.

In section 3 we review some results on locally automorphic Lie systems and give a procedure to find some of their invariants. We also work out two examples to better illustrate these concepts. The first is the so-called generalised Darboux–Brioschi–Halphen system, which appears in the study of triply orthogonal surfaces and vacuum Einstein equations for hyper-Kähler Bianchi-IX metrics [23, 24, 35]. Additionally, it also occurs when reducing the self-dual Yang–Mills equations corresponding to an infinite-dimensional gauge group of diffeomorphisms of a three-dimensional sphere [23]. The second example is a first-order control system on  $\mathbb{R}^5$  with two controls [53, 58].

Section 4 devises a new reduction procedure for multisymplectic Lie systems. We begin by working out a couple of introductory examples: the Schwarz equation and dissipative quantum harmonic oscillators. Having studied some examples, we develop a multisymplectic Lie systems reduction theory. These results are applied to the control system introduced in the previous section and to quantum harmonic oscillators with a spin-magnetic term.

A reconstruction procedure for multisymplectic Lie systems is developed in section 5. This reconstruction is achieved by combining several different reductions in an appropriate way and allows us to recover the initial multisymplectic Lie system. This procedure is applied to the example of quantum harmonic oscillators with a spin-magnetic term introduced in the previous section.

## 2. Some basic concepts and notations

Let us assume some general statements to hold throughout the work unless otherwise explicitly stated. All mathematical objects are smooth, real, and globally defined. This allows us to avoid non-relevant technical problems while stressing the main ideas of our theory. Hereafter,  $N$  will represent an  $n$ -dimensional connected manifold. All manifolds are considered connected. The sum over crossed repeated indices is understood.

### 2.1. Generalised distributions and $t$ -dependent vector fields

Consider a Lie algebra  $V$  with a Lie bracket  $[\cdot, \cdot]$ . Given two subsets  $\mathcal{A}, \mathcal{B} \subset V$ , we denote by  $[\mathcal{A}, \mathcal{B}]$  the linear subspace generated by the Lie brackets between elements of  $\mathcal{A}$  and  $\mathcal{B}$ . Meanwhile,  $\text{Lie}(\mathcal{B})$  stands for the smallest Lie subalgebra of  $V$  containing  $\mathcal{B}$ .

A *Stefan–Sussmann* (or *generalised*) *distribution* on a manifold  $N$  is a correspondence  $\mathcal{D}$  associating each  $x \in N$  with a linear subspace  $\mathcal{D}_x \subset T_x N$ . The dimension of  $\mathcal{D}_x$  is called the *rank* of  $\mathcal{D}$  at  $x$ . It is said that  $\mathcal{D}$  is *regular at*  $x \in N$  when its rank is constant on a neighbourhood of  $x$ . Meanwhile, the generalised distribution  $\mathcal{D}$  is called *regular* when its rank is constant on the whole  $N$ . Regular generalised distributions are called *distributions*. A vector field  $Y \in \mathfrak{X}(N)$  takes values in  $\mathcal{D}$ , let us write it  $Y \in \mathcal{D}$ , when  $Y_x \in \mathcal{D}_x$  for all  $x \in N$ .

We write  $\mathfrak{X}(N)$  for the  $C^\infty(N)$ -module of vector fields on  $N$ . Meanwhile,  $\Omega(N)$  and  $\Omega^k(N)$  stand for the  $C^\infty(N)$ -modules of differential forms and differential  $k$ -forms on  $N$ , respectively. A set  $\mathcal{V} \subset \mathfrak{X}(N)$  of vector fields on  $N$  allows us to define a generalised distribution  $\mathcal{D}^\mathcal{V}$  on  $N$  mapping each  $x \in N$  to the linear span of all the values of its vector fields at  $x$ , i.e.  $\mathcal{D}_x^\mathcal{V} = \text{span}\{X_x | X \in \mathcal{V}\}$ .

A  *$t$ -dependent vector field* on  $N$  is a mapping  $X : (t, x) \in \mathbb{R} \times N \mapsto X(t, x) \in TN$  so that  $\tau_N \circ X = \pi_2$ , where  $\pi_2 : (t, x) \in \mathbb{R} \times N \mapsto x \in N$  and  $\tau_N : TN \rightarrow N$  is the canonical tangent bundle projection. A  $t$ -dependent vector field  $X$  on  $N$  amounts to a  $t$ -parameterised family of vector fields  $\{X_t\}_{t \in \mathbb{R}}$  on  $N$ , with  $X_t : x \in N \mapsto X(t, x) \in TN$  [21]. This enables us to relate  $t$ -dependent vector fields to the following geometric structures.

An integral curve of  $X$  is a curve  $\gamma : \mathbb{R} \rightarrow N$  satisfying that

$$\frac{d\gamma}{dt}(t) = X(t, \gamma(t)), \quad \forall t \in \mathbb{R}. \tag{2.1}$$

Hence,  $\tilde{\gamma} : t \in \mathbb{R} \mapsto (t, \gamma(t)) \in \mathbb{R} \times N$  becomes an integral curve of the *autonomisation*  $\tilde{X}$  of  $X$ , namely the vector field  $\tilde{X} = \partial/\partial t + X$  on  $\mathbb{R} \times N$  [1, 21]. On the contrary, if  $\tilde{\gamma} : \mathbb{R} \rightarrow \mathbb{R} \times N$  is an integral curve of the autonomisation  $\tilde{X}$  and a section of the bundle  $\pi_1 : (t, x) \in \mathbb{R} \times N \mapsto t \in \mathbb{R}$ , then  $\gamma = \pi_2 \circ \tilde{\gamma}$  is an integral curve of  $X$ . Note that  $\tilde{\gamma}$  is a section of the bundle  $\pi_1$  if, and only if,  $\pi_1 \circ \tilde{\gamma} = \text{Id}_{\mathbb{R}}$ . It is worth noting, for clarity, that  $\sigma(t) = (t + 1, \gamma(t + 1))$ , where  $\gamma(t)$  is a solution to (2.1), is not a section of  $\pi_1$ , but it is an integral curve of  $\tilde{X}$ .

Every  $t$ -dependent vector field  $X$  on a manifold  $N$  gives rise to a unique system (2.1) on  $N$  describing its integral curves. Conversely, a system (2.1) describes the integral curves  $\tilde{\gamma} : t \in \mathbb{R} \mapsto (t, \gamma(t)) \in \mathbb{R} \times N$  of the autonomisation of a unique  $t$ -dependent vector field  $X$  on  $N$ . This allows us to use  $X$  to identify both a  $t$ -dependent vector field on  $N$  and its associated system (2.1). This identification will simplify the terminology of our paper without leading to any misunderstanding.

**Definition 2.1.** The *smallest Lie algebra* of a  $t$ -dependent vector field  $X$  on  $N$  is the smallest (in the sense of inclusion) Lie algebra over the reals,  $V^X$ , including the vector fields  $\{X_t\}_{t \in \mathbb{R}}$ , namely  $V^X = \text{Lie}(\{X_t\}_{t \in \mathbb{R}})$ . The *associated distribution* of  $X$ , denoted by  $\mathcal{D}^{V^X}$ , is the generalised distribution on  $N$  spanned by the elements of the Lie algebra of vector fields  $V^X$ .

It can only be ensured that the rank of  $\mathcal{D}^{V^X}$  is constant on the connected components of an open and dense subset of  $N$ , where  $\mathcal{D}^{V^X}$  becomes regular, involutive, and integrable (see [22]). The most relevant instance for us is when  $V^X$  is a finite-dimensional Lie algebra and, therefore, the generalised distribution  $\mathcal{D}^{V^X}$  becomes integrable on the whole  $N$  in the sense of Stefan–Sussmann (see [54, p 63] for details).

### 2.2. Lie systems

Let us survey some fundamental notions in the theory of Lie systems to be used hereafter (see [21] for details).

**Definition 2.2.** A *superposition rule* depending on  $m$  particular solutions for a system  $X$  in  $N$  is a function  $\Phi : N^m \times N \rightarrow N$ ,  $x = \Phi(x_{(1)}, \dots, x_{(m)}; \lambda)$ , so that the general solution,  $x(t)$ , of  $X$  can be written as  $x(t) = \Phi(x_{(1)}(t), \dots, x_{(m)}(t); \lambda)$ , where  $x_{(1)}(t), \dots, x_{(m)}(t)$  is any generic family of particular solutions of  $X$  and  $\lambda$  is a point in  $N$  that is related to the initial conditions. A *Lie system* is a non-autonomous system of first-order ordinary differential equations that admits a superposition rule.

The *Lie–Scheffers theorem* [19, 47] states the conditions that ensure that a system  $X$  possesses a superposition rule:

**Theorem 2.3.** A system  $X$  on  $N$  possesses a superposition rule if, and only if,  $X = \sum_{\alpha=1}^r b_{\alpha}(t)X_{\alpha}$ , for a family  $X_1, \dots, X_r$  of vector fields on  $N$  spanning an  $r$ -dimensional Lie algebra and a set  $b_1(t), \dots, b_r(t)$  of  $t$ -dependent functions.

In other words,  $X$  possesses a superposition rule if, and only if,  $\dim V^X < \infty$ . Then, a Lie system can always be written as  $X = \sum_{\alpha=1}^r b_{\alpha}(t)X_{\alpha}$ , where  $X_1, \dots, X_r$  span a Lie algebra  $V$  that may strictly contain  $V^X$ . Then,  $V$  is called a *Vessiot–Guldberg Lie algebra* of  $X$ . It should be noted that  $V$  may not be univocally defined for a system  $X$ , while  $V^X$  is always characterised by  $X$  [21].

Previous comments allow us to hereafter denote a *Lie system* as a triple  $(N, X, V)$ , where  $N$  is a manifold and  $V$  is a Vessiot–Guldberg Lie algebra of a  $t$ -dependent vector field,  $X$ , on  $N$ .

Let us show how the integration of a Lie system on a manifold can be reduced to integrate one of its Vessiot–Guldberg Lie algebras to a Lie group action and to know a particular solution of a Lie system on a Lie group of a very specific type, a so-called automorphic Lie system [18, 60]. With this aim, let us recall some basic results regarding Lie group actions.

Let  $\varphi : D \subset G \times N \rightarrow N$  be a local (left) Lie group action and define  $\varphi_x : g \in D_G \subset G \mapsto \varphi(g, x) \in N$  so that  $D_G = \{g \in G | (g, x) \in D\}$  [57]. For every  $\xi \in T_e G$ , we set the fundamental vector field  $\xi_N \in \mathfrak{X}(N)$  to be  $\xi_N(x) = T_e \varphi_x(\xi)$ . Let us denote by  $\xi^L$  and  $\xi^R$  the respective invariant vector fields associated with  $\xi \in T_e G$ . Hence, there are linear isomorphisms  $\xi \in T_e G \mapsto \xi^L \in \mathfrak{X}_L(G)$  and  $\xi \in T_e G \mapsto \xi^R \in \mathfrak{X}_R(G)$ , where  $\mathfrak{X}_L(G)$  and  $\mathfrak{X}_R(G)$  are the linear spaces of left- and right-invariant vector fields on  $G$ . The space  $\mathfrak{g} = T_e G$  inherits a Lie algebra structure from  $\mathfrak{X}_L(G)$  via the identification given by the linear isomorphism  $\xi \in T_e G \mapsto \xi^L \in \mathfrak{X}_L(G)$ . Then (cf [44, ch 20]),

- (a) The map  $\widehat{\varphi} : \mathfrak{g} \rightarrow \mathfrak{X}(N)$ ,  $\xi \mapsto \xi_N$ , is a Lie algebra antihomomorphism (the *infinitesimal generator* of  $\varphi$ ).
- (b) For every  $x \in N$ , the *right*-invariant vector field  $\xi^R$  and the fundamental vector field  $\xi_N$  are  $\varphi_x$ -related [44, p 182], namely  $T_g \varphi_x(\xi^R(g)) = \xi_N(\varphi_x(g))$  for every  $g \in G$ .

In general, an antihomomorphism  $\mathfrak{g} \rightarrow \mathfrak{X}(N)$  is called a (left) Lie algebra action, and, when  $\mathfrak{g}$  is finite-dimensional, it can be integrated to a Lie group action. More precisely:

**Theorem 2.4.** *Let  $\mathfrak{g}$  be a finite-dimensional Lie algebra, and let  $G$  be a Lie group with Lie algebra  $\mathfrak{g}$ . Given a Lie algebra morphism  $\widehat{\varphi} : \mathfrak{g} \rightarrow \mathfrak{X}(N)$ , there exists a local Lie group action  $\varphi : G \times N \rightarrow N$  whose infinitesimal generator is  $\widehat{\varphi}$ . If  $G$  is connected and simply connected and the vector fields  $\widehat{\varphi}(\mathfrak{g})$  are complete, then  $\varphi$  becomes a global Lie group action.*

The proof of these results, and other related facts, can be found, for instance, in [57, p 58] and [9, p 207]. This Lie group action is the device that relates the Lie system on  $N$  to a Lie system on  $G$ :

**Theorem 2.5.** *Let  $(N, X, V)$  be a Lie system given by  $X = \sum_{\alpha=1}^r b_\alpha(t)X_\alpha$ , where  $X_1, \dots, X_r$  is a basis of the Vessiot–Guldberg Lie algebra  $V$  and  $b_1(t), \dots, b_r(t)$  are arbitrary  $t$ -dependent functions. Let  $G$  be a Lie group whose Lie algebra is isomorphic to  $V$ , and let  $\varphi : D_G \subset G \times N \rightarrow N$  be a local Lie group action as given by theorem 2.4. If  $X_\alpha^R$  is the right-invariant vector field on  $G$  related to the vector field  $X_\alpha$  through  $\varphi$ , with  $\alpha = 1, \dots, r$ , then*

- (a) The triple  $(G, X^G, V^G)$ , where

$$X^G(t, g) = \sum_{\alpha=1}^r b_\alpha(t)X_\alpha^R(g), \quad \forall g \in G, \forall t \in \mathbb{R}, \tag{2.2}$$

and  $V^G = \mathfrak{X}_R(G)$ , is a Lie system on  $G$ .

- (b) For every  $x_0 \in N$  and  $t \in \mathbb{R}$ , the vector field  $X_t^G$  is  $\varphi_{x_0}$ -related with  $X_t$ ; namely, the  $t$ -dependent vector fields  $X^G$  and  $X$  are  $\varphi_{x_0}$ -related.
- (c) If  $g(t)$  is the integral curve of  $X^G$  with  $g(0) = e$ , and  $x_0 \in N$ , then  $x(t) = \varphi(g(t), x_0)$  is the integral curve of  $X$  with  $x(0) = x_0$ .

The proof of this result is almost immediate: the vector fields  $X_\alpha^R$  span the Lie algebra  $\mathfrak{X}_R(G)$ , the  $t$ -dependent vector field  $X^G$  is  $\varphi_{x_0}$ -related with  $X$ , and this implies that  $\varphi_{x_0}$  maps integral curves of  $X^G$  to integral curves of  $X$  (see [18, 21] for details).



Hence, knowing the explicit form of  $\varphi$  allows us to obtain the solutions to  $X$  via a single particular solution to (2.2). The other way around, the general solution of  $X$  determines the integral curve of (2.2) with  $g(0) = e$  using an algebraic system of equations determined by  $\varphi$ .

Lie systems (2.2) are frequently called *automorphic Lie systems* [8, 60]. Due to their specific structure, automorphic Lie systems possess invariant differential forms with respect to their evolution (see [33] for details). From now on, we say that the automorphic Lie system  $(G, X^G, \mathfrak{X}_R(G))$  is the automorphic Lie system *related* to  $(N, X, V)$ .

The automorphic Lie system related to a  $t$ -dependent right-invariant vector field (2.2) is invariant relative to right multiplications  $R_g : g' \in G \rightarrow g'g \in G$ . Then, if  $g_1(t)$  is the particular solution to the system  $X^G$  with an initial condition  $g_1(0) \in G$ , then  $R_g g_1(t)$  is the particular solution of  $X^G$  with the initial condition  $g_1(0)g$ . Consequently, the general solution to  $X^G$  can be brought into the form  $g(t) = \Phi^G(g_1(t); g) := R_g g_1(t)$ , which leads to the definition of the superposition rule  $\Phi^G : (g_1; g) \in G \times G \mapsto R_g g_1 \in G$ , which depends on a unique particular solution [21].

### 2.3. Multisymplectic manifolds and Lie systems

This section reviews the fundamental properties of multisymplectic manifolds to be used hereafter (see [13, 14, 25] for details).

A differential  $k$ -form  $\omega$  on  $N$  is called *one-nondegenerate* if, and only if, the vector bundle morphism

$$\begin{aligned} \omega^\flat: \quad TN &\rightarrow \Lambda^{k-1}T^*N \\ X_p &\mapsto \iota_{X_p}\omega_p, \end{aligned}$$

where  $X_p \in T_pN$  and  $\iota_{X_p}\omega_p$  stands for an inner contraction, is injective. If this is the case, the induced morphism of  $C^\infty(N)$ -modules  $\widehat{\omega} : \mathfrak{X}(N) \rightarrow \Omega^{k-1}(N), X \mapsto \iota_X\omega$ , is also injective.

**Definition 2.6.** A *multisymplectic  $k$ -form* on an  $n$ -dimensional manifold  $N$  is a one-nondegenerate and closed differential  $k$ -form  $\Theta \in \Omega^k(N)$ . A *multisymplectic manifold* of degree  $k$  is a pair  $(N, \Theta)$ , where  $\Theta$  is a multisymplectic  $k$ -form on  $N$ .

The multisymplectic two-forms are the symplectic forms. Multisymplectic  $n$ -forms on  $N$  are the volume forms on  $N$ . We hereafter assume that  $\dim N \geq 2$  and, thus, every multisymplectic  $k$ -form has degree  $k \geq 2$ . The ‘multisymplectic’ term can be misleading and it should not be confused with other structures, such as  $k$ -symplectic and polysymplectic [45, p 21], where several differential forms appear. Anyhow, we will stick to the standard nomenclature, which has been established for almost fifty years now [40, 41].

**Definition 2.7.** Let  $(N, \Theta)$  be a multisymplectic manifold of degree  $k$ . A vector field  $X$  on  $N$  is *locally Hamiltonian* if  $\iota_X\Theta$  is closed. A vector field  $X$  is *Hamiltonian* if  $\iota_X\Theta$  is exact, i.e.  $\iota_X\Theta = d\Upsilon$  for a differential  $(k - 2)$ -form  $\Upsilon_X$  on  $N$ . In previous cases,  $\iota_X\Theta$  and  $\Upsilon_X$  are called *Hamiltonian*  $(k - 1)$ - and  $(k - 2)$ -forms associated with  $X$ , respectively.

A vector field is locally Hamiltonian if, and only if,  $\Theta$  is invariant by  $X$ , that is,

$$\mathcal{L}_X\Theta = 0,$$

where  $\mathcal{L}_X\Theta$  stands for the Lie derivative of  $\Theta$  relative to  $X$ .

To keep the notation simple, we will frequently call  $\iota_X\Omega$  and  $\Upsilon_X$  Hamiltonian forms associated with  $X$ . Note that the degree is obvious from the context and does not need to be detailed.

In analogy with symplectic geometry, it is possible to define certain brackets on the spaces of Hamiltonian forms on multisymplectic manifolds. Notwithstanding, their properties generally differ significantly from the properties of the brackets defined in symplectic geometry.

**Definition 2.8.** Let  $(N, \Theta)$  be a multisymplectic manifold of degree  $k$ .

- A bracket between Hamiltonian  $(k - 1)$ -forms can be set as follows. Let  $\xi, \zeta \in \text{Im } \widehat{\Theta} \subset \Omega^{k-1}(N)$ , and let  $X, Y \in \mathfrak{X}(N)$  be the unique vector fields such that  $\iota_X \Theta = \xi$  and  $\iota_Y \Theta = \zeta$ . The bracket between  $\xi$  and  $\zeta$  is defined by

$$\{\xi, \zeta\} = \iota_{[Y, X]} \Theta \in \text{Im } \widehat{\Theta}. \tag{2.3}$$

This bracket satisfies the Jacobi identity and, therefore, becomes a Lie bracket.

- The bracket between Hamiltonian  $(k - 2)$ -forms will be now defined. Let  $X, Y$  be Hamiltonian vector fields and let  $\Upsilon_X, \Upsilon_Y \in \Omega^{k-2}(N)$  be Hamiltonian forms of  $X$  and  $Y$ , respectively. Then,

$$\{\Upsilon_X, \Upsilon_Y\} = \iota_Y \iota_X \Theta$$

gives rise to a bracket on  $(k - 2)$ -forms. It can be proved that the bracket of Hamiltonian  $(k - 2)$ -forms needs not be a Lie bracket for  $k > 2$  [13].

Although the brackets for Hamiltonian  $(k - 1)$  and  $(k - 2)$ -forms have been denoted in the same way, this will not lead to confusion and will simplify the notation.

Note that

$$d\{\Upsilon_X, \Upsilon_Y\} = d\iota_Y \iota_X \Theta = \iota_{[Y, X]} \Theta = \{d\Upsilon_X, d\Upsilon_Y\}. \tag{2.4}$$

The equality  $d\{\Upsilon_X, \Upsilon_Y\} = \iota_{[Y, X]} \Theta$ , shows that  $[Y, X]$  is a Hamiltonian vector field admitting a Hamiltonian form  $\{\Upsilon_X, \Upsilon_Y\}$ . Consequently, the space of (locally) Hamiltonian vector fields ( $\text{Ham}_{\text{loc}}(N)$ )  $\text{Ham}(N)$  becomes a Lie algebra, and the map attaching a (locally) Hamiltonian vector field  $X$  to  $\iota_X \Theta$  is an injective Lie algebra anti-homomorphism.

Let us recall the notion of a multivector field, which is needed in our reduction procedure for multisymplectic Lie systems (see [26, 30] for more details). An  $\ell$ -multivector field on  $N$  is a section of the bundle  $\Lambda^\ell(TN)$ . An  $\ell$ -multivector field  $Y$  on  $N$  is said to be *decomposable* if there exists a family of vector fields  $Y_1, \dots, Y_\ell \in \mathfrak{X}(N)$  such that  $Y = Y_1 \wedge \dots \wedge Y_\ell$ . We will denote by  $\mathfrak{X}^\ell(N)$  the set of  $\ell$ -multivector fields.

Let  $(N, \Theta)$  be a multisymplectic manifold of degree  $k$ . An  $\ell$ -multivector field  $Y$  is *Hamiltonian* (with respect to  $\Theta$ ) if there exists a  $(k - \ell - 1)$ -form  $\theta$  such that  $\iota_Y \Theta = d\theta$ . This notion allows us to generalise the notion of Hamiltonian vector fields to multivector fields. Furthermore,  $Y$  is *locally Hamiltonian* or *multisymplectic* if  $\mathcal{L}_Y \Theta = 0$  (see [13, 14]).

Finally, let us recall the notion of a multisymplectic Lie system [33].

**Definition 2.9.** A (locally) multisymplectic Lie system is a triple  $(N, \Theta, X)$ , where  $X$  is a Lie system whose smallest Lie algebra  $V^X$  is a finite-dimensional real Lie algebra of (locally) Hamiltonian vector fields relative to a multisymplectic structure  $\Theta$  on  $N$ .

#### 2.4. Unimodular Lie algebras

This section surveys the fundamentals of unimodular Lie algebras and Lie groups to be used hereafter.



Let  $G$  be a Lie group with Lie algebra  $\mathfrak{g} = T_e G$ . A (left) *Haar measure* on  $G$  is a left-invariant volume form on  $G$  [39]. Every Lie group admits a Haar measure given by a left-invariant volume form, and it is unique up to a non-zero multiplicative constant (cf [12]).

Let  $\{X_1^L, \dots, X_r^L\}$  be a basis of the Lie algebra  $\mathfrak{X}_L(G)$  of left-invariant vector fields on  $G$  and let  $\{\eta_1^L, \dots, \eta_r^L\}$  be its dual basis of left-invariant differential one-forms. Then, any left-invariant volume form on  $G$  is a non-zero scalar multiple of

$$\Theta = \eta_1^L \wedge \dots \wedge \eta_r^L.$$

If  $X^L$  is a left-invariant vector field on  $G$ , then [33]:

$$\mathcal{L}_{X^L} \Theta = -\text{Tr}(\text{ad}_{X^L}) \Theta; \tag{2.5}$$

where  $\text{Tr}$  stands for the trace of an endomorphism, and  $\text{ad} : v \in \mathfrak{g} \mapsto \text{ad}_v \in \text{End}(\mathfrak{g})$ , is the adjoint representation of a Lie algebra  $\mathfrak{g}$  and given by  $\text{ad}_v w = [v, w]$ .

A Lie group is called *unimodular* if its Haar measure is *bi-invariant*, namely it is left- and right-invariant [52]. For instance, Abelian, compact, and semi-simple Lie groups, are all unimodular [62]. This work focusses on the Lie algebras of unimodular Lie groups, whose most important properties (for our purposes) are detailed in the definition and proposition below (for details, see [33]).

**Definition 2.10.** A finite-dimensional Lie algebra  $\mathfrak{g}$  is called *unimodular* when the maps  $\text{ad}_v \in \text{End}(\mathfrak{g})$  are traceless—we say then that the adjoint representation is traceless.

**Proposition 2.11.** A (connected) Lie group  $G$  is unimodular if, and only if, its Lie algebra is unimodular.

**Remark 2.12.** It is worth noting that each left-invariant vector field  $X$  on  $G$  has a flow of the form  $\phi : t \in \mathbb{R} \mapsto R_{\exp(tX_e)} \in \text{Diff}(G)$ . Then, a vector field  $Y$  on a connected Lie group  $G$  is right-invariant if, and only if, it commutes with *every* left-invariant vector field  $X^L$ , namely  $\mathcal{L}_{X^L} Y = 0$ . This also applies to tensor fields on  $G$ .

**Definition 2.13.** Let  $\mathfrak{g}$  be an  $n$ -dimensional Lie algebra. We write  $\mathcal{G}(r, \mathfrak{g})$  for the set of decomposable  $r$ -multivectors spanning  $r$ -dimensional unimodular Lie subalgebras. The set  $\mathcal{G}(\mathfrak{g}) = \bigcup_{r=0}^n \mathcal{G}(r, \mathfrak{g})$  is called the *unimodular Grassmannian* of  $\mathfrak{g}$ .

### 3. Locally automorphic Lie systems and invariant forms

In this section, we recall conditions that ensure the existence of a multisymplectic form  $\Theta$  invariant with respect to the elements of a Vessiot–Guldberg Lie algebra  $V$ .

In general, it is difficult to find multisymplectic forms compatible with a Lie system  $X$  admitting a Vessiot–Guldberg Lie algebra  $V$ , because this requires finding adequate solutions  $\Theta$  to a system of partial differential equations  $\mathcal{L}_Y \Theta = 0$  for every  $Y \in V$ . However, we can develop several simpler methods to find compatible invariant forms for a special class of Lie systems with many relevant physical applications: the so-called locally automorphic Lie systems.

#### 3.1. Locally automorphic Lie systems

**Definition 3.1.** A *locally automorphic Lie system* is a triple  $(N, X, V)$ , where  $X$  is a Lie system on the manifold  $N$  with a Vessiot–Guldberg Lie algebra  $V$  such that  $\dim V = \dim N$  and  $\mathcal{D}^V = TN$ .

The following result, whose proof can be found in [33], shows that locally automorphic Lie systems are locally diffeomorphic to automorphic Lie systems, thus motivating the previous definition.

**Theorem 3.2.** *Consider a locally automorphic Lie system  $(N, X, V)$ . Let  $G$  be a Lie group whose Lie algebra is isomorphic to  $V$ , let  $\varphi$  be a local action of  $G$  on  $N$  obtained from the integration of  $V$ , and let  $(G, X^G, V^G)$  be the corresponding automorphic Lie system on  $G$  given by theorem 2.5. Then, for every  $x \in N$ , the map  $\varphi_x = \varphi(\cdot, x)$  is a local diffeomorphism such that  $T_g\varphi_x(X_t^G(g)) = X_t(\varphi_x(g))$  for every  $t \in \mathbb{R}$ .*

Recall that the action  $\varphi$  can be ensured to be globally defined if, and only if,  $G$  is simply connected and the Lie algebra  $V$  consists of complete vector fields [57]. From now on, we assume that the action  $\varphi$  is defined globally.

The mapping  $\varphi$  allows us not only to find local diffeomorphisms  $\varphi_x : G \rightarrow N$ , with  $x \in N$ , but also maps certain geometric structures related to the locally automorphic Lie system  $(N, X, V)$  with the associated automorphic Lie system  $(G, X^R, V^R)$ .

As a consequence of theorem 3.2, we have the following corollaries.

**Corollary 3.3.** *Consider a locally automorphic Lie system  $(N, X, V)$ . Then,  $X$  admits a superposition rule that depends only on one particular solution of  $X$ .*

**Corollary 3.4.** *Let  $(N, X, V^X)$  be a locally automorphic Lie system on a (connected) manifold. Then all  $t$ -independent constants of motion of  $X$  are constants.*

The existence of the local diffeomorphism  $\varphi_x$  allows us to obtain theoretical properties of locally automorphic Lie systems, such as corollary 3.3. However, its use to map them into automorphic Lie systems is quite limited due to the locality of  $\varphi_x$  and the difficulties in obtaining an explicit expression, which must be obtained by solving a system of nonlinear ordinary differential equations determined by the Lie algebra  $V$ .

This can be seen in the next two examples of locally automorphic Lie systems.

**Example 3.5 (The generalised Darboux–Brioschi–Halphen (DBH) system [24]).**

Consider the system of ordinary differential equations given by

$$\begin{cases} \frac{dw_1}{dt} = w_3w_2 - w_1w_3 - w_1w_2 + \tau^2, \\ \frac{dw_2}{dt} = w_1w_3 - w_2w_1 - w_2w_3 + \tau^2, \\ \frac{dw_3}{dt} = w_2w_1 - w_3w_2 - w_3w_1 + \tau^2, \end{cases} \quad (3.1)$$

where

$$\tau^2 = \alpha_1^2(w_1 - w_2)(w_3 - w_1) + \alpha_2^2(w_2 - w_3)(w_1 - w_2) + \alpha_3^2(w_3 - w_1)(w_2 - w_3),$$

and  $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$ . The DBH system with  $\tau = 0$  appears in the description of triply orthogonal surfaces and the vacuum Einstein equations for hyper-Kähler Bianchi-IX metrics [23, 24, 35]. Furthermore, the generalised DBH system for  $\tau \neq 0$  is a reduction of the self-dual Yang–Mills equations corresponding to an infinite-dimensional gauge group of diffeomorphisms of a three-dimensional sphere [23].

Even though the DBH system is autonomous, it is useful, e.g. to obtain its Lie symmetries [28], to view it as a Lie system related to a Vessiot–Guldberg Lie algebra  $V^{\text{DBH}}$  ( $V$  to simplify

the notation) spanned by the vector fields

$$\begin{aligned} X_1 &= \frac{\partial}{\partial w_1} + \frac{\partial}{\partial w_2} + \frac{\partial}{\partial w_3}, \\ X_2 &= w_1 \frac{\partial}{\partial w_1} + w_2 \frac{\partial}{\partial w_2} + w_3 \frac{\partial}{\partial w_3}, \\ X_3 &= -(w_3 w_2 - w_1(w_3 + w_2) + \tau^2) \frac{\partial}{\partial w_1} - (w_1 w_3 - w_2(w_1 + w_3) + \tau^2) \frac{\partial}{\partial w_2} \\ &\quad - (w_2 w_1 - w_3(w_2 + w_1) + \tau^2) \frac{\partial}{\partial w_3}. \end{aligned}$$

One can check that the commutation relations are,

$$[X_1, X_2] = X_1, \quad [X_1, X_3] = 2X_2, \quad [X_2, X_3] = X_3.$$

Thus, we have  $V \simeq \mathfrak{sl}_2$ ,  $\dim V = \dim \mathcal{O}$  and  $X_1 \wedge X_2 \wedge X_3 \neq 0$  on an open submanifold  $\mathcal{O}$  of  $\mathbb{R}^3$ . Hence,  $(\mathcal{O}, X, V)$  is a locally automorphic Lie system. To obtain a local diffeomorphism mapping the DBH system into an automorphic one, one needs to integrate the vector fields of  $V$ . In view of their analytic form, it is clear that it is very difficult to provide such a local diffeomorphism.

**Example 3.6 (A control system [53, 58]).** Consider the following system of ordinary differential equations on  $\mathbb{R}^5$ :

$$\begin{aligned} \frac{dx_1}{dt} &= b_1(t), & \frac{dx_2}{dt} &= b_2(t), & \frac{dx_3}{dt} &= b_2(t)x_1, & \frac{dx_4}{dt} &= b_2(t)x_1^2, \\ \frac{dx_5}{dt} &= 2b_2(t)x_1x_2, \end{aligned} \tag{3.2}$$

where  $b_1(t)$  and  $b_2(t)$  are arbitrary  $t$ -dependent functions.

This control system is defined by the  $t$ -dependent vector field  $X^{\text{CS}} = b_1(t)X_1 + b_2(t)X_2$  on  $\mathbb{R}^5$ , where the vector fields

$$\begin{aligned} X_1 &= \frac{\partial}{\partial x_1}, & X_2 &= \frac{\partial}{\partial x_2} + x_1 \frac{\partial}{\partial x_3} + x_1^2 \frac{\partial}{\partial x_4} + 2x_1x_2 \frac{\partial}{\partial x_5}, \\ X_3 &= \frac{\partial}{\partial x_3} + 2x_1 \frac{\partial}{\partial x_4} + 2x_2 \frac{\partial}{\partial x_5}, & X_4 &= \frac{\partial}{\partial x_4}, & X_5 &= \frac{\partial}{\partial x_5} \end{aligned} \tag{3.3}$$

are such that their only non-vanishing commutation relations are

$$[X_1, X_2] = X_3, \quad [X_1, X_3] = 2X_4, \quad [X_2, X_3] = 2X_5. \tag{3.4}$$

It is remarkable that the initial system of differential equation (3.2) can be described via a  $t$ -dependent vector field that can be written as a linear combination of the vector fields  $X_1, X_2$  with  $t$ -dependent coefficients. Nevertheless, to prove that  $X$  is a Lie system, one has to add to  $X_1, X_2$  as many vector fields as necessary to ensure that all such vector fields close a Vessiot–Guldberg Lie algebra. In fact, the vector fields  $X_3, X_4, X_5$  appear as successive Lie brackets of the vector fields  $X_1, X_2$  and give rise, all of them, to a basis of a Vessiot–Guldberg Lie algebra.

Thus,  $X_1, \dots, X_5$  span a five-dimensional nilpotent Lie algebra  $V^{\text{CS}}$ . Since  $X^{\text{CS}}$  takes values in  $V^{\text{CS}}$ , we have that  $(\mathbb{R}^5, X^{\text{CS}}, V^{\text{CS}})$  is a Lie system [58]. The vector fields of  $V^{\text{CS}}$  span a distribution  $\mathcal{D}^{V^{\text{CS}}} = \mathbb{T}\mathbb{R}^5$  and  $\dim V^{\text{CS}} = \dim \mathbb{R}^5$ . Then,  $(\mathbb{R}^5, X^{\text{CS}}, V^{\text{CS}})$  becomes a locally automorphic Lie system.

### 3.2. Invariants for locally automorphic Lie systems

A Lie symmetry of a Lie system  $(N, X, V)$  is a vector field  $Y \in \mathfrak{X}(N)$  such that  $\mathcal{L}_Y Z = 0$  for every vector field  $Z \in V$ . If  $V$  is spanned by the vector fields  $X_1, \dots, X_r$ , this condition is equivalent to saying that  $Y$  has to fulfil the following system of partial differential equations:

$$\mathcal{L}_{X_i} Y = 0, \quad i = 1, \dots, r. \tag{3.5}$$

Note that the space of Lie symmetries of  $(N, X, V)$  just depends on the Vessiot–Guldberg Lie algebra  $V$ , which motivates denoting it by  $\text{Sym}(V)$ . Moreover,  $\text{Sym}(V)$  is a Lie algebra. In what follows, we study  $\text{Sym}(V)$  for the case of locally automorphic Lie systems  $(N, X, V)$ .

Consider a locally automorphic Lie system  $(N, X, V)$ . Then, each mapping  $\varphi_x$  maps it to an automorphic Lie system  $(G, X^R, V^R)$ . It is clear that  $\text{Sym}(V^R) = V^L$ . Since  $\varphi_x$  is a local diffeomorphism mapping  $V$  onto  $V^R$ , then it also maps  $\text{Sym}(V)$  onto  $V^L$ . Thus, we have the following lemma, whose implications are shown in example 3.8.

**Lemma 3.7.** *The Lie algebra of Lie symmetries of  $(N, X, V)$ , namely  $\text{Sym}(V)$ , is isomorphic to  $V$ .*

**Example 3.8.** Consider again example 3.6, where we studied the control system given by (3.2).

Solving the linear system of partial differential equation (3.5) in the coefficients of a vector field  $Y$  in the basis  $\partial/\partial x_1, \dots, \partial/\partial x_5$ , which demands a very long and tedious calculation, one sees that every Lie symmetry  $Y$  of an arbitrary control system of the form (3.2) has to be a linear combination with constant coefficients of the vector fields

$$\begin{aligned} Y_1 &= \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial x_3} + 2x_3 \frac{\partial}{\partial x_4} + x_2^2 \frac{\partial}{\partial x_5}, & Y_2 &= \frac{\partial}{\partial x_2} + 2x_3 \frac{\partial}{\partial x_5}, \\ Y_3 &= \frac{\partial}{\partial x_3}, & Y_4 &= \frac{\partial}{\partial x_4}, & Y_5 &= \frac{\partial}{\partial x_5}. \end{aligned} \tag{3.6}$$

It is easy to see that the vector fields  $-Y_i$ , where  $i = 1, \dots, 5$ , span a Lie algebra with the same structure constants as  $V^{\text{CS}} = \langle X_1, \dots, X_5 \rangle$ .

From the fact that every locally automorphic Lie system  $(N, X, V)$  is locally diffeomorphic to an automorphic Lie system  $(G, X^R, V^R)$ , we find that every differential form on  $N$  that is invariant with respect to the Lie derivative of elements of  $V$  must be locally diffeomorphic to a left-invariant differential form on  $G$ . Since  $\text{Sym}(V)$  is also diffeomorphic to  $V^L$ , and taking into account remark 2.12, one gets the following theorem:

**Theorem 3.9.** *Consider a locally automorphic Lie system  $(N, X, V)$ . Let  $Y_1, \dots, Y_r$  be a basis of  $\text{Sym}(V)$ , with a dual frame  $\nu^1, \dots, \nu^r$ . Then, a differential form on  $N$  is invariant with respect to the Lie algebra  $V$  if, and only if, it is a linear combination with real coefficients of the exterior products of  $\nu^1, \dots, \nu^r$ .*

**Remark 3.10.** Let us consider a result on the invariants of general multisymplectic Lie systems. Let  $(N, \Theta, X)$  be a multisymplectic Lie system  $X$  with a Vessiot–Guldberg Lie algebra

$V^X$ . A Lie algebra of vector fields  $W$  on  $N$  is a Lie algebra of symmetries of the Lie system  $X$  if, and only if,  $[V^X, W] = 0$ . Let  $Y_1, \dots, Y_n$  be a basis of  $W$ . Then,

$${}_{\iota}Y_1 \wedge \dots \wedge Y_n \ominus$$

are constants of the motion of our system.

#### 4. Reduction of multisymplectic Lie systems

This section introduces a reduction procedure for multisymplectic Lie systems by infinitesimal symmetries of the associated Lie system and its compatible multisymplectic form. On the one hand, this simplifies the study of certain multisymplectic Lie systems by transforming them into new ones on manifolds of smaller dimension. On the other hand, the multisymplectic form of the reduced multisymplectic Lie system enables one to use the methods described in [33] to study its properties. To introduce our multisymplectic Lie system reduction theory, we illustrate a reduction procedure based upon some examples.

##### 4.1. Introductory examples

The following examples show how multisymplectic Lie systems can be reduced to new multisymplectic Lie systems on manifolds of smaller dimension.

**4.1.1. Schwarz equations.** Consider a Schwarz equation of the form [7, 56]

$$\frac{d^3x}{dt^3} = \frac{3}{2} \left( \frac{dx}{dt} \right)^{-1} \left( \frac{d^2x}{dt^2} \right)^2 + 2b_1(t) \frac{dx}{dt}, \tag{4.1}$$

where  $b_1(t)$  is a non-constant function. The relevance of (4.1) is due to its appearance in the study of Ermakov systems [42] and the Schwarz derivative (see [17] and references therein).

The differential equation (4.1) is known to be a higher-order Lie system [16]. This means that the associated system of first-order differential equations obtained by adding the variables  $v := dx/dt$  and  $a := d^2x/dt^2$ , i.e.

$$\frac{dx}{dt} = v, \quad \frac{dv}{dt} = a, \quad \frac{da}{dt} = \frac{3}{2} \frac{a^2}{v} + 2b_1(t)v \tag{4.2}$$

is a Lie system. Indeed, (4.2) is associated with the  $t$ -dependent vector field on  $\mathcal{O} := \{(x, v, a) \in \mathbb{T}^2\mathbb{R} | v \neq 0\}$  of the form  $X^S = X_3 + b_1(t)X_1$ , where the vector fields

$$X_1 = 2v \frac{\partial}{\partial a}, \quad X_2 = v \frac{\partial}{\partial v} + 2a \frac{\partial}{\partial a}, \quad X_3 = v \frac{\partial}{\partial x} + a \frac{\partial}{\partial v} + \frac{3}{2} \frac{a^2}{v} \frac{\partial}{\partial a} \tag{4.3}$$

satisfy the commutation relations

$$[X_1, X_2] = X_1, \quad [X_1, X_3] = 2X_2, \quad [X_2, X_3] = X_3. \tag{4.4}$$

Consequently,  $X_1, X_2, X_3$  span a three-dimensional Lie algebra of vector fields  $V^S$  isomorphic to  $\mathfrak{sl}_2$  and  $X^S$  becomes a  $t$ -dependent vector field taking values in  $V^S$ , i.e.  $X^S$  is a Lie system.

In [33] it was proved that the vector fields of  $V^S$  are Hamiltonian relative to the multisymplectic structure  $\Theta_S$  on  $\mathcal{O}$  given by  $\Theta_S = \frac{1}{2v^3} da \wedge dv \wedge dx$  satisfying that

$$\mathcal{L}_{X_\alpha} \Theta_S = 0, \quad \alpha = 1, 2, 3. \tag{4.5}$$

Therefore,  $(\mathcal{O}, \Theta_S, X^S)$  is a multisymplectic Lie system. Moreover,

$$\begin{aligned} \iota_{X_1} \Theta_S &= \frac{1}{v^2} dv \wedge dx = d\left(-\frac{1}{v} dx\right) = d\theta_1, \\ \iota_{X_2} \Theta_S &= \frac{1}{v} \left(\frac{a}{v^2} dv - \frac{1}{2v} da\right) \wedge dx = d\left(-\frac{a}{2v^2} dx + \frac{dv}{2v}\right) = d\theta_2, \\ \iota_{X_3} \Theta_S &= -\frac{3a^2}{4v^4} dx \wedge dv - \frac{a}{2v^3} da \wedge dx + \frac{1}{2v^2} da \wedge dv \\ &= d\left(-\frac{a^2}{4v^3} dx + \frac{a}{v^2} dv - \frac{1}{2v} da\right) = d\theta_3. \end{aligned} \tag{4.6}$$

Hence,  $X_1, X_2, X_3$  are Hamiltonian vector fields with respect to the multisymplectic manifold  $(\mathcal{O}, \Theta_S)$ , with Hamiltonian one-forms

$$\theta_1 = -\frac{dx}{v}, \quad \theta_2 = -\frac{a}{2v^2} dx + \frac{dv}{2v}, \quad \theta_3 = -\frac{a^2}{4v^3} dx + \frac{a}{v^2} dv - \frac{1}{2v} da.$$

Consequently, no matter what the  $t$ -dependent coefficient  $b_1(t)$  in (4.2) is, the evolution of  $X^S$  preserves the volume form  $\Theta_S$ . Since  $\mathcal{D}^{V^S} = T\mathcal{O}$ , and in view of (4.5), the value of  $\Theta_S$  at a point  $o \in \mathcal{O}$  determines the value of  $\Theta_S$  on the connected component of  $o$  in  $\mathcal{O}$ . Moreover,  $\Theta_S$  is, up to a multiplicative constant on each connected component of  $\mathcal{O}$ , the only volume form satisfying equation (4.5). Since any one-form or two-form on a three-dimensional manifold is one-degenerate, the system under study has a unique, up to a non-zero proportional constant, multisymplectic form which is invariant under the action of  $V^S$ .

The Schwarz equation, whose first-order system  $X^S$  is given by (4.2), admits a Lie algebra of Lie symmetries,  $\text{Sym}(V^S)$ , spanned by (see [43, 55])

$$Y_1 = \frac{\partial}{\partial x}, \quad Y_2 = x \frac{\partial}{\partial x} + v \frac{\partial}{\partial v} + a \frac{\partial}{\partial a}, \quad Y_3 = x^2 \frac{\partial}{\partial x} + 2vx \frac{\partial}{\partial v} + 2(ax + v^2) \frac{\partial}{\partial a}. \tag{4.7}$$

Since  $X^S$  is a locally automorphic Lie system,  $\text{Sym}(V^S)$  is the Lie algebra of Lie symmetries given by lemma 3.7 and  $[X_\alpha, Y_\beta] = 0$ , for  $\alpha, \beta = 1, 2, 3$ . Therefore,  $W := \langle X_1, X_2, X_3, Y_1, Y_2, Y_3 \rangle$  is a Lie algebra isomorphic to  $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$ .

A short computation shows that

$$\begin{aligned} \iota_{Y_1} \Theta_S &= d\left(\frac{1}{4v^2} da\right), & \iota_{Y_2} \Theta_S &= d\left(-\frac{a}{2v} d\left(\frac{x}{v}\right)\right), \\ \iota_{Y_3} \Theta_S &= d\left(-\frac{x}{v} dv - \frac{a}{2v} d\left(\frac{x^2}{v}\right)\right). \end{aligned}$$

Since the Vessiot–Guldberg Lie algebra for  $X^S$ , namely (4.3), consists also of Hamiltonian vector fields relative to  $\Theta_S$ , the Lie algebra  $W$  is made of Hamiltonian vector fields relative to  $\Theta_S$  and  $\mathcal{L}_Z \Theta_S = 0$  for every  $Z \in W$ .

We denote by  $G_2$  the one-parameter group of diffeomorphisms generated by the flow of the Lie symmetry  $Y_2$  and the inversion  $(x, v, a) \mapsto (-x - v, -a)$  on  $\mathcal{O}$ . Define a projection  $\pi : (x, v, a) \in \mathcal{O} \mapsto (\bar{x} = x/v, \bar{a} = a/v) \in \mathcal{O}/G_2$ , where  $\mathcal{O}/G_2$  is the space of orbits of the action of  $G_2$  on  $\mathcal{O}$ , which admits an immediate manifold structure as it is diffeomorphic to  $\mathbb{R}^2$ . Since



$Y_2$  is a Lie symmetry of  $X^S$ , the system  $X^S$  can be reduced onto  $\mathcal{O}/G_2$ . In coordinates, the projections of  $X_1, X_2, X_3$  onto  $\mathcal{O}/G_2$  read

$$\bar{X}_1 := 2 \frac{\partial}{\partial \bar{a}}, \quad \bar{X}_2 := -\bar{x} \frac{\partial}{\partial \bar{x}} + \bar{a} \frac{\partial}{\partial \bar{a}}, \quad \bar{X}_3 := (1 - \bar{x}\bar{a}) \frac{\partial}{\partial \bar{x}} + \frac{\bar{a}^2}{2} \frac{\partial}{\partial \bar{a}},$$

respectively. They form a basis of the projection,  $V_R^S$ , of the Lie algebra  $V^S = \langle X_1, X_2, X_3 \rangle$  onto  $\mathcal{O}/G_2$ . In fact,

$$[\bar{X}_1, \bar{X}_2] = \bar{X}_1, \quad [\bar{X}_1, \bar{X}_3] = 2\bar{X}_2, \quad [\bar{X}_2, \bar{X}_3] = \bar{X}_3. \tag{4.8}$$

Since  $V^S$  is simple and  $V_R^S \neq \{0\}$ , one gets  $V_R^S \simeq V^S \simeq \mathfrak{sl}_2$ .

The vector fields  $\bar{X}_1, \bar{X}_2, \bar{X}_3$  are Hamiltonian relative to the symplectic form defined by

$$\bar{\Theta} := \frac{1}{2} d\bar{x} \wedge d\bar{a}. \tag{4.9}$$

In fact,

$$\iota_{\bar{X}_1} \bar{\Theta} = -d\bar{x}, \quad \iota_{\bar{X}_2} \bar{\Theta} = -\frac{1}{2} d(\bar{x}\bar{a}), \quad \iota_{\bar{X}_3} \bar{\Theta} = \frac{1}{2} d\left(\bar{a} - \frac{\bar{x}\bar{a}^2}{2}\right).$$

Moreover,  $\bar{\Theta}$  is the only two-form in  $\mathcal{O}/G_2 \cong \mathbb{R}^2$  such that  $\pi^* \bar{\Theta} = \iota_{Y_2} \Theta_S$ .

Thus, the multisymplectic Lie system  $(\mathcal{O}, \Theta_S, X^S)$  reduces to a new multisymplectic Lie system  $(\mathcal{O}/G_2, \bar{\Theta}, X_R^S)$  with a Vessiot–Guldberg Lie algebra  $V_R^S$ , where  $X_R^S$  is the projection of  $X^S$  onto  $\mathcal{O}/G_2$ . In particular, this reduced Lie system is a Lie–Hamilton system (the multisymplectic form is a symplectic form).

To illustrate the simplification obtained by our multisymplectic reduction, let us comment on the particular reduced system  $X_R^S = \bar{X}_3 - 1/4\bar{X}_1$ , whose integral curves satisfy the system of differential equations

$$\frac{d\bar{x}}{dt} = 1 - \bar{x}\bar{a}, \quad \frac{d\bar{a}}{dt} = \frac{\bar{a}^2}{2} - \frac{1}{2}. \tag{4.10}$$

Note that this system is a Hamiltonian system relative to the symplectic form (4.9). As a symplectic Hamiltonian system on the plane, many of its properties, such as its superposition rules, have been studied elsewhere [3, 5]. Let us comment on other properties that have not been studied so far, e.g. its equilibrium points. System (4.10) has equilibrium points at two points given by

$$\bar{x} = \bar{a} = \pm 1.$$

It is notable that the system on  $\mathcal{O}$  of the form

$$\frac{dx}{dt} = v, \quad \frac{dv}{dt} = a, \quad \frac{da}{dt} = \frac{3}{2} \frac{a^2}{v} - \frac{v}{2}, \tag{4.11}$$

that projects onto (4.10) has no equilibrium point since  $v \neq 0$  on  $\mathcal{O}$ . The points in  $\mathcal{O}$  that project onto equilibrium points of (4.10) are called *relative equilibrium points* [2, 50]. Note that, although we have assumed  $b_1(t)$  to be non-constant in (4.2), we have also succinctly commented which parts of our above analysis hold true for every, possibly constant, function  $b_1(t)$ . In particular, (4.11) leads to a multisymplectic Lie system  $(\mathcal{O}, \Theta_S, X^S)$  which can be reduced, but it does not give rise to a locally automorphic Lie system when  $b_1(t)$  is a constant.

Although analytical solutions for (4.11) can be obtained using mathematical manipulation software, the solutions obtained in this way are, as far as we were able to analyse them, too complicated to be easily studied. This justifies the execution of our reduction procedure. More precisely, the solutions to (4.10) are of the form

$$\begin{aligned} \bar{x}(t) &= (2c_2 - 1)(1 + \cosh(t + 2c_1)) + \sinh(t + 2c_1), \\ \bar{a}(t) &= \frac{1 - e^{2c_1+t}}{e^{2c_1+t} + 1}, \quad c_1, c_2 \in \mathbb{R}, \end{aligned}$$

for solutions with  $|\bar{a}(t)| < 1$  and

$$\begin{aligned} \bar{x}(t) &= -1 - 2c_2 + e^{t+2c_1}c_2 + e^{-t-2c_1}(1 + c_2), \\ \bar{a}(t) &= \frac{1 + e^{2c_1+t}}{1 - e^{2c_1+t}}, \quad c_1, c_2 \in \mathbb{R}, \end{aligned}$$

for solutions for  $|\bar{a}(t)| > 1$ . Meanwhile, the solutions for  $\bar{a}(t) = \pm 1$  read

$$\bar{x}(t) = \pm 1 + e^{\mp t}c_2, \tag{4.12}$$

respectively.

It can be established whether two Lie systems in the plane related to a Vessiot–Guldberg Lie algebra isomorphic to  $\mathfrak{sl}_2$ , like (4.10), are locally diffeomorphic via the so-called *Casimir tensor fields* (see [5] for details). In particular, it was proved in [5, p 5] that the Casimir tensor field for a Vessiot–Guldberg Lie algebra on the plane isomorphic to  $\mathfrak{sl}_2$  in a basis with structure constants (4.8) is given by

$$\begin{aligned} G &= X_1 \otimes X_3 + X_3 \otimes X_1 - 2X_2 \otimes X_2 \\ &= -2\bar{x}^2 \frac{\partial}{\partial \bar{x}} \otimes \frac{\partial}{\partial \bar{x}} + 2\frac{\partial}{\partial \bar{x}} \otimes \frac{\partial}{\partial \bar{a}} + 2\frac{\partial}{\partial \bar{a}} \otimes \frac{\partial}{\partial \bar{x}}. \end{aligned}$$

It was also shown in [5, lemma 4.1] that the determinant of the coefficients of  $G$ , namely  $\det G = -4$ , allows us to determine that there exists a local diffeomorphism on  $\mathbb{R}^2$  mapping (4.10) onto a Lie system on the plane related to a Vessiot–Guldberg Lie algebra


$$\left\langle \frac{\partial}{\partial x} + \frac{\partial}{\partial y}, x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y}, x^2 \frac{\partial}{\partial x} + y^2 \frac{\partial}{\partial y} \right\rangle,$$

the so-called  $I_4$ . It was proved in [46, table 1], that this Lie algebra is a Lie algebra of Killing vector fields relative to a pseudo-Riemannian metric on  $\mathbb{R}^2$ . Physically, it is also notable that (4.10) is locally diffeomorphic to a Milne–Pinney equation [5, lemma 4.1 and proposition 6.1], which appears in many physical problems (see [20, 21, 48] and references therein).

It is possible to see in figure 1 that equilibrium points of (4.10) are unstable (solutions close to them move away from them as time passes by).

There is a last remark about (4.10) to be considered. Since it is a Hamiltonian system, its evolution conserves the area of any set of solutions. Despite this, system (4.10) is still unstable around its equilibrium points.

**4.1.2. Dissipative quantum harmonic oscillators.** Let us define  $G^{\text{do}} := \text{SL}_2 \ltimes \text{H}_3$ , where  $\text{SL}_2$  is the Lie group of unimodular real  $2 \times 2$  matrices with real entries,  $\text{H}_3$  is the Heisenberg group of upper real triangular unimodular  $3 \times 3$  matrices with ones in the diagonal, and  $\ltimes$  represents the semidirect product of  $\text{SL}_2$  acting on the normal subgroup  $\text{H}_3$  of  $G^{\text{do}}$  via a Lie group morphism



**Figure 1.** The first three diagrams represent some integral curves of the system (4.11) on the three-dimensional manifold  $\mathcal{O}$  from different points of view, illustrating the fact that it has no equilibrium points, as already commented. The figure on the second line depicts the integral curves of the reduced multisymplectic system (4.10) close to the equilibrium points  $(1, 1)$  and  $(-1, -1)$ .

$\Phi : A \in \text{SL}_2 \mapsto \Phi_A \in \text{Aut}(\text{H}_3)$ , where  $\text{Aut}(\text{H}_3)$  is the space of Lie group automorphisms on  $\text{H}_3$ . Note that the Lie group product in  $G^{\text{do}}$  can be written as

$$(A, B)(A', B') = (AA', B\Phi_A(B')), \quad A, A' \in \text{SL}_2, \quad B, B' \in \text{H}_3.$$

Note that the elements of  $\text{SL}_2$  in  $\text{SL}_2 \times \text{H}_3$  can be then written as  $(A, I_3)$ , where  $A \in \text{SL}_2$  and  $I_3$  is the neutral element of  $\text{H}_3$ . Meanwhile, the elements of  $\text{H}_3$  in  $\text{SL}_2 \times \text{H}_3$  can be brought into the form  $(I_2, B)$  for some  $B \in \text{H}_3$  and the neutral element  $I_2$  in  $\text{SL}_2$ . Then,

$$(A, I_3)(I_2, B)(A^{-1}, I_3) = (A, \Phi_A(B))(A^{-1}, I_3) = (I_2, \Phi_A(B)), \quad A \in \text{SL}_2, \quad B \in \text{H}_3.$$

Hence,  $\Phi_A(B)$  can be considered as the inner automorphism induced by  $A$  acting on  $B$  within  $\text{SL}_2 \times \text{H}_3$ . It is standard to consider  $G^{\text{do}}$  as a matrix Lie group, at least locally around its neutral element. In this case, the elements of  $\text{SL}_2$  and  $\text{H}_3$  can be considered as subgroups of matrices in a certain  $\text{GL}(n, \mathbb{R})$  and one can write  $\Phi_A(B) = ABA^{-1}$  for every  $A \in \text{SL}_2$  and  $B \in \text{H}_3$ . The Lie algebra  $\mathfrak{g}^{\text{do}}$  of  $G^{\text{do}}$  satisfies that  $\mathfrak{g}^{\text{do}} \simeq \mathfrak{sl}_2 \oplus_{\mathcal{S}} \mathfrak{h}_3$ , where  $\oplus_{\mathcal{S}}$  represents a semi-direct sum of the Lie algebra  $\mathfrak{sl}_2 \simeq \text{T}_e\text{SL}_2$  with the ideal  $\mathfrak{h}_3 \simeq \text{T}_e\text{H}_3$  of  $\text{T}_eG^{\text{do}}$ .

Let us consider the automorphic Lie system on  $G^{\text{do}}$  of the form

$$\frac{dg}{dt} = \sum_{\alpha=1}^6 b_{\alpha}(t)X_{\alpha}^R(g), \quad g \in G^{\text{do}}, \quad t \in \mathbb{R}, \tag{4.13}$$

where  $b_1(t), \dots, b_6(t)$  are arbitrary  $t$ -dependent functions, and  $\{X_1^R, \dots, X_6^R\}$  form a basis of the Lie algebra  $V^{\text{do}}$  of right-invariant vector fields on  $G^{\text{do}}$  so that  $\{X_1^R, X_2^R, X_3^R\}$  is a basis of the Lie algebra  $V_{\mathfrak{sl}} := \langle X_1^R, X_2^R, X_3^R \rangle \simeq \mathfrak{sl}_2$  and  $\{X_4^R, X_5^R, X_6^R\}$  stands for a basis of the ideal  $V_h := \langle X_4^R, X_5^R, X_6^R \rangle \simeq \mathfrak{h}_3$  of  $V^{\text{do}}$ . Geometrically, system (4.13) amounts to the  $t$ -dependent vector field on  $G^{\text{do}}$  given by

$$X^{\text{do}}(t, g) := \sum_{\alpha=1}^6 b_{\alpha}(t)X_{\alpha}^R(g), \quad g \in G^{\text{do}}, \quad t \in \mathbb{R}. \tag{4.14}$$

Since the right-hand side of (4.14) is a  $t$ -dependent vector field  $X^{\text{do}}$  on  $G^{\text{do}}$  taking values in the finite-dimensional Lie algebra  $V^{\text{do}}$ , the system  $X^{\text{do}}$  is a Lie system.

The relevance of system (4.13) is due to the fact that it appears in the study of quantum harmonic oscillators with dissipation (see [21, equation (3.9)] for details). More specifically, they appear in methods to solve  $t$ -dependent Hamilton operators of the form

$$\widehat{H}(t) = \alpha(t)\frac{\widehat{x}^2}{2} + \beta(t)\frac{\widehat{p}^2}{2} + \gamma(t)\frac{\widehat{x}\widehat{p} + \widehat{p}\widehat{x}}{4} + \delta(t)\widehat{p} + \epsilon(t)\widehat{x} + \phi(t)\widehat{\text{Id}}, \tag{4.15}$$

where  $\alpha(t), \beta(t), \gamma(t), \delta(t), \epsilon(t), \phi(t)$  are arbitrary  $t$ -dependent functions, while  $\widehat{x}, \widehat{p}, \widehat{\text{Id}}$ , are the position, moment, and identity operators on the space of complex square-integrable functions on  $\mathbb{R}$ , respectively. It is evident that (4.15) has many relevant applications in many quantum mechanical problems.

Although the relevant properties of  $X^{\text{do}}$  are better understood using a geometric approach, a representation in coordinates of  $X^{\text{do}}$  will also be useful to describe some of its features. More specifically, let us use the canonical coordinates of the second kind  $\{v_1, \dots, v_6\}$  on  $G^{\text{do}}$  induced by (see [21] for details)

$$\begin{aligned} \phi : \sum_{i=1}^6 v_i a^i \in \mathfrak{g} &\mapsto \exp(-v_4 a^4) \exp(-v_5 a^5) \exp(-v_6 a^6) \exp(-v_1 a^1) \\ &\times \exp(-v_2 a^2) \exp(-v_3 a^3) \in G^{\text{do}}, \end{aligned} \tag{4.16}$$

where  $\{a^1, \dots, a^6\}$  is a basis of  $T_e G^{\text{do}} \simeq \mathfrak{g}^{\text{do}}$  spanning the commutation relations given in [21, p 70] and  $X_i^R(e) = -a^i$  for  $i= 1, \dots, 6$ . This allows one to write system (4.13) in local coordinates as

$$\begin{cases} \frac{dv_1}{dt} = b_1(t) + b_2(t)v_1 + b_3(t)v_1^2, & \frac{dv_4}{dt} = b_4(t) + \frac{1}{2}b_2(t)v_4 + b_1(t)v_5, \\ \frac{dv_2}{dt} = b_2(t) + 2b_3(t)v_1, & \frac{dv_5}{dt} = b_5(t) - b_3(t)v_4 - \frac{1}{2}b_2(t)v_5, \\ \frac{dv_3}{dt} = e^{v_2}b_3(t), & \frac{dv_6}{dt} = b_6(t) - b_5(t)v_4 + \frac{1}{2}b_3(t)v_4^2 - \frac{1}{2}b_1(t)v_5^2. \end{cases} \tag{4.17}$$

Comparing the system  $X^{\text{do}}$  in intrinsic form (4.13) with the above, we obtain

$$\begin{aligned} X_1^{\text{R}} &= \frac{\partial}{\partial v_1} + v_5 \frac{\partial}{\partial v_4} - \frac{1}{2} v_5^2 \frac{\partial}{\partial v_6}, & X_2^{\text{R}} &= v_1 \frac{\partial}{\partial v_1} + \frac{\partial}{\partial v_2} + \frac{1}{2} v_4 \frac{\partial}{\partial v_4} - \frac{1}{2} v_5 \frac{\partial}{\partial v_5}, \\ X_3^{\text{R}} &= v_1^2 \frac{\partial}{\partial v_1} + 2v_1 \frac{\partial}{\partial v_2} + e^{v_2} \frac{\partial}{\partial v_3} - v_4 \frac{\partial}{\partial v_5} + \frac{1}{2} v_4^2 \frac{\partial}{\partial v_6}, & X_4^{\text{R}} &= \frac{\partial}{\partial v_4}, \\ X_5^{\text{R}} &= \frac{\partial}{\partial v_5} - v_4 \frac{\partial}{\partial v_6}, & X_6^{\text{R}} &= \frac{\partial}{\partial v_6}. \end{aligned} \tag{4.18}$$

The commutation relations between the above vector fields read

$$\begin{aligned} [X_1^{\text{R}}, X_2^{\text{R}}] &= X_1^{\text{R}}, \\ [X_1^{\text{R}}, X_3^{\text{R}}] &= 2X_2^{\text{R}}, \quad [X_2^{\text{R}}, X_3^{\text{R}}] = X_3^{\text{R}}, \\ [X_1^{\text{R}}, X_4^{\text{R}}] &= 0, \quad [X_2^{\text{R}}, X_4^{\text{R}}] = -\frac{1}{2} X_4^{\text{R}}, \quad [X_3^{\text{R}}, X_4^{\text{R}}] = X_5^{\text{R}}, \\ [X_1^{\text{R}}, X_5^{\text{R}}] &= -X_4^{\text{R}}, \quad [X_2^{\text{R}}, X_5^{\text{R}}] = \frac{1}{2} X_5^{\text{R}}, \quad [X_3^{\text{R}}, X_5^{\text{R}}] = 0, \quad [X_4^{\text{R}}, X_5^{\text{R}}] = -X_6^{\text{R}}, \\ [X_1^{\text{R}}, X_6^{\text{R}}] &= 0, \quad [X_2^{\text{R}}, X_6^{\text{R}}] = 0, \quad [X_3^{\text{R}}, X_6^{\text{R}}] = 0, \quad [X_4^{\text{R}}, X_6^{\text{R}}] = 0, \quad [X_5^{\text{R}}, X_6^{\text{R}}] = 0. \end{aligned} \tag{4.19}$$

Therefore, the vector fields given in (4.18) span a six-dimensional real Lie algebra  $V^{\text{do}}$  isomorphic to the semidirect sum of Lie algebras  $\mathfrak{sl}_2 \oplus \mathfrak{sh}_3$ .

Following the method described in [33], one can construct a multisymplectic form on  $G^{\text{do}}$  turning the elements of  $V^{\text{do}}$  into locally Hamiltonian vector fields. Since  $(X_1^{\text{R}} \wedge \dots \wedge X_6^{\text{R}})(g) \neq 0$  on each  $g \in G^{\text{do}}$ , the vector fields  $X_1^{\text{R}}, \dots, X_6^{\text{R}}$  admit a family of differential one-forms  $\eta_1^{\text{R}}, \dots, \eta_6^{\text{R}}$  on  $G^{\text{do}}$  such that  $\eta_\alpha^{\text{R}}(X_\beta^{\text{R}}) = \delta_{\alpha\beta}$  for  $\alpha, \beta = 1, \dots, 6$ , where  $\delta_{\alpha\beta}$  is the Kronecker delta. More specifically, the local expressions of these differential one-forms are

$$\begin{aligned} \eta_1^{\text{R}} &= dv_1 - v_1 dv_2 + v_1^2 e^{-v_2} dv_3, & \eta_2^{\text{R}} &= dv_2 - 2v_1 e^{-v_2} dv_3, & \eta_3^{\text{R}} &= e^{-v_2} dv_3, \\ \eta_4^{\text{R}} &= -v_5 dv_1 + \left( v_1 v_5 - \frac{1}{2} v_4 + \frac{1}{2} v_4 v_5 \right) dv_2 \\ &\quad + (v_1 v_4 - v_1^2 v_5 - v_1 v_4 v_5 + v_4^2) e^{-v_2} dv_3 + dv_4 + v_4 dv_5, \\ \eta_5^{\text{R}} &= \frac{1}{2} v_5 dv_2 + (v_4 - v_1 v_5) e^{-v_2} dv_3 + dv_5, \\ \eta_6^{\text{R}} &= \frac{1}{2} v_5 dv_1 - \frac{1}{2} v_1 v_5 dv_2 + \frac{1}{2} (v_1^2 v_5 - v_4^2) e^{-v_2} dv_3 + dv_6. \end{aligned}$$

As  $\eta_1^{\text{R}}, \dots, \eta_6^{\text{R}}$  are right-invariant differential forms, it makes sense to denote  $V^{\text{do}*} := \langle \eta_1^{\text{R}}, \dots, \eta_6^{\text{R}} \rangle$ . This enables us to define the volume form

$$\Theta^{\text{do}} := \eta_1^{\text{R}} \wedge \dots \wedge \eta_6^{\text{R}} = e^{-v_2} dv_1 \wedge \dots \wedge dv_6. \tag{4.20}$$

Since  $\Theta^{\text{do}}$  is a volume form, it is clearly a multisymplectic one.

Let us prove that the vector fields  $X_1^{\text{R}}, \dots, X_6^{\text{R}}$  are locally Hamiltonian relative to  $\Theta^{\text{do}}$ . This is indeed a consequence of the geometric construction of  $\Theta^{\text{do}}$  and the Lie algebra structure of  $\mathfrak{g}^{\text{do}}$ . We have that

$$\begin{aligned}
 (\mathcal{L}_{X^R} \eta^R)(\bar{X}^R) &= \mathcal{L}_{X^R}(\iota_{\bar{X}^R} \eta^R) - \eta^R([X^R, \bar{X}^R]) = -\eta^R([X^R, \bar{X}^R]), \\
 &\forall X^R, \bar{X}^R \in V^{\text{do}}, \forall \eta^R \in V^{\text{do}*}.
 \end{aligned}$$

Since all the  $\bar{X}^R$  span the tangent bundle to  $G^{\text{do}}$ , it follows that

$$\mathcal{L}_{X^R} \eta^R = -\text{ad}_{X^R}^* \eta^R, \quad \forall X^R \in V^{\text{do}}, \forall \eta^R \in V^{\text{do}*}.$$

The adjoint representation  $\text{ad} : X^R \in V^{\text{do}} \mapsto \text{ad}_{X^R} := [X^R, \cdot] \in \text{End}(V^{\text{do}})$  is such that  $\text{ad}(V^{\text{do}})$  consists of traceless operators on  $V^{\text{do}}$ . Hence, the adjoint representation of  $V^{\text{do}}$  is traceless and  $V^{\text{do}}$  is a unimodular Lie algebra. Then,  $V^{\text{do}}$  acts on  $V^{\text{do}*}$  through the coadjoint Lie algebra representation  $\text{coad} : X^R \in V^{\text{do}*} \mapsto -\text{ad}_{X^R}^* \in \text{End}(V^{\text{do}*})$ . Since  $\text{ad}$  is traceless, the representation  $\text{coad}$  is also traceless. Thus,

$$\mathcal{L}_{X^R} \Theta^{\text{do}} = -\text{Tr}(\text{ad}_{X^R}^*) \Theta^{\text{do}} = 0, \quad \forall X^R \in V^{\text{do}}.$$

As a consequence, the elements of  $V^{\text{do}}$  are locally Hamiltonian vector fields with respect to the multisymplectic volume form  $\Theta^{\text{do}}$ . The above construction gives rise to a Lie system on  $G^{\text{do}}$  admitting a Vessiot–Guldberg Lie algebra of locally Hamiltonian vector fields  $X_1^R, \dots, X_6^R$  relative to the multisymplectic form  $\Theta^{\text{do}}$ , that is  $(G^{\text{do}} = \text{SL}_2 \ltimes \text{H}_3, \Theta^{\text{do}}, X^{\text{do}})$  is a multisymplectic Lie system.

The multisymplectic form,  $\Theta^{\text{do}}$ , given in (4.20), and the elements of  $V^{\text{do}}$ , detailed in (4.18), are right-invariant. Hence, for every left-invariant vector field  $Y^L$  on  $G^{\text{do}}$ , one has that  $\mathcal{L}_{Y^L} \Theta^{\text{do}} = 0$  and  $\mathcal{L}_{Y^L} X^{\text{do}} = 0$ . Then,  $Y^L$  is a Lie symmetry of  $V^{\text{do}}$ , that is,  $Y^L \in \text{Sym}(V^{\text{do}})$ . Lemma 3.7 yields that  $V_L^{\text{do}} := \text{Sym}(V^{\text{do}})$  is isomorphic to  $V^{\text{do}}$ . As shown next, this will lead to the construction of multisymplectic reductions of the multisymplectic Lie system  $(G^{\text{do}}, \Theta^{\text{do}}, X^{\text{do}})$  by unimodular Lie subalgebras of  $V_L^{\text{do}} \simeq \mathfrak{sl}_2 \oplus \mathfrak{sh}_3$ .

Let  $\{X_1^L, \dots, X_6^L\}$  be a basis of left-invariant vector fields on  $G^{\text{do}}$  satisfying that

$$X_\alpha^L(e) = X_\alpha^R(e), \quad \alpha = 1, \dots, 6. \tag{4.21}$$

A tedious calculation yields the local coordinate expression of these vector fields:

$$\begin{aligned}
 X_1^L &= e^{v_2} \frac{\partial}{\partial v_1} + 2v_3 \frac{\partial}{\partial v_2} + v_3^2 \frac{\partial}{\partial v_3}, & X_2^L &= \frac{\partial}{\partial v_2} + v_3 \frac{\partial}{\partial v_3}, & X_3^L &= \frac{\partial}{\partial v_3}, \\
 X_4^L &= e^{-v_2/2} (e^{v_2} - v_1 v_3) \frac{\partial}{\partial v_4} - e^{-v_2/2} v_3 \frac{\partial}{\partial v_5} - e^{-v_2/2} (e^{v_2} - v_1 v_3) v_5 \frac{\partial}{\partial v_6}, & & & & \\
 X_5^L &= v_1 e^{-v_2/2} \frac{\partial}{\partial v_4} + e^{-v_2/2} \frac{\partial}{\partial v_5} - v_1 v_5 e^{-v_2/2} \frac{\partial}{\partial v_6}, & X_6^L &= \frac{\partial}{\partial v_6}.
 \end{aligned} \tag{4.22}$$

Note that  $V_L^{\text{do}} = \langle X_1^L, \dots, X_6^L \rangle$ . The above properties will be useful in the following results.

As an example of reduction for the multisymplectic Lie system  $(G^{\text{do}}, \Theta^{\text{do}}, X^{\text{do}})$ , we consider the unimodular Lie subalgebra  $V_{\mathfrak{sl}_2}^L := \langle X_1^L, X_2^L, X_3^L \rangle \simeq \mathfrak{sl}_2$  and an associated Noether invariant  $\Theta_d^{\text{sl}} := \iota_{X_1^L \wedge X_2^L \wedge X_3^L} \Theta^{\text{do}}$  of the Lie system  $X^{\text{do}}$ . The idea is to obtain a new multisymplectic Lie system  $(\tilde{G}, \tilde{\Theta}_d^{\text{sl}}, \tilde{X}_d^{\text{sl}})$  such that the following conditions hold:

- (a) If  $G^{\text{do}}/\text{SL}_2$  stands for the manifold of left cosets relative to the subgroup  $\text{SL}_2$  of  $G^{\text{do}}$ , then the canonical projection  $\pi^{\text{sl}} : G^{\text{do}} \rightarrow \tilde{G} := G^{\text{do}}/\text{SL}_2$  is a submersion and  $\tilde{G}$  becomes a manifold, which does not need to have a Lie group structure.
- (b) The vector fields of the Vessiot–Guldberg Lie algebra  $V^{\text{do}}$  project onto a Lie algebra  $\tilde{V}_d^{\text{sl}} := (\pi^{\text{sl}})_*(V^{\text{do}})$  of vector fields on  $\tilde{G}$ .



(c) Finally,  $(\pi^{sl})^*\tilde{\Theta}_d^{sl} = \Theta_d^{sl} = \iota_{X_1^L \wedge X_2^L \wedge X_3^L} \Theta^{do}$ .

The action of  $SL_2$  on  $G^{do}$  on the right has fundamental vector fields  $\langle X_1^L, X_2^L, X_3^L \rangle$ . The orbits of this action are the left cosets of  $SL_2$  in  $G^{do}$ . Furthermore, the quotient space  $G^{do}/SL_2$  is diffeomorphic to  $H_3$ . Thus, the first condition is satisfied.

Let us verify the second condition. Since the elements of  $V^{do}$  commute with left-invariant vector fields on  $G^{do}$ , they are all  $\pi^{sl}$ -projectable vector fields onto  $G^{do}/SL_2$ . As  $V^{do} = \langle X_1^R, \dots, X_6^R \rangle$  is a Lie algebra whose distribution,  $\mathcal{D}^{V^{do}}$ , is such that  $\mathcal{D}_g^{V^{do}} = T_g G^{do}$  for every  $g \in G^{do}$ , the projections of  $X_1^R, \dots, X_6^R$  onto  $G^{do}/SL_2 \simeq H_3$  span  $T(G^{do}/SL_2)$ . Hence, the initial system  $X^{do}$  can be projected onto  $G^{do}/SL_2$  giving rise to a Lie system on  $G^{do}/SL_2$  with a Vessiot–Guldberg Lie algebra  $\tilde{V}_d^{sl} = (\pi^{sl})_*(V^{do})$ . The morphism  $(\pi^{sl})_*|_{V^{do}} : V^{do} \rightarrow \tilde{V}_d^{sl}$  is a surjective Lie algebra morphism.

Note that  $v_1, \dots, v_6$  form a canonical coordinate system of the second kind defined locally in  $G^{do}$  whilst, in view of our parametrisation (4.16), the  $v_4, v_5, v_6$  are, indeed global, coordinates in  $G^{do}/SL_2$ . In coordinates  $v_1, \dots, v_6$ , we have the canonical projection  $\pi^{sl} : (v_1, \dots, v_6) \in G^{do} \mapsto (v_4, v_5, v_6) \in G^{do}/SL_2 \simeq H_3$ . The projections of the elements of the basis (4.18) of the Vessiot–Guldberg Lie algebra  $V^{do}$  of  $X^{do}$  onto  $G^{do}/SL_2$  read

$$\begin{aligned} \tilde{X}_1^{sl} &= v_5 \frac{\partial}{\partial v_4} - \frac{1}{2} v_5^2 \frac{\partial}{\partial v_6}, & \tilde{X}_2^{sl} &= \frac{1}{2} v_4 \frac{\partial}{\partial v_4} - \frac{1}{2} v_5 \frac{\partial}{\partial v_5}, & \tilde{X}_3^{sl} &= -v_4 \frac{\partial}{\partial v_5} + \frac{1}{2} v_4^2 \frac{\partial}{\partial v_6}, \\ \tilde{X}_4^{sl} &= \frac{\partial}{\partial v_4}, & \tilde{X}_5^{sl} &= \frac{\partial}{\partial v_5} - v_4 \frac{\partial}{\partial v_6}, & \tilde{X}_6^{sl} &= \frac{\partial}{\partial v_6}. \end{aligned} \tag{4.23}$$

A simple calculation shows that

$$[\tilde{X}_\alpha^{sl}, \tilde{X}_\beta^{sl}] = c_{\alpha\beta}^\gamma \tilde{X}_\gamma^{sl}, \quad \alpha, \beta, \gamma = 1, \dots, 6,$$

where the  $c_{\alpha\beta}^\gamma$  are again the same constants given in (4.19) for  $V^{do}$ . This shows that  $\tilde{V}_d^{sl} \simeq V^{do} \simeq \mathfrak{g}^{do}$ . This gives rise to a Lie system  $X_{sl}^{G^{do}} := \sum_{\alpha=1}^6 b_\alpha(t) \tilde{X}_\alpha^{sl}$  on the quotient space  $G^{do}/SL_2$ .

Finally, let us prove that there exists a multisymplectic form  $\tilde{\Theta}_d^{sl}$  on  $G^{do}/SL_2$  turning the elements of  $\tilde{V}_d^{sl}$  into locally Hamiltonian vector fields. Since  $V_{sl}^L$  is a Lie subalgebra, it makes sense to define the restriction  $\text{ad}_{X_\alpha^L}|_{V_{sl}^L}$  of each  $\text{ad}_{X_\alpha^L} : V_L^{do} \rightarrow V_L^{do}$ , with  $\alpha = 1, \dots, 6$ , to  $V_{sl}^L$ . In view of the commutation relations (4.19) of the vector fields of  $V^{do}$ , it follows that  $\text{Tr}(\text{ad}_{X_\alpha^L}|_{V_{sl}^L}) = 0$ . Therefore,

$$\mathcal{L}_{X_\alpha^L} \left( \iota_{X_1^L \wedge X_2^L \wedge X_3^L} \Theta^{do} \right) = \text{Tr} \left( \text{ad}_{X_\alpha^L}|_{V_{sl}^L} \right) \iota_{X_1^L \wedge X_2^L \wedge X_3^L} \Theta^{do} = 0, \quad \alpha = 1, 2, 3. \tag{4.24}$$

Moreover,  $\Theta_d^{sl} := \iota_{X_1^L \wedge X_2^L \wedge X_3^L} \Theta^{do}$  satisfies

$$(\ker \Theta_d^{sl})(g) = \langle X_1^L(g), X_2^L(g), X_3^L(g) \rangle, \quad \forall g \in G^{do}. \tag{4.25}$$

Expressions (4.24) and (4.25) ensure that there exists a unique well-defined differential form  $\tilde{\Theta}_d^{sl}$  on  $G^{do}/SL_2$  such that  $(\pi^{sl})^*\tilde{\Theta}_d^{sl} = \Theta_d^{sl}$  relative to the canonical projection  $\pi^{sl} : G^{do} \rightarrow G^{do}/SL_2$ . As the vector fields of  $\tilde{V}_d^{sl}$  span the tangent bundle to  $G^{do}/SL_2$  and  $\tilde{\Theta}_d^{sl}$  is invariant under the action by Lie derivatives of the elements of  $\tilde{V}_d^{sl}$  on it, then  $\tilde{\Theta}_d^{sl}$  is the only multisymplectic form, up to a non-zero proportionality constant, turning the vector fields of  $\tilde{V}_d^{sl}$  into locally Hamiltonian vector fields relative to  $\tilde{\Theta}_d^{sl}$ . Since  $\Theta^{do}$  is a volume form on a six-dimensional manifold, then  $\tilde{\Theta}_d^{sl}$  is a differential three-form. Therefore,  $\tilde{\Theta}_d^{sl}$  is a volume form

on  $G^{\text{do}}/\text{SL}_2$  and  $d\tilde{\Theta}_d^{sl} = 0$ . In view of (4.25), it follows that  $\tilde{\Theta}_d^{sl}$  is nondegenerate. Hence,  $(G^{\text{do}}/\text{SL}_2, \tilde{\Theta}_d^{sl})$  is a multisymplectic manifold.

The differential form  $\tilde{\Theta}_d^{sl}$  can be explicitly derived by taking into account that it is, up to a non-zero constant, invariant by the above projected vector fields (4.23) spanning the Lie algebra  $\tilde{V}_d^{sl}$ , namely  $\tilde{\Theta}_d^{sl} = \lambda dv_4 \wedge dv_5 \wedge dv_6$  for a certain  $\lambda \in \mathbb{R} \setminus \{0\}$  and  $\iota_{X_4^L \wedge X_5^L \wedge X_6^L}(\pi^{sl})^* \tilde{\Theta}_d^{sl} = 1$  when  $v_4 = v_5 = v_6 = 0$ . Hence,

$$\tilde{\Theta}_d^{sl} = dv_4 \wedge dv_5 \wedge dv_6.$$

It is worth noting that the multisymplectic reduction process has been carried out assuming  $V_{sl}^L$  to be unimodular.

#### 4.2. Multisymplectic reduction of multisymplectic Lie systems

After studying the above examples, the following question arises: Can the above process be repeated for other multisymplectic Lie subalgebras of  $V^{\text{do}}$ ? To answer this question, we define the momentum map

$$\begin{aligned} \mathcal{J}: \quad G \times \mathcal{G}(\mathfrak{g}) &\longrightarrow \mathring{\wedge} T^*G \\ (g, X_1^L \wedge \dots \wedge X_p^L) &\longmapsto (\iota_{X_1^L \wedge \dots \wedge X_p^L} \Theta)(g), \end{aligned}$$

where  $\mathcal{J}(g, X_1^L \wedge \dots \wedge X_p^L) \in \mathring{\wedge} T_g^*G$ , the set  $\mathcal{G}(\mathfrak{g})$  is the unimodular Grassmannian defined in definition 2.13 and  $\Theta$  is a multisymplectic form on  $G$ .

In the example in section 4.1.1, we considered the space  $\mathcal{J}(g, Y_2)$ , while  $\mathcal{J}(g, X_1^L \wedge X_2^L \wedge X_3^L)$  was used in the multisymplectic reduction of the example in section 4.1.2. In both cases,  $\mathfrak{g}$  is the abstract Lie algebra isomorphic to the Lie algebra of Lie symmetries  $\text{Sym}(V^X)$ , where  $V^X$  is the Vessiot–Guldberg Lie algebra of the multisymplectic Lie system. Furthermore, it can be shown that in both cases

$$\mathcal{L}_X \mathcal{J} = 0, \quad \forall X \in V^X.$$

This section aims to use the examples of reduction of multisymplectic Lie systems already studied to provide a general multisymplectic reduction procedure for multisymplectic Lie systems.

**4.2.1. Multisymplectic reduction.** Let  $(N, \Theta, X)$  be a multisymplectic Lie system, where  $(N, \Theta)$  is a multisymplectic manifold and  $X$  denotes a Lie system on  $N$ . Let  $V^X$  be a Vessiot–Guldberg Lie algebra of  $X$  and denote by  $\text{Sym}(V^X)$  the finite-dimensional Lie algebra of Lie symmetries of a Lie system  $(N, X, V^X)$ . Consider  $\mathfrak{g}$  to be the abstract Lie algebra isomorphic to  $\text{Sym}(V^X)$ .

To obtain a general reduction procedure for multisymplectic Lie systems, one considers the map

$$\begin{aligned} \mathcal{J}: \quad N \times \mathcal{G}(\mathfrak{g}) &\longrightarrow \mathring{\wedge} T^*N \\ (x, w) &\longmapsto (\iota_w \Theta)(x), \end{aligned}$$

where  $(\iota_w \Theta)(x) \in \mathring{\wedge} T_x^*N$  for every  $x \in N$  and  $w \in \mathcal{G}(\mathfrak{g})$ .

Let  $V$  be an  $r$ -dimensional unimodular Lie subalgebra of  $\text{Sym}(V^X)$  and let  $\mathcal{D}^V$  be the distribution generated by the vector fields of  $V$ . We consider the reduction of the original system by  $V$ . The reduced system is a multisymplectic Lie system if:

- (a) The quotient space  $N/\mathcal{D}^V$  is a manifold and the canonical projection  $\pi : N \rightarrow N/\mathcal{D}^V$  is a submersion.
- (b) The elements of a basis of  $V$  are  $\pi$ -projectable.
- (c) There exists a multisymplectic form  $\tilde{\Theta}$  on  $N/\mathcal{D}^V$  such that  $(\pi^*\tilde{\Theta})(x) = \mathcal{J}(x, X_1 \wedge \cdots \wedge X_r)$ , for every  $x \in N$  and where  $X_1 \wedge \cdots \wedge X_r$  is the unique (up to an irrelevant multiplicative non-zero constant)  $r$ -vector field generated by the elements of  $V$ .

The idea is to give conditions in terms of  $\Theta$ ,  $V$ , and  $\mathcal{J}$  so that these three properties are fulfilled. The examples in the previous section can be considered as an application of the following reduction procedure.

**Theorem 4.1.** *Let  $\Theta$  be an  $(r + 1)$ -nondegenerate multisymplectic form on  $N$  and let  $G$  be an  $r$ -dimensional unimodular Lie group. We consider a proper and free unimodular Lie group action  $\Phi : G \times N \rightarrow N$  and denote by  $V_\Phi$  the Lie algebra of fundamental vector fields of  $\Phi$ . If  $w \in \Lambda^r V_\Phi \setminus \{0\}$  is a Hamiltonian  $r$ -vector field with respect to the multisymplectic form  $\Theta$ , then there exists a unique multisymplectic form  $\tilde{\Theta}$  on  $N/G$  given by*

$$\pi^*\tilde{\Theta} = \iota_w \Theta,$$

where  $\pi : N \rightarrow N/G$  is the quotient projection of  $G$  onto  $N/G$ .

**Proof.** The differential form  $\iota_w \Theta$  can be projected onto  $N/G$  if its contraction with the elements of  $V_\Phi$  vanishes and  $\iota_w \Theta$  is invariant relative to the elements of  $V_\Phi$ . The first condition is ensured by the fact that  $w \in \Lambda^r V_\Phi$  and therefore  $\iota_X \iota_w \Theta = \iota_{w \wedge X} \Theta = 0$  for all  $X \in V_\Phi$  because  $X \wedge w = 0$ . The second condition follows from the following considerations. By assumption, the vector fields  $X_1, \dots, X_r$  are  $\Theta$ -Hamiltonian, i.e.  $\mathcal{L}_X \Theta = 0$  for every  $X \in V_\Phi$ , and  $V_\Phi$  is a unimodular Lie algebra of vector fields, namely  $\mathcal{L}_X w = 0$  for all  $X \in V_\Phi$ . This implies that

$$\mathcal{L}_X \iota_w \Theta = (-1)^r \iota_w \mathcal{L}_X \Theta + \iota_{\mathcal{L}_X w} \Theta = 0, \quad \forall X \in V_\Phi.$$

In consequence,  $\mathcal{L}_X \Theta = 0$  and  $\iota_X \iota_w \Theta = 0$  for all  $X \in V_\Phi$ , which implies that there exists a differential form  $\tilde{\Theta}$  on  $N/G$  such that  $\pi^*\tilde{\Theta} = \iota_w \Theta$ . The differential form  $\tilde{\Theta}$  is also unique because  $\pi^* : \Omega^\bullet(N/G) \rightarrow \Omega^\bullet(N)$  is an injection.

Moreover,  $\tilde{\Theta}$  is closed since the assumption that  $w$  is Hamiltonian enables us to write that

$$0 = \mathcal{L}_w \Theta = (d\iota_w + (-1)^r \iota_w d)\Theta = d\iota_w \Theta = d\pi^*\tilde{\Theta} = \pi^*d\tilde{\Theta} \implies d\tilde{\Theta} = 0.$$

Since  $\Theta$  is  $(r + 1)$ -nondegenerate and  $\Phi$  is effective, then  $w \neq 0$  and the form  $\iota_w \Theta$  is one-nondegenerate. Hence,  $\tilde{\Theta}$  is also one-nondegenerate. This turns  $(N/G, \tilde{\Theta})$  into a multisymplectic manifold.  $\square$

**Definition 4.2.** A pair  $(\Theta, \Phi)$  satisfying the conditions of the theorem 4.1 is called a *multisymplectic reduction scheme*.

Once a reduction scheme is provided, it is mandatory to show how this can be applied to the reduction of a multisymplectic Lie system.

**Theorem 4.3.** *Let  $(N, \Theta, X)$  be a multisymplectic Lie system and let  $(\Theta, \Phi)$  be a multisymplectic reduction scheme whose  $\{\Phi_g\}_{g \in G}$  are symmetries of the elements of  $V^X$ . Then,  $X$  can be*

projected onto  $N/G$  giving rise to a new multisymplectic Lie system  $(N/G, \tilde{\Theta}, \pi_*X)$ , where  $\tilde{\Theta}$  is the multisymplectic form on  $N/G$  induced by the multisymplectic reduction scheme  $(\Theta, \Phi)$ .

**Proof.** If the  $\Phi_g$ , with  $g \in G$ , are symmetries of the elements of  $V^X$ , in particular they are symmetries of  $X$  because it takes values in  $V^X$ . Hence,  $X$  can be projected onto  $N/G$ . If  $\pi : N \rightarrow N/G$  is the quotient projection, then  $\pi_*|_{V^X} : V^X \rightarrow \mathfrak{X}(N/G)$  is a Lie algebra morphism and  $\pi_*(V^X)$  is therefore finite-dimensional turning  $\pi_*X$  into a Lie system. Since  $(N, \Theta, X)$  is a multisymplectic Lie system and the elements of  $V^X$  commute with the fundamental vector fields of the Lie group action  $\Phi$ , one has that  $\mathcal{L}_{X \wedge X_1 \wedge \dots \wedge X_r} \Theta = 0$  and, as  $X$  and  $\Theta$  are projectable onto  $N/G$ , then  $\mathcal{L}_{\pi_*X} \tilde{\Theta} = 0$ . Therefore  $(N/G, \tilde{\Theta}, \pi_*X)$  is a multisymplectic Lie system.  $\square$

It is frequently accessible to find Lie symmetries for a Lie system by using Lie theory methods. Nevertheless, the crux is to find a multisymplectic form satisfying the conditions provided in theorem 4.3.

**4.2.2. Reductions of a control system.** Consider again the control system given in example 3.6. Let us now apply our multisymplectic reduction to this control system. Recall that the vector fields  $X_1, \dots, X_5$  span, in view of their unique non-vanishing commutation relations (3.4), a unimodular Lie algebra  $V$ . Moreover,  $\mathcal{D}^V = \mathbb{T}\mathbb{R}^5$ . In consequence, this control system is a locally automorphic one and the application to it of the techniques devised in [33] gives rise to a multisymplectic Lie system  $(\mathbb{R}^5, \Theta^B, X)$ . For example, we can construct the multisymplectic structure

$$\Theta^B = \eta_1 \wedge \eta_2 \wedge \eta_3 \wedge \eta_4 \wedge \eta_5,$$

where  $\eta_1, \dots, \eta_5$  is a dual basis to  $X_1, \dots, X_5$  given by

$$\begin{aligned} \eta_1 &= dx_1, & \eta_2 &= dx_2, & \eta_3 &= -x_1 dx_2 + dx_3, \\ \eta_4 &= x_1^2 dx_2 - 2x_1 dx_3 + dx_4, & \eta_5 &= -2x_2 dx_3 + dx_5. \end{aligned}$$

In coordinates,

$$\Theta^B = dx_1 \wedge \dots \wedge dx_5.$$

Let us use the Lie symmetries of this system to accomplish several of its possible reductions. Recall that the Lie symmetries of the control systems (3.2) are given in (3.6). Indeed, after a long but simple calculation, we obtain a Lie algebra  $V^S$  of Lie symmetries spanned by

$$\begin{aligned} Y_1 &= \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial x_3} + 2x_3 \frac{\partial}{\partial x_4} + x_2^2 \frac{\partial}{\partial x_5}, & Y_2 &= \frac{\partial}{\partial x_2} + 2x_3 \frac{\partial}{\partial x_5}, \\ Y_3 &= \frac{\partial}{\partial x_3}, & Y_4 &= \frac{\partial}{\partial x_4}, & Y_5 &= \frac{\partial}{\partial x_5}. \end{aligned}$$

Recall also that

$$Y_\alpha(0) = X_\alpha(0), \quad \alpha = 1, \dots, 5$$

and the only non-vanishing commutation relations between the previous vector fields read

$$[Y_1, Y_2] = -Y_3, \quad [Y_1, Y_3] = -2Y_4, \quad [Y_2, Y_3] = -2Y_5.$$

Hence,  $V^S = \langle Y_1, \dots, Y_6 \rangle \simeq V$ . Let  $\mathfrak{g}_C$  be the abstract Lie algebra isomorphic to  $V$ . This leads us to define a momentum map

$$\begin{aligned} \mathcal{J}: \mathbb{R}^5 \times \mathcal{G}(\mathfrak{g}_C) &\longrightarrow \overset{\bullet}{\wedge} T^*\mathbb{R}^5 \\ (\xi, w) &\longmapsto \iota_w \Theta^B(\xi). \end{aligned}$$

Consider the unimodular subalgebra  $V_{45}^L := \langle Y_4, Y_5 \rangle$  and the associated Noether invariant of  $X^G$ :

$$\bar{\Theta}_{45} := \iota_{Y_4 \wedge Y_5} \Theta^B = dx_1 \wedge dx_2 \wedge dx_3.$$

The Lie algebra  $\langle Y_4, Y_5 \rangle \simeq \mathbb{R}^2$  can be considered as the Lie algebra of fundamental vector fields of a proper and free unimodular Lie group action

$$\Phi_{45} : (\lambda_1, \lambda_2; x_1, x_2, x_3, x_4, x_5) \in \mathbb{R}^2 \times \mathbb{R}^5 \longmapsto (x_1, x_2, x_3, x_4 + \lambda_1, x_5 + \lambda_2) \in \mathbb{R}^5.$$

Moreover, the space  $\Lambda^2 V_\Phi = \langle Y_4 \wedge Y_5 \rangle$  is Hamiltonian. Since  $\Theta^B$  is five-nondegenerate, the pair  $(\Theta^B, \Phi_{45})$  becomes a reduction scheme and theorem 4.1 can be applied to reduce this multisymplectic structure to  $\mathbb{R}^5/\mathbb{R}^2$ . More specifically, the orbits of the action  $\Phi$  take the form  $\mathcal{O}_{[x_1, x_2, x_3, x_4, x_5]} = \{x_1\} \times \{x_2\} \times \{x_3\} \times \mathbb{R}^2$  and the quotient space is  $\mathbb{R}^5/\mathbb{R}^2 \simeq \mathbb{R}^3$ . The variables  $x_1, x_2, x_3$  give rise to a coordinate system in the quotient and one can define the submersion

$$\pi_{45} : (x_1, \dots, x_5) \in \mathbb{R}^5 \longmapsto (x_1, x_2, x_3) \in \mathbb{R}^3.$$

Additionally, theorem 4.1 ensures that there exists a unique multisymplectic form  $\widehat{\Theta}_{45}$  on  $\mathbb{R}^3$  whose pull-back to  $\mathbb{R}^5$  is  $\bar{\Theta}_{45}$ , namely  $\widehat{\Theta}_{45} = dx_1 \wedge dx_2 \wedge dx_3$ .

Since  $(\Theta^B, \Phi_{45})$  is a reduction scheme such that the  $\Phi_g$ , with  $g \in \mathbb{R}^2$ , are symmetries of the Lie system of the multisymplectic Lie system  $(\mathbb{R}^5, \Theta^B, X)$ , theorem 4.3 ensures that we can reduce the multisymplectic Lie system. In fact, the Vessiot–Guldberg Lie algebra for the control system (3.2) project onto  $\mathbb{R}^3$  giving rise to a new Lie algebra  $V_{45}$  spanned by vector fields

$$X_1^{45} = \frac{\partial}{\partial x_1}, \quad X_2^{45} = \frac{\partial}{\partial x_2} + x_1 \frac{\partial}{\partial x_3}, \quad X_3^{45} = \frac{\partial}{\partial x_3}.$$

Hence, also the system  $X^{CS}$  can be projected onto  $\mathbb{R}^3$  to give rise to a Lie system  $X_{45}$  on  $\mathbb{R}^3$ . Moreover, the Lie algebra  $V_{45}$  consists of Hamiltonian vector fields relative to  $\widehat{\Theta}_{45}$ , which turns the triple  $(\mathbb{R}^3, \widehat{\Theta}_{45}, X_{45})$  into a new multisymplectic Lie system.

#### 4.2.3. Systems on Lie groups and quantum harmonic oscillators with a spin-magnetic term.

Let us consider a Lie system on  $G^{os} := \text{SL}_2 \times \text{SO}_3$  related to a quantum mechanical system given by a  $t$ -dependent harmonic oscillator with a spin-magnetic term. This example will be useful in the next section to illustrate how several reductions of a multisymplectic Lie system can be used to reconstruct the dynamics of the original one.

The Lie algebra of  $G^{os}$  takes the form  $\mathfrak{g}^{os} := \mathfrak{sl}_2 \oplus \mathfrak{so}_3$  where  $\oplus$  represents the direct sum of the Lie algebra  $\mathfrak{sl}_2$  and  $\mathfrak{so}_3$ .

Let us define the Lie system on  $G^{os}$  of the form

$$\frac{dg}{dt} = \sum_{\alpha=1}^r b_\alpha(t) X_\alpha^R(g) =: X^{os}(t, g), \quad g \in G^{os}, \quad t \in \mathbb{R}, \tag{4.26}$$

where  $b_1(t), \dots, b_6(t)$  are arbitrary  $t$ -dependent functions and  $\{X_1^R, \dots, X_6^R\}$  form a basis of the Lie algebra  $V^R$  of right-invariant vector fields on a Lie group  $G^{\text{os}}$  in such a way that  $\{X_1^R, X_2^R, X_3^R\}$  forms a basis of the Lie algebra  $V_{\mathfrak{sl}} := \langle X_1^R, X_2^R, X_3^R \rangle \simeq \mathfrak{sl}_2$  and  $\{X_4^R, X_5^R, X_6^R\}$  stands for a basis of the Lie algebra  $V_{\mathfrak{so}} := \langle X_4^R, X_5^R, X_6^R \rangle \simeq \mathfrak{so}_3$ .

The relevance of the Lie system (4.26) is due to the fact that it appears in the solution of quantum harmonic oscillators with a spin-magnetic term [21]. More specifically, particular cases of (4.26) appear as the automorphic Lie systems associated with the  $t$ -dependent Schrödinger equation related to a quantum  $t$ -dependent Hamiltonian operator of the form

$$\widehat{H}_1(t) = \frac{\widehat{p}_x^2 + \widehat{p}_y^2 + \widehat{p}_z^2}{2} + \omega^2(t) \frac{\widehat{x}^2 + \widehat{y}^2 + \widehat{z}^2}{2} + B_x(t)\widehat{S}_x + B_y(t)\widehat{S}_y + B_z(t)\widehat{S}_z, \tag{4.27}$$

where  $\omega(t)$  is any  $t$ -dependent frequency,  $B_x(t), B_y(t), B_z(t)$  are the coordinates of a magnetic field and  $\widehat{S}_x, \widehat{S}_y, \widehat{S}_z$  can be assumed to be, for instance, the Pauli matrices up to a proportional constant. Note that  $\widehat{H}_1(t)$  acts on  $\mathcal{L}^2(\mathbb{R}^3) \otimes \mathbb{C}^2$ , where  $\mathcal{L}^2(\mathbb{R}^3)$  is the space of complex square-integrable functions on  $\mathbb{R}^3$ . It is remarkable that (4.26) is also related to other  $t$ -dependent Hamiltonian operators, e.g.

$$\widehat{H}_2(t) = \frac{\widehat{p}_x^2 + \widehat{p}_y^2 + \widehat{p}_z^2}{2} + \omega^2(t) \frac{\widehat{x}^2 + \widehat{y}^2 + \widehat{z}^2}{2} + B_x(t)\widehat{L}_x + B_y(t)\widehat{L}_y + B_z(t)\widehat{L}_z, \tag{4.28}$$

acting on  $\mathcal{L}^2(\mathbb{R}^3)$ , where  $\widehat{L}_x, \widehat{L}_y, \widehat{L}_z$  are the angular momentum operators relative to the axes  $OX, OY, OZ$ , respectively. In both previous cases, the key to understand the relation between (4.27) and (4.28) is the fact that  $i\widehat{H}_j(t)$ , with  $j = 1, 2$ , can be written as a linear combination with  $t$ -dependent constants of a family of skew-Hermitian operators closing a real Lie algebra isomorphic to  $\mathfrak{g}^{\text{os}}$  (cf [21]).

The above examples show that we can define a momentum map of the form

$$\mathcal{J}^{\text{os}} : G^{\text{os}} \times \mathcal{G}(\mathfrak{g}^{\text{os}}) \rightarrow \dot{\wedge} T^*G^{\text{os}}.$$

We call  $\mathcal{G}(\mathfrak{g}^{\text{os}})$  a *traceless Grassmannian*, i.e. the space of traceless subalgebras of  $\mathfrak{g}^{\text{os}}$ . This mapping allows us to gather all previous reduction processes. Although the relevant properties of the system are better understood by using a geometric approach, we hereafter provide a local coordinate expression for this system by using canonical coordinates of the second kind on  $G^{\text{os}}$  of the form

$$\phi : (v_1, \dots, v_6) \in \mathfrak{g}^{\text{os}} \mapsto \exp(-v_1 e_1) \times \dots \times \exp(-v_6 e_6) \in G^{\text{os}},$$

for  $X_\alpha^R(e) = -e_\alpha$  for  $\alpha = 1, \dots, 6$ , namely

$$\begin{cases} \frac{dv_1}{dt} = b_1(t) + b_2(t)v_1 + b_3(t)v_1^2, & \frac{dv_4}{dt} = b_4(t) + \tan v_5(b_5(t)\sin v_4 + b_6(t)\cos v_4), \\ \frac{dv_2}{dt} = b_2(t) + 2b_3(t)v_1, & \frac{dv_5}{dt} = b_5(t)\cos v_4 - b_6(t)\sin v_4, \\ \frac{dv_3}{dt} = e^{v_2} b_3(t), & \frac{dv_6}{dt} = \frac{b_5(t)\sin v_4 + b_6(t)\cos v_4}{\cos v_5}. \end{cases} \tag{4.29}$$



We have now the following coordinate expressions for the vector fields

$$\begin{aligned}
 X_1^R &= \frac{\partial}{\partial v_1}, & X_4^R &= \frac{\partial}{\partial v_4}, \\
 X_2^R &= v_1 \frac{\partial}{\partial v_1} + \frac{\partial}{\partial v_2}, & X_5^R &= \sin v_4 \tan v_5 \frac{\partial}{\partial v_4} + \cos v_4 \frac{\partial}{\partial v_5} + \frac{\sin v_4}{\cos v_5} \frac{\partial}{\partial v_6}, \\
 X_3^R &= v_1^2 \frac{\partial}{\partial v_1} + 2v_1 \frac{\partial}{\partial v_2} + e^{v_2} \frac{\partial}{\partial v_3}, & X_6^R &= \cos v_4 \tan v_5 \frac{\partial}{\partial v_4} - \sin v_4 \frac{\partial}{\partial v_5} + \frac{\cos v_4}{\cos v_5} \frac{\partial}{\partial v_6}.
 \end{aligned}
 \tag{4.30}$$

Recall that right-invariant vector fields must be defined everywhere. The singularity in the above local coordinate expression when  $v_5 = \pi/2 + k\pi$  for  $k \in \mathbb{Z}$  is only due to the fact that canonical coordinates of the second kind are defined, generically, only on an open neighbourhood of the neutral element of  $G^{\text{os}}$ . A simple change of coordinates would remove the singularity in the expression of right-invariant vector fields when the previous second canonical coordinates fail to be well defined.

Next, it will be useful to describe the commutation relations among the above vector fields

$$\begin{aligned}
 [X_1^R, X_2^R] &= X_1^R, & [X_2^R, X_3^R] &= X_3^R, & [X_3^R, X_4^R] &= 0, & [X_4^R, X_5^R] &= X_6^R, & [X_5^R, X_6^R] &= X_4, \\
 [X_1^R, X_3^R] &= 2X_2^R, & [X_2^R, X_4^R] &= 0, & [X_3^R, X_5^R] &= 0, & [X_4^R, X_6^R] &= -X_5^R, \\
 [X_1^R, X_4^R] &= 0, & [X_2^R, X_5^R] &= 0, & [X_3^R, X_6^R] &= 0, \\
 [X_1^R, X_5^R] &= 0, & [X_2^R, X_6^R] &= 0, \\
 [X_1^R, X_6^R] &= 0.
 \end{aligned}$$

Thus, the vector fields given in (4.30) span a real six-dimensional Lie algebra  $V^{\text{os}}$  isomorphic to  $\mathfrak{sl}_2 \oplus \mathfrak{so}_3$ . Their dual forms are given by

$$\begin{aligned}
 \eta_1^R &= dv_1 - v_1 dv_2 + v_1^2 e^{-v_2} dv_3, & \eta_4^R &= dv_4 - \sin v_5 dv_6, \\
 \eta_2^R &= dv_2 - 2v_1 e^{-v_2} dv_3, & \eta_5^R &= \cos v_4 dv_5 + \sin v_4 \cos v_5 dv_6, \\
 \eta_3^R &= e^{-v_2} dv_3, & \eta_6^R &= -\sin v_4 dv_5 + \cos v_4 \cos v_5 dv_6.
 \end{aligned}$$

Following the method presented in [33], we can build up a multisymplectic form on  $G^{\text{os}} = \text{SL}_2 \times \text{SO}_3$  such that the elements of  $V^{\text{os}}$  become Hamiltonian vector fields with respect to it. Indeed, the obtained multisymplectic form reads

$$\Theta^{\text{os}} = \eta_1^R \wedge \dots \wedge \eta_6^R = e^{-v_2} \cos v_5 dv_1 \wedge dv_2 \wedge dv_3 \wedge dv_4 \wedge dv_5 \wedge dv_6.
 \tag{4.31}$$

We will use the Lie symmetries of the multisymplectic Lie system  $(G^{\text{os}}, \Theta^{\text{os}}, X^{\text{os}})$  to reduce it. Recall that the Lie symmetries of a Lie system are given by the system of partial differential equation (3.5). In this case, the Lie algebra  $V^{\mathcal{S}} = \text{Sym}(V^{\text{os}})$  is spanned by the vector fields

$$\begin{aligned}
 Y_1^L &= e^{v_2} \frac{\partial}{\partial v_1} + 2v_3 \frac{\partial}{\partial v_2} + v_3^2 \frac{\partial}{\partial v_3}, & Y_4^L &= \frac{\cos v_6}{\cos v_5} \frac{\partial}{\partial v_4} - \sin v_6 \frac{\partial}{\partial v_5} + \cos v_6 \tan v_5 \frac{\partial}{\partial v_6}, \\
 Y_2^L &= \frac{\partial}{\partial v_2} + v_3 \frac{\partial}{\partial v_3}, & Y_5^L &= \frac{\sin v_6}{\cos v_5} \frac{\partial}{\partial v_4} + \cos v_6 \frac{\partial}{\partial v_5} + \sin v_6 \tan v_5 \frac{\partial}{\partial v_6}, \\
 Y_3^L &= \frac{\partial}{\partial v_3}, & Y_6^L &= \frac{\partial}{\partial v_6},
 \end{aligned}
 \tag{4.32}$$

where the only non-vanishing commutation relations are

$$\begin{aligned}
 [Y_1^L, Y_2^L] &= -Y_1^L, & [Y_2^L, Y_3^L] &= -Y_3^L, & [Y_4^L, Y_5^L] &= -Y_6^L, & [Y_5^L, Y_6^L] &= -Y_4^L, \\
 [Y_1^L, Y_3^L] &= -2Y_2^L, & [Y_4^L, Y_6^L] &= Y_5^L.
 \end{aligned}$$

Hence,  $V^S = \langle Y_1^L, Y_2^L, Y_3^L, Y_4^L, Y_5^L, Y_6^L \rangle \simeq V^{\text{os}}$  via the Lie algebra isomorphism mapping  $X_\alpha^R \mapsto -Y_\alpha^L$  for  $\alpha = 1, \dots, 6$ .

Let us consider the momentum map

$$\begin{aligned}
 \mathcal{J}^{\text{os}}: G^{\text{os}} \times \mathcal{G}(\mathfrak{g}^{\text{os}}) &\longrightarrow \overset{\bullet}{\wedge} T^*G^{\text{os}} \\
 (\xi, w) &\longmapsto \iota_w \Theta^{\text{os}}(\xi).
 \end{aligned}$$

- First, consider the reduction of  $\Theta^{\text{os}}$  by  $Y_1^L \wedge Y_2^L \wedge Y_3^L$ . We have the quotient map projection

$$\pi_{123}: (v_1, \dots, v_6) \in G^{\text{os}} \longmapsto (v_4, v_5, v_6) \in G/\text{SL}_2 \simeq \text{SO}_3.$$

The projections of the elements of the basis (4.30) of the Vessiot–Guldberg Lie algebra  $V^{\text{os}}$  onto  $G^{\text{os}}/\text{SL}_2$  read  $\tilde{X}_1^{sl} = \tilde{X}_2^{sl} = \tilde{X}_3^{sl} = 0$ , while

$$\begin{aligned}
 \tilde{X}_4^{sl} &= \frac{\partial}{\partial v_4}, \\
 \tilde{X}_5^{sl} &= \sin v_4 \tan v_5 \frac{\partial}{\partial v_4} + \cos v_4 \frac{\partial}{\partial v_5} + \frac{\sin v_4}{\cos v_5} \frac{\partial}{\partial v_6}, \\
 \tilde{X}_6^{sl} &= \cos v_4 \tan v_5 \frac{\partial}{\partial v_4} - \sin v_4 \frac{\partial}{\partial v_5} + \frac{\cos v_4}{\cos v_5} \frac{\partial}{\partial v_6}.
 \end{aligned}$$

It is clear that  $\tilde{V}^{sl} = \langle \tilde{X}_4^{sl}, \tilde{X}_5^{sl}, \tilde{X}_6^{sl} \rangle$  is a three-dimensional Lie algebra isomorphic to  $\mathfrak{so}_3$ . This gives rise to a Lie system

$$X^{sl} = \sum_{\alpha=4}^6 b_\alpha(t) \tilde{X}_\alpha^{sl}$$

on the quotient space  $G_1^{\text{os}} = G^{\text{os}}/\text{SL}_2$ . This is a multisymplectic Lie system with respect to the multisymplectic three-form  $\Theta_{123}^{\text{os}}$  determined by

$$\pi_{123}^* \Theta_{123}^{\text{os}} = \iota_{Y_1^L \wedge Y_2^L \wedge Y_3^L} \Theta^{\text{os}} = \cos v_5 \, dv_4 \wedge dv_5 \wedge dv_6$$

on  $G_1^{\text{os}}$ .

- On the other hand, consider the reduction of  $\Theta^{\text{os}}$  by  $Y_4^L \wedge Y_5^L \wedge Y_6^L$ . We have the projection

$$\pi_{456}: (v_1, \dots, v_6) \in G^{\text{os}} \longmapsto (v_1, v_2, v_3) \in G^{\text{os}}/\text{SO}_3 \simeq \text{SL}_2.$$

The projections of the elements of the basis (4.30) of the Vessiot–Guldberg Lie algebra  $V^{os}$  onto  $G^{os}/SO_3$  read

$$\tilde{X}_1^{so} = \frac{\partial}{\partial v_1}, \quad \tilde{X}_2^{so} = v_1 \frac{\partial}{\partial v_1} + \frac{\partial}{\partial v_2}, \quad \tilde{X}_3^{so} = v_1^2 \frac{\partial}{\partial v_1} + 2v_1 \frac{\partial}{\partial v_2} + e^{v_2} \frac{\partial}{\partial v_3},$$

and  $\tilde{X}_4^{so} = \tilde{X}_5^{so} = \tilde{X}_6^{so} = 0$ . It is clear that  $\tilde{V}^{so} = \langle \tilde{X}_1^{so}, \tilde{X}_2^{so}, \tilde{X}_3^{so} \rangle$  is three-dimensional Lie algebra isomorphic to  $\mathfrak{sl}_2$ . This gives rise to a Lie system

$$X^{so} = \sum_{\alpha=1}^3 b_{\alpha}(t)\tilde{X}_{\alpha}^{so}$$

on the quotient space  $G_2^{os} = G^{os}/SO_3$ . This is a multisymplectic Lie system on  $G_2^{os}$  with respect to the multisymplectic three-form  $\Theta_{456}^{os}$  induced by

$$\pi_{456}^* \Theta_{456}^{os} = \iota_{Y_4^L \wedge Y_5^L \wedge Y_6^L} \Theta^{os} = -e^{-v_2} dv_1 \wedge dv_2 \wedge dv_3.$$

### 5. Multisymplectic reconstruction

This section gives criteria for determining a multisymplectic Lie system  $(M, \Theta, X)$  from several of its multisymplectic reductions. Let us start by giving a particular example that illustrates our more general techniques to be deployed later on.

Let us go back to the multisymplectic Lie system  $(G^{os}, \Theta^{os}, X^{os})$  related to (4.26) and associated with quantum harmonic oscillators with a spin-magnetic term. Recall that  $(G^{os}, \Theta^{os}, X^{os})$  gives rise, by using multisymplectic reductions, to two different multisymplectic Lie systems

$$(G_1^{os}, \Theta_{123}^{os}, X^{sl}), \quad (G_2^{os}, \Theta_{456}^{os}, X^{so}).$$

Recall that  $\Theta_{123}^{os}$  is a multisymplectic form on  $G_1^{os}$  such that  $\pi_{123}^* \Theta_{123}^{os} = \iota_{Y_1^L \wedge Y_2^L \wedge Y_3^L} \Theta^{os}$ . As

$$Y_1^L \wedge Y_2^L \wedge Y_3^L = e^{v_2} \frac{\partial}{\partial v_1} \wedge \frac{\partial}{\partial v_2} \wedge \frac{\partial}{\partial v_3}, \tag{5.1}$$

and  $\Theta^{os}$  is six-nondegenerate because it is a volume form on  $G^{os}$ , the original multisymplectic form is uniquely determined by  $\Theta_{123}^{os}$ , namely  $\Theta_{123}^{os} = \cos v_5 dv_4 \wedge dv_5 \wedge dv_6$ , and it must take the form (4.31). Alternatively, the knowledge of  $\Theta_{456}^{os}$ , namely  $\Theta_{456}^{os} = -e^{-v_2} dv_1 \wedge dv_2 \wedge dv_3$ , and the fact that it was obtained as a multisymplectic reduction via the contraction of  $\Theta^{os}$  with

$$Y_4^L \wedge Y_5^L \wedge Y_6^L = \frac{1}{\cos v_5} \frac{\partial}{\partial v_4} \wedge \frac{\partial}{\partial v_5} \wedge \frac{\partial}{\partial v_6},$$

determines again the value of  $\Theta^{os}$ , which matches the value obtained previously via the previous multisymplectic reduction.

Let us now focus on retrieving the  $t$ -dependent vector field  $X^{os}$  on  $G^{os}$ , whose reductions gave rise to the  $t$ -dependent vector fields on  $G_1^{os}$  and  $G_2^{os}$ , given by

$$X^{sl} = \sum_{\alpha=4}^6 b_{\alpha}(t)\tilde{X}_{\alpha}^{sl}, \quad X^{so} = \sum_{\alpha=1}^3 b_{\alpha}(t)\tilde{X}_{\alpha}^{so}, \tag{5.2}$$

respectively. In this case, we know that  $\pi_{123}^* X^{os} = X^{sl}$ , which determines  $X^{os}$  up to a vector field on  $G^{os}$  taking values in  $\ker T\pi_{123}$ . Similarly,  $\pi_{456}^* X^{os} = X^{so}$ , which retrieves the value of

$X^{os}$  up to a vector field on  $G^{os}$  taking values in  $\ker T\pi_{456}$ . Hence, the value of  $X^{os}$  is determined at a point  $g \in G^{os}$  up to an element of

$$\ker T_p\pi_{123} \cap \ker T_p\pi_{456} = \{0\}.$$

In other words,  $X^{os}$  is uniquely determined by (5.2). Note that this example shows how to determine the original multisymplectic Lie system on a Lie group, related to a multisymplectic volume form, via two of its multisymplectic reductions.

The previous example illustrates how to set sufficient conditions to retrieve our initial multisymplectic Lie system from its multisymplectic reductions. For instance, we can consider the following proposition.

**Proposition 5.1.** *Let  $G$  be a Lie group with certain Lie subgroups  $G_1, \dots, G_k$ . Let  $(G/G_i, \Theta^i, X^i)$ , with  $i = 1, \dots, k$ , be a set of multisymplectic reductions of a multisymplectic Lie system  $(G, \Theta, X)$ , let  $\pi_i : G \rightarrow G/G_i$  and  $\iota_{Y_i}\Theta$ , with  $1, \dots, k$ , be the projections and Noether invariants related to the multisymplectic reductions induced by the multivector fields  $Y_1, \dots, Y_k$  on  $G$ . Assume that*

$$\bigcap_{i=1}^k \ker T_g\pi_i = \{0\}, \quad \forall g \in G,$$

and suppose that  $\Theta$  is a volume form on  $G$ . Then, the multisymplectic Lie system  $(G, \Theta, X)$  is univocally determined from its multisymplectic reductions.

**Proof.** Since by construction of the multisymplectic reductions, one has that  $X_{\pi_i(g)}^i = T_g\pi_i X_g$  for  $i = 1, \dots, k$  and every  $g \in G$ , then the value of  $X$  at  $g$ , let us say  $X_g$ , is determined by  $X^1, \dots, X^k$  up to an element of  $\bigcap_{i=1}^k \ker T_g\pi_i$ , which is, by assumption, equals to zero. Hence,  $X$  is determined univocally by  $X^1, \dots, X^k$ .

Let us now study the determination of the multisymplectic form  $\Theta$  on  $G$  via the Noether invariants  $\iota_{Y_i}\Theta$  associated with the multisymplectic reductions and the induced multisymplectic forms on the different quotients  $G/G_i$ . Given one of them, let us say  $\Theta^i$ , one has that

$$\iota_{Y_i}\Theta = \pi_i^*\Theta^i.$$

Since  $\Theta$  is a volume form and  $\pi$  is a surjective submersion,  $\Theta$  is univocally determined by  $\Theta^i$ , the multivector field  $Y_i$ , and the above relation.  $\square$

To understand the generality of the assumptions of the previous proposition, it is worth noting that multisymplectic volume forms are not as restrictive as it may seem at first. In two-dimensional manifolds, every multisymplectic form is a volume form. On three-dimensional manifolds, it occurs the same way as differential two- and one-forms are degenerate. In four-dimensional manifolds, differential three-forms are degenerate because they can all be written, locally, as the contraction of a volume form with a vector field. Moreover, differential one-forms are also degenerate and the only multisymplectic forms on a four-dimensional manifold are volume forms or symplectic ones. It is worth stressing that we are mainly interested in multisymplectic forms that are not symplectic ones, as only in this case the multisymplectic formalism will provide information that is not available by means of other theories. Hence, non-volume multisymplectic forms, which are not symplectic forms, appear only on manifolds of dimension five or higher.

Anyhow, proposition 5.1 can be modified to deal with multisymplectic Lie systems relative to a class of non-volume multisymplectic forms. As shown next, these multisymplectic forms need not be  $k$ -nondegenerate for  $k > 1$ , and this leads to new features. Let us provide an example to illustrate the new geometric aspects introduced in this case.

Consider a multisymplectic form on  $\mathbb{R}^8$  of the form

$$\Omega^S = \sum_{1 \leq i_1 < i_2 < i_3 < i_4 < i_5 < i_6 \leq 8} dx^{i_1} \wedge dx^{i_2} \wedge dx^{i_3} \wedge dx^{i_4} \wedge dx^{i_5} \wedge dx^{i_6},$$

where  $\{x^1, \dots, x^8\}$  are global coordinates on  $\mathbb{R}^8$ . In fact,  $\Omega^S$  is closed and, since the mappings  $\Omega_x^{S\sharp} : T_x M \rightarrow \wedge^5 T_x M$  are injections for every  $x \in M$ , one-nondegenerate. Moreover,  $\Omega^S$  is also two- and three-nondegenerate, i.e. if  $\iota_Z \Omega^S = 0$  for  $Z \in \mathfrak{X}^2(\mathbb{R}^8)$  or  $Z \in \mathfrak{X}^3(\mathbb{R}^8)$ , then  $Z = 0$ . In fact,  $\iota_Z \Omega^S = 0$  can be seen as an algebraic equation in the coefficients of  $Z \in \mathfrak{X}^s(\mathbb{R}^8)$  in a certain basis of differential  $s$ -forms. In particular, when  $Z \in \mathfrak{X}^2(\mathbb{R}^8)$ , the algebraic system  $\iota_Z \Omega^S = 0$  can be seen as a system of  $\binom{8}{4}$  linear equations on the  $\binom{8}{2}$  coefficients of  $Z$ , which, after a simple but tedious calculation, turns out to have a unique trivial  $Z = 0$  solution. Moreover, if  $Z \in \mathfrak{X}^3(\mathbb{R}^8)$ , the algebraic system  $\iota_Z \Omega^S = 0$  can be seen as a system of  $\binom{8}{3}$  linear equations on the  $\binom{8}{3}$  coefficients of  $Z$ , which, as in the previous case, has only a unique zero trivial solution. Note that this last fact also implies that  $\Omega^S$  must be one- and two-nondegenerate.

Consider the Lie system on  $\mathbb{R}^8$  of the form

$$X^S = \sum_{i=1}^8 b_i(t) \frac{\partial}{\partial x^i},$$

where  $b_1(t), \dots, b_8(t)$  are arbitrary  $t$ -dependent functions. It is immediate that the vector fields  $\{\partial/\partial x^1, \dots, \partial/\partial x^8\}$  are Hamiltonian relative to  $\Omega^S$ . Then,  $(\mathbb{R}^8, \Omega^S, X^S)$  becomes a multisymplectic Lie system.

Consider the Lie group action  $\Phi : (\lambda^1, \dots, \lambda^8, x^1, \dots, x^8) \in \mathbb{R}^8 \times \mathbb{R}^8 \mapsto (\lambda^1 + x^1, \dots, \lambda^8 + x^8) \in \mathbb{R}^8$ , which admits a momentum map

$$\mathcal{J} : \mathbb{R}^8 \times \mathcal{G}(\mathbb{R}^8) \rightarrow \dot{\wedge} T^* \mathbb{R}^8.$$

Given the Hamiltonian bivector fields on  $\mathbb{R}^8$  of the form

$$Z_a = \frac{\partial}{\partial x^1} \wedge \frac{\partial}{\partial x^2}, \quad Z_b = \frac{\partial}{\partial x^4} \wedge \frac{\partial}{\partial x^5}, \quad Z_c = \frac{\partial}{\partial x^7} \wedge \frac{\partial}{\partial x^8},$$

let us analyse the multisymplectic reductions of  $\Omega^S$  related to them. In particular,

$$\begin{aligned} \iota_{Z_a} \Omega^S &= \sum_{3 \leq i_1 < i_2 < i_3 < i_4 \leq 8} dx^{i_1} \wedge dx^{i_2} \wedge dx^{i_3} \wedge dx^{i_4}, \\ \iota_{Z_c} \Omega^S &= \sum_{1 \leq i_1 < i_2 < i_3 < i_4 \leq 6} dx^{i_1} \wedge dx^{i_2} \wedge dx^{i_3} \wedge dx^{i_4}, \\ \iota_{Z_b} \Omega^S &= \sum_{1 \leq i_1 < i_2 < i_3 < i_4 \leq 8} \epsilon_{45i_1 i_2 i_3 i_4} dx^{i_1} \wedge dx^{i_2} \wedge dx^{i_3} \wedge dx^{i_4}, \end{aligned}$$

where  $\epsilon_{45i_1 i_2 i_3 i_4}$  is the Levi-Civita symbol relative to the sequence of indices  $4, 5, i_1, i_2, i_3, i_4$ . The above Noether invariants are basic forms relative to the canonical projections  $\pi_a : (x^1, \dots, x^8) \in \mathbb{R}^8 \mapsto (x^3, \dots, x^8) \in \mathbb{R}^6$ ,  $\pi_b : (x^1, \dots, x^8) \in \mathbb{R}^8 \mapsto (x^1, x^2, x^3, x^6, x^7, x^8) \in \mathbb{R}^6$ ,

and  $\pi_c : (x^1, \dots, x^8) \in \mathbb{R}^8 \mapsto (x^1, x^2, x^3, x^4, x^5, x^6) \in \mathbb{R}^6$  associated with the multisymplectic reductions of  $\Omega^S$  by  $Z_a, Z_b$ , and  $Z_c$ , respectively. In other words,  $\pi_a^* \Omega_a = \iota_{Z_a} \Omega^S$ ,  $\pi_b^* \Omega_b = \iota_{Z_b} \Omega^S$ ,  $\pi_c^* \Omega_c = \iota_{Z_c} \Omega^S$  for three unique differential four-forms  $\Omega_a, \Omega_b$ , and  $\Omega_c$  on  $\mathbb{R}^6$ , which are therefore closed. They are also one-nondegenerate because  $\Omega^S$  is three-nondegenerate.

On the other hand, the value of  $\pi_a^* \Omega_a$  does not allow us to characterise univocally  $\Omega^S$  via the equality  $\iota_{Z_a} \Omega^S = \pi_a^* \Omega_a$  since there are many differential six-forms,  $\Omega$ , satisfying that  $\iota_{Z_a} \Omega = 0$ . Therefore,  $\pi_a^* \Omega_a = \iota_{Z_a} \Omega^S$  determines  $\Omega^S$  up to a differential six-form, let us say  $\Omega_{aa}$ , whose contraction with  $Z_a$  vanishes. Moreover, the differential six-forms  $\Omega^S + \epsilon \Omega_{aa}$ , for small  $\epsilon$ , are one-nondegenerate and  $\iota_{Z_a} (\Omega^S + \epsilon \Omega_{aa}) = \pi_a^* \Omega_a$ . Hence, to determine every possible multisymplectic six-form on  $\mathbb{R}^8$ , let us say  $\Omega$ , satisfying  $\iota_{Z_a} \Omega = \pi_a^* \Omega_a$ , it is interesting to define the following notion.

**Definition 5.2.** Let  $w$  be a  $r$ -tangent vector at  $x \in M$ , i.e.  $w \in \wedge^r T_x M$ . The  $\ell$ -annihilator of  $w$ , denoted by  $w^{\circ, \ell}$ , is the space of skew-symmetric  $\ell$ -forms,  $\Omega_x$ , on  $T_x M$  such that  $\iota_w \Omega_x = 0$ .

In the previous example, the knowledge of  $\pi_a^* \Omega_a$  determines the value of  $\Omega^S$  up to a differential six-form vanishing on  $Z_a$ . Meanwhile,

$$Z_a^{\circ, 6} \ni dx^{i_1} \wedge dx^{i_2} \wedge dx^{i_3} \wedge dx^{i_4} \wedge dx^{i_5} \wedge dx^{i_6}$$

if, and only if,  $\{1, 2\} \not\subset \{i_1, \dots, i_6\}$ , which determines  $Z_a^{\circ, 6}$ . Similarly, one can determine  $Z_b^{\circ, 6}$  and  $Z_c^{\circ, 6}$ . From this, it follows that  $Z_a^{\circ, 6} \cap Z_b^{\circ, 6} \cap Z_c^{\circ, 6} = \{0\}$  and  $\Omega^S$  is the unique differential six-form satisfying that  $\iota_{Z_a} \Omega^S = \pi_a^* \Omega_a$ ,  $\iota_{Z_b} \Omega^S = \pi_b^* \Omega_b$ , and  $\iota_{Z_c} \Omega^S = \pi_c^* \Omega_c$ . Hence, it is immediate that proposition 5.1 can be generalised as follows.

**Proposition 5.3.** Let  $G$  be a finite-dimensional Lie group with certain Lie subgroups  $G_1, \dots, G_k$ . Let  $(G/G_i, \Theta^i, X^i)$ , with  $i = 1, \dots, k$ , be a set of multisymplectic reductions of a multisymplectic Lie system  $(G, \Theta, X)$ ; where  $\Theta$  is an  $(s + 1)$ -nondegenerate multisymplectic  $\ell$ -form on  $G$ . Let  $\pi_i : G \rightarrow G/G_i$  and  $\Theta^i = \iota_{Z_i} \Theta$ , with  $1, \dots, k$  and  $Z_i \in \mathfrak{X}^s(G)$ , be the projections and Noether invariants related to the multisymplectic reductions. If

$$\bigcap_{i=1}^k \ker T_g \pi_i = \{0\} \quad \text{and} \quad \bigcap_{i=1}^k Z_i^{\circ, \ell}(g) = \{0\}, \quad \forall g \in G,$$

then, the multisymplectic Lie system  $(G, \Theta, X)$  is univocally determined from its multisymplectic reductions  $(G/G_i, \Theta^i, X^i)$ , with  $i = 1, \dots, k$ .

## 6. Conclusions and outlook

This work presents a reduction procedure for multisymplectic Lie systems, which were first introduced in [33], by using a suitable momentum map. We have also developed a reconstruction procedure which, in favourable cases, allows us to recover the original Lie system from its reductions. These results are illustrated by working out several examples, including the Schwarz equation, dissipative quantum harmonic oscillators, a control system, and a quantum harmonic oscillator with a spin-magnetic term.

In the future, it is planned to study new physical and mathematical applications of our methods, e.g. to Schwarz derivatives [48]. It would also be interesting to investigate multisymplectic Lie systems that are not of locally transitive type. In those cases, the Lie system is not locally diffeomorphic to an automorphic Lie system on a Lie group and we cannot use the structures

on Lie groups to obtain multisymplectic forms, Lie symmetries, reductions and so on. This also raises the question of the existence of a multisymplectic form that turns a Lie system into a multisymplectic one.

Note that shown results about Schwarz equations highlight that multisymplectic structures may allow for the study of relative equilibrium points for systems that are not Hamiltonian ones relative to any symplectic or Poisson structure. It would be interesting to study the extension of energy–momentum methods [51] to Hamiltonian systems relative to multisymplectic forms. The energy–momentum method is based heavily on a Marsden–Weinstein reduction procedure (cf [50, 51]), which has been depicted here for multisymplectic structures. Therefore, our work solves one of the main drawbacks in developing such a method. We hope to follow these lines of research in upcoming works.

Finally, it would be interesting to develop an analogue of our methods to study the reductions of Lie systems admitting a Vessiot–Guldberg Lie algebra of Hamiltonian vector fields relative to other geometric structures, e.g.  $k$ -cosymplectic [45] or  $k$ -contact [31] structures.

## Acknowledgments

The authors acknowledge fruitful discussions and comments from our colleague Miguel-C Muñoz-Lecanda. J de Lucas acknowledges partial financial support from the Project MINIATURA 5 of the Polish National Science Centre (NCN) under Grant Number Nr 2021/05/X/ST1/01797. X Gràcia, X Rivas, N Román-Roy, and S Vilariño acknowledge partial financial support from the Spanish Ministerio de Ciencia e Innovación and Gobierno de Aragón projects PGC2018-098265-B-C31, PGC2018-098265-B-C33, RED2018-102541-T, and E48\_20R. We also thank the anonymous referees, whose interesting comments and suggestions have helped us improve our article.

## Data availability statement

No new data were created or analysed in this study.

## ORCID iDs

Javier de Lucas  <https://orcid.org/0000-0001-8643-144X>

Xavier Gràcia  <https://orcid.org/0000-0003-1006-4086>

Xavier Rivas  <https://orcid.org/0000-0002-4175-5157>

Narciso Román-Roy  <https://orcid.org/0000-0003-3663-9861>

Silvia Vilariño  <https://orcid.org/0000-0003-0404-1427>

## References

- [1] Abraham R and Marsden J E 1978 *Foundations of Mechanics* (New York: Benjamin-Cummings)
- [2] Ames W F 1978 Nonlinear superposition for operator equations *Nonlinear Equations in Abstract Spaces* (New York: Academic) pp 43–66 Proc. Int. Symp. Nonlinear Equations in Abstract Spaces, University of Texas Arlington, Texas 1977
- [3] Ballesteros A, Blasco A, Herranz F J, de Lucas J and Sardón C 2015 Lie–Hamilton systems on the plane: properties, classification, and applications *J. Differ. Equ.* **258** 2873–907



- [4] Ballesteros A, Cariñena J F, Herranz F J, de Lucas J and Sardón C 2013 From constants of motion to superposition rules for Lie–Hamilton systems *J. Phys. A: Math. Theor.* **46** 285203
- [5] Ballesteros A, Campoamor-Stursberg R, Fernandez-Saiz E, Herranz F J and de Lucas J 2018 Poisson–Hopf algebra deformations of Lie–Hamilton system *J. Phys. A: Math. Theor.* **51** 065202
- [6] Berkovich L M 2007 Method of factorization of ordinary differential operators and some of its applications *Appl. Anal. Discret. Math.* **1** 122–49
- [7] Blacker C 2021 Reduction of multisymplectic manifolds *Lett. Math. Phys.* **111** 64
- [8] Blacker Casey, Miti Antonio Michele and Ryvkin Leonid 2022 Reduction of  $L_\infty$ -algebras of observables on multisymplectic manifolds arXiv [2206.03137](https://arxiv.org/abs/2206.03137)
- [9] Blasco A, Herranz J F, de Lucas J and Sardón C 2015 Lie–Hamilton systems on the plane: applications and superposition rules *J. Phys. A: Math. Theor.* **48** 345202
- [10] Blázquez Sanz D 2008 Differential Galois Theory and Lie–Vessiot Systems *PhD Thesis* Universitat Politècnica de Catalunya, Barcelona
- [11] Bourbaki N 1968 *Groupes et algèbres de Lie* (Paris: Hermann)
- [12] Bump D 2013 *Lie Groups (Graduate Texts in Mathematics vol 225)* (Berlin: Springer)
- [13] Cantrijn F, Ibort L A and de León M 1996 Hamiltonian structures on multisymplectic manifolds *Rend. Sem. Mat. Univ. Politec. Torino* **54** 225–36
- [14] Cantrijn F, Ibort A and De León M 1999 On the geometry of multisymplectic manifolds *J. Aust. Math. Soc. A* **66** 303–30
- [15] Cariñena J F, Clemente-Gallardo J, Jover-Galtier J A and de Lucas J 2019 Application of Lie systems to quantum mechanics: superposition rules *Proc. 60 Years Alberto Ibort Fest Classical and Quantum Physics: Geometry, Dynamics and Control Springer Procs. Phys.* (Berlin: Springer)
- [16] Cariñena J F, Grabowski J and de Lucas J 2012 Superposition rules for higher-order differential equations, and their applications *J. Phys. A: Math. Theor.* **45** 185202
- [17] Cariñena J F, Grabowski J, de Lucas J and Sardón C 2014 Dirac–Lie systems and Schwarzian equations *J. Differ. Equ.* **257** 2259–752
- [18] Cariñena J F, Grabowski J and Marmo G 2000 *Lie–Scheffers Systems: A Geometric Approach* (Naples: Bibliopolis)
- [19] Cariñena J F, Grabowski J and Marmo G 2007 Superposition rules, Lie theorem and partial differential equations *Rep. Math. Phys.* **60** 237–58
- [20] Cariñena J F and de Lucas J 2009 Applications of Lie systems in dissipative Milne–Pinney equations *Int. J. Geom. Methods Mod. Phys.* **6** 683–99
- [21] Cariñena J F and Lucas J d 2011 Lie systems: theory, generalisations, and applications *Dissertationes Math.* **479** 1–162
- [22] Cariñena J F, de Lucas J and Sardón C 2013 Lie–Hamilton systems: theory and applications *Int. J. Geom. Methods Mod. Phys.* **10** 1350047
- [23] Chakravarty S and Halburd R 2003 First integrals of a generalized Darboux–Halphen systems *J. Math. Phys.* **44** 1751–62
- [24] Darboux G 1878 Sur la théorie des coordonnées curvilignes et les systèmes orthogonaux *Ann. Sci. Ecole Norm. Sup.* **7** 101–50
- [25] Echeverría-Enríquez A, Ibort A, Ibort A, Muñoz-Lecanda C M and Román-Roy N 2012 Invariant forms and automorphisms of locally homogeneous multisymplectic manifolds *J. Geom. Mech.* **4** 397–419
- [26] Echeverría-Enríquez A, Muñoz-Lecanda M C and Román-Roy N 1998 Multivector fields and connections: setting Lagrangian equations in field theories *J. Math. Phys.* **39** 4578–603
- [27] Echeverría-Enríquez A, Muñoz-Lecanda M C and Román-Roy N 2018 Remarks on multisymplectic reduction *Rep. Math. Phys.* **81** 415–24
- [28] Estévez P G, Herranz F J, de Lucas J and Sardón C 2016 Lie symmetries for Lie systems: applications to systems of ODEs and PDEs *Appl. Math. Comput.* **273** 435–52
- [29] Flores Espinoza R 2011 Periodic first integrals for Hamiltonian systems of Lie type *Int. J. Geom. Methods Mod. Phys.* **8** 1169–77
- [30] Forger M and Römer H 2001 A Poisson bracket on multisymplectic phase space *Rep. Math. Phys.* **48** 211–8
- [31] Gaset J, Gràcia X, Muñoz-Lecanda M C, Rivas X and Román-Roy N 2020 A contact geometry framework for field theories with dissipation *Ann. Phys., NY* **414** 168092
- [32] Grabowski J and de Lucas J 2013 Mixed superposition rules and the Riccati hierarchy *J. Differ. Equ.* **254** 179–98

- [33] Gràcia X, de Lucas J, Muñoz-Lecanda M C and Vilariño S 2019 Multisymplectic structures and invariant tensors for Lie systems *J. Phys. A: Math. Theor.* **52** 215201
- [34] Grundland A M, Martina L and Rideau G 1997 Partial differential equations with differential constraint *Advances in Mathematical Sciences-CRM's 25 Anniversary 11 (Procs. Lecture Notes)* (Providence, RI: American Mathematical Society) pp 135–54
- [35] Halphen G 1881 Sur un système d'équations différentielles *C.R. Acad. Sci. Paris* **92** 1101–3
- [36] Herranz F J, de Lucas J and Sardón C 2015 Jacobi–Lie systems: fundamentals and low-dimensional classification *The 10th AIMS Conf. Dynamical Systems, Differential Equations and Applications*
- [37] Herranz F J, De Lucas J and Tobolski M 2017 Lie–Hamilton systems on curved spaces: a geometrical approach *J. Phys. A: Math. Theor.* **50** 495201
- [38] Ibragimov N H 2009 Integration of systems of first-order equations admitting nonlinear superposition *J. Nonlinear Math. Phys.* **16** 137–47
- [39] James I M 1996 Reflections of the history of topology *Seminario Mat. Fis. Milano* **66** 87–96
- [40] Kijowski J 1973 A finite-dimensional canonical formalism in the classical field theory *Commun. Math. Phys.* **30** 99–128
- [41] Kijowski J and Tulczyjew W M 1979 *A Symplectic Framework for Field Theories (Lect. Notes Phys. 107)* (Berlin: Springer)
- [42] Leach P G L and Andriopoulos S K 2008 The Ermakov equation: a commentary *Appl. Anal. Discrete Math.* **2** 146–57
- [43] Leach P G L and Govinder K S 1999 On the uniqueness of the Schwarzian and linearisation by nonlocal contact transformation *J. Math. Anal. Appl.* **235** 84–107
- [44] Lee J M 2013 *Introduction to Smooth Manifolds* 2nd edn (Berlin: Springer)
- [45] de León M, Salgado M and Vilariño S 2015 *Methods of Differential Geometry in Classical Field Theories* (Singapore: World Scientific)
- [46] Lewandowski M M and de Lucas J 2017 Geometric features of Vessiot–Guldberg Lie algebras of conformal and Killing vector fields on  $\mathbb{R}^2$  *Banach Center Publ.* **113** 243–62
- [47] Lie S and Scheffers G 1893 *Vorlesungen über kontinuierliche Gruppen mit geometrischen und anderen Anwendungen* (Leipzig: Teubner)
- [48] de Lucas J and Sardón C 2020 *A Guide to Lie Systems with Compatible Geometric Structures* (Singapore: World Scientific)
- [49] de Lucas J and Vilariño S 2015  $k$ -symplectic Lie systems: theory and applications *J. Differ. Equ.* **258** 2221–55
- [50] de Lucas J and Zawora B M 2021 A time-dependent energy–momentum method *J. Geom. Phys.* **170** 104364
- [51] Marsden J E and Simo J C 1988 The energy momentum method *Act. Acad. Sci. Tau.* **1** 245–68
- [52] Milnor J 1976 Curvatures of left invariant metrics on Lie groups *Adv. Math.* **21** 293–329
- [53] Nikitin S 2000 Control synthesis for Čaplygin polynomial systems *Acta Appl. Math.* **60** 199–212
- [54] Ortega J P and Ratiu T S 2004 *Momentum Maps and Hamiltonian Reduction (Prog. Math. 222)* (Boston, MA: Birkhäuser)
- [55] Ovsienko V and Tabachnikov S 2005 *Projective Differential Geometry Old and New: From the Schwarzian Derivative to Cohomology of Diffeomorphism Groups* (Cambridge: Cambridge University Press)
- [56] Ovsienko V and Tabachnikov S 2009 What is the Schwarzian derivative *Not. AMS* **56** 34–6
- [57] Palais R S 1957 A global formulation of the Lie theory of transformation groups *Mem. Am. Math. Soc.* **22** 1–123
- [58] Ramos A 2006 New links and reductions between the Brockett nonholonomic integrator and related systems *Rend. Semin. Mat. Univ. Politec. Torino* **64** 39–54
- [59] Temple G 1960 A superposition principle for ordinary nonlinear differential equations *Lectures on Topics in Nonlinear Differential Equations* Report 1415 (Carderock: David Taylor Model Basin) pp 1–15
- [60] Vessiot E 1904 Sur la théorie de Galois et ses diverses généralisations *Ann. Sci. Ecole Norm. Sup.* **21** 9–85
- [61] Winternitz P 1983 Lie groups and solutions of nonlinear differential equations *Nonlinear Phenomena (Lecture Notes Phys. 189)* (Berlin: Springer) pp 263–331
- [62] Yoo W 2015 The automorphisms of a Lie algebra *Appl. Math. Sci.* **9** 121–7