

The Equidistant Dimension of Graphs

A. González¹ **D** · C. Hernando² **D** · M. Mora² **D**

Received: 23 July 2021 / Revised: 22 March 2022 / Accepted: 13 April 2022 © The Author(s) 2022

Abstract

A subset *S* of vertices of a connected graph *G* is a *distance-equalizer set* if for every two distinct vertices $x, y \in V(G) \setminus S$ there is a vertex $w \in S$ such that the distances from x and y to w are the same. The *equidistant dimension* of *G* is the minimum cardinality of a distance-equalizer set of G. This paper is devoted to introduce this parameter and explore its properties and applications to other mathematical problems, not necessarily in the context of graph theory. Concretely, we first establish some bounds concerning the order, the maximum degree, the clique number, and the independence number, and characterize all graphs attaining some extremal values. We then study the equidistant dimension of several families of graphs (complete and complete multipartite graphs, bistars, paths, cycles, and Johnson graphs), proving that, in the case of paths and cycles, this parameter is related to 3-AP-free sets. Subsequently, we show the usefulness of distance-equalizer sets for constructing doubly resolving sets.

Keywords Distance-equalizer set \cdot Equidistant dimension \cdot Resolving set \cdot Doubly resolving set \cdot Metric dimension

Mathematics Subject Classification 05C12 · 05C69

Communicated by Rosihan M. Ali.

A. González gonzalezh@us.es

> C. Hernando carmen.hernando@upc.edu

M. Mora merce.mora@upc.edu

¹ Department of Didactics of Mathematics, Universidad de Sevilla, Seville, Spain

² Departament de Matemàtiques, Universitat Politècnica de Catalunya, Barcelona, Spain

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No. 734922.

1 Introduction

The notion of resolving set, also known as locating set, was introduced by Slater [40] and, independently, by Harary and Melter [28]. This concept arises in diverse areas, including location problems in networks of different nature (see [11]). For example, in order to locate a failure in a computer network modeled as a graph, we are interested in a subset of vertices S such that every vertex of the underlying graph might be uniquely determined by its vector of distances to the vertices of S. Such a set is called a *resolving set* of the graph, and the *metric dimension* of that graph is the minimum cardinality of a resolving set.

Resolving sets and several related sets, such as identifying codes, locatingdominating sets or watching systems, have been widely studied during the last decades (see [3, 10, 18, 35]), as well as doubly resolving sets, a type of subset of vertices more restrictive than resolving sets with multiple applications in different areas [11, 12, 15, 26, 31-33]. However, many recent papers [13, 16, 17, 24, 41, 43, 44] have turned their attention precisely in the opposite direction to resolvability, thus trying to study anonymization problems in networks instead of location aspects. For instance, the need to ensure privacy and anonymity in social networks makes necessary to develop graph tools such as the concepts of antiresolving set and metric antidimension, introduced by Trujillo-Rasua and Yero [41]. Indeed, a subset of vertices A is a 2-antiresolving set if, for every vertex $v \notin A$, there exists another different vertex $w \notin A$ such that v and w have the same vector of distances to the vertices of A. The 2-metric antidimension of a graph is the minimum cardinality among all its 2-antiresolving sets. With the same spirit, this paper introduces new graph concepts that can also be applied to anonymization problems in networks: distance-equalizer set and equidistant dimension. Furthermore, we shall see that these concepts have concrete applications in mathematical problems, such as obtaining new bounds on the size of doubly resolving sets of graphs, as well as a new formulation in terms of graphs of a classical problem of number theory.

The paper is organized as follows. In Sect. 2, we define distance-equalizer sets and the equidistant dimension and show bounds in terms of other graph parameters: order, diameter, maximum degree, independence number, and clique number. Section 3 is devoted to characterize all graphs attaining some extremal values of the equidistant dimension. In Sect. 4 we study this parameter for some families of graphs: complete and complete multipartite graphs, bistars, paths, cycles, and Johnson graphs. For the particular cases of paths and cycles, we show that this parameter is related to 3-AP-free sets. In Sect. 5, we obtain bounds for general graphs and trees on the minimum cardinality of doubly resolving sets in terms of the equidistant dimension. Finally, we present some conclusions and open problems in Sect. 6.

All graphs considered in this paper are connected, undirected, simple, and finite. The vertex set and the edge set of a graph G are denoted by V(G) and E(G), respectively.

The order of G is |V(G)|. For any vertex $v \in V(G)$, its open neighborhood is the set $N(v) = \{w \in V(G) : vw \in E(G)\}$ and its closed neighborhood is $N[v] = N(v) \cup \{v\}$.



Fig. 1 Black vertices form a distance-equalizer set of minimum size for P_8

The *degree* of a vertex v, denoted by deg(v), is defined as the cardinality of N(v). If deg(v) = 1, then we say that v is a *leaf*, in which case the only vertex adjacent to v is called its *support vertex*.

When deg(v) = |V(G)| - 1, we say that v is *universal*. The *maximum degree* of G is $\Delta(G) = \max \{ \deg(v) : v \in V(G) \}$ and its *minimum degree* is $\delta(G) = \min \{ \deg(v) : v \in V(G) \}$. The *distance* between two vertices $v, w \in V(G)$ is denoted by d(v, w), and the *diameter* of G is $D(G) = \max\{d(v, w) : v, w \in V(G)\}$. A *clique* is a subset of pairwise adjacent vertices and the *clique number* of G, denoted by $\omega(G)$, is the maximum cardinality of a clique of G. An *independent set* of G is a subset of pairwise non-adjacent vertices and the *independence number* of G, denoted by $\alpha(G)$, is the maximum cardinality of an independent set of G. For undefined terms we refer the reader to [42].

For every integer $n \ge 1$, let $[n] = \{1, 2, ..., n\}$. We denote by P_n the *path* of order n with vertex set [n] and edges ij with j = i + 1 and $i \in [n-1]$, and by C_n the *cycle* of order $n, n \ge 3$, with vertex set [n] and the same edge set as P_n together with the edge 1n. Also, for every $r, s \ge 3$, we denote by $K_{1,r}$ the *star* on r + 1 vertices with vertex set $\{v\} \cup [r]$, vertex v being called the *center* of the star, and edge set $\{vi : i \in [r]\}$. A *bistar*, denoted by $K_2(r, s)$, is a graph obtained by joining the centers of two stars $K_{1,r-1}$ and $K_{1,s-1}$.

2 Distance-Equalizer Sets and Equidistant Dimension

Let x, y, w be vertices of a graph G. We say that w is equidistant from x and y if d(x, w) = d(y, w). A subset S of vertices is called a *distance-equalizer set* for G if for every two distinct vertices $x, y \in V(G) \setminus S$ there exists a vertex $w \in S$ equidistant from x and y. The equidistant dimension of G, denoted by eqdim(G), is the minimum cardinality of a distance-equalizer set of G. For example, if v is a universal vertex of a graph G, then $S = \{v\}$ is a minimum distance-equalizer set of G, and so eqdim(G) = 1. Also, a distance-equalizer set of P_8 is shown in Fig. 1, and it can be easily checked that P_8 has no distance-equalizer set of size at most 4, so $eqdim(P_8) = 5$.

The following results are immediate but make it easier to prove subsequent results.

Lemma 1 Let G be a graph. If S is a distance-equalizer set of G and v is a support vertex of G, then S contains v or all leaves adjacent to v. Consequently,

$$eqdim(G) \ge | \{v \in V(G) : v \text{ is a support vertex} \} |$$

Proof No vertex is equidistant from a leaf and its support vertex, since every path from a leaf to any other vertex goes through its support vertex. Hence, if v is not in S, then all leaves hanging from v must be in S.

Recall that a graph G is *bipartite* whenever V(G) can be partitioned into two independent sets, say A, B, which are called its *partite sets*.

Proposition 1 Let G be a bipartite graph with partite sets A and B. If S is a distanceequalizer set of G, then $A \subseteq S$ or $B \subseteq S$. Consequently, $eqdim(G) \ge min\{|A|, |B|\}$.

Proof The distance between two vertices in the same partite set is even, while the distance between vertices of different partite sets is odd. Hence, there is no vertex equidistant from two vertices belonging to different partite sets. Therefore, $A \subseteq S$ or $B \subseteq S$.

If G is a graph of order n, with $n \ge 2$, then any set of vertices of cardinality n - 1 is obviously a distance-equalizer set. Hence, n - 1 is an immediate upper bound on the equidistant dimension of nontrivial graphs. We next prove some upper bounds involving classical graph parameters.

Proposition 2 For every graph G of order $n \ge 2$, the following statements hold.

- (i) $eqdim(G) \le n \Delta(G)$ and the bound is tight if $\Delta(G) \ge n/2$;
- (ii) $eqdim(G) \le n \omega(G)$, if $G \ncong K_n$, and the bound is tight if $\omega(G) \le n/2$;
- (iii) $eqdim(G) \leq \frac{n(D(G)-1)+1}{D(G)}$, and the bound is tight if D(G) = 2;
- (iv) $eqdim(G) \le n \alpha(G)$, whenever D(G) = 2, and the bound is tight if $\alpha(G) \ge n/2$.
- **Proof** (i) Let v be a vertex of degree $\Delta(G)$. It is easy to see that the set $S = V(G) \setminus N(v)$ is a distance-equalizer set of cardinality $n \Delta(G)$, and so $eqdim(G) \le n \Delta(G)$.

To prove tightness, let $H_{a,b}$ with $a \ge 1$ and $0 \le b < a$ be the graph with vertex set $\{v, v_1, \ldots, v_a, u_1, \ldots, u_b\}$ and edge set $\{vv_i : i \in [a]\} \cup \{v_iu_i : i \in [b]\}$. This graph has order a + b + 1 and maximum degree a, so $eqdim(H_{a,b}) \le b + 1$ as we have just seen. Moreover, this graph is bipartite with partite sets $A = \{v, u_1, \ldots, u_b\}$ and $B = \{v_1, \ldots, v_a\}$, and so by Proposition 1 we have that $eqdim(H_{a,b}) \ge min\{|A|, |B|\} = b + 1$ since b < a. Hence, $H_{a,b}$ is a graph of order a+b+1 and maximum degree a, with $a \ge \frac{a+b+1}{2}$ and $eqdim(H_{a,b}) = b+1$, showing that the given the bound is tight.

(ii) Suppose that G is a non complete graph of order n. Let W be a clique of maximum size ω(G). Thus, |W| = ω(G) < n. Since G is connected, there is at least one vertex u ∉ W adjacent to a vertex w ∈ W. Hence, Δ(G) ≥ deg(w) ≥ |W| = ω(G) and, by item (i), eqdim(G) ≤ n − Δ(G) ≤ n − ω(G).

To prove that the bound is tight for $\omega(G) \le n/2$, consider for example the graph $G_{r,s}$ of order n = r+s, for every $r \ge 2$ and $1 \le s \le r$, obtained by attaching a leaf to *s* vertices of a complete graph of order *r*. Then, by Lemma 1, $eqdim(G_{r,s}) \ge s$. Since the set formed by all except one leaves together with the support vertex of the remaining leaf is a distance-equalizer set, we have $eqdim(G_{r,s}) = s = n - r = n - \omega(G_{r,s})$.

(iii) Let v be a vertex of G for which there exists another vertex at distance D(G). For every $i \in [D(G)]$, let V_i be the set of vertices at distance i from v, and observe that all vertices in V_i are equidistant from v. Also, there must exist a set V_{i_0} , with $1 \le i_0 \le D$, having at least $\frac{n-1}{D(G)}$ vertices. Therefore, $V(G) \setminus V_{i_0}$ is a distanceequalizer set, and consequently $eqdim(G) \le n - \frac{n-1}{D(G)} = \frac{n(D(G)-1)+1}{D(G)}$. To prove that the bound is tight, consider the complete bipartite graph $K_{r,r}$,

To prove that the bound is tight, consider the complete bipartite graph $K_{r,r}$, for every $r \ge 2$. These graphs have diameter 2 and, as we will see later, $eqdim(K_{r,r}) = r$. Hence, $eqdim(K_{r,r}) = r = \lfloor \frac{2r(2-1)+1}{2} \rfloor$.

(iv) Let W be an independent set of cardinality α(G). Then, the set S = V\W is a distance-equalizer set. Indeed, note that the vertices not in S are the vertices in W. If v, w ∈ W, then d(v, w) = 2, because v, w are not adjacent and D(G) = 2. Hence, there is a vertex u ∉ W such that d(u, v) = d(u, w) = 1. Thus, S is a distance-equalizer set and we obtain eqdim(G) ≤ |S| = n - α(G). Eigenvertex and the vertice area before when a set of the vertice of the vertice of the vertex with a ≥ n. Then K we have

Finally, consider any complete bipartite graph $K_{r,s}$, with $s \ge r$. Then, $K_{r,s}$ has order n = r + s, the independence number is $\alpha(K_{r,s}) = s$ and, as we will see later, $eqdim(K_{r,s}) = r$. Hence, $eqdim(K_{r,s}) = n - \alpha(K_{r,s})$.

The bound given in Proposition 2(i) is not tight for all values of $\Delta(G)$ and n, for example when $\Delta(G) = 2$ and $n \ge 7$. Indeed, the only graphs satisfying $\Delta(G) = 2$ are paths and cycles and, as it will be seen below, the equidistant dimension of paths and cycles of order $n \ge 7$ is at most n - 3.

3 Extremal Values

In this section we characterize all nontrivial graphs achieving extremal values for the equidistant dimension, concretely, graphs *G* of order $n \ge 2$ such that $eqdim(G) \in \{1, 2, n-2, n-1\}$. We also derive a Nordhaus–Gaddum type bound for the equidistant dimension.

Theorem 1 For every graph G of order $n \ge 2$, the following statements hold.

(i) eqdim(G) = 1 if and only if $\Delta(G) = n - 1$;

- (ii) eqdim(G) = 2 if and only if $\Delta(G) = n 2$.
- **Proof** (i) If $\Delta(G) = n 1$, then $eqdim(G) \le n (n 1) = 1$ by Proposition 2(i), and so eqdim(G) = 1. Conversely, if eqdim(G) = 1, then there exists a vertex vsuch that $S = \{v\}$ is a distance-equalizer set of G. We claim that v has degree n - 1. Indeed, suppose on the contrary that there is a vertex u not adjacent to v. Then, since there is at least one vertex w adjacent to v, we have $d(v, w) = 1 \ne d(v, u)$. Hence, $\{v\}$ is not a distance-equalizer set of G, a contradiction. Therefore, G has maximum degree n - 1.
- (ii) If G has maximum degree n 2, then eqdim(G) ≤ n (n 2) = 2 by Proposition 2(i), and eqdim(G) ≠ 1 by item (i). Hence, eqdim(G) = 2. Conversely, suppose that eqdim(G) = 2 and let S = {u, v} be a distance-equalizer set of G. We first prove that N(u)\{v} ⊆ N(v)\{u} or N(v)\{u} ⊆ N(u)\{v}. Suppose on the contrary that N(u)\{v} ⊈ N(v)\{u} and N(v)\{u} ⊈ N(u)\{v}. Then, there exist vertices x, y such that x ∈ N(u)\{v} and x ∉ N(v)\{u}, y ∈ N(v)\{u} and y ∉ N(u)\{v}. Thus, vertices x, y verify that d(x, u) = 1 ≠ d(y, u) and

 $d(y, v) = 1 \neq d(x, v)$, contradicting that S is a distance-equalizer set. Without loss of generality, we assume $N(v) \setminus \{u\} \subseteq N(u) \setminus \{v\}$.

Next, we prove that $V(G) = N[u] \cup \{v\}$. Suppose on the contrary that there is a vertex $z \notin N[u] \cup \{v\}$. Thus, $d(z, u) \ge 2$ and $d(z, v) \ge 2$. If $N(v) \setminus \{u\}$ is nonempty, then for any vertex $x \in N(v) \setminus \{u\}$ we have $d(x, u) = 1 \neq d(z, u) \ge 2$ and $d(x, v) = 1 \neq d(z, v) > 2$, contradicting that *S* is a distance-equalizer set.

Otherwise $N(v)\setminus\{u\}$ is empty, and so v is a leaf with u as its support vertex. Thus, for any $y \in N(u)\setminus\{v\}$, we have $d(y, u) = 1 \neq d(z, u) \geq 2$ and $d(y, v) = 2 \neq d(z, v) = d(z, u) + 1 \geq 3$, contradicting again that S is distance-equalizer.

Finally, we have that $\Delta(G) \neq n-1$ by the preceding item. Hence, $v \notin N(u)$ and eqdim(G) = deg(u) = n-2.

Theorem 2 For any graph G of order n, the following statements hold.

- (i) If $n \ge 2$, then eqdim(G) = n 1 if and only if G is a path of order 2.
- (ii) If $n \ge 3$, then eqdim(G) = n 2 if and only if $G \in \{P_3, P_4, P_5, P_6, C_3, C_4, C_5\}$.
- **Proof** (i) It is obvious that $eqdim(P_2) = 1$. Conversely, if G is a graph with eqdim(G) = n 1, then $\Delta(G) = 1$ by Proposition 2(i), and the only connected graph with maximum degree equal to 1 is the path of order 2.
- (ii) A straightforward computation shows that the graphs P_3 , P_4 , P_5 , P_6 , C_3 , C_4 , and C_5 have equidistant dimension equal to the order minus 2. Conversely, if *G* is a graph with eqdim(G) = n 2, then $\Delta(G) \le 2$ by Proposition 2(i). As we have seen above, the path of order 2 is the only connected graph with maximum degree 1. Hence, $\Delta(G) = 2$, that is, *G* is a path or a cycle of order at least 3.

It is easy to see that in both cases the set $[n]\setminus\{1, 3, 7\}$ is a distance-equalizer set whenever $n \ge 7$, and so $eqdim(G) \le n-3$ but eqdim(G) = n-2. Therefore, $3 \le n \le 6$ and consequently $G \in \{P_3, P_4, P_5, P_6, C_3, C_4, C_5\}$ since $eqdim(C_6) = 3 \ne 6-2$.

Corollary 1 If G is a graph of order $n \ge 7$, then $1 \le eqdim(G) \le n - 3$.

Now, we provide a Nordhaus–Gaddum type bound on the equidistant dimension. Nordhaus–Gaddum type inequalities establish bounds on the sum of a parameter for a graph and its complement. Recall that the *complement* of a graph G, denoted by \overline{G} , is the graph on the same vertices as G and two vertices are adjacent in \overline{G} if and only if they are not adjacent in G. Also, a graph G is *doubly connected* if both G and \overline{G} are connected. Note that nontrivial doubly connected graphs have order at least 4.

The following result is a direct consequence of Proposition 2(i).

Proposition 3 If G is a doubly connected graph, then $eqdim(\overline{G}) \leq \delta(G) + 1$.

Theorem 3 If G is a doubly connected graph of order $n \ge 4$, then $4 \le eqdim(G) + eqdim(\overline{G}) \le n + 1$. Moreover, these bounds are tight.

Proof First observe that a graph G satisfying eqdim(G) = 1 is not doubly connected. Indeed, in such a case, by Theorem 1(i), it contains a universal vertex v that is an isolated vertex in \overline{G} . Hence, $eqdim(\overline{G}) \ge 2$ and $eqdim(\overline{G}) \ge 2$, whenever G is doubly connected, and the lower bound follows. The family of bistars $G = K_2(2, n - 2)$, $n \ge 4$, provides examples of graphs attaining the lower bound for every $n \ge 4$. As we will see in Theorem 4(iii) below, these graphs satisfy eqdim(G) = 2, and it is easy to check that $eqdim(\overline{G}) = 2$.

The upper bound is a direct consequence of Propositions 2(i) and 3, because

$$eqdim(G) + eqdim(\overline{G}) \le n - \Delta(G) + \delta(G) + 1 \le n + 1.$$

The cycle C_5 attains the upper bound, since $\overline{C_5} = C_5$ and, by Theorem 2(ii), $eqdim(C_5) = 3$.

4 Equidistant Dimension of Some Families of Graphs

In this section we study the equidistant dimension of some families of graphs, concretely of complete, complete bipartite and complete multipartite graphs, bistars, paths, cycles, and Johnson graphs.

4.1 Complete Graphs, Complete Multipartite Graphs, and Bistars

Recall that, for every positive integer *n*, the *complete graph* K_n is the graph of order *n* in which every pair of vertices is connected by an edge. Also, the *complete bipartite graph* $K_{r,s}$, with *r*, *s* positive integers, is the bipartite graph with partite sets *A*, *B* such that |A| = r and |B| = s, and edge set given by all pairs vu with $v \in A$ and $u \in B$. More generally, a *complete p-partite* graph, denoted by $K_{n_1,...,n_p}$, is a graph with set of vertices $A_1 \cup \cdots \cup A_p$ such that A_1, \ldots, A_p , which are called its *partite sets*, are pairwise disjoint, verify $|A_i| = n_i \ge 1$, and two vertices are adjacent if and only if they belong to A_i and A_j , respectively, with $i \ne j$. Note that complete bipartite graphs.

Theorem 4 Let $n, r, s, p, n_1, ..., n_p$ be positive integers such that $n \ge 2$, $s \ge r$, $p \ge 3$ and $n_p \ge \cdots \ge n_1 \ge 1$. Then, the following statements hold.

- (i) $eqdim(K_n) = 1;$
- (ii) $eqdim(K_{r,s}) = r;$
- (iii) $eqdim(K_2(r, s)) = r;$
- (iv) $eqdim(K_{n_1,...,n_p}) = \min\{n_1, 3\}.$

Proof (i) It is a direct consequence of Theorem 1(i).

- (ii) By Proposition 1, $eqdim(K_{r,s}) \ge r$. Since both partite sets of $K_{r,s}$ are distanceequalizer sets, we have $eqdim(K_{r,s}) = r$.
- (iii) Note that $K_2(r, s)$ is a bipartite graph with partite sets of cardinality r and s. Hence, by Proposition 1, $eqdim(K_2(r, s)) \ge r$. Moreover, if A is the partite set

of cardinality r, then A is a distance-equalizer set because it contains a vertex at distance 1 from every vertex not in A.

(iv) If $n_1 = 1$, then $K_{n_1,...,n_p}$ has a universal vertex and $eqdim(K_{n_1,...,n_p}) = 1$, by Theorem 1(i).

If $n_1 = 2$, then $K_{n_1,...,n_p}$ has maximum degree equal to the order minus 2 and $eqdim(K_{n_1,...,n_p}) = 2$, by Theorem 1(ii).

Otherwise $n_1 \ge 3$, and so the maximum degree of $K_{n_1,...,n_p}$ is at most the order minus 3 and $eqdim(K_{n_1,...,n_p}) \ge 3$, by Theorem 1. Moreover, it is very easy to verify that any set consisting of 3 vertices from different partite sets is distance-equalizer. Thus, we conclude that $eqdim(K_{n_1,...,n_p}) = 3$.

4.2 Paths

We next show that distance-equalizer sets and the equidistant dimension of paths are related to 3-AP-free sets and the function r(n) introduced by Erdös and Turán [23]. A subset $S \subseteq [n]$ is 3-AP-free if $a + c \neq 2b$, for every distinct terms $a, b, c \in S$. The largest cardinality of a 3-AP-free subset of [n] is denoted by r(n).

We begin by introducing some preliminary results. A subset of [n] is called *even-sum* if all its elements have the same parity.

Proposition 4 Let $S \subseteq [n]$ for some integer n. Then, S is a distance-equalizer set of P_n if and only if $[n] \setminus S$ is a 3-AP-free even-sum set.

Proof Let us denote by *A* the set of vertices of P_n labeled with even numbers and by *B* the set of vertices labeled with odd numbers. (Note that P_n is a bipartite graph and *A*, *B* are its partite sets). Also, let *S* be a distance-equalizer set of P_n with |S| = r. By Proposition 1, either $A \subseteq S$ or $B \subseteq S$. Thus, if $T = [n] \setminus S = \{t_1, \ldots, t_{n-r}\}$, then t_1, \ldots, t_{n-r} have the same parity, that is, *T* is an even-sum set. Moreover, $(t_i + t_j)/2$ is the only vertex of P_n equidistant from t_i and t_j . Hence $(t_i + t_j)/2 \in S$, that is, $(t_i+t_j)/2 \notin T$. Then, *T* is a 3-*AP*-*free* set. Conversely, suppose that $T = \{t_1, \ldots, t_{n-r}\}$ is a 3-*AP*-free even-sum set. Then, for all pair of vertices t_i, t_j of *T*, we have $(t_i+t_j)/2 \in [n] \setminus T$. Hence, $S = [n] \setminus T$ is a distance-equalizer set of P_n .

Corollary 2 For every positive integer n, it holds that

 $eqdim(P_n) = n - max\{|T| : T \text{ is a 3-AP-free even-sum subset of } [n]\}.$

Proposition 5 [19] Let k_1, \ldots, k_r , *n* be different positive integers. Then, one of the sets $\{2k_1 - 1, 2k_2 - 1, \ldots, 2k_r - 1\}$ or $\{2k_1, 2k_2, \ldots, 2k_r\}$ is a 3-AP-free even-sum set of [n] if and only if $\{k_1, \ldots, k_r\}$ is a 3-AP-free subset of $\lceil n/2 \rceil$].

The equidistant dimension of a path is derived from the results above.

Theorem 5 For every positive integer n, it holds that

$$eqdim(P_n) = n - r\left(\left\lceil \frac{n}{2} \right\rceil\right).$$

Hence, obtaining the equidistant dimension of paths amounts to computing the function r(n), which has been widely studied [5, 7–9, 22, 25, 27, 29, 37–39]. In fact, many papers are devoted to obtain the values of r(n) in some specific cases ($n \le 23$ and n = 41 [23]; $n \le 27$ and $41 \le n \le 43$ [39]; $n \le 123$ [21]), which allows us to compute $eqdim(P_n)$ in all those cases (see Table 1 for $n \le 20$ and n = 50). Also, other works [5, 21, 37] provide bounds on r(n) that are useful to approach $eqdim(P_n)$, such as

$$n^{1-c/\sqrt{\log n}} < r(n) < \frac{cn}{\log \log n}.$$

Besides its relationship with the function r(n), the equidistant dimension of paths is also related to a problem concerning covering squares of a chessboard by queens proposed by Cockayne and Hedetniemi [19]. Indeed, the authors are interested in determining the minimum number of queens needed to be placed on the major diagonal of a chessboard in order to reach all the remaining squares with a single chess movement. More formally, a subset $K \subseteq [n]$ is a *diagonal dominating set* if its |K| queens placed in position $\{(k, k) | k \in K\}$ on the black major diagonal of an $n \times n$ chessboard cover the entire board. The minimum cardinality of a diagonal dominating set is denoted by diag(n). It is proved in [19] that diagonal dominating sets are precisely the complements of 3-AP-free even-sum sets, which combined with Proposition 4 leads us to see that the distance-equalizer sets of P_n are the diagonal dominating sets in [n], and consequently $eqdim(P_n) = diag(n)$.

Finally, we do not know the exact value of the equidistant dimension of trees. However, in this family of graphs, it looks that paths are those graphs needing more vertices to construct a distance-equalizer set. Indeed, it is easily seen that, for every pair of vertices of a path, there is at most one equidistant vertex. Hence, we believe that the following conjecture holds true.

Conjecture 1 If T is a tree of order n, then $eqdim(T) \le eqdim(P_n)$.

4.3 Cycles

In this section, the equidistant dimension of cycles of even order is completely determined, while for cycles of odd order, lower and upper bounds in terms of r(n) are given.

Theorem 6 For every positive integer $n \ge 3$, the following statements hold.

(i)
$$eqdim(C_n) = \begin{cases} \frac{n}{2}, & \text{for } n \text{ even, } n \neq 0 \mod 4; \\ \frac{3n}{4} - 1, \text{ for } n \text{ even, } n \equiv 0 \mod 4. \end{cases}$$

(ii) $\frac{n-1}{2} \le eqdim(C_n) \le n - r\left(\left\lceil \frac{n+1}{4} \right\rceil\right), \text{ for } n \text{ odd.}$

Proof Throughout this proof, for every $i, j \in [n]$, we use the expression $\frac{i+j+n}{2}$ to represent the only integer in [n] modulo n whenever $\frac{i+j+n}{2}$ is an integer. Thus, for

every pair of vertices i, j of C_n , the vertices equidistant from them are $\frac{i+j}{2}$ and $\frac{i+j+n}{2}$, whenever these values are integers. Hence, there is exactly one vertex equidistant from i and j, when n is odd; there is no equidistant vertex from i and j, whenever n is even and i, j have distinct parity; and there are exactly two vertices equidistant from i and j, if n even and i, j have the same parity. Moreover, in this last case, the vertices equidistant from i and j are antipodal.

- (i) Let n be an even integer, and let us denote by A (resp., B) the set of vertices of C_n labeled with odd (resp., even) numbers. As n is even, C_n is a bipartite graph and, by Proposition 1, for every distance-equalizer set S, either A ⊆ S or B ⊆ S. Hence, |S| ≥ n/2. We distinguish two subcases.
 - (a) Case $n \neq 0 \mod 4$. We claim that *A* is a distance-equalizer set (see an example in Fig. 2a). Indeed, for every $i, j \in [n] \setminus A$, the numbers $\frac{i+j}{2}$ and $\frac{i+j+n}{2}$ are integers of different parity, because *n* is even but $n \neq 0 \mod 4$. Thus, either $\frac{i+j}{2}$ or $\frac{i+j+n}{2}$ belongs to *A*. Hence, *A* is a distance-equalizer set and $eqdim(C_n) = n/2$.
 - (b) Case n ≡ 0 mod 4. Let S be a distance-equalizer set, and let us assume, relabeling the vertices if necessary, that A ⊆ S. Thus, [n]\S ⊆ B. First, we suppose that there is a pair of antipodal vertices in [n]\S ⊆ B. We can assume without loss of generality that these vertices are n/2 and n. For every i ∈ {2, 4, ..., n/2 2}, the only vertices equidistant from i and n i are n/2 and n. Since n/2 and n are not in S, we derive that at least one of the vertices i or n i must be in S, for every i ∈ {2, 4, ..., n/2 2}. Therefore, besides the vertices from A, the set S contains at least net not negative from B. Therefore, begins a for the vertices from B. Therefore, network of the vertices from B. T

$$eqdim(C_n) \ge \frac{n}{2} + \frac{n/2 - 2}{2} = \frac{3n}{4} - 1.$$

It is straightforward that the same bound holds if we suppose that there is no pair of antipodal vertices in $[n] \setminus S \subseteq B$.

Now, we consider the set $S = [n] \setminus \{2, 4, 6, ..., n/2 + 2\}$ of size $|S| = \frac{3n}{4} - 1$. It is easy to check that every pair of vertices not in *S* has a vertex in $\{n/2 + 3, n/2 + 4, ..., n, 1\} \subseteq S$ equidistant from them. Thus, we conclude that *S* is a distance-equalizer set of minimum cardinality (see an example in Fig. 2b), and so $eqdim(C_n) = \frac{3n}{4} - 1$.

(ii) Let *n* be an odd integer, and let *S* be a distance-equalizer set of minimum size. Since any set of cardinality *n*−1 is a distance-equalizer set, we can assume without loss of generality that *n* ∉ *S*. As *n* is an odd integer, *n* is the only vertex of *C_n* equidistant from each pair of vertices *i*, *n*−*i*, with *i* ∈ {1, ..., (*n*−1)/2}, and so at least one of them must be in *S*. Therefore, eqdim(*C_n*) ≥ (*n*−1)/2. To prove the upper bound, let *S*₁ = {*i* : (*n*+1)/2 < *i* ≤ *n*} and consider a distance-equalizer set *S*₂ of *P*_{(*n*+1)/2}. We claim that *S* = *S*₁ ∪ *S*₂ is a distance-equalizer set *S*₂ of *P*_{(*n*+1)/2}.

equalizer set of C_n (see an example in Fig. 2c). Indeed, any two vertices *i*, *j* not in *S* belong to [(n + 1)/2], and there is a vertex in S_2 equidistant from then, since

Table 1 Equidistant dimensionof some paths and cycles



Fig. 2 The set of black vertices form a distance-equalizer set of the cycles **a** C_{10} , **b** C_{12} , and **c**, **d** C_{13} . In cases (**a**, **b**, **d**), the given distance-equalizer set has minimum cardinality

n	3	4	5	6	7	8	9	10	11	12
$r(\lceil n/2 \rceil)$	2	2	2	2	3	3	4	4	4	4
$eqdim(P_n)$	1	2	3	4	4	5	5	6	7	8
$eqdim(C_n)$	1	2	3	3	4	5	5	5	7	8
n	13	14	1:	5	16	17	18	19	20	50
$r(\lceil n/2 \rceil)$	4	4	4		4	5	5	5	5	10
$eqdim(P_n)$	9	10	1	1	12	12	13	14	15	40
$eqdim(C_n)$	9	7	1	1	11	12	9	13	14	25

 S_2 is a distance-equalizer set of $P_{(n+1)/2}$. Hence,

$$eqdim(C_n) \le |S_1| + |S_2| = \frac{n-1}{2} + eqdim(P_{\frac{n+1}{2}}) = n - r\left(\left\lceil \frac{n+1}{4} \right\rceil\right).$$

Note that the distance-equalizer set constructed in the proof of the preceding theorem for odd cycles is not necessarily of minimum cardinality. In Fig. 2c, d, the distance-equalizer set described in the proof of the theorem and a distance-equalizer set of minimum cardinality for C_{13} are shown.

In Table 1 the values of the equidistant dimension of C_n for $n \le 20$ and n = 50 are given. Note that some of these values have been obtained with computer.

4.4 Johnson Graphs

Johnson graphs are important because of their connections with other combinatorial structures such as projective planes and symmetric designs [3]. Furthermore, there exist different studies about geometric versions of these graphs because of their multiple applications in network design (see for instance [4]). Due to these facts, among others, properties of Johnson graphs have been widely studied in the literature: spectra [34], induced subgraphs [2], connectivity [1], colorings [6], distances [20], automorphisms [36], and metric dimension [3]. In this subsection we study the equidistant dimension of Johnson graphs, obtaining an upper bound for several cases.

The Johnson graph J(n, k), with $n > k \ge 1$, has as vertex set the k-subsets of a *n*-set and two vertices are adjacent if their intersection has size k - 1. Thus, it can be easily seen that the distance between any two vertices X, Y is given by

$$d(X, Y) = |X \setminus Y| = |Y \setminus X| = k - |X \cap Y|.$$

Consequently, a vertex $U \in V(J(n, k))$ is equidistant from vertices X and Y if and only if $|U \cap X| = |U \cap Y|$.

Proposition 6 For any positive integer k, it holds that

$$eqdim(J(n,k)) \leq n$$

whenever $n \in \{2k - 1, 2k + 1\}$ or $n > 2k^2$.

Proof Consider the vertices of J(n, k) as k-subsets of the n-set $W = \{0, ..., n - 1\}$. For each positive integer *i*, let $S_i = \{i, i + 1, ..., i + k - 1\} \in V(J(n, k))$ where sums are taken modulo *n* (thus $S_{i+rn} = S_i$ for any integer *r*). We claim that the set $S = \{S_0, ..., S_{n-1}\}$ is a distance-equalizer set of J(n, k). Suppose on the contrary the existence of two vertices $X, Y \in V(J(n, k)) \setminus S$ such that $|S_i \cap X| \neq |S_i \cap Y|$ for every $i \in \{0, 1, ..., n - 1\}$.

First, we assume that $n > 2k^2$. For j, r integers, let $T(j, r) = \{j, j + 1, ..., j + r\} \subseteq W$, where the sums are also taken modulo n. Let $\mathcal{T} = \{T_1, ..., T_s\}$ be the family of sets T(j, r) satisfying $T(j, r) \cap (X \cup Y) = \emptyset$, $j - 1 \in X \cup Y$ and $j + r + 1 \in X \cup Y$. Note that \mathcal{T} is a partition of $W \setminus (X \cup Y)$ with at most 2k parts, by construction. Moreover, $|T_i| \leq k - 1$ for every $i \in \{1, ..., s\}$, otherwise, if $T_i = T(j, r)$, where $r \geq k$, then $|S_j \cap X| = |S_j \cap Y| = 0$, which contradicts our hypothesis. Therefore,

$$n = |W| = |X \cup Y| + |W \setminus (X \cup Y)| \le 2k + 2k(k-1) = 2k^2,$$

contradicting our assumption on n.

Now, suppose $n \in \{2k - 1, 2k + 1\}$. Let $u = (u_0, \ldots, u_{n-1})$ be the vector of $\{-1, 1, 0\}^n$ such that $u_i = 1$ if $i \in X \setminus Y$; $u_i = -1$ if $i \in Y \setminus X$; and $u_i = 0$ otherwise. Observe that u has at most 2k nonzero components, and the same number of 1's and -1's. Hence, $\sum_{i=0}^{n-1} u_i = 0$. Let $s_i = \sum_{j=i}^{i+k-1} u_j$. Observe that $s_i = |S_i \cap X| - |S_i \cap Y|$, for every $i \in \{0, 1, \ldots, n-1\}$. Hence, $s_i \neq 0$, for every $i \in \{0, 1, \ldots, n-1\}$, because no set S_i is equidistant from X and Y. Next, we prove that $s_i s_{i+k} < 0$ for every $i \in \{0, \ldots, n-1\}$. Indeed, we have that

$$s_i + s_{i+k} + u_{i+2k} = \sum_{i=0}^{n-1} u_i = 0$$
, when $n = 2k + 1$;
 $s_i + s_{i+k} - u_i = \sum_{i=0}^{n-1} u_i = 0$, when $n = 2k - 1$.

🖉 Springer

Therefore, for n = 2k + 1,

$$s_{i+k} = -s_i - u_{i+2k} \le -1 - u_{i+2k} \le 0, \text{ if } s_i > 0;$$

$$s_{i+k} = -s_i - u_{i+2k} \ge 1 - u_{i+2k} \ge 0, \text{ if } s_i < 0$$

and for n = 2k - 1,

$$s_{i+k} = -s_i + u_i \le -1 + u_i \le 0$$
, if $s_i > 0$;
 $s_{i+k} = -s_i + u_i \ge 1 + u_i \ge 0$, if $s_i < 0$.

Hence, for every $i \in \{0, ..., n-1\}$, we have $s_i s_{i+k} < 0$ since $s_{i+k} \neq 0$, and it can be derived that $s_i s_{i+rk} < 0$, for r odd, and $s_i s_{i+rk} > 0$, for r even. Then, $s_i s_{i+nk} < 0$, since n is odd, which is a contradiction since $s_{i+nk} = s_i$.

5 Using Distance-Equalizer Sets for Constructing Doubly Resolving Sets

We now explore different relationships among distance-equalizer sets and doubly resolving sets. To do this, we first need to formally define resolving sets. Indeed, a subset S of vertices is a resolving set of a graph G if, for every pair of vertices $x, y \in V(G)$, there exists a vertex $v \in S$ such that $d(v, x) \neq d(v, y)$. The metric dimension of G, denoted by dim(G), is the minimum cardinality of a resolving set of G. Observe that, on the one hand, a set of vertices can be at the same time resolving set and distance-equalizer set. For example, it is easy to check that any independent set S of cardinality three of a cycle of order 6 is both resolving and distance-equalizer. However, in this case S is a distance-equalizer set of minimum cardinality, but S is not a resolving set of minimum cardinality, since $eqdim(C_6) = 3$ and $dim(C_6) = 3$ 2. On the other hand, there are graphs satisfying dim(G) = eqdim(G) with no minimum resolving set being a distance-equalizer set. For example, the cycle or order 4 satisfies $eqdim(C_4) = dim(C_4) = 2$, but there is no set of cardinality two that is both resolving and distance-equalizer because a resolving set on two vertices is formed by two adjacent vertices and distance-equalizer sets of cardinality two are formed by two non-adjacent vertices.

Doubly resolving sets were introduced in [11] as a tool for computing the metric dimension of Cartesian products of graphs. Furthermore, different authors have provided interesting applications of doubly resolving sets on source location [15, 31], algorithmic studies and relations with other graph parameters [12, 26, 32, 33].

We say that two vertices u, v doubly resolve a pair of vertices x, y of G (or that $\{x, y\}$ is doubly resolved by u, v) if $d(u, x) - d(u, y) \neq d(v, x) - d(v, y)$. A set $S \subseteq V(G)$ is a doubly resolving set of G if every pair $\{x, y\} \subseteq V(G)$ is doubly resolved by two vertices of S (it is said that S doubly resolves $\{x, y\}$), and the minimum cardinality of such a set is denoted by $\psi(G)$. Observe that a doubly resolving set is also a resolving set, and so $dim(G) \leq \psi(G)$.

Proposition 7 For every graph G, it holds that

 $\psi(G) \leq dim(G) + 2 \ eqdim(G).$

Proof Let A be a resolving set, and let B be a distance-equalizer set. We next construct a set C of vertices satisfying $0 \le |C| \le eqdim(G)$ and such that for every pair of vertices $x, y \in V(G)$, there exist $u, v \in A \cup B \cup C$ doubly resolving x and y.

First, notice that if $x, y \in B$ then x and y are doubly resolved by themselves, and if $x, y \notin B$, then x and y are doubly resolved by vertices u and v, where $u \in A$ is a vertex resolving x and y, and $v \in B$ is equidistant from x and y. Hence, in both cases, x and y are doubly resolved by a pair of vertices in $A \cup B$.

Now, suppose that $x \in B$ and $y \notin B$. We claim that, for every $x \in B$ there is at most one vertex $y_x \notin B$ such that x, y_x are not doubly resolved by $A \cup B$ and besides, in such a case, $d(u, x) + d(x, y_x) = d(u, y_x)$ for every $u \in A \cup B$. Indeed, suppose that there exists $y' \notin B$ such that the pair x, y' is not doubly resolved by $A \cup B$. Then, for all $u \in A \cup B$, the pair of vertices $x, u \in A \cup B$ does not doubly resolve x, y'. Hence, d(u, x) - d(u, y') = -d(x, y'). In a similar way, if y'' is a vertex such that $y'' \notin B, y'' \neq y'$ and the pair x, y'' is not doubly resolved by $A \cup B$, then for all $u \in A \cup B$ we have d(u, x) - d(u, y'') = -d(x, y''). Therefore, d(x, y') - d(x, y'') = d(u, y') - d(u, y''). Thus, for every pair of vertices $u, v \in A \cup B$ we obtain d(u, y') - d(u, y'') = d(x, y') - d(x, y'') = d(v, y') - d(v, y''), implying that $y', y'' \notin B$ are not doubly resolved by $A \cup B$, which is not possible as we have seen in the former paragraph. Consider the (possibly empty) set

 $C = \{y_x : x \in B, y_x \notin B \text{ and } x, y_x \text{ are not doubly resolved by } A \cup B\}.$

Then, $0 \le |C| \le |B|$ and, by construction, $A \cup B \cup C$ doubly resolves x and y whenever $x \in B$ and $y \notin B$. Therefore, the set $S = A \cup B \cup C$ is a doubly resolving set for G and, consequently, $|S| \le dim(G) + 2 \ eqdim(G)$.

We think that the preceding bound can be improved as follows.

Conjecture 2 For every graph G, it holds that $\psi(G) \leq dim(G) + eqdim(G)$.

A graph attaining the upper bound given in the preceding conjecture is shown in Fig. 3.

Although we have no proof of Conjecture 2, we next prove that it holds true for trees.

Theorem 7 For every tree T, it holds that

$$\psi(T) \le \dim(T) + eqdim(T).$$

Proof Let S be the set of all support vertices of T. For every $v \in S$, consider the sets of vertices $L_v = \{z : z \text{ is a leaf adjacent to } v\}$ and $T_v = \{v\} \cup L_v$, and observe that





the sets T_v are pairwise disjoint. First, note that it is well known that the set of leaves of a tree T is the unique minimum doubly resolving set of T (see [11]). Hence,

$$\psi(T) = \sum_{v \in S} |L_v|.$$

Also, note that if W is a resolving set of T, then $|W \cap L_v| \ge |L_v| - 1$ for every $v \in S$ (see [30]).

Let W' be the union of a minimum resolving set and a minimum distance-equalizer set of T. We claim that $|W' \cap T_v| \ge |L_v|$, for every $v \in S$. Indeed, $|W' \cap L_v| \ge |L_v| - 1$, since W' is a resolving set, and $v \in W'$ or $L_v \subseteq W'$, by Lemma 1. In any case, $|W' \cap T_v| \ge |L_v|$.

Then,

$$\psi(T) = \sum_{v \in \mathcal{S}} |L_v| \le \sum_{v \in \mathcal{S}} |W' \cap T_v| \le |W'| \le dim(T) + eqdim(T).$$

We finish this section by analyzing lower and upper bounds on dim(G)+eqdim(G). Concretely, we are interested in the minimum and maximum value of dim(G) + eqdim(G) for graphs of order *n*. First, note that for any nontrivial graph *G* of order *n*,

$$2 \le \dim(G) + eqdim(G) \le 2(n-1). \tag{1}$$

The lower bound in (1) is attained only by the paths P_2 and P_3 , by Theorem 1(i), and the upper bound, only by the path P_2 , by Theorem 2(i). Hence, for every graph G of order at least 4,

$$3 \le \dim(G) + eqdim(G) \le 2n - 3.$$

In order to study this question, we consider the following functions defined for integers $n \ge 4$:

$$\Sigma(n) := \max\{dim(G) + eqdim(G) : |V(G)| = n\}$$



Fig. 4 The graph G_3

 $\sigma(n) := \min\{dim(G) + eqdim(G) : |V(G)| = n\}.$

Proposition 8 For every integer $n \ge 4$, the following statements hold.

(i) $\Sigma(n) \ge \frac{3n}{2} - 3;$ (ii) $\sigma(n) \le \log_2(n) + 2.$

- **Proof** (i) Consider the complete bipartite graph $G = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$, for which dim(G) = n 2 (see [14]), and $eqdim(G) = \lfloor n/2 \rfloor$, by Theorem 4(ii). Hence, $dim(G) + eqdim(G) = n 2 + \lfloor n/2 \rfloor \ge 3n/2 3$.
- (ii) For every $k \ge 1$, consider the graph G_k , with $V(G_k) = A \cup B \cup C$, where $A = \{v\}, B = \{1, ..., k\}, C = \{w : w \text{ is a binary word of length } k\}$ and two different vertices x and y are adjacent in G_k if and only if one of the following conditions hold (see an example in Fig. 4):
 - one of the vertices is v;
 - one of the vertices belongs to C, say x = w ∈ C, and the other one belongs to B, say y = j ∈ B, and w has the digit 1 in the jth position.

Then, G_k is a graph of order $n = 2^k + k + 1$. Moreover, A is a distance-equalizer set since v is a universal vertex, and it is easy to check that $A \cup B$ is a resolving set. Hence, $dim(G_k) + eqdim(G_k) \le k + 2 \le \log_2(n) + 2$.

6 Conclusions and Open Problems

In this paper, the notion of equidistant dimension as a parameter to evaluate sameness in graphs is introduced. The value of this invariant in several families of graphs and relations with other parameters has been provided. In Table 2, the equidistant dimension, metric dimension and minimum cardinality of doubly resolving sets of some families of graphs are given. Also, all graphs reaching some extremal values of the equidistant dimension have been characterized.

As future work, besides solving Conjecture 1 about the equidistant dimension of trees, and Conjecture 2 about the relation of the equidistant dimension with doubly

G	Constraints	eqdim(G)	dim(G)	$\psi(G)$
P _n	$n \ge 2$	$n-r\left(\left\lceil \frac{n}{2} \right\rceil\right)$	1	2
	$n = 4k \ge 4$	$\frac{3n}{4} - 1$	2	3
C_n	$n = 4k + 2 \ge 6$	$\frac{n}{2}$	2	3
	$n = 2k + 1 \ge 5$	$\leq n - r\left(\left\lceil \frac{n+1}{4} \right\rceil\right)$	2	2
K _n	$n \ge 3$	1	n - 1	n-1
$K_{r,s}$	$2 \le r \le s = n - r$	r	n-2	n-2
$K_{1,n-1}$	$n \ge 4$	1	n-2	n - 1
$K_2(r,s)$	$3 \le r \le s = n - r$	r	n-4	n-2
K_{n_1,\ldots,n_p}	$p \ge 3, n_1 + \dots + n_p = n$	$\min\{3, n_1, \ldots, n_p\}$	n-p	n-p

Table 2 Equidistant dimension and related parameters of some families of graphs

resolving sets, it would be interesting to relate distance-equalizer sets to other types of sets of vertices such as dominating sets, cut sets or determining sets, for example. Also, it could be of interest to find other graph families whose equidistant dimension connects with other problems, thus producing similar results as the relationship between the computation of this parameter in paths and AP-3-free sequences. Finally, we could perform new techniques that allow us to compute exact values for the equidistant dimension of Johnson graphs.

Acknowledgements A. González is supported by "VI Plan Propio de Investigación y Transferencia" of the Universidad de Sevilla (Spain) and by the Research Group in Mathematics Education FQM-226 of the Junta de Andalucía (Spain). C. Hernando and M. Mora are supported by project PID2019-104129GB-100/AEI/10.13039/501100011033 of the Spanish Ministry of Science and Innovation and by project Gen. Cat. DGR 2017SGR1336. M. Mora is also supported by project H2020-MSCA-RISE project 734922 - CONNECT.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- 1. Alspach, B.: Johnson graphs are Hamilton-connected. Ars Math. Contemp. 6, 21–23 (2013)
- Aslam, M., Ali, A.: Some results on induced subgraphs of Johnson graphs. Int. Math. Forum 7(9–12), 445–454 (2012)
- Bailey, R.F., Cáceres, J., Garijo, D., González, A., Márquez, A., Meagher, K., Puertas, M.L.: Resolving sets for Johnson and Kneser graphs. Eur. J. Comb. 34(4), 736–751 (2013)

- Bautista-Santiago, C., Cano, J., Fabila-Monroy, R., Flores-Peñaloza, D., González-Aguilar, H., Lara, D., Sarmiento, E., Urrutia, J.: On the connectedness and diameter of a geometric Johnson graph. Discrete Math. Theor. Comput. Sci. 15(3), 21–30 (2013)
- Behrend, F.A.: On sets of integers which contain no three in arithmetic progression. Proc. Natl. Acad. Sci. 23, 331–332 (1946)
- Bitan, S., Etzion, T.: On the chromatic number, colorings, and codes of the Johnson graph. Discrete Appl. Math. 70(2), 163–175 (1996)
- 7. Bloom, T.F.: A quantitative improvement for Roth's theorem on arithmetic progressions. J. Lond. Math. Soc. Second Ser. **93**(3), 643–663 (2016)
- Bloom, T.F., Sisask, O.: Logarithmic bounds for Roths theorem via almost-periodicity. Discrete Anal. 4, 20 (2019)
- 9. Bourgain, J.: On triples in arithmetic progression. Geom. Funct. Anal. 9(5), 968–984 (1999)
- Cáceres, J., Hernando, C., Mora, M., Pelayo, I.M., Puertas, M.L., Seara, C.: On the metric dimension of infinite graphs. Discrete Appl. Math. 160(18), 2618–2626 (2012)
- Cáceres, J., Hernando, C., Mora, M., Pelayo, I.M., Puertas, M.L., Seara, C., Wood, D.R.: On the metric dimension of Cartesian products of graphs. SIAM J. Discrete Math. 21(2), 423–441 (2007)
- Čangalović, M., Kratica, J., Vujčič, V.K., Stojanovič, M.: Minimal double resolving sets of prism graphs. Optimization 62(8), 1037–1043 (2013)
- Casas-Roma, J., Herrera-Joancomartí, J., Torra, V.: A survey of graph-modification techniques for privacy-preserving on networks. Artif. Intell. Rev. 47(3), 341–366 (2017)
- Chartrand, G., Eroh, L., Johnson, M.A., Oellermann, O.R.: Resolvability in graphs and the metric dimension of a graph. Discrete Appl. Math. 105, 99–113 (2000)
- Chen, X., Wang, C.: Approximability of the minimum weighted doubly resolving set problem. Comput. Comb. LNCS 8591, 357–368 (2014)
- 16. Chester, S., Srivastava, G.: Social network privacy for attribute disclosure attacks. In: 2011 International Conference on Advances in Social Networks Analysis and Mining, pp. 445–449. IEEE (2011)
- Chester, S., Kapron, B.M., Srivastava, G., Venkatesh, S.: Complexity of social network anonymization. Soc. Netw. Anal. Min. 3(2), 151–166 (2013)
- Claverol, M., García, A., Hernández, G., Hernando, C., Maureso, M., Mora, M., Tejel, J.: Metric dimension of maximal outerplanar graphs. Bull. Malays. Math. Sci. Soc. 44(4), 2603–2630 (2021)
- Cockayne, E.J., Hedetniemi, S.T.: On the diagonal queens domination problem. J. Comb. Theory Ser. A 42(1), 137–139 (1986)
- Dabrowski, A., Moss, L.S.: The Johnson graphs satisfy a distance extension property. Combinatorica 20(2), 295–300 (2000)
- Dybizbański, J.: Sequences containing no 3-term arithmetic progressions. Electron. J. Comb. 19(2), P15 (2012)
- 22. Elkin, M.: An improved construction of progression-free sets. Isr. J. Math. 184, 93-128 (2011)
- 23. Erdös, P., Turán, P.: On some sequences of integers. J. Lond. Math. Soc. 11, 261-264 (1936)
- 24. Feder, T., Nabar, S., Terzi, E.: Anonymizing graphs (2008). CoRR arXiv:0810.5578
- Gasarch, W., Glenn, J., Kruskal, C.: Finding large 3-free sets. I. The small n case. J. Comput. Syst. Sci. 74(4), 628–655 (2008)
- González, A., Hernando, C., Mora, M.: Metric-locating-dominating sets of graphs for constructing related subsets of vertices. Appl. Math. Comput. 332, 449–456 (2018)
- 27. Gowers, W.: A new proof of Szeméredi's theorem. Geom. Funct. Anal. 11, 465-588 (2001)
- 28. Harary, F., Melter, R.A.: On the metric dimension of a graph. Ars Comb. 2, 191-195 (1976)
- Heath-Brown, D.R.: Integer sets containing no arithmetic progressions. J. Lond. Math. Soc. (2) 35, 385–394 (1987)
- Hernando, C., Mora, M., Pelayo, I.M., Seara, C., Wood, D.R.: Extremal graph theory for metric dimension and diameter. Electron. J. Comb. 17, R30 (2010)
- Jiang, J., Wen, S., Yu, S., Xiang, Y., Zhou, W.: Identifying propagation sources in networks: state-ofthe-art and comparative studies. IEEE Commun. Surv. Tutor. 19(1), 465–481 (2017)
- Kratica, J., Čangalović, M., Kovačević-Vujčić, V.: Computing minimal doubly resolving sets of graphs. Comput. Oper. Res. 36(7), 2149–2159 (2009)
- Kratica, J., Kovačević-Vujčić, V., Čangalović, M., Stojanović, M.: Minimal doubly resolving sets and the strong metric dimension of some convex polytopes. Appl. Math. Comput. 218(19), 9790–9801 (2012)

- Krebs, M., Shaheen, A.: On the spectra of Johnson graphs. Electron. J. Linear Algebra 17, 154–167 (2008)
- Lobstein, A.: Watching systems, identifying, locating-dominating and discriminating codes in graphs. http://www.infres.enst.fr/lobstein/debutBIBidetlocdom.pdf
- Ramras, M., Donovan, E.: The automorphism group of a Johnson graph. SIAM J. Discrete Math. 25, 267–270 (2011)
- 37. Roth, K.F.: On certain sets of integers. J. Lond. Math. Soc. 28(1), 104-109 (1953)
- 38. Sanders, T.: On Roth's theorem on progressions. Ann. Math. (2) 174(1), 619–636 (2011)
- Sharma, A.: Sequences of integers avoiding 3-term arithmetic progressions. Electron. J. Comb. 19, P27 (2012)
- 40. Slater, P.J.: Leaves of trees. Congr. Numer. 14, 549-559 (1975)
- Trujillo-Rasua, R., Yero, I.G.: k-metric antidimension: a privacy measure for social graphs. Inf. Sci. 328, 403–417 (2016)
- 42. West, D.B.: Introduction to Graph Theory, Second Edition Prentice Hall, Upper Saddle River (2001)
- Zheleva, E., Getoor, L.: Privacy in social networks: a survey. In: Aggarwal, C.C. (ed.) Social Network Data Analytics, 1st edn., pp. 277–306. Springer, Berlin (2011)
- Zhou, B., Pei, J., Luk, W.S.: A brief survey on anonymization techniques for privacy preserving publishing of social network data. SIGKDD Explor. Newsl. 10(2), 12–22 (2008)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.