



Microbial community and antimicrobial resistance in fecal samples from wild and domestic ruminants in Maiella National Park, Italy

Camilla Smoglica^{a,*}, Simone Angelucci^{a,b}, Muhammad Farooq^a, Antonio Antonucci^b, Fulvio Marsilio^a, Cristina E. Di Francesco^a

^a University of Teramo, Faculty of Veterinary Medicine, Loc. Piano D'Accio, 64100 Teramo, Italy

^b Wildlife Research Center, Maiella National Park, via del vivaio, 65023 Caramanico Terme (PE), Italy

ARTICLE INFO

Keywords:

Wild animals
Bacteria
Resistance genes
mcr-1
Wild-livestock interface

ABSTRACT

This study aimed to provide new insights about antimicrobial resistance genes abundance and microbial communities of wild and domestic ruminants in wildlife-livestock interface. In total, 88 fecal samples were recovered from Apennine chamois, red deer, goat, cattle and sheep, and were collected in pools. The populations under study were selected based on ecological data useful to define sympatric and non-sympatric populations. Samples were screened for commonly used in farms under study or critically important antimicrobial resistance genes (*aadA2*, *TetA*, *TetB*, *TetK*, *TetM*, *mcr-1*). The microbial community composition was found to be different based on the species and land use of animals under study. Indeed, it was mostly characterized by phyla *Firmicutes* in bovine, *Bacteroidota* in chamois and *Proteobacteria* in red deer. Additionally, positive correlations between antibiotic resistance genes and microbial taxa (e.g., *Tet* genes correlated with *Firmicutes* and *Patescibacteria*) were described. Of the antimicrobials investigated, the abundance of *mcr-1* gene suggests the importance of monitoring the wildlife in order to detect the emerging resistance genes contamination in environment. This study provides new data that highlight the importance of multidisciplinary and uncultured study in order to describe the spreading of antimicrobial resistance and related contamination in the environment.

1. Introduction

Antimicrobial resistance (AMR) is a multifaced issue considered one of the the 21st century most serious global threats [1]. The spread of AMR involves many habitats which include environments characterized by low or no human impact [1]. The mechanisms of resistance spreading are mostly related to mobile resistance genes and to selection and maintenance of multi-resistant bacteria in the environment based on the horizontal transfer [2]. Resistant microorganisms and their antimicrobial resistance genes (ARGs) may remain and eventually persist into environmental microbial community [3]. In this view the ARGs have been considered as emerging environmental contaminants [3]. Indeed, the 60–90% of antimicrobials used at farm level may be released in the environment through animal urine and feces [2], leading to the selection of resistance bacteria by means of their long-time persistence in environment.

Previous studies have mainly focused on the detection of ARGs

starting from cultured selected bacteria, but it is noteworthy that most microbial species are non-culturable [4,5]. In this regard, alternative PCR- based approaches without preliminary culture have been suggested as a powerful method to detect potential antibiotic resistance in total microbial DNA [6]. This approach resulted mainly applied for soil, residues and water-related samples analysis [7,8] and during the last years has been also realized in wildlife [4,9–11]. Despite the wildlife has been suggested as an indicator for the spread of ARGs [6], there is non-univocal published data about its role in antibiotic resistance dynamics. Species living in different ecological niches are considered to have different roles in AMR dispersion. Indeed, the presence of AMR in wildlife results to be associated to habitat features, trophic characteristics, and anthropic pressure. Sympatric wild animals (that use anthropized habitats as feeding or refuge sites) have higher possibilities of harboring and spreading resistant bacteria [12]. Therefore, the wildlife has been used as sentinels of AMR and related ARGs in the environment and the species mostly investigated resulted wild birds and

* Corresponding author.

E-mail addresses: csmoglica@unite.it (C. Smoglica), simone.angelucci@parcomajella.it (S. Angelucci), mfarooq@unite.it (M. Farooq), antonio.antonucci@parcomajella.it (A. Antonucci), fmarsilio@unite.it (F. Marsilio), cedifrancesco@unite.it (C.E. Di Francesco).

<https://doi.org/10.1016/j.oneht.2022.100403>

Received 1 March 2022; Received in revised form 17 May 2022; Accepted 17 May 2022

Available online 21 May 2022

2352-7714/© 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

small mammals [13].

In this view, the aim of this study is to determine the fecal microbial community and occurrence of ARGs of wild and domestic ruminants with different levels of wildlife-livestock interface in the protected area of Maiella National Park (MNP), Italy.

2. Material and methods

2.1. Study area and sampling design

During October and November 2019, were collected a total of 22 fecal pools from red deer (*Cervus elaphus*), Apennine chamois (*Rupicapra pyrenaica ornata*) and extensive livestock, living in MNP (Italy). The Apennine chamois is a rare subspecies of chamois living in defined areas of Central Italy. The population size of the MNP is approximately 1300 individuals, and these data are the result of the reintroduction programs carried out in the past years. This species coexists in the territories of the Park with other widespread wild ungulates as the red deer population, of approximately 1500 individuals, and domestic cattle, sheep and goats traditionally raised in small farms by extensive grazing systems. The distribution of wild and domestic animals was previously determined by monitoring programs and georeferencing data useful to define the level of grazing lands sharing [14]. In this way, a group of 100 Apennine chamois sympatric with a farm of 120 goats, and 50 red deer coexisting with a farm of 300 sheep has been selected. Additionally, the non-sympatric populations composed by 70 cattle, 210 goats, 100 Apennine chamois and 20 red deer living in different areas of the Park were




Fig. 1. Distribution of sympatric and non-sympatric animals investigated in grazing lands of the MNP [14].

included (Fig. 1).

As previously reported, the samples were collected gathering the freshly deposited feces and following the groups of grazing animals without disturbance of the animal activities [14]. The samples were stored at refrigeration temperature and analyzed by the laboratory of the Faculty of Veterinary Medicine of the University of Teramo within 3 h from the collection.

2.2. DNA extraction

Twenty-five grams of sample were homogenized in 250 ml of Phosphate Buffered Saline (PBS) and filtered using a sterile Filtra-bag (280 μ m pore size) by Stomacher® for 2 min. After centrifugation at 2903 G for 10 min at 4 °C; total DNA was extracted from 200 mg of the resulting pellet using GeneAll Exgene™ Stool DNA mini kit. DNA quality and quantity were assessed using the spectrophotometer and fluorometer Denovix DS-11 FX (Wilmington, USA).

2.3. 16S rRNA gene sequencing and data analysis

The V3 and V4 regions of 16S rRNA gene were amplified using the primer V3(5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG-3') and V4 (5'-GTCTCGTGGGCTCGGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC-3') according to Klindworth et al. [15] with Illumina adapter overhang nucleotide sequences. The PCR products were pooled and indexed using the Nextera XT Index kit (Illumina, USA) following the 16S Metagenomic Sequencing Library Preparation Guide protocol (Illumina, USA). Libraries were sequenced using the Illumina MiSeq sequencing platform (San Diego, California, USA) with a 2 × 300 bp paired-end approach. The 16S rRNA data analysis was carried out using DADA2 package within the Quantitative Insights into Microbial Ecology 2 (QIIME2 version 2019.4) software [16,17]. The taxonomy categories were assigned by a Naive Bayes classifier and the q2-feature-classifier plugin, and identified using SILVA-Naive Bayes sklearn trained database [18]. The raw sequence reads have been deposited in the NCBI Short Read Archive under the accession number PRJNA832550.

The microbial community characterization and α - and β -diversity statistics were performed by the software Calypso (<http://cgenome.net/wiki/index.php/Calypso>) [19]. The microbial community composition and quantification within each sample group and among different animals were analyzed as previously described [1].

2.4. Determination of antimicrobial resistance genes (ARGs) by quantitative PCR (qPCR) analysis

The panel of 6 ARGs (*aadA2*, *TetA*, *TetB*, *TetK*, *TetM*, *mcr-1*) was selected based on the main antimicrobial classes used in livestock of study areas (tetracycline and streptomycin) along with the colistin, which is considered a critically important antibiotic for human medicine, and it has been particularly monitored in Europe by official surveillance program during the last years [20]. These genes along with the 16S rRNA gene were analyzed by SYBR Green® qPCR as previously described [7,9] in order to normalize the abundance of ARGs in fecal samples [21]. Each run was realized in triplicate. The copy number of target genes was calculated including an 8-point calibration curve obtained using ten-fold diluted positive controls and a good correlation coefficient ($0.993 > R^2 > 0.999$). The normalized copy number of ARGs was calculated as the ratio of ARG copy number to 16S rRNA gene copy number [21]. The sensitivity of the technique was determined using a baseline threshold of -7 . Any value ≤ -8 was considered negative as described by [21].

2.5. Statistical analysis

The student's *t*-test of ARGs copies and Spearman correlation analysis

to assess the association between the relative abundance of microbial taxa at phylum levels with ARGs, species or animal's group, were calculated using standard statistical software packages, Stata [22].

3. Results

3.1. General description of DNA sequences

Quality analysis and trimming returned a total of 8432 sequences with 8432 different features, with an average of 746,182 sequences per individual sample based on the analysis at operational taxonomy unit (OTU) level. The sequencing depth was good as indicated by the rarefaction curves for samples (Supplementary data 1).

3.2. Composition of bacterial communities

The microbial community composition of each sample was defined by means of the 16S rRNA sequencing analysis. At phylum level, member of the *Firmicutes* and *Bacteroidota* were dominant. In detail, *Firmicutes* phylum was highly abundant in bovine with the average of relative frequency of 56.3%, while a higher percentage of *Bacteroidota* (48%) respect to *Firmicutes* (38.7%) was identified in chamois. *Proteobacteria* phylum (26.3%) was more abundant in red deer samples, while *Verrucomicrobiota* (bovine: 5.2%; goat: 8.6%; sheep: 4.3%) was mostly detected in domestic animals (Fig. 2). The distribution at class level is consistent with the phylum frequency (Fig. 3). Indeed, the *Bacteroidia* class (*Bacteroidota* phylum) and *Clostridia* class (*Firmicutes* phylum) resulted prominent. *Clostridia* class (54.3%) was highly abundant in bovine, while a higher percentage of *Bacteroidia* (47.9%) respect to *Clostridia* (30.7%) was described in chamois. *Gammaproteobacteria* (*Proteobacteria* phylum) (30.7%) was more predominant in red deer, while *Verrucomicrobiae* (*Verrucomicrobiota* phylum) was mostly reported in domestic animals (bovine:3.4%; goat: 5.1%; sheep: 1.7%).

A strong positive correlation was observed between domestic species and phyla *Cyanobacteria* ($r = 0.81p = 0.001$), *Fibrobacteria* ($r = 0.83 p = 0.001$), *Spirochaetota* ($r = 0.85 p = 0.001$), *Verrucomicrobiota* ($r = 0.77 p = 0.001$), while a moderate correlation was described between domestic animals and *Desulfomicrobiota* ($r = 0.66 p = 0.0005$) or *Elusimicrobiota* ($r = 0.66 p = 0.0005$). Additionally, a moderate positive correlation resulted between wild animals and phyla *Patescibacteria* ($r = 0.41 p = 0.04$) and *Proteobacteria* ($r = 0.51 p = 0.001$).

Concerning the group of animals, it was described a moderate positive correlation between sympatric animals and *Proteobacteria* ($r = 0.48 p = 0.01$) or *Patescibacteria* ($r = 0.68 p = 0.0003$), while it was reported a moderate positive correlation between non-sympatric animals and *Firmicutes* ($r = 0.59 p = 0.003$).

B: bovine; C: chamois; G: goat; D: red deer; S: sheep.

B: bovine; C: chamois; G: goat; D: red deer; S: sheep.

3.3. Bacterial community α -diversity

The α -diversity was evaluated at OTU level using Shannon index (Supplementary data 2) highlighting a significant difference ($p < 0.01$) between the species. The lowest alpha diversity was observed in Apennine chamois and red deer while the highest diversity was noticed in bovine, goats and sheep. Finally, α -diversity appeared comparable between sympatric and non-sympatric populations.

3.4. Comparison among bacterial communities (β -diversity)

The analysis of β -diversity at OTU level was realized with principal coordinate analysis.

(PCoA) along with permutational multi-variable analysis of variance (PERMANOVA) showing a significant difference ($p < 0.05$) of microbial community in sympatric animals compared to non-sympatric animals. This finding was supported by PCoA graphs, in which the Bray-Curtis distance was used as a distance metrics of β -diversity, and a difference in bacterial communities' distribution between sympatric and non-sympatric animal samples (Fig. 4) was reported. Additionally, the differences in microbial community composition between species were significant ($p < 0.05$) except for goat and sheep or chamois and red deer, and PCoA graph confirmed these observations (Fig. 4).

3.5. Determination of antimicrobial resistance genes (ARGs) by quantitative PCR (qPCR) analysis

All investigated samples were found positive to at least one ARG.

The *aadA2*, *TetA* and *TetK* were detected in all the samples. The gene *TetB* was not amplified in samples from sympatric and non-sympatric goats and in non-sympatric red deer, while only one samples from sympatric sheep was resulted negative for *TetM* gene. Finally, the gene *mcr-1* was not reported in one sample from sympatric chamois.




Fig. 2. Distribution of relative frequency of bacterial phyla in wild and domestic animals.




Fig. 3. Heatmap of microbial community distribution in wild and domestic animals at class level.

The total abundance of ARGs normalized to 16S rRNA ranged from 4.59×10^{-7} to 4.30×10^1 as reported in Table 1.

The highest concentration of ARG was found in a sample of goat for *mcr-1* (2.16×10^1 copies/16SRNA copies) in the domestic animal's group and for *TetM* (4.30×10^1 ARGs copies/16SRNA copies) in Apennine chamois in wild animals.

Between the sympatric animals, the highest concentration of gene was identified in Apennine chamois for *mcr-1* (7.51×10^{-1}) and in non-sympatric animals in goat for *mcr-1* (4.47×10^1 ARGs copies/16SRNA copies).

The statistical analysis showed that the means of ARGs of *TetA* ($p = 0.04$), *TetK* ($p = 0.01$), *TetM* ($p = 0.001$) and *mcr-1* ($p = 0.01$) were highly significantly ($p < 0.05$) different in sympatric respect to non-sympatric animals. No significant difference of ARGs mean was identified between wild and domestic animals.

3.6. Co-occurrence between ARGs and bacterial taxa

The Spearman rank sum correlation analysis allowed to achieve different results. In detail, one moderate and two strong positive correlations resulted between phylum *Patescibacteria* and *TetA* ($r = 0.46$ $p = 0.02$), *TetK* ($r = 0.65$ $p = 0.0006$) and *TetM* ($r = 0.66$ $p = 0.0005$). Additionally, a moderate positive correlation was described between *Firmicutes* and *TetK* ($r = 0.42$ $p = 0.04$) or *TetM* ($r = 0.40$ $p = 0.04$). Considering the gene *mcr-1*, a moderate correlation with *Myxococcota* ($r = 0.42$ $p = 0.04$) was reported and finally a moderate correlation was described for *Actinobacteriota* and *adaA2* ($r = 0.55$ $p = 0.005$), and *Bdellovibrionota* and *TetA* ($r = 0.40$ $p = 0.04$).

4. Discussion

This study represents the first analysis of microbial communities and

ARGs in wildlife-livestock interface in Apennines, particularly relevant for the sampling design realized in a protected area and for the rare species involved such as Apennine chamois. Based on these features, the study did not allow to collect a large number of samples, but the research constitutes a preliminary step for further and more accurate investigations.

Similar methodologies were recently carried out on wildlife or environment in north-eastern Germany [11], in Poland [23] and Italy [24] but no ecological features (i.e., geographical distribution, size population and land use) of investigated animals were evaluated.

Additionally, Lanconi et al. [25] focused on the microbial community and AMR in agricultural soils fertilized with livestock manure in Italy, without considering the potential influence of wild species co-existing in the same territories.

The results obtained in our study showed that *Bacteroidota* and *Firmicutes*, along with *Proteobacteria* in wild ruminants, as reported by previous studies, are the dominant colonizing bacterial groups of the digestive system in domestic ruminants [26,27]. The *Firmicutes/Bacteroidota* ratio in gut microbiota appears to evolve during the life of animals and humans and it results in diverse digestive ability [28]. Several factors related to individual features and geographical localization may influence the microbiota in terms of quantity and diversity [28]. For examples, in dogs a food supplementation with fiber was associated to the increase of *Firmicutes* and *Bacteroidota*, and it was described as determinant for digestion of carbohydrates and other high molecular nutrients [28]. In wildlife *Proteobacteria* was described as most predominant phylum in giant panda and wild sika deer, resulting useful to optimize the degradation of lignin [29]. Finally, this phylum was also related to catabolism of various components of bovine fodder [30]. This evidence suggests that the composition of dietary may influence the relative abundance of these phyla in animals. Additionally, significant differences between wild and domestic animals and sympatric and non-




Fig. 4. β -diversity between sympatric or non-sympatric animals (A) and between species (B) by means of PCoA analysis according to Bray Curtis distances.

sympatric populations were observed in our study. The correlation analysis showed results in line with previous studies. In detail, the association between phyla and domestic animals confirmed the common bacterial phyla of fecal microbiota in domestic ruminants, as previously

described by other authors [31]. In addition, the correlation between *Proteobacteria* and wild animals was described according to the data previously reported [27]. Finally, the correlation between non-sympatric animals with *Firmicutes* and sympatric animals with *Proteobacteria* or *Patescibacteria*, may be related to the different activities and diets of animals. Indeed, other authors have associated *Firmicutes* to the feedlot animals and *Patescibacteria* to grazing animals [31].

These results suggest that the diversity of microbial community may be influenced by species (wild or domestic animals) and diets as assumed by other authors [28] and that the use and level of sharing of the land play a key role to define the composition of fecal microbiota.

Regarding the ARGs distribution, a significant difference of *TetA*, *TetK*, *TetM* and *mcr-1* genes was observed between sympatric and non-sympatric animals, highlighting one more time the potential influence of the environment to define the ARGs profile. Indeed, different habitats may accumulate different ARGs [32].

Different correlations between phyla and ARGs have already been described. In detail, the *Proteobacteria* phylum was previously identified as particularly abundant of ARGs [33] and this evidence may explain the correlation detected in this study between ARGs and the proteobacterial classes, recently renamed as *Myxococcota* and *Bdellovibrionota* phyla [34]. The correlation between *aadA2* and *Actinobacteria* reported in this study, were previously justified by other authors considering that many aminoglycosides have been isolated by bacteria of this phylum [35]. The correlation of *Tet* genes with *Firmicutes* and *Patescibacteria* was observed and appears to be in line with previous results that include the tetracycline genes in the top 15 ARGs accumulated in different habitats (i.e., hospitals, farms, soil, water, and wastewater) [32].

Tetracycline resistance-encoding genes were the most prevalent, as described in other similar studies in Brazilian wild birds [1], in tortoises in the Galapagos Archipelago, in Ecuador [9], in guignas [10] and Andean foxes in Chile [4], and in wild boar and foxes in Poland and Germany [11,23]. The high tetracycline resistance occurrence is not surprising, considering that these antibiotics have been widely employed for treatments in human and veterinary medicine as well as in agriculture [1]. Indeed, as reported by technical staff of MNP, this antibiotic was widely used in livestock of extensive farm situated in the protected area (Angelucci S. personal communication). Noteworthy, the 75–80% of the doses of tetracyclines are excreted in urine and feces with a long-term persistence of 578 days in soil that returns in a selection of resistant bacteria [36].

Table 1

Normalized copies of ARGs detected in each sample of wild and domestic animals.

Group	ID sample	Animal	<i>aadA2</i>	<i>TetA</i>	<i>TetB</i>	<i>TetK</i>	<i>TetM</i>	<i>mcr-1</i>	
Sympatric	S1	Sheep	1.56×10^{-4}	2.47×10^{-3}	1.96×10^{-3}	4.79×10^{-5}	4.28×10^{-7}	6.36×10^{-2}	
	S2	Sheep	1.36×10^{-4}	7.97×10^{-4}	2.24×10^{-3}	1.52×10^{-5}	Neg	9.38×10^{-3}	
	S3	Sheep	1.02×10^{-3}	2.07×10^{-3}	3.42×10^{-3}	3.63×10^{-5}	4.59×10^{-7}	1.50×10^{-1}	
	D1	Red deer	3.65×10^{-4}	8.35×10^{-3}	3.62×10^{-3}	8.24×10^{-6}	3.54×10^{-7}	3.05×10^{-3}	
	D5	Red deer	3.15×10^{-1}	3.66×10^{-4}	1.92×10^{-3}	1.40×10^{-5}	1.47×10^{-6}	1.97×10^{-2}	
	D6	Red deer	4.72×10^{-4}	2.24×10^{-3}	4.47×10^{-4}	2.25×10^{-5}	1.79×10^{-7}	6.42×10^{-4}	
	G1	Goats	2.31×10^{-4}	3.16×10^{-2}	Neg	2.26×10^{-5}	8.14×10^{-6}	5.52×10^{-1}	
	G2	Goats	1.75×10^{-4}	4.14×10^{-2}	Neg	1.99×10^{-5}	2.74×10^{-6}	5.11×10^{-2}	
	G3	Goats	9.93×10^{-5}	2.44×10^{-2}	Neg	1.16×10^{-4}	6.41×10^{-6}	2.85×10^{-1}	
	C1	Chamois	8.81×10^{-2}	1.41×10^{-1}	1.20×10^{-2}	7.41×10^{-4}	2.94×10^{-6}	7.51×10^{-1}	
	C2	Chamois	1.67×10^{-3}	1.87×10^{-2}	1.01×10^{-3}	1.24×10^{-4}	1.13×10^{-5}	Neg	
	Non-sympatric	D2	Red deer	4.50×10^{-4}	2.40×10^{-2}	Neg	2.79×10^{-4}	2.22×10^{-5}	4.83×10^{-1}
		D3	Red deer	5.05×10^{-3}	6.52×10^{-2}	Neg	1.21×10^{-3}	1.25×10^{-4}	2.04×10^{-1}
D4		Red deer	1.02×10^{-3}	3.22×10^{-2}	Neg	2.72×10^{-4}	2.95×10^{-5}	1.29×10^1	
G4		Goats	1.13×10^{-3}	2.87×10^{-2}	Neg	3.15×10^{-5}	7.58×10^{-5}	4.47×10^1	
G5		Goats	7.98×10^{-4}	1.29×10^{-2}	Neg	6.97×10^{-5}	3.03×10^{-4}	5.53×10^{-1}	
G6		Goats	4.38×10^{-4}	1.85×10^{-2}	Neg	5.42×10^{-4}	2.69×10^{-4}	2.16×10^1	
B1		Cattle	2.61×10^{-6}	8.13×10^{-3}	1.08×10^{-2}	3.65×10^{-5}	3.81×10^{-1}	8.07×10^{-2}	
B2		Cattle	4.12×10^{-7}	1.64×10^{-2}	9.51×10^{-3}	1.02×10^{-4}	2.74×10^{-1}	3.44×10^{-1}	
B3		Cattle	2.94×10^{-4}	7.36×10^{-3}	3.72×10^{-3}	3.18×10^{-5}	2.62×10^{-1}	8.67×10^{-2}	
C3		Chamois	6.10×10^{-8}	1.43×10^{-1}	1.22×10^{-2}	3.89×10^{-4}	4.30×10^1	5.73×10^{-1}	
C4		Chamois	2.03×10^{-8}	1.00×10^{-1}	1.01×10^{-2}	2.35×10^{-4}	1.44×10^1	3.39×10^{-1}	
C5		Chamois	1.55×10^{-6}	3.33×10^{-2}	1.63×10^{-3}	7.83×10^{-5}	5.93×10^{-1}	4.46×10^{-2}	

Neg: negative sample was defined as any value ≤ -8

Therefore, the soil resistome derived from pathogenic and non-pathogenic bacteria present in the environment may consist of ARGs relevant to both human and veterinary medicine [36].

In this view, the colistin resistance gene *mcr-1* gene resulted the most abundant target observed not only in domestic animals but also in wild sympatric group, highlighting the importance of studies focused on the environmental contamination of AMR. The worldwide presence of colistin resistance gene, previously reported in human [37], animals [38,39], animal food-products [40], and the environment [41], is an alarming indication of the inevitable progression to pan-drug resistance [37]. The *mcr-1* gene evidence of this study confirms the important role of wildlife as possible sentinels of this trend in the human-wildlife-environment interface [42–45]. Similar reports carried out on wild animals showed the *mcr-1* in free-ranging seabirds, such as gulls in Europe [45], North America [46,47], Oceania [48], and South America [1,49], while in wild mammals phenotypic and genetic colistin resistance was reported in isolates recovered from fallow deer in Europe [50], barbary macaques in Africa [51] and Père David's deer in Asia [52]. In Central Italy, similar studies are available only for isolates from hunted wild boar and free-ranging Apennine chamois [20,42,53].

To date, the investigations regarding animal and environment context are mainly focused on selected pathogens and ARGs of public health concerns [54]. Conversely, the results of this study suggest that new measures to understand drug resistance dynamics are necessary. In this regard, the environmental monitoring, the risk assessment, and the quantification of AMR should be realized involving wildlife sources and by means of conventional microbiological analysis improved by culture-independent strategy [54].

5. Conclusions

The ARGs differences found in this study were linked to species' gut microbiome composition and ecological relationship between animals, providing for the first-time new data concerning this topic in protected area in Italy and at wildlife-livestock interface.

These preliminary results represent an attempt to overcome the limitation of culture-based methods and opportunistically sampling used to study AMR, applying a multidisciplinary approach targeted on the territory.

Funding

This work was supported by the Project “Demetra” (Dipartimento di Eccellenza 2018–2022, CUP_C46C18000530001) funded by the Italian Ministry for Education, University and Research.

Ethical approval

Not required.

CRediT authorship contribution statement

Camilla Smoglica: Conceptualization, Methodology, Software, Data curation, Formal analysis, Software, Validation, Writing – original draft, Writing – review & editing. **Simone Angelucci:** Visualization, Investigation, Writing – review & editing. **Muhammad Farooq:** Data curation, Formal analysis. **Antonio Antonucci:** Visualization, Investigation, Writing – review & editing. **Fulvio Marsilio:** Writing – review & editing, Funding acquisition, Supervision. **Cristina E. Di Francesco:** Conceptualization, Methodology, Software, Software, Validation, Writing – original draft, Writing – review & editing.

Declaration of Competing Interest

None declared.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.onehlt.2022.100403>.

References

- [1] A.C. Ewbank, F. Esperón, C. Sacristán, I. Sacristán, R. Krul, E. Cavalcante de Macedo, O. Calatayud, I. Bueno, R. de Francisco Strefezzi, J.L. Catão-Dias, Seabirds as anthropization indicators in two different tropical biotopes: a One Health approach to the issue of antimicrobial resistance genes pollution in oceanic islands, *Sci. Total Environ.* 754 (2021), 142141, <https://doi.org/10.1016/j.scitotenv.2020.142141>.
- [2] H. Heuer, H. Schmitt, K. Smalla, Antibiotic resistance gene spread due to manure application on agricultural fields, *Curr. Opin. Microbiol.* 4 (3) (2011) 236–243, <https://doi.org/10.1016/j.mib.2011.04.009>.
- [3] S. Petrin, I. Patuzzi, A. Di Cesare, A. Tiengo, G. Sette, G. Biancotto, G. Corno, M. Drigo, C. Losasso, V. Cibin, Evaluation and quantification of antimicrobial residues and antimicrobial resistance genes in two Italian swine farms, *Environ. Pollut.* 255 (Pt 1) (2019), 113183, <https://doi.org/10.1016/j.envpol.2019.113183>.
- [4] A. Cevdanes, F. Esperón, S. Di Cataldo, E. Neves, N. Sallaberry-Pincheira, J. Millán, Antimicrobial resistance genes in Andean foxes inhabiting anthropized landscapes in central Chile, *Sci. Total Environ.* 724 (2020), 138247, <https://doi.org/10.1016/j.scitotenv.2020.138247>.
- [5] J.C. Lagier, G. Dubourg, M. Million, F. Cadoret, M. Bilen, F. Fenollar, A. Levasseur, J.M. Rolain, P.E. Fournier, D. Raoult, Culturing the human microbiota and culturomics, *Nat. Rev. Microbiol.* 16 (2018) 540–550, <https://doi.org/10.1038/s41579-018-0041-0>.
- [6] H.K. Allen, J. Donato, H.H. Wang, K. Cloud-Hansen, J. Davies, J. Handelsman, Call of the wild: antibiotic resistance genes in natural environments, *Nat. Rev. Microbiol.* 8 (2010) 251–259, <https://doi.org/10.1038/nrmicro2312>.
- [7] L. Jiang, X. Hu, T. Xu, H. Zhang, D. Sheng, D. Yin, Prevalence of antibiotic resistance genes and their relationship with antibiotics in the Huangpu River and the drinking water sources, Shanghai, China, *Sci. Total Environ.* 458–460 (2013) 267–272, <https://doi.org/10.1016/j.scitotenv.2013.04.038>.
- [8] F. Esperón, C. Sacristán, M. Carballo, A. de la Torre, Antimicrobial resistance genes in animal manure, manure-amended and non anthropogenically impacted soils in Spain, *Adv. Biosci. Biotechnol.* 9 (2018) 469–480, <https://doi.org/10.4236/abb.2018.99032>.
- [9] A. Nieto-Claudin, S.L. Deem, C. Rodríguez, S. Cano, N. Moity, F. Cabrera, F. Esperón, Antimicrobial resistance in Galapagos tortoises as an indicator of the growing human footprint, *Environ. Pollut.* 284 (2021), 117453, <https://doi.org/10.1016/j.envpol.2021.117453>.
- [10] I. Sacristán, F. Esperón, F. Acuña, E. Aguilar, S. García, M.J. López, A. Cevdanes, E. Neves, J. Cabello, E. Hidalgo-Hermoso, J. Millán, C. Napolitano, Antibiotic resistance genes as landscape anthropization indicators: using a wild felid as sentinel in Chile, *Sci. Total Environ.* 703 (2020), 134900, <https://doi.org/10.1017/CBO9781107415324.004>.
- [11] T. Homeier-Bachmann, A.K. Schütz, S. Dreyer, J. Glanz, K. Schaufner, F.J. Conraths, Genomic analysis of ESBL-producing *E. coli* in wildlife from North-Eastern Germany, *Antibiotics* 11 (2022) 123, <https://doi.org/10.3390/antibiotics11020123>.
- [12] A.M. Ramey, C.A. Ahlstrom, Antibiotic resistant bacteria in wildlife: perspectives on trends, acquisition and dissemination, data gaps, and future directions, *J. Wildl. Dis.* 56 (1) (2020) 1–15 (PMID: 31567035).
- [13] R.T. Torres, J. Carvalho, M.V. Cunha, E. Serrano, J.D. Palmeira, C. Fonseca, Temporal and geographical research trends of antimicrobial resistance in wildlife - a bibliometric analysis, *One Health* 11 (2020), 100198, <https://doi.org/10.1016/j.onehlt.2020.100198>.
- [14] C. Smoglica, A. Vergara, S. Angelucci, A.R. Festino, A. Antonucci, F. Marsilio, C. E. Di Francesco, Evidence of linezolid resistance and virulence factors in *Enterococcus* spp. isolates from wild and domestic ruminants, Italy, *Antibiotics* 11 (2022) 223, <https://doi.org/10.3390/antibiotics11020223>.
- [15] A. Klindworth, E. Pruesse, T. Schweer, J. Peplies, C. Quast, M. Horn, et al., Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies, *Nucleic Acids Res.* 41 (2013), e1, <https://doi.org/10.1093/nar/gks808>.
- [16] E. Bolyen, J.R. Rideout, M.R. Dillon, N.A. Bokulich, C.C. Abnet, G.A. Al-Ghalith, et al., Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2, *Nat. Biotechnol.* 37(8) (2019) 852–857. doi:10.1038/s41587-019-0209-9, Erratum in: *Nat. Biotechnol.* 37 (9) (2019) 1091.
- [17] B.J. Callahan, P.J. McMurdie, M.J. Rosen, A.W. Han, A.J. Johnson, S.P. Holmes, DADA2: high-resolution sample inference from Illumina amplicon data, *Nat. Methods* 13 (7) (2016) 581–583, <https://doi.org/10.1038/nmeth.3869>.
- [18] P. Yilmaz, L.W. Parfrey, P. Yarza, J. Gerken, E. Pruesse, C. Quast, T. Schweer, J. Peplies, W. Ludwig, F.O. Glöckner, The SILVA and “all-species living tree project (LTP)” taxonomic frameworks, *Nucleic Acids Res.* 42 (Database issue) (2014) D643–D648, <https://doi.org/10.1093/nar/gkt1209>.
- [19] M. Zakrzewski, C. Proietti, J.J. Ellis, S. Hasan, M.J. Brion, B. Berger, L. Krause, Calypso: a user-friendly web-server for mining and visualizing microbiome-environment interactions, *Bioinformatics* 33 (5) (2017) 782–783, <https://doi.org/10.1093/bioinformatics/btw725>.
- [20] C. Smoglica, A. Vergara, S. Angelucci, A.R. Festino, A. Antonucci, L. Moschetti, M. Farooq, F. Marsilio, C.E. Di Francesco, Resistance patterns, *mcr-4* and *OXA-48*

- genes, and virulence factors of *Escherichia coli* from Apennine chamois living in sympatry with domestic species, Italy, *Animals* (Basel). 12 (2) (2022) 129, <https://doi.org/10.3390/ani12020129>.
- [21] C.M. McCann, B. Christgen, J.A. Roberts, J.Q. Su, K.E. Arnold, N.D. Gray, Y.G. Zhu, D.W. Graham, Understanding drivers of antibiotic resistance genes in High Arctic soil ecosystems, *Environ. Int.* 125 (2019) 497–504, <https://doi.org/10.1016/j.envint.2019.01.034>.
- [22] Stata Statistical Software: Release 16. (2019) College Station, TX: StataCorp LLC.
- [23] M. Skarżyńska, P. Leekitcharoenphon, R.S. Hendriksen, F.M. Aarestrup, D. Wasyl, A metagenomic glimpse into the gut of wild and domestic animals: quantification of antimicrobial resistance and more, *PLoS One* 15 (12) (2020), e0242987, <https://doi.org/10.1371/journal.pone.0242987>.
- [24] P. Monaco, F. Divino, G. Naclerio, A. Buccì, Microbial community analysis with a specific statistical approach after a record breaking snowfall in southern Italy, *Ann. Microbiol.* 70 (2020) 63, <https://doi.org/10.1186/s13213-020-01604-6>.
- [25] A. Laconi, L. Mughini-Gras, R. Tolosi, G. Grilli, A. Trocino, L. Carraro, F. Di Cesare, P. Cagnardi, A. Piccirillo, Microbial community composition and antimicrobial resistance in agricultural soils fertilized with livestock manure from conventional farming in northern Italy, *Sci. Total Environ.* 760 (2021), 143404, <https://doi.org/10.1016/j.scitotenv.2020.143404>.
- [26] Y. Guan, H. Yang, S. Han, L. Feng, T. Wang, J. Ge, Correction to: comparison of the gut microbiota composition between wild and captive sika deer (*Cervus nippon hortulorum*) from feces by high-throughput sequencing, *AMB Express* 8(1) (2018) 15. doi:10.1186/s13568-018-0535-1, Erratum for: *AMB Express* 7 (1) (2017) 212.
- [27] J. Lima, M.D. Auffret, R.D. Stewart, R.J. Dewhurst, C.A. Duthie, T.J. Snelling, A. W. Walker, T.C. Freeman, M. Watson, R. Roehe, Identification of rumen microbial genes involved in pathways linked to appetite, growth and feed conversion efficiency in cattle, *Front. Genet.* 10 (701) (2019) 1–18, [10.3389/fgene.2019.00701](https://doi.org/10.3389/fgene.2019.00701).
- [28] N. Szeligowska, P. Cholewińska, K. Czyż, K. Wojnarowski, M. Janczak, Inter and intraspecies comparison of the level of selected bacterial phyla in in cattle and sheep based on feces, *BMC Vet. Res.* 17 (1) (2021) 224, <https://doi.org/10.1016/10.1186/s12917-021-02922-w>.
- [29] W. Fang, Z. Fang, P. Zhou, F. Chang, Y. Hong, X. Zhang, H. Peng, Y. Xiao, Evidence for lignin oxidation by the giant panda fecal microbiome, *PLoS One* 7 (11) (2017), e50312, <https://doi.org/10.1371/journal.pone.0050312>.
- [30] N.J. Evans, J.M. Brown, R.D. Murray, B. Getty, R.J. Birtles, C.A. Hart, S.D. Carter, Characterization of novel bovine gastrointestinal tract *Trimonema* isolates and comparison with bovine digital dermatitis *trimonemes*, *Appl. Environ. Microbiol.* 77 (1) (2011) 138–147, <https://doi.org/10.1128/AEM.00993-10>.
- [31] Z. Zhang, L. Yang, Y. He, X. Luo, S. Zhao, X. Jia, Composition of fecal microbiota in grazing and feedlot Angus beef cattle, *Animals* (Basel). 11 (11) (2021) 3167, <https://doi.org/10.3390/ani11113167>.
- [32] M. Zhuang, Y. Achmon, Y. Cao, X. Liang, L. Chen, H. Wang, B.A. Siame, K.Y. Leung, Distribution of antibiotic resistance genes in the environment, *Environ. Pollut.* 285 (2021), 117402, <https://doi.org/10.1016/j.envpol.2021.117402>.
- [33] Y. Hu, X. Yang, J. Li, N. Lv, F. Liu, J. Wu, I.Y. Lin, N. Wu, B.C. Weimer, G.F. Gao, Y. Liu, B. Zhu, The bacterial mobile resistome transfer network connecting the animal and human microbiomes, *Appl. Environ. Microbiol.* 82 (22) (2016) 6672–6681, <https://doi.org/10.1128/AEM.01802-16>.
- [34] D.W. Waite, M. Chuvochina, C. Pelikan, D.H. Parks, P. Yilmaz, M. Wagner, A. Loy, T. Naganuma, R. Nakai, W.B. Whitman, M.W. Hahn, J. Kuever, P. Hugenholz, Proposal to reclassify the proteobacterial classes *Deltaproteobacteria* and *Oligoflexia*, and the phylum *Thermodesulfobacteria* into four phyla reflecting major functional capabilities, *Int. J. Syst. Evol. Microbiol.* 70 (11) (2020) 5972–6016, <https://doi.org/10.1099/ijsem.0.004213>.
- [35] M. Fatahi-Bafghi, Antibiotic resistance genes in the Actinobacteria phylum, *Eur. J. Clin. Microbiol. Infect. Dis.* 38 (9) (2019) 1599–1624, <https://doi.org/10.1007/s10096-019-03580-5>.
- [36] M. Cycoń, A. Mroziak, Z. Piotrowska-Seget, Antibiotics in the soil environment-degradation and their impact on microbial activity and diversity, *Front. Microbiol.* 10 (2019) 338, <https://doi.org/10.3389/fmicb.2019.00338>.
- [37] Y.Y. Liu, Y. Wang, T.R. Walsh, L.X. Yi, R. Zhang, J. Spencer, Y. Doi, G. Tian, B. Dong, X. Huang, L.F. Yu, D. Gu, H. Ren, X. Chen, L. Lv, D. He, H. Zhou, Z. Liang, J.H. Liu, J. Shen, Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study, *Lancet Infect. Dis.* 16 (2016) 161–168, [https://doi.org/10.1016/S1473-3099\(15\)00424-7](https://doi.org/10.1016/S1473-3099(15)00424-7).
- [38] M.F. Anjum, N.A. Duggett, M. AbuOun, L. Randall, J. Nunez-Garcia, R.J. Ellis, J. Rogers, R. Horton, C. Brena, S. Williamson, F. Martelli, R. Davies, C. Teale, Colistin resistance in *Salmonella* and *Escherichia coli* isolates from a pig farm in Great Britain, *J. Antimicrob. Chemother.* 71 (8) (2016) 2306–2313, <https://doi.org/10.1093/jac/dkw149>.
- [39] X.F. Zhang, Y. Doi, X. Huang, H.Y. Li, L.L. Zhong, K.J. Zeng, Y.F. Zhang, S. Patil, G. B. Tian, Possible transmission of mcr-1-harboring *Escherichia coli* between companion animals and human, *Emerg. Infect. Dis.* 22 (9) (2016) 1679–1681, <https://doi.org/10.3201/eid2209.160464>.
- [40] D.F. Monte, A. Mem, M.R. Fernandes, L. Cerdeira, F. Esposito, J.A. Galvão, B.D.G. M. Franco, N. Lincopan, M. Landgraf, Chicken meat as a reservoir of Colistin-resistant *Escherichia coli* strains carrying mcr-1 genes in South America, *Antimicrob. Agents Chemother.* 61 (5) (2017), <https://doi.org/10.1128/AAC.02718-16> e02718-16.
- [41] A.G. Sacramento, M.R. Fernandes, F.P. Sellera, M.E. Muñoz, R. Vivas, S. S. Dolabella, N. Lincopan, Genomic analysis of MCR-1 and CTX-M-8 co-producing *Escherichia coli* ST58 isolated from a polluted mangrove ecosystem in Brazil, *J Glob Antimicrob Resist.* 15 (2018) 288–289, <https://doi.org/10.1016/j.jgar.2018.10.024>.
- [42] G. Cilia, B. Turchi, F. Fratini, V.V. Ebani, L. Turini, D. Cerri, F. Bertelloni, Phenotypic and genotypic resistance to colistin in *E. coli* isolated from wild boar (*Sus scrofa*) hunted in Italy, *Eur. J. Wildl. Res.* 67 (2021) 57, <https://doi.org/10.1007/s10344-021-01501-6>.
- [43] J. Bonnedahl, M. Drobni, M. Gauthier-Clerc, J. Hernandez, S. Granholm, Y. Kayser, A. Melhus, G. Kahlmeter, J. Waldenström, A. Johansson, B. Olsen, Dissemination of *Escherichia coli* with CTX-M type ESBL between humans and yellow-legged gulls in the south of France, *PLoS One* 4 (6) (2009), e5958, <https://doi.org/10.1371/journal.pone.0005958>.
- [44] C.A. Ahlstrom, J. Bonnedahl, H. Woksepp, J. Hernandez, B. Olsen, A.M. Ramey, Acquisition and dissemination of cephalosporin-resistant *E. coli* in migratory birds sampled at an Alaska landfill as inferred through genomic analysis, *Sci. Rep.* 8 (1) (2018) 7361, <https://doi.org/10.1038/s41598-018-25474-w>.
- [45] M. Ruzauskas, L. Vaskeviciute, Detection of the mcr-1 gene in *Escherichia coli* prevalent in the migratory bird species *Larus argentatus*, *J. Antimicrob. Chemother.* 71 (8) (2016) 2333–2334, <https://doi.org/10.1093/jac/dkw245>.
- [46] C. Atterby, A.M. Ramey, G.G. Hall, J. Järhult, S. Börjesson, J. Bonnedahl, Increased prevalence of antibiotic-resistant *E. coli* in gulls sampled in southcentral Alaska is associated with urban environments, *Infect Ecol Epidemiol.* 6 (2016) 32334, <https://doi.org/10.3402/iee.v6.32334>.
- [47] C.A. Ahlstrom, J. Bonnedahl, H. Woksepp, J. Hernandez, J.A. Reed, L. Tibbitts, B. Olsen, D.C. Douglas, A.M. Ramey, Satellite tracking of gulls and genomic characterization of faecal bacteria reveals environmentally mediated acquisition and dispersal of antimicrobial-resistant *Escherichia coli* on the Kenai Peninsula, Alaska, *Mol Ecol.* 28 (10) (2019) 2531–2545, <https://doi.org/10.1111/mec.15101>.
- [48] S. Mukerji, M. Stegger, A.V. Truswell, T. Laird, D. Jordan, R.J. Abraham, A. Harb, M. Barton, M. O’Dea, S. Abraham, Resistance to critically important antimicrobials in Australian silver gulls (*Chroicocephalus novaehollandiae*) and evidence of anthropogenic origins, *J. Antimicrob. Chemother.* 74 (9) (2019) 2566–2574, <https://doi.org/10.1093/jac/dkz242>.
- [49] A. Liakopoulos, D.J. Mevius, B. Olsen, J. Bonnedahl, The colistin resistance mcr-1 gene is going wild, *J. Antimicrob. Chemother.* 71 (8) (2016) 2335–2336, <https://doi.org/10.1093/jac/dkw262>.
- [50] R.T. Torres, M.V. Cunha, D. Araujo, H. Ferreira, C. Fonseca, J.D. Palmeira, Emergence of colistin resistance genes (mcr-1) in *Escherichia coli* among widely distributed wild ungulates, *Environ. Pollut.* 291 (2021), 118136, <https://doi.org/10.1016/j.envpol.2021.118136>.
- [51] T. Bachiri, R. Lalaoui, S. Bakour, M. Allouache, N. Belkebla, J.M. Rolain, A. Touati, First report of the plasmid-mediated Colistin resistance gene mcr-1 in *Escherichia coli* ST405 isolated from wildlife in Bejaia, Algeria, *Microb. Drug Resist.* 24 (7) (2018) 890–895, <https://doi.org/10.1089/mdr.2017.0026>.
- [52] X. Lu, X. Xiao, Y. Liu, S. Huang, R. Li, Z. Wang, Widespread prevalence of plasmid-mediated colistin resistance Genemcr-1 in *Escherichia coli* from Père David’s Deer in China, *mSphere* 5 (6) (2020), <https://doi.org/10.1128/mSphere.01221-20> e01221-20.
- [53] M. Rega, I. Carmosino, P. Bonilauri, V. Frascolla, A. Vismarra, C. Bacci, Prevalence of ESBL, AmpC and Colistin-Resistant *E. coli* in meat: a comparison between pork and wild boar, *Microorganisms* 9 (2021) 214, <https://doi.org/10.3390/microorganisms9020214>.
- [54] D.G.J. Larsson, C.F. Flach, Antibiotic resistance in the environment, *Nat. Rev. Microbiol.* 4 (2021) 1–13, <https://doi.org/10.1038/s41579-021-00649-x>.