

Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Response to Comment on "Minimal and Maximal Models to Quantitate Glucose Metabolism: Tools to Measure, to Simulate and to Run in Silico Clinical Trials"

Original Citation:

Availability:

This version is available at: 11577/3454724 since: 2022-09-01T16:44:53Z

Publisher:

SAGE

Published version:

DOI: 10.1177/19322968211060069

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

Response to Comment on “Minimal and Maximal Models to Quantitate
Glucose Metabolism: Tools to Measure, to Simulate and to
Run in Silico Clinical Trials”

Cobelli C., Ph.D¹ and Dalla Man C., Ph.D²

Author Affiliations: ¹Department of Woman and Child’s Health
University of Padova, Padova, Italy; ²Department of Information
Engineering, University of Padova, Padova, Italy

Claudio Cobelli
Department of Woman
and Child’s Health,
University of Padova
Via N. Giustiniani, 3
35128 Padova, Italy
cobelli@dei.unipd.it
+39 335 6055945

Chiara Dalla Man
Department of Information
Engineering,
University of Padova
Via Gradenigo 6B
35131 Padova, Italy
dallaman@dei.unipd.it
+39 0498277713

Keywords: oral minimal model, structural identifiability, numerical identifiability, glucose effectiveness, insulin sensitivity

Corresponding Author: Claudio Cobelli, Department of Woman and Child’s Health, University of Padova, Via N. Giustiniani, 3 35128 Padova, Italy. email address: cobelli@dei.unipd.it.

Funding Source: This study was supported by MIUR (Italian Minister for Education), Departments of Excellence, Law 232/2016 and by the European Commission HORIZON2020 - FET FORGETDIABETES, EU951933.

Conflict-of-Interest Disclosure: None

Acknowledgements: None

We thank Eichenlaub and coworkers for their interesting letter to the editor entitled *Comment on “Minimal and Maximal Models to Quantitate Glucose Metabolism: Tools to Measure, to Simulate and to Run in Silico Clinical Trials”*. When developing the original model (1), we acknowledged that the need of fixing S_G (fractional glucose effectiveness) to a population value was an important limitation of the method. To test the implication of this assumption on the model-derived insulin sensitivity, S_I , we performed an extensive validation work against independent techniques, including glucose clamp (2) and multiple tracer experiment (3), and we were able to prove that S_I is well correlated with model-independent indices. Eichenlaub and coworkers’ overlook the S_I validation studies (2,3) that fully address their concerns: the correlation was 0.81 $p < 0.001$ with clamp in 10 normal and 11 impaired glucose tolerant subjects (2) and 0.86, $p < 0.0001$ with the multiple tracer experiment in 88 healthy individuals (3). Therefore, their critiques cannot be sustained upfront such a large body of validation evidence (2, 3).

In addition, Eichelelaub and coworkers have only considered the *a priori (structural)* identifiability, which is a necessary but not sufficient condition of the identification problem: it is mandatory to consider also the *a posteriori (numerical)* identifiability. In fact, parameter S_G can only be estimated with poor precision during an oral challenge. To overcome this problem one has to use a priori knowledge either by fixing S_G to a population value or by using a Bayesian prior on GEZI (glucose effectiveness at zero insulin) as detailed in the Appendix of (4). Initially, the population value strategy was successfully used as documented in the validation studies (2, 3). More recently, we adopted a novel strategy, which allows to avoid to choose a population value by introducing the parameter GEZI and estimating it with a Bayesian Maximum a Posteriori estimator, which we propose as the recommended method.

References

1. Dalla Man C, Caumo A, Cobelli C. The oral glucose minimal model: estimation of insulin sensitivity from a meal test. *IEEE Trans Biomed Eng.* 2002;49(5):419-429.
2. Dalla Man C, Yarasheski KE, Caumo A, Robertson H, Toffolo G, Polonsky KS, Cobelli C. Insulin sensitivity by oral glucose minimal models: validation against clamp. *Am J Physiol Endocrinol Metab* 289: E954–E959, 2005.
3. Dalla Man C, Caumo A, Basu R, Rizza R, Toffolo G, Cobelli C. Minimal model estimation of glucose absorption and insulin sensitivity from an oral test: validation with a tracer method. *Am J Physiol Endocrinol Metab* 287: E637–E643, 2004.
4. Basu A, Dalla Man C, Basu R, Toffolo G, Cobelli C, Rizza RA. Effects of Type 2 diabetes on insulin secretion, insulin action, glucose metabolism. *Diabetes Care* 32:866-72, 2009.