

Article

Effects of Garlic Powder and Salt on Meat Quality and Microbial Loads of Rabbit Burgers

Simone Mancini ^{1,*}, Simona Mattioli ², Roberta Nuvoloni ^{1,3}, Francesca Pedonese ^{1,3},
Alessandro Dal Bosco ² and Gisella Paci ^{1,3}

¹ Department of Veterinary Sciences, University of Pisa, Viale delle Piagge 2, 56124 Pisa, Italy; roberta.nuvoloni@unipi.it (R.N.); francesca.pedonese@unipi.it (F.P.); gisella.paci@unipi.it (G.P.)

² Department of Agricultural, Food and Environmental Science, University of Perugia, Borgo XX Giugno 74, 06100 Perugia, Italy; simona.mattioli@hotmail.it (S.M.); alessandro.dalboscio@unipg.it (A.D.B.)

³ Interdepartmental Research Center “Nutraceuticals and Food for Health”, University of Pisa, Via del Borghetto 80, 56124 Pisa, Italy

* Correspondence: simone.mancini@unipi.it

Received: 9 July 2020; Accepted: 28 July 2020; Published: 31 July 2020



Abstract: The aim of the research study was to evaluate the effects of a common culinary spice such as garlic powder and salt addition on the quality and microbial shelf life of rabbit meat burgers. Rabbit burgers were evaluated for pH, the colour parameters, the water holding capacity and microbial loads during storage time of seven days at 4 °C. Four different formulations of burgers ($n = 180$ in total) were tested as control samples (only meat, C), burgers with garlic powder (at 0.25%, G), burgers with salt (at 1.00%, S) and burgers with both garlic powder and salt (0.25% and 1.00%, respectively, GS). As results, it was highlighted that garlic powder and salt addition significantly affected pH, water holding capacity and some colour parameters of burgers. In particular, salt affected the pH of the raw burgers, leading to lower values that partially influenced all the colour parameters with higher a^* values of S burgers. The mix of garlic powder and salt (GS burgers) showed mixed effects even if more closed to the G burgers than S ones. Salt expressed its properties of binding water molecules reducing drip and cooking losses in S and GS burgers. No variations in microbial loads were highlighted in relation to the formulations. Storage time affected all the parameters, highlighting a deterioration of the burgers' quality and an increase of the microbial loads.

Keywords: spice; ingredient; colour; ready-to-cook; meat preparation

1. Introduction

Decreasing rabbit meat consumption is spreading around the world, and the Mediterranean basin in particular, where rabbit meat was historically consumed, displays a decreasing per-capita consumption [1,2]. Other kinds of meat and the low amount of time available for cooking are affecting rabbit meat consumption, mostly among young consumers looking for more approachable kinds of food [3]. Indeed, rabbits are normally sold as pre-packed whole carcasses or cut-up (such as hind legs and loin) [4,5]. Production of meat products with rabbit meat could be a response to increase consumers' willingness to purchase this product.

Burgers are extensively consumed as fast meals and are recognizable and well known worldwide. Burgers represent an appetizing and easy to cook protein food that does not require long cooking processes or culinary preparation. This popularity and high rate of consumption have driven the lifestyle changes that drive consumers to also prefer ready-to-cook products in meat sector.

Burgers, and in general meat products, could represent a way to reintroduce rabbit meat to daily consumption. Indeed, in the last years attention has been paid to several different types of meat

used as a basis for protein burgers as this type of product could be a solution to meet the consumers' willingness to purchase. Moreover, these combinations raise rabbit meat to a right market level of importance also in relation to its nutritional value, especially for children and the elderly [2].

Normally, burgers are sold as a ready-to-cook product, and several different recipes are available on market shelves to attract consumers' attention. Burgers are sold as meat mixed with spices and other ingredients among which salt is always present. Spices and herbs are used as flavours, colours and aroma enhancers and also as preservatives due to their phytochemicals [6]. Salt (NaCl, sodium chloride) due to its chemical properties, mostly the ability to reduce water activity, can also play a role as a bacteriostatic in meat products that contain a low level of salt, typically 1–3%, increasing the overall safety of the food product [7].

In this study we tested garlic (*Allium sativum* L.), a widely used spice in the Mediterranean basin, in rabbit recipes. Garlic, besides its importance as flavour carrier and sensory characteristics, can also play an important role as an antimicrobial [8,9]. In this work burgers added with salt alone and garlic powder and salt mixes were tested in order to identify more products that could reliably meet consumers' liking. Although rabbit burgers are already on the market in some European countries where cuniculture and rabbit meat consumption have a historical tradition, such as Italy and Spain, the products have a low market penetration.

Therefore, the aim of this study was to evaluate different effects of garlic powder and salt, alone and in combinations, on burgers quality and microbial loads during refrigerated storage in order to increase our knowledge about rabbit meat products.

2. Materials and Methods

2.1. Experimental Design and Burgers Manufacture

Frozen rabbit hind legs (−20 °C, 1 month of frozen storage), derived from hybrid rabbits reared under intensive conditions (ninety day-old, 2.7 ± 0.30 kg) and fed commercial pelleted feed, were thawed for 18 h at 4 °C and then deboned. Meat was finely ground with a DN30323 meat mincer (DiNa Professional, Catania, Italy) into twelve meat batches, as previously reported [10,11]. Four different formulations (three batches per formulation) were performed as: control (C, only meat); meat supplemented with 0.25% of garlic powder (G); meat supplemented with 1.00% of salt (S); meat supplemented with 0.25% of garlic powder and 1.00% of salt (GS). Garlic powder and salt were purchased as food ingredients (garlic powder produced by Drogheria e Alimentari S.p.A., Florence, Italy, batch number: L010545; salt was sea salt type, NaCl).

Each batch was hand mixed and fifteen burgers of 100 g were formatted with a DN8097 forming machine (DiNa Professional, Catania, Italy; diameter 100 mm), for a total of forty-five burgers per formulation and 180 burgers in total. Burgers were then packaged in single Styrofoam trays, overwrapped with polyethylene film and stored raw at 4 ± 0.5 °C. At the fixed storage times (day 0, 4 and 7 of storage; D0, D4 and D7) burgers from each batch were tested as raw samples and as cooked samples.

Raw and cooked burgers were analysed at D0, D4 and D7 for the determination of the pH, the colour parameters, the water holding capacity and microbial loads (performed only on raw samples).

2.2. pH

The pH was measured using a pH meter (Eutech pH2700 Meter, Eutech Instruments Pte Ltd., Singapore) equipped with a XS Sensor Standard S7 (XS Sensor, Modena, Italy) and an automatic temperature compensator. pH meter was calibrated before each session with buffer solutions at pH 4.01 and 7.01 (HI7004L and HI7007L Hanna Instruments, Padova, Italy).

2.3. Colour

Colour was expressed as L^* (lightness), a^* (redness), and b^* (yellowness) according to the CIElab system [12]. Colour parameters were measured using a Minolta CR300 chroma meter (Minolta, Osaka, Japan) with an aperture size of 8 mm, illuminant D65 and incidence angle of 0° . Before each session, the colorimeter was calibrated with a white tile ($L^* = 98.14$, $a^* = -0.23$ and $b^* = 1.89$). Each data point was the mean of three replications measured on the surface of the burgers at randomly selected locations. Chroma (C^*) and Hue angle (h^*) were calculated as function of a^* and b^* following the formulas:

$$C^* = \sqrt{(a^{*2} + b^{*2})}$$

$$h^* = \tan^{-1}\left(\frac{b^*}{a^*}\right)$$

The total colour difference (ΔE) was calculated as proposed by Sharma and Bala [13] between two different formulations at the same storage time or between two different storage times for the same formulation following the formula:

$$\Delta E_{\alpha-\beta} = \sqrt{(L_{\alpha}^* - L_{\beta}^*)^2 + (a_{\alpha}^* - a_{\beta}^*)^2 + (b_{\alpha}^* - b_{\beta}^*)^2}$$

where α and β subscripts of L^* , a^* and b^* referred to two different formulations at the same storage time or two different storage times for the same formulation. Cooking effect on colour were determined for each F at each storage time (ST). As proposed by Sharma and Bala [13], the threshold of a human noticeable difference was fixed at 2.3 points.

2.4. Water Holding Capacity

Drip loss was calculated as proposed by Lundström and Malmfors [14] within the F between D0-D4 and D0-D7. Cooking loss was calculated as percentage of the decrease of weight before and after cooking in a preheated oven at 163°C to an internal temperature of 71°C and were turned every 4 min to prevent excess surface crust formation [15].

2.5. Microbial Quantifications

Ten grams of sample were aseptically removed and homogenised in a Stomacher 400 Circulator Lab Blender (Seward, Worthing, UK) with 90 mL of 0.1% peptone salt solution. Further serial dilutions were made in the same diluent and used for standard plate enumerations.

Total aerobic mesophilic and psychrotrophic bacteria were determined on Plate Count Agar (pour plate method) with incubation at 30°C for 72 h, and 7°C for 10 days, respectively; Enterobacteriaceae on Violet Red Bile Glucose Agar at 37°C for 24 h; *Escherichia coli* on Tryptone Bile X-Glucuronide Medium (TBX) at 44°C for 24 h, lactic acid bacteria on MRS Agar (pour plate method) in anaerobiosis (Anaerogen 2.5L) at 30°C for 72 h; *Brochothrix thermosphacta* on Streptomycin Thallous Acetate (STA) agar with STA selective supplement at 25°C for 48 h; *Pseudomonas* spp. on Pseudomonas Agar base with CFC supplement at 25°C for 72 h; yeasts and moulds on Yeast Extract Glucose Chloramphenicol Agar (pour plates method) after incubation at 25°C for 5 days. Where not specified, spread plate method was used. All cultural media and supplements were from Oxoid (Basingstoke, UK). The bacterial counts were expressed as log Colony-Forming Units (CFU) per gram of sample.

2.6. Statistical Analysis

The effects of the formulation (F), of the storage time (ST) and their interaction ($F \times ST$) on the burger parameters were analysed through a two-way ANOVA using the R software Version 1.2.5019 (R Core Team, The R Foundation for Statistical Computing, Vienna, Austria) [16]. The significance

level was set at 5% (statistically significant for $p < 0.05$), and if statistical significance was found, the differences were assessed using Tukey's test ($p < 0.05$). When the interaction $F \times ST$ was not significant the results are reported as the mean of the fixed effects F and ST ; the variability is expressed as root mean square error (RMSE).

3. Results and Discussion

Results of the pH, the colour parameters and the water holding capacity of the raw burgers are reported in Table 1. The formulation (F) significantly effected all the tested parameters. Furthermore, also storage time influenced quite all the parameters, indeed, only L^* and b^* coordinates did not show significant differences for ST ($p = 0.099$ and $= 0.066$, respectively). The addition of salt lead to lower pH values in average, as evidenced by the pH of S and GS formulations in relation to C and G burgers ($p = 0.002$). These differences in pH values might played a role in the colour values; indeed L^* of S and GS burgers were lower than C and G ones ($p < 0.001$), as pH and lightness are linked by a negative correlation. Additions of garlic powder modified the redness (a^*) and yellowness (b^*) coordinates ($p < 0.001$ and $p = 0.013$, respectively). Natural pigments in garlic powder lead to a pale-yellow raw product than when processed turn to a strong yellow colour turning more vivid. Due to degradation processes the garlic pigments could produce a green-yellow tone [17–19]. Garlic natural pigments showed their effect in G and GS burgers with the reduction of redness value, as well as an increase of yellowness value. Additions of ingredients, both salt and garlic powder and its mix, decreased the chroma index in relation to the C burgers, beside S and GS burgers showed a greater decrement in chroma as salt addition affected negatively both a^* and b^* coordinates ($p < 0.001$). On the other hand, garlic induced an increase in h^* leading to light-yellow burgers in colour due to the increase of b^* value at the expense of a^* value ($p < 0.001$).

Table 1. pH, colour parameters and water holding capacity of raw rabbit burgers.

Item	Formulation (F)				Storage Time (ST, Days)			p-Value			RMSE
	C	G	S	GS	D0	D4	D7	F	ST	F × ST	
pH	5.97 ^a	5.96 ^a	5.87 ^b	5.86 ^b	5.89 ^y	5.89 ^y	5.97 ^x	0.002	0.002	0.360	0.058
L^*	57.24 ^a	57.65 ^a	51.31 ^b	51.04 ^b	55.38	53.80	53.75	<0.001	0.099	0.861	2.769
a^*	5.86 ^a	4.62 ^b	5.53 ^a	4.09 ^b	5.57 ^x	5.34 ^x	3.97 ^y	<0.001	<0.001	0.343	0.759
b^*	6.07 ^b	7.43 ^a	5.93 ^b	6.41 ^{ab}	7.04	5.97	6.37	0.013	0.066	0.252	0.938
C^*	10.34 ^a	8.83 ^{ab}	8.29 ^b	7.71 ^b	10.19 ^x	8.10 ^y	8.07 ^y	<0.001	<0.001	0.234	1.209
h^*	40.70 ^b	58.46 ^a	48.19 ^b	58.70 ^a	44.22 ^y	49.34 ^y	60.97 ^x	<0.001	<0.001	0.224	6.250
Drip loss%	0.63 ^a	0.77 ^a	0.40 ^b	0.44 ^b	0.00 ^z	0.73 ^y	0.95 ^x	0.016	<0.001	0.316	0.835
Cooking loss%	24.33 ^a	22.00 ^a	17.83 ^b	16.15 ^b	21.07 ^x	21.17 ^x	17.99 ^y	<0.001	0.032	0.828	0.024

C: control; G: control + 0.25% of garlic powder; S: control + 1% of salt; GS: control + 0.25% of garlic powder + 1% salt. ^{a,b} Different letters in the same row indicate significant differences at $p < 0.05$ for F . ^{x,y,z} Different letters in the same row indicate significant differences at $p < 0.05$ for ST .

During storage time a slight increase in pH was revealed at D7 ($p = 0.002$), that might be imputable to an alkalisation of meat resulting from an increase in ammoniacal nitrogen levels and to the degradation of proteins and amino acids by Gram-negative bacteria [20–22]. Also the reduction of a^* and C^* and the increase of h^* during storage time ($p < 0.001$ for all indexes) could be ascribed to bacteria metabolism actions and due to the formation of metmyoglobin produced by myoglobin oxidation [23,24].

Water holding capacity was affected principally from salt addition as both S and GS burgers showed lowest drip and cooking losses ($p = 0.016$ and $p < 0.001$, respectively), indeed salt due to its chemical properties contributes to water and fat binding in meat products and this property is enhanced by mincing processing [25–27]. As expected during storage time drip loss increased due to the natural water release, furthermore, the loss of water by the raw product during the storage lead to day 7 at more dry samples with a consequence of a lowest cooking loss.

Cooking flattered the formulation differences, with statistical evidence only on L^* and h^* parameters (Table 2). After cooking the presence of salt in the samples affected the L^* value with lower values of S and GS than C and G samples ($p < 0.001$), following the trend reported in raw samples. Even if a^* and

b* coordinates were not affected by F ($p = 0.065$ and $p = 0.278$, respectively) the h* index revealed a difference in colour between C and the other formulation that appeared lighter in colours ($p = 0.004$). During storage time a rise in b* value was highlighted ($p < 0.001$) as degradation of the pale pink colour of rabbit meat and the formation of yellowness complex. As consequence of b* value rise also both C* and h* increased their values meaning a lighter vivid yellowness samples at D7 than D0 ($p < 0.001$ and $p = 0.032$, respectively).

Table 2. pH and colour parameters of cooked rabbit burgers.

Item	Formulation (F)				Storage Time (ST, Days)			p-Value			RMSE
	C	G	S	GS	D0	D4	D7	F	ST	F × ST	
pH	6.13	6.11	6.07	6.06	6.10	6.07	6.11	0.218	0.053	0.893	0.102
L*	69.70 ^a	70.60 ^a	66.43 ^b	65.31 ^b	67.76	68.61	67.66	<0.001	0.452	0.201	2.278
a*	7.19	6.48	6.09	6.18	6.29	6.21	6.95	0.065	0.567	0.084	0.959
b*	14.77	14.27	15.23	16.14	14.09 ^y	14.58 ^y	16.64 ^x	0.278	<0.001	0.522	1.341
C*	16.43	15.70	16.42	17.29	15.44 ^y	15.86 ^y	18.09 ^x	0.567	<0.001	0.291	1.436
H*	64.09 ^b	65.98 ^a	68.02 ^a	68.97 ^a	65.90 ^y	66.97 ^x	67.42 ^x	0.004	0.032	0.102	2.563

C: control; G: control + 0.25% of garlic powder; S: control + 1% of salt; GS: control + 0.25% of garlic powder + 1% salt. ^{a,b} Different letters in the same row indicate significant differences at $p < 0.05$ for F. ^{x,y} Different letters in the same row indicate significant differences at $p < 0.05$ for ST.

Cooking might affect several chemical, physical, and even biological characteristics of the products. As burger require a cooking section to be eaten it is important how this final step is performed in order to maintain the chemical and nutritional properties added via formulation and physical properties related to sensory acceptance by consumers [28,29].

Colour differences (ΔE s) within F between ST are reported in Table 3; colour differences (ΔE s) within ST between F are reported in Table 4. All the F changed in a noticeable way the overall colour during the 7 days of storage (D0–D7). The addition of the sole garlic powder induced a strong variation colour after 4 days as reported by the ΔE value of D0–D4 period. That might be related to the rapid oxidation of the garlic compounds and the formation of a green-yellow hue that mitigate the pink rabbit meat colour. This modification in raw burgers affected also the ΔE between cooked burgers at D0 versus D4 as G formulation was the only one to reported values over the threshold of 2.3 points. On the contrary C, GS S cooked burgers showed higher variation in colour between D4 and D7.

Table 3. Colour differences (ΔE) within Formulation (F) between Storage Time (ST).

Formulation (F)	Storage Time (ST, Days)								
	Raw Samples			Cooked Samples			Raw–Cooked Samples		
	D0–D4	D4–D7	D0–D7	D0–D4	D4–D7	D0–D7	D0	D4	D7
C	2.21	3.09*	5.18*	0.97	3.38*	4.32*	13.65*	14.93*	17.42*
G	4.66*	2.51*	3.57*	3.69*	6.96*	3.58*	12.63*	18.39*	14.51*
GS	2.72*	2.57*	5.18*	2.78*	3.06*	4.50*	15.76*	19.30*	20.05*
S	2.48*	2.24	4.33*	1.96	3.11*	2.37*	15.27*	16.57*	20.34*

* Value over the threshold (2.3 points) with a noticeable difference in colour between the samples. C: control; G: control + 0.25% of garlic powder; S: control + 1% of salt; GS: control + 0.25% of garlic powder + 1% salt.

Table 4. Colour differences (ΔE) within Storage Time (ST) between Formulation (F).

Formulation (F)	Storage Time (ST, Days)											
	D0				D4				D7			
	C	G	S	GS	C	G	S	GS	C	G	S	GS
C		2.81*	8.21*	7.21*		4.84*	7.29*	7.39*		4.94*	5.37*	4.94*
G	0.67		8.27*	6.64*	4.03*		6.15*	5.27*	0.51		7.02*	8.10*
S	5.77*	5.40*		1.73	2.67*	6.26*		3.70*	2.84*	2.41*		1.15
GS	5.34*	4.82*	2.25		6.21*	9.93*	1.88		3.53*	3.06*	1.68	

* Value over the threshold (2.3 points) with a noticeable difference in colour between the samples. C: control; G: control + 0.25% of garlic powder; S: control + 1% of salt; GS: control + 0.25% of garlic powder + 1% salt. In the columns of ΔE s within ST between F the values in italic or bold refer respectively to raw and cooked samples.

Colour differences at the same storage time between F highlighted that salt addition strongly changed the burgers' colour at D0 in both raw and cooked samples, reaching high ΔE s between S and GS in relation to C and G formulations. This tendency was maintained also at D4 even though also G burgers increase their colour distances from the control burgers. After cooking no noticeable differences was reported between S and GS burgers, countering the garlic effect on colour. After 7 days of storage the cooked burgers showed the lowest ΔE values between the formulations highlighting that the oxidation process occurred by the time affected all the samples inducing a general colour variation.

In Table 5 are reported the microbial loads of the raw burgers. No statistical differences were highlighted for the F main factor; thus, no effect of salt or garlic addition was evidenced on the microbial load. *Escherichia coli* and *Brochothrix thermosphacta* were not detected in the burgers both in relation to the formulation and the storage time. All the detected bacterial loads increased during storage time ($p < 0.001$ for all the parameters), mostly with differences between each fixed day of analysis. Only the total aerobic psychrotrophic bacteria showed to reach the highest value at D4 and to maintain it at D7.

Table 5. Microbial determinations on raw rabbit burgers.

Item	Formulation (F)				Storage Time (ST, Days)			p-Value			RMSE
	C	G	S	GS	D0	D4	D7	F	ST	F × ST	
Enterobacteriaceae	3.80	3.94	3.77	3.94	1.59 ^z	3.96 ^y	6.04 ^x	0.824	<0.001	0.691	0.445
<i>Pseudomonas</i> spp.	5.41	5.31	4.91	4.96	2.23 ^z	5.73 ^y	7.48 ^x	0.507	<0.001	0.984	0.748
Lactic acid bacteria	3.49	3.79	3.42	3.99	2.21 ^z	3.36 ^y	5.44 ^x	0.633	<0.001	0.969	0.923
Yeast and moulds	3.69	3.44	3.19	3.09	1.99 ^z	3.34 ^y	4.73 ^x	0.355	<0.001	0.665	0.686
Total aerobic mesophilic bacteria	6.62	6.49	6.05	6.21	4.22 ^z	6.73 ^y	8.08 ^x	0.293	<0.001	0.692	0.603
Total aerobic psychrotrophic bacteria	6.29	5.71	5.95	5.53	3.61 ^y	6.42 ^x	7.58 ^x	0.703	<0.001	0.794	1.299

C: control; G: control + 0.25% of garlic powder; S: control + 1% of salt; GS: control + 0.25% of garlic powder + 1% salt. ^{x,y,z} Different letters in the same row indicate significant differences at $p < 0.05$ for ST.

Spices and other ingredients added into minced meat products, such as burgers, might affect in different ways the growth of microorganisms. Turmeric and ginger powders showed a bacteriostatic effect against several different bacteria in rabbit burgers stored at 4 °C [20,23]. Similarly, pork burgers/patties supplemented with different plant products, such as passion fruit co-products and tea or grape extracts, showed a lower bacterial growth than the respective control treatment [30–32]. Ingredients' activity against microorganisms' growth might be also related to the physical form and technological transformations, to the employed concentrations or to the meat used. Indeed, Sallam et al. [33] reported activities against aerobic plate count of fresh garlic (30 g/kg) and garlic powder (9 g/kg) added in chicken sausages stored at 3 °C up to 21 days. Also Aydin et al. [34] reported activities of fresh garlic (10%) in ground beef refrigerated for 24 h in terms of total aerobic mesophilic bacteria and coliform bacteria.

4. Conclusions

The additions of garlic powder and salt to rabbit meat could bring several characteristics modifications to burgers along with culinary perceptions. Both garlic powder and salt also played a role in the colour changes in relation to the storage time. No effects on the microbial loads suggest that higher concentrations of garlic powder or salt are needed if their use is to be also intended as bacteriostatic additives. Different garlic products such as fresh minced or extracts could produce higher/lower beneficial effects, thus further studies are needed to better the potential application of this spice and how addition of salt and/or garlic could affect burgers' flavour and consumers' acceptance.

Author Contributions: Conceptualization, S.M. (Simone Mancini), R.N. and G.P.; methodology, S.M. (Simone Mancini) and G.P.; formal analysis, S.M. (Simone Mancini), R.N. and F.P.; writing—original draft preparation, S.M. (Simone Mancini); writing—review and editing, S.M. (Simone Mancini), S.M. (Simona Mattioli), R.N., F.P., A.D.B. and G.P.; supervision, G.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kallas, Z.; Gil, J.M. A dual response choice experiments (DRCE) design to assess rabbit meat preference in Catalonia: A heteroscedastic extreme-value model. *Br. Food J.* **2012**, *114*, 1394–1413. [[CrossRef](#)]
2. Petracci, M.; Soglia, F.; Baldi, G.; Balzani, L.; Mudalal, S.; Cavani, C. Technical note: Estimation of real rabbit meat consumption in Italy. *World Rabbit Sci.* **2018**, *26*, 91–96. [[CrossRef](#)]
3. González-Redondo, P.A.; Contreras-Chacón, G.M. Perceptions among university students in seville (SPAIN) of the rabbit as livestock and as a companion animal. *World Rabbit Sci.* **2012**, *20*, 155–162. [[CrossRef](#)]
4. Petracci, M.; Cavani, C. Rabbit meat processing: Historical perspective to future directions. *World Rabbit Sci.* **2013**, *21*, 217–226. [[CrossRef](#)]
5. Zotte, A.D. Rabbit farming for meat purposes. *Anim. Front.* **2014**, *4*, 62–67. [[CrossRef](#)]
6. Embuscado, M.E. Spices and herbs: Natural sources of antioxidants—A mini review. *J. Funct. Foods* **2015**, *18*, 811–819. [[CrossRef](#)]
7. Betts, G.; Everis, L.; Betts, R. Microbial issues in reducing salt in food products. In *Reducing Salt in Foods*; Elsevier: Amsterdam, The Netherlands, 2007; pp. 174–200, ISBN 9781845690182.
8. Casella, S.; Leonardi, M.; Melai, B.; Fratini, F.; Pistelli, L. The role of diallyl sulfides and dipropyl sulfides in the in vitro antimicrobial activity of the essential oil of garlic, *Allium sativum* L., and leek, *Allium porrum* L. *Phyther. Res.* **2013**, *27*, 380–383. [[CrossRef](#)]
9. Chung, L.Y. The antioxidant properties of garlic compounds: Allyl cysteine, alliin, allicin, and allyl disulfide. *J. Med. Food* **2006**, *9*, 205–213. [[CrossRef](#)]
10. Mancini, S.; Nuvoloni, R.; Pedonese, F.; Paci, G. Effects of garlic powder and salt additions in rabbit meat burgers: Preliminary evaluation. *J. Food Process. Preserv.* **2019**, *43*, e13894. [[CrossRef](#)]
11. Mancini, S.; Mattioli, S.; Nuvoloni, R.; Pedonese, F.; Dal Bosco, A.; Paci, G. Effects of garlic powder and salt additions on fatty acids profile, oxidative status, antioxidant potential and sensory properties of raw and cooked rabbit meat burgers. *Meat Sci.* **2020**, *169*, 108226. [[CrossRef](#)]
12. Commission Internationale de l’Eclairage. *Official Recommendations on Uniform Colour Spaces, Colour Differences Equations and Metric Colour Terms*; CIE: Paris, France, 1976.
13. Sharma, G.; Bala, R. *Digital Color Imaging Handbook*; Sharma, G., Ed.; Electrical Engineering & Applied Signal Processing Series; CRC Press: Boca Raton, FL, USA, 2002; Volume 11, ISBN 978-0-8493-0900-7.
14. Lundström, K.; Malmfors, G. Variation in light scattering and water-holding capacity along the porcine Longissimus dorsi muscle. *Meat Sci.* **1985**, *15*, 203–214. [[CrossRef](#)]
15. AMSA. *Research Guidelines for Cookery, Sensory Evaluation and Instrumental Tenderness Measurements of Fresh Meat*; National Live Stock and Meat Board: Chicago, IL, USA, 1995.
16. R Core Team. *R: A Language and Environment for Statistical Computing*; The R Foundation for Statistical Computing: Vienna, Austria, 2015.
17. Lee, E.-J.; Cho, J.-E.; Kim, J.-H.; Lee, S.-K. Green pigment in crushed garlic (*Allium sativum* L.) cloves: Purification and partial characterization. *Food Chem.* **2007**, *101*, 1677–1686. [[CrossRef](#)]
18. Wang, D.; Yang, X.; Wang, Z.; Hu, X.; Zhao, G. Isolation and identification of one kind of yellow pigments from model reaction systems related to garlic greening. *Food Chem.* **2009**, *117*, 296–301. [[CrossRef](#)]
19. Wang, D.; Li, X.; Zhao, X.; Zhang, C.; Ma, Y.; Zhao, G. Characterization of a new yellow pigment from model reaction system related to garlic greening. *Eur. Food Res. Technol.* **2010**, *230*, 973–979. [[CrossRef](#)]
20. Mancini, S.; Preziuso, G.; Dal Bosco, A.; Roscini, V.; Szendrő, Z.; Fratini, F.; Paci, G. Effect of turmeric powder (*Curcuma longa* L.) and ascorbic acid on physical characteristics and oxidative status of fresh and stored rabbit burgers. *Meat Sci.* **2015**, *110*, 93–100. [[CrossRef](#)]
21. Kılıç, B.; Şimşek, A.; Claus, J.R.; Atilgan, E. Encapsulated phosphates reduce lipid oxidation in both ground chicken and ground beef during raw and cooked meat storage with some influence on color, pH, and cooking loss. *Meat Sci.* **2014**, *97*, 93–103. [[CrossRef](#)] [[PubMed](#)]
22. Rodríguez-Calleja, J.M.; García-López, M.-L.; Santos, J.A.; Otero, A. Development of the aerobic spoilage flora of chilled rabbit meat. *Meat Sci.* **2005**, *70*, 389–394. [[CrossRef](#)]
23. Mancini, S.; Preziuso, G.; Fratini, F.; Torracca, B.; Nuvoloni, R.; Dal Bosco, A.; Paci, G. Qualitative improvement of rabbit burgers using *Zingiber officinale* Roscoe powder. *World Rabbit Sci.* **2017**, *25*, 367. [[CrossRef](#)]

24. Choe, J.-H.; Jang, A.; Lee, E.-S.; Choi, J.-H.; Choi, Y.-S.; Han, D.-J.; Kim, H.-Y.; Lee, M.-A.; Shim, S.-Y.; Kim, C.-J. Oxidative and color stability of cooked ground pork containing lotus leaf (*Nelumbo nucifera*) and barley leaf (*Hordeum vulgare*) powder during refrigerated storage. *Meat Sci.* **2011**, *87*, 12–18. [[CrossRef](#)]
25. Fernández-Ginés, J.M.; Fernández-Lopez, J.; Sayas-Barbera, E.; Perez-Alvarez, J.A. Meat Products as Functional Foods: A Review. *J. Food Sci.* **2005**, *70*, R37–R43. [[CrossRef](#)]
26. Mariutti, L.R.B.; Bragagnolo, N. Influence of salt on lipid oxidation in meat and seafood products: A review. *Food Res. Int.* **2017**, *94*, 90–100. [[CrossRef](#)] [[PubMed](#)]
27. Overholt, M.F.; Mancini, S.; Galloway, H.O.; Preziuso, G.; Dilger, A.C.; Boler, D.D. Effects of salt purity on lipid oxidation, sensory characteristics, and textural properties of fresh, ground pork patties. *LWT Food Sci. Technol.* **2016**, *65*, 890–896. [[CrossRef](#)]
28. King, N.J.; Whyte, R. Does it look cooked? A review of factors that influence cooked meat color. *J. Food Sci.* **2006**, *71*, R31–R40. [[CrossRef](#)]
29. Rodriguez-Estrada, M.T.; Penazzi, G.; Caboni, M.F.; Bertacco, G.; Lercker, G. Effect of different cooking methods on some lipid and protein components of hamburgers. *Meat Sci.* **1997**, *45*, 365–375. [[CrossRef](#)]
30. López-Vargas, J.H.; Fernández-López, J.; Pérez-Álvarez, J.Á.; Viuda-Martos, M. Quality characteristics of pork burger added with albedo-fiber powder obtained from yellow passion fruit (*Passiflora edulis* var. *flavicarpa*) co-products. *Meat Sci.* **2014**, *97*, 270–276. [[CrossRef](#)] [[PubMed](#)]
31. Lorenzo, J.M.; Sineiro, J.; Amado, I.R.; Franco, D. Influence of natural extracts on the shelf life of modified atmosphere-packaged pork patties. *Meat Sci.* **2014**, *96*, 526–534. [[CrossRef](#)]
32. Mancini, S.; Paci, G.; Fratini, F.; Torracca, B.; Nuvoloni, R.; Dal Bosco, A.; Roscini, V.; Preziuso, G. Improving pork burgers quality using *Zingiber officinale* Roscoe powder (ginger). *Meat Sci.* **2017**, *129*, 161–168. [[CrossRef](#)]
33. Sallam, K.I.; Ishioroshi, M.; Samejima, K. Antioxidant and antimicrobial effects of garlic in chicken sausage. *LWT Food Sci. Technol.* **2004**, *37*, 849–855. [[CrossRef](#)]
34. Aydin, A.; Bostan, K.; Erkan, M.E.; Bingöl, B. The Antimicrobial Effects of Chopped Garlic in Ground Beef and Raw Meatball (Çiğ Köfte). *J. Med. Food* **2007**, *10*, 203–207. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).