

Editorial



## **Biological Activity of Essential Oils**

## Francesca Mancianti 1,2,\* and Valentina Virginia Ebani 1,2

- <sup>1</sup> Dipartimento di Scienze Veterinarie, Università di Pisa, Viale delle Piagge, 2 I 56124 Pisa, Italy; valentina.virginia.ebani@unipi.it
- <sup>2</sup> Centro Interdipartimentale di Ricerca Nutraceutica e Alimentazione per la Salute "Nutrafood", Università di Pisa, Via del Borghetto, 80 I 56124 Pisa, Italy
- \* Correspondence: francesca.mancianti@unipi.it

Received: 28 January 2020; Accepted: 05 February 2020; Published: 5 February 2020

Essential oils (EOs) have for a long time been recognized to possess several different biological activities. Several among these secondary plant metabolites exhibit marked antimicrobial effects that have made their use as an antiseptic and/or preservative in food well known, since the ancient times.

In this Special Issue, several oils extracted from plants growing in different environments and frequently employed in ethnomedicine were investigated in 30 original research and two review papers. This issue encompasses both in vivo and in vitro studies.

Some novel approaches are reported in both selecting and delivering them. In particular, machine learning algorithms were applied to analyze the chemical composition of the EOs to understand how it can be related to the anti-biofilm potencies [1]. On the other hand, the use of biomaterials with properties for improving antimicrobial activity is up to date, and coating hydroxyapatite with peppermint EO would significantly enhance its antibacterial effect [2].

The composition of the EOs within the same botanical species is influenced by various parameters such as time of harvest, mode of extraction, and conservation [3]. Therefore, in the present issue, a strong correlation between the chemical composition and environmental parameters (geographical position, bioclimate, and nature of soils) of individual oil samples extracted from aerial parts of *Eryngium campestre* was reported [4].

The varying composition of EOs makes it difficult to compare their antimicrobial activities, however, there is a need to understand the antimicrobial effects of EOs to gain insights into their application potential [5]. In this framework, a critical, excellent review of the antimicrobial activity of EOs is provided by Winska et al. [6], with the view to promote further investigations into targeting EOs and microorganisms.

Based on this information, in this Special Issue, most EOs have been screened for their in vitro antibacterial/antifungal activity, and in some selected papers, the micro-organisms were chosen from the perspective of a practical application to face different situations in the field. *Clausena lausium* EO showed potent activity against *Candida* yeasts, corroborating the efficacy of these traditional remedies in Chinese folk medicine [7], while *Cymbopogon flexuosum*, *Litsea cubeba*, and *Citrus bergamia* EOs appeared active against *Saprolegnia parasitica* [8]. However, attention should be given to in vivo administration of such products, being that some EOs have been proven to exert strong antimicrobial effects in animal practice where they would show toxic effects damaging the liver, kidney or gastrointestinal tissues [9].

Although the use of EOs as antimicrobials has been reported to not induce relevant changes in the sensitivity to antibacterial drugs [10], a novel insight into the mechanisms of action and *Staphylococcus aureus* resistance against active individual constituents such as carvacrol, citral, and (+)-limonene oxide was provided by Berdejo et al.[11]. Selected microorganisms, exposed to subinhibitory doses of such compounds showed a higher tolerance to lethal treatments by individual constituents or heat. This investigation suggests the need for further studies aiming to evaluate whether the emergence of resistant strains in the presence of EOs and their main components is a general phenomenon. EOs have shown remarkable healing effects in wounds both in vivo and in vitro models [14] and a contributor to the potential of wound healing by *Eugenia dysenterica* EO has also been reported [15]. Although EOs have antiviral actions against animal/human viruses [16,17], Vuko et al. [18] reported a further study on the anti-phytoviral activity of *Micromeria croatica* EO in inhibiting satellite RNA Associated Cucumber Mosaic Virus Infection.

component, curlone, appeared as a promising anti-trypanosomal candidate [13].

In a green approach to crop protection, EOs from *Coriandrum sativum* (Sumalan et al., 2019) [19] and *Solidago canadensis* appeared active against several phytopathogenic bacteria and fungi [20] as well as geranium and rosewood EOs versus *Fusarium graminearum* [21].

EOs can also act as allelochemicals, in order to defend the plants against predators and other competing plants. *Xanthium strumarium* leaves EO was successfully tested as a green bioherbicide [22], while the phytotoxic activity of *Heracleum mantegazzianum*, an alien botanical species, was assessed on monocot and dicot plants species [23].

Among other less investigated biological effects, there is an anticancer potential of some EOs, checked in vitro and/or in vivo. EO obtained from Gannanzao orange showed a significant inhibition in the proliferation and migration of hepatoma and colorectal cancer cells [24]. The latter were sensitive to EO from *Origanum onites*, which was also able to inhibit in vivo the growth of syngeneic CT26 colon tumors when used for prophylactic oral administration in a murine experimental model [25]. Of interest also is the study by de Lima et al. [26] regarding the in vivo anticancer effect of EO from the leaves of *Croton matourensis*, a medicinal plant from Brazil.

Tea seed EO showed a marked anti-obesity effect in a murine model, which was able to prevent obesity, reduce physical fatigue, and improve exercise performance, suggesting a possible use for menopause-related metabolic syndromes [27].

In a perspective to investigate the therapeutic potential of organosulfur compounds for various disorders with immune and/or inflammatory mechanisms, garlic EO and some organosulfur compounds could serve as biological response modifiers by augmenting phagocyte functions. They were in fact able to activate neutrophil functional activity, suggesting that neutrophil stimulation by organosulfur components from *Allium* spp. might enhance resistance to infection [28].

Finally, thee antispasmodic effects of EOs from 39 botanical species were reviewed by Codrura Heghes et al, [29], indicating such compounds as promising alternatives to conventional antispasmodic drugs.

**Acknowledgments:** The Guest Editor wishes to thank all the authors for their contributions to this Special Issue, all the reviewers for their work in evaluating the submitted articles, and the editorial staff of Molecules, for their kind help in making this Special Issue.

Conflicts of Interest: The author declares no conflicts of interest.

## References

- Patsilinakos, A.; Artini, M.; Papa, R.; Sabatino, M.; Bozovic, M.; Garzoli, S.; Vrenna, G.; Buzzi, R.; Manfredini, S.; Selan, L.; Ragno, R. Machine learning analyses on data including essential oil chemical composition and in vitro experimental antibiofilm activities against *Staphylococcus* species. *Molecules* 2019, 24, 890.
- Badea, M.L.; Iconaru, S.L.; Groza, A.; Chifiriuc, M.C.; Beuran, M.; Predoi, D. Peppermint essential oil-doped hydroxyapatite nanoparticles with antimicrobial properties *Molecules* 2019, 24,
- Yuan, Y.; Huang, M.; Pang, Y.X.; Yu, F.L.; Chen, C.; Liu, L.W.; Chen, Z.X.; Zhang, Y.B.; Chen, X.L.; Hu, X. Variations in essential oil yield, composition, and antioxidant activity of different plant organs from *Blumea balsamifera* (L.) DC. at different growth times. *Molecules* 2016, *21*, 1024.

- Medbouhi, A.; Benbelaid, F.; Djabou, N.; Beaufay, C.; Bendahou, M.; Quetin-Leclercq, J.; Tintaru, A.; Costa, J.; Muselli, A. Essential oil of algerian *Eryngium campestre*: Chemical variability and evaluation of biological activities. *Molecules* 2019, 24, 2575.
- 5. Van de Vel, E.; Sampers, I.; Raes, K. A review on influencing factors on the minimum inhibitory concentration of essential oils. *Crit. Rev. Food Sci. Nutr.* **2019**, *59*, 357–378.
- Winska, K.; Maczka, W.; Lyczko, J.; Grabarczyk, M.; Czubaszek, A.; Szumny, A. Essential oils as antimicrobial agents-myth or real alternative? *Molecules* 2019, 24, 2130.
- He, X.; Lantong, Z.; Chen, J.; Sui, J.; Yi, G.; Wu, J.; Ma, Y. Correlation between chemical composition and antifungal activity of *Clausena lansium* essential oil against *Candida* spp. *Molecules* 2019, 24, 1394.
- Nardoni, S.; Najar, B.; Fronte, B.; Pistelli, L. In vitro activity of essential oils against Saprolegnia parasitica. Molecules 2019, 24, 1270
- 9. Horky, P.; Skalickova, S.; Smerkova, K.; Skladanka, J. Essential oils as additives: Pharmacokinetics and potential toxicity in monogastric animals. *Animals* **2019**, *9*, 352.
- Leite de Souza, E. The effects of sublethal doses of essential oils and their constituents on antimicrobial susceptibility and antibiotic resistance among food related bacteria: A review. *Trends Food Sci Technol.* 2016 56, 1–12.
- Berdejo, D.; Checa, B.; Pagan E.; Renzoni, A.; Kelley, W.L.; Pagan, R.; Garcia-Gonzalo, D. Sub inhibitory doses of individual constituents of essential oils can select for *Staphylococcus aureus* resistant mutants. *Molecules* 2019, 24, 170.
- Margatto Rottini, M.; Fernandes Amaral, A.; Pinto Ferreira, J.L.; Souza Coelho Oliveira, E.; Rocha de Andrade Silva, J.; Nosomi Taniwaki, N.; dos Santos, AR.; Almeida Souza, F.; da Silva Freitas de Souza, C.; da Silva Calabrese, K. *Endlicheria* bracteolata (Meisn.) essential oil as a weapon against *Leishmania amazonensis*: In vitro assay. *Molecules* 2019, 24, 2525
- Binh Le, T.; Beaufay, C.; Nghiem, D.T.; Pham, T.A.; Mingeot-Leclercq M.P.; Quetin Leclercq, J. Evaluation of anti trypanosomal activity of vietnamese essential oils, with emphasis on *Curcuma longa* L. and its components. *Molecules* 2019, 24, 1158
- Perez-Recalde, M.; Ruiz Arias, I.E.; Hermida, E.B. Could essential oils enhance biopolymers performance for wound healing? A systematic review. *Phytomedicine* 2018, 38, 57–65.
- Mazutti da Silva, S.M.; Rezende Costa, C.R.; Martins Gelfuso, G.; Silva Guerra, E.N.; De Medeiros Nóbrega, Y.K.; Gomes, S.M.; Pic-Taylor, A.; Fonseca-Bazzo, Y.M.; Silveira, D.; Magalhães, P.O. Wound Healing Effect of Essential Oil Extracted from *Eugenia dysenterica* DC (Myrtaceae) Leaves. *Molecules* 2019, 24, 2.
- Schnitzler, P. Essential Oils for the Treatment of Herpes Simplex Virus Infections. *Chemotherapy*. 2019, 64, 1– 7. doi: 10.1159/000501062
- Tariq, S.; Wani, S.; Rasool, W.; Shafi, K.; Bhat, M.A.; Prabhakar, A.; Shalla, A.H.; Rather, M.A. A comprehensive review of the antibacterial, antifungal and antiviral potential of essential oils and their chemical constituents against drug-resistant microbial pathogens. *Microb Pathog.* 2019, 134, 103580.
- Vuko, E.; Rusak, G.; Dunkic, V.; Kremer, D.; Kosalec, I.; Rada, B.; Bezic, N. Inhibitionof satellite RNA associated cucumber mosaic virus infection by essential oil of *Micromeria croatica* (Pers.) Schott. *Molecules* 2019, 24, 1342
- Sumalan, R.M.; Alexa, E.; Popescu, I.; Negrea, M.; Radulov, I.; Obistiolu, D.; Cocan, I. Exploring ecological alternatives for crop protection using *Coriandrum sativum* essential oil. *Molecules* 2019, 24, 2040.
- Elshafie, Hs.; Grul'ova, D.; Baranova, B.; Caputo, L.; De Martino, L.; Sedlak, V.; Camele, I.; De Feo, V. Antimicrobial activity and chemical composition of essential extracted from *Solidago canadensis* L. *Molecules* 2019, 24, 1206.
- 21. Krzysko-Lupicka, T.; Walkowlak, W.; Bialon, M. Comparison of the fungistatic activity of selected essential oils relative to *Fusarium graminearum* isolates. *Molecules* **2019**, *24*, 311.
- El Gawad, A.A.; Elshamy A.; Elo Gendy A.E.N.; Gaara A.; Assaeed, A. Volatiles profiling, allelopathic activity, and antioxidant potentiality of *Xanthium strumarium* leaves essential oil from Egypt: Evidence from chemometrics analysis. *Molecules* 2019, 24, 584.
- Matouskova, M.; Jurova, J.; Grul'ova, D., Wajs-Bonikowska, A.; Renco, M.; Sedlak, V.; Poracova, J.; Gogal'ova, Z.; Kalemba, D. Phytotoxic effect of invasive *Heracleum mantegazzianum* essential oil on dicot and monocot species. *Molecules* 2019, 24, 425.
- Liu, K.; Deng, W.; Hu, W.; Cao, S.; Zhong, B.; Chun, J. Extraction of 'Gannanzao' orange peel essential oil by response surface methodology and its effect on cancer cell proliferation and migration. *Molecules* 2019, 24, 499.
- Spyridopoulou, K.; Fitsiou, E.; Bouloukosta, E.; Tiptiri-Kourpeti, A.; Vamvakias, M.; Oreopoulou A.; Papavassilopoulou E.; Pappa, A.; Chilchia, K. Extraction, chemical composition, and anticancer potential of *Origanum onites* L. essential oil. *Molecules* 2019, 24, 2612

- De Lima, E.J.S.P.; Alves R.G.; D' Elia, G.M.A.; da Annunciacao T.A.; Silva, V.R.; de S. Santos, L.; Soares M.B.P.; Cardozo N.M.D.; Costa, E.V.; da Silva F.M.A.; Koolen H.H.F.; Bezerra, D.P. Antitumour effect of the essential oil from the leaves of *Croton matourensis* Aubl. (Euphorbiaceae). *Molecules* 2019, 24, 2974.
- Tung, Y.T.; Hsu, Y.J.; Chien, Y.W.; Huang , C.C.; Huang, W.C.; Chiu, W.C. Tea seed oil prevents obesity, reduces physical fatigue, and improves exercise performance in high fat diet induced obese ovariectomized mice. *Molecules* 2019, 24, 980.
- Schepetkin, I.A.; Kirpotina, L.N.; Khlebnikov, A.I.; Balasubramanian, N.; Quinn, M.T. Neutrophil immunomodulatory activity of of natural organosulfur compounds. *Molecules* 2019, 24, 1809
- 29. Codruta Heghes, S.; Vostinaru, O.; Rus, L.M.; Mogosan, C.; Iuga, C.A.; Filip, L. Antispasmodic effect of essential oils and their constituents: A review. *Molecules* **2019**, *24*, 1675.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).