

# **The paleopathological evidence on the origins of human tuberculosis: a review**

Ileana Buzic<sup>1</sup>, Valentina Giuffra<sup>1</sup>

<sup>1</sup>Division of Paleopathology, Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Italy

**Keywords:** tuberculosis, paleopathology, history, Neolithic, and

**Running title:** On the origins of TB: a paleopathological review

**Correspondence:** Ileana Buzic, MA, Doctoral School of History - Specialisation Archaeology, University “1 Decembrie 1918” of Alba Iulia, Romania. Email: ileana.buzic@gmail.com

## **Abstract**

Tuberculosis (TB) has been one of the most important infectious diseases affecting mankind and still represents a plague on a global scale. In this narrative review the origins of tuberculosis are outlined, according to the evidence of paleopathology. In particular the first cases of human TB in ancient skeletal remains are presented, together with the most recent discoveries resulting from the paleomicrobiology of the tubercle bacillus, which provide innovative information on the history of TB. The paleopathological evidence of TB attests the presence of the disease starting from Neolithic times. Traditionally, it was thought that TB has a zoonotic origin, being acquired by humans from cattle during the Neolithic revolution. However, the biomolecular studies proposed a new evolutionary scenario demonstrating that human TB has a human origin. The researches show that the disease was present in the early human populations of Africa at least 70000 years ago and that it expanded following the migrations of *Homo sapiens* out of Africa, adapting to the different human groups. The demographic success of TB during the Neolithic period was due to the growth of density and size of the human host population, and not the zoonotic transfer from cattle, as previously hypothesized. These data demonstrate a long coevolution of the disease and its human host. Understanding the changes of TB through time thanks to the advances in the field of paleopathology can help to solve the present problems and understand the future evolution of TB.

## Introduction

Tuberculosis (TB) is an infectious disease caused by bacteria of the *Mycobacterium tuberculosis* complex (MTBC), which includes *M. tuberculosis*, *M. bovis*, *M. canettii*, *M. africanum*, *M. pinnipedii*, *M. microti*, *M. caprae*, *M. mungi* and *M. orygis*. The first two strains are most frequently responsible for TB in humans [1]. Despite the efforts to reduce the burden of TB worldwide and the progress in care and prevention, it is still a major global cause of disease and death [2].

Starting from the industrial era, TB became particularly diffuse, favoured by the increased population density and the poor living conditions. Thanks to the improvement of health status, alimentation and hygiene during the 20<sup>th</sup> century in developed countries, the disease began to decrease its incidence; the introduction of the BCG (Bacillus Calmette-Guérin) vaccine in 1921 and the use of chemotherapeutic drugs efficacious against the infection, such as streptomycin and isoniazid in 1943 and 1952 respectively, further decreased the incidence. However, despite being considered in the 80s of the 20<sup>th</sup> century as a conquered infection [3], TB still remains a plague on a global scale and is the top infectious killer globally by being present today in both developed and developing countries; it is estimated that a quarter of the world population (1.7 billion people) is infected with *M. tuberculosis*, 5–15% of which will develop the TB disease during their lifetime [4]. Statistics show that in 2016 TB was the underlying cause of 1.3 million deaths among HIV-negative people and furthermore was a contributing cause of 374,000 deaths among people with HIV [4]. Due to historical, social and geopolitical determinants, the regions with the highest rates of TB are Sub-Saharan Africa, South-East Asia and East Europe [5, 6]. The only available vaccine (BCG) is obtained from attenuated strains of *M. bovis*, but is not extensively effective; no other vaccines had been produced in the meantime [7]. Finally, the appearance of TB-strains that are resistant to antibiotics, in particular multi-drug and even totally drug-resistant strains, represents a real challenge in the struggle against the disease [8].

Paleopathology, the interdisciplinary field of research that focuses on the development of ancient diseases and their impact and distribution in past populations, together with ancient DNA analyses,

can provide important contributions to our knowledge of this major pathogen through the study of how TB evolved through time.

### **Early paleopathological and biomolecular evidences for TB in ancient skeletal remains**

The paleopathological evidence for TB is based on primary sources, such as the analysis of skeletal remains and mummified soft tissues. The diagnosis of TB can be considered reliable on the basis of specific skeletal features; these include characteristic changes to the spine, consisting in lytic lesions affecting the vertebral bodies with resulting ankylosis, body collapse and kyphosis (Pott's disease); extraspinal unifocal lytic lesions with absence of new bone formation; single joint ankylosis, specially localised in the hip, knee and wrist; and new bone formation on the internal surface of the ribs [9, 10]. In the last decades, the development of the new field of palaeomicrobiology permitted in several cases further corroboration of the diagnosis through the detection of TB ancient DNA (aDNA); the latest technologies of high-throughput sequencing and metagenomics allowed us to obtain a complete picture of the pathogen in ancient human remains [10].

However, modern clinical data demonstrated that only 1-5% of patients with pulmonary TB develops skeletal lesions [11]; therefore, it should be considered that the detection of TB in paleopathology is largely underestimated [12].

The earliest human skeletal remains with paleopathological changes of TB date back to 8000-10000 years ago, corresponding to the Neolithic revolution, and come from the Near East. Before this period, a single controversial diagnosis of Leptomeningitis tuberculosa, in relation to non specific endocranial changes of the frontal bone of a fossil, has been attributed to a Homo erectus from Turkey dating from the middle Pleistocene (490000–510000 years BP) [13]; this could represent the most ancient example of TB in a human fossil, but this interpretation was questioned by other authors [14]. The most ancient animal case, which was confirmed by morphological and biomolecular analyses, is represented by a ca. 17000-year-old late Pleistocene long-horned extinct bison (*Bison antiquus*) enclosed sediments from the Natural Trap Cave from Wyoming (United States). This fossil

documents that TB was present in North America at least 20000 years BP, long before the domestication [15].

The most notable of the first TB traces in human skeletal remains are concentrated in the Near East and Europe.

As for the Near East, five cases from two sites located in the Fertile Crescent belonging to the Pre-Pottery Neolithic B (PPNB) period provide evidence for the presence of human TB. Four individuals with lesions consistent with TB were brought to light from Dja'de el Mughara, situated on the Euphrates river in Northern Syria and dating to the pre-domestication phase (8800-8300 BC). An individual from Tell Aswad from Southern Syria, belonging to the early domestication phase (8200-7600 BC), displayed features typical of Hypertrophic Pulmonary Osteoarthropathy (HPOA), that is caused by chronic pulmonary diseases such as TB. In addition to the morphological features, multidisciplinary analyses such as morphological examination, MicroCT scan, lipid biomarkers and molecular analyses were also performed with positive diagnostic results on selected specimens from these archaeological sites, confirming the presence of human TB in the pre-domestication and early domestication phases of the Neolithic [16].

Another three cases evoking skeletal human TB were reported in the Neolithic village of Ain Ghazal in Jordan, dated to 7250 BC and also located in the cradle of agriculture; these cases, whose diagnosis was only morphological as no molecular analyses were performed, suggest the presence of human TB before and/or during the introduction of agriculture and animal domestication [17].

From the same geographical area, in the ancient Levantine village of Atlit-Yam now submerged (Israel), there is a report of two cases of TB on the basis of scarce paleopathological changes involving an adult individual and an adolescent, presumed to be mother and son, as they were buried together. The site is dated from 6200-5500 BC, corresponding to the last phase of the Pre-Pottery Neolithic period, when the transition to agriculture and animal domestication was fully accomplished. Lipid biomarkers and molecular analyses confirmed the presence of TB in the two individuals [18, 19].

Lipid biomarkers examination consists in the extraction, derivatisation and high-performance liquid chromatography (HPLC) analysis of mycobacterial cell wall mycolic acids [18-19].

As for Europe, among the earliest cases of TB in skeletal evidence there are cases of tuberculous spondylitis (Pott's disease) in two individuals from the Early Neolithic belonging to the Linear Pottery culture (5400–4800 BC) from three sites (Halberstadt, Derenburg and Karsdorf) from Saxony-Anhalt in central Germany. Molecular analyses detected the presence of pathogens belonging to the MTC in skeletal remains from all the three sites taken in examination [20].

Other cases with morphological evidences of TB were unearthed from several Neolithic sites dated back to 5000 BC, including Heidelberg, Germany [22], Złota, Poland [20], Hódmezővásárhely-Gorzsa, Hungary (confirmed by lipid biomarker analyses and ancient DNA analysis) [23], Alsónyék-Bátaszék, Hungary (confirmed by molecular analyses) [24, 25], attesting to the diffusion of the infection in different areas of Europe.

A particular concentration of TB was observed in the Finalese area in Liguria (Northwestern Italy) belonging to the Middle Neolithic period (4000-3500 BC), where three cases were discovered in three important caves that are in close proximity to each other: Arene Candide [26], Arma dell'Aquila [27], and Pollera [28] caves. The diagnosis was based on typical skeletal lesions; biomolecular analyses using polymerase chain reaction (PCR) were performed on the Pollera case, but did not confirm the presence of the MTBC ancient DNA.

As for Africa, 13 cases from the Upper Egyptian site of Nagada (4500-3000 BC) suggest that the earliest evidence of TB in Egypt could be dated back to 4500 BC [3]. The first Egyptian cases of TB confirmed by molecular analyses date back to the predynastic period (3500-2650 BC) and suggest that infection with *M. tuberculosis* was relatively recurrent in this period [29-33].

Outside the European landscape, a possible Neolithic case of TB was observed in an adult individual from Shanghai, China, associated with the Songze culture (3900-3200 BC), at the beginning of the wet rice agriculture [34].

From the late Neolithic onward the paleopathological evidence of TB becomes more frequent.

Finally, in the prehistoric New World, macroscopic and molecular analyses suggest that the earliest evidence of human TB in skeletal remains and mummified soft tissues is attested in prehistoric South America in Peru by 700 AD, while in North America it is attested by 900 AD in the Southwest, where large permanent agricultural settlements are concentrated [35, 36].

### **The biomolecular advancements on the origins of TB**

As highlighted in the previous section, the paleopathological evidence of human TB attests the presence of the disease starting from Neolithic times; no sure skeletal cases are dated before this period [37]. However, absence of evidence does not mean evidence of absence. Did TB affect humans in more ancient times? When did TB become a human pathogen? Did human TB originate from animal TB? The new next-generation sequencing technologies tried to answer these questions [38, 39].

According to the traditional theory, formulated before the advent of the biomolecular studies, humans acquired TB from cattle during the Neolithic revolution due to the zoonotic transfer from the newly domesticated animals [40-42]. As several infectious diseases have a zoonotic origin, it was thought that TB was also transferred to humans from animals; in particular, *M. bovis* would have infected humans, then adapting and evolving into *M. tuberculosis*. This theory was supported by the fact that the oldest human remains presenting paleopathological changes of TB were found in the Neolithic period, while the findings in animal remains have suggested a more ancestral existence of TB in cattle [15].

The evolutionary history of TB was dramatically changed by recent biomolecular studies that challenged the old theory, proposing an African origin for the MTBC long pre-dating the Neolithic [43-44].

Phylogenetic studies have clarified that within MTB seven lineages, associated with different areas of the world, can be identified [46]: the Lineage 1 (East Africa, Philippines), the modern Euroasian lineages 2 (which includes the Beijing family), 3, 4 that are in close phylogenetic relationship, African

lineages 5 and 6 with *M. africanum* phylogeographically distributed in West Africa, and finally, the newly described lineage 7 [47] collected from patients with TB from Ethiopia. Animal species, including *M. bovis*, *M. microti* and *M. pinnipedii*, represent monophyletic lineages, and *M. canettii* has an ancestral position in the MTBC. The chronological positions of the lineages are determined by the presence or absence of an MTB-specific deletion (TbD1); the presence makes them ancestral (TbD1<sup>+</sup>), the absence makes them modern (TbD1<sup>-</sup>) [48, 49]. On this basis, lineages 1, 5, and 6, the animal lineages and *M. canettii* represent the ancestral MTBC group, whereas lineages 2, 3 and 4 are defined as modern from an evolutionary point of view, because they are thought to have diffused more recently; lineage 7 resulted as intermediate between ancient and modern MTB lineages. These findings suggest that the animal lineages and *M. africanum* have diverged from the progenitor of the ancestral MTB lineages and that the tubercle bacillus was originally a human pathogen [49], confirming previous studies which suggested that animal adapted MTBC strains diverged from the major MTBC strains [48, 43-45]. Therefore, the new evolutionary scenario proposed that the human TB did not derive from *M. bovis*, but that, on the contrary, MTB has a human origin.

Furthermore, a recent work seems to confirm this theory with phylogenetic studies based on whole-genome data of MTCB. This study is based on the sequencing of a global collection of 259 MTBC modern clinical strains representative of the diversity of the global MTBC for reconstructing the phylogenetic relationship between them [50]. The overall results of three independent phylogenetic analyses for the determination of the most recent common ancestor of TB using a Bayesian approach suggested East and West Africa as the most probable geographic origin of MTBC. The results of this research would show the common origin in Africa of *Homo sapiens* and MTBC and that TB was present in the early human populations of Africa at least 70000 years ago, then infecting human for thousands of years [50].

The study of Comas and colleagues [50] also would demonstrate that MTBC evolved in parallel with its human host. The disease expanded in correspondence with the migrations of *Homo sapiens* out of Africa. It has been hypothesized that the two *M. africanum* lineages, *M. canettii* and lineage 7 might



have remained in Africa, whereas the others could have reached Europe and Asia, following the colonization of these areas, and finally becoming endemic worldwide. This work would suggest that the different lineages have adapted to different human groups. On the basis of this study a correlation between the divergence of the Lineage 1 of MTBC about 67000 years ago and the first human wave of migration out of Africa that was directed around the Indian Ocean was observed. Another split of MTBC that results occurring 46000 years ago corresponds instead to the second great migration of *Homo sapiens* from Africa, which was directed to Eurasia [50]. With the rise of Neolithic Demographic Transition 10000±2000 years ago, TB raised its rate of diffusion together with the increase in the human population density following the advance of agriculture and animal domestication. These results would suggest that the demographic success of TB during the Neolithic period was due to the growth of density and size of the human host population, and not to the zoonotic transfer from cattle, as previously hypothesized [50].

Also, of interest is the study of the dispersal of the modern Lineage 2, which includes the “Beijing” strains that have an increased virulence and transmission potential, a shorter latency phase than ancient lineages, furthermore being particularly successful in geographical diffusion, and are associated with antibiotic resistance [8]. The study has evidenced that the dating of Lineage 2 in Asia coincides with the arrival of the anatomically modern humans in East Asia about 42000-32000 years ago confirmed by the archaeological evidence, presents a first expansion about 11000-6000 years ago corresponding to the development of agriculture in China about 8000 years ago and, later, a further expansion about 5000-3000 years ago, when agriculture was introduced in the regions near China [50]. These results seem to show that tuberculosis had various moments of expansion other than the increase during the Neolithic in the Mediterranean region.

Furthermore, these data could explain why TB presents an adaptation to both low and high host population densities. In fact, the high level of asymptomatic latency that allows reactivation of TB and the slow progression to disease are an indication of adaptation to low host population density that is consistent with the ancestral emergence of MTBC in Africa before the Neolithic revolution. On the

other hand, the modality of aerosol transmission and the high virulence in order to guarantee the maximum level of transmission are typical of a high-density population environment. This double adaptation would have ensured to the bacterium a large diffusion in both high- and low-density areas, resulting in the fact that one-third of the human population is infected but remains asymptomatic, representing an extraordinary reservoir for the bacterium [8].

The study of the origins and patterns of change in the evolution of MTBC would demonstrate that migration and demography of the anatomically modern humans affected the evolution and spread of TB. The major recent molecular studies on the evaluation of its demography and timing of propagation seem to show a long coevolution of the disease and its human host, demonstrating that TB represents an exemplary model of adaptation to humans [8]. They also suggest that the palaeomicrobiology of TB can be fully understood through an integrated approach in correlation-association with the human host genome sequencing, considering the relation between the evolution of TB genomes and their human hosts.

## **Conclusions**

TB was traditionally thought to be a zoonosis, but the advancements of biomolecular analyses performed by various teams of researchers seem to suggest that *M. tuberculosis* did not evolve from *M. bovis*, but has a complex evolutionary history that reflects human evolution. In order to understand the evolution of the bacterium and the current epidemiology of TB it is necessary to investigate the history of the disease in parallel with that of humans. In fact, MTBC has evolved with humans during thousands of years, influencing reciprocally their evolution. Understanding the changes of TB through time thanks to the advent of the next-generation sequencing technology can aid the problems of the present and future evolution of TB in relationship with its human host.

## **Acknowledgments**

Funding sources: this research did not receive any specific grant from funding agencies in the public, commercial, or non-profit sectors.

## **Conflicts of interest statement**

None declared.

## **Authors' contributions**

IB performed the bibliographic research and wrote the text; VG conceived the design, partially wrote the text and revised the manuscript. All authors critically revised the manuscript. All authors have read and approved the latest version of the manuscript.

## **References**

- [1] Brites D, Gagneux S. Co-evolution of Mycobacterium tuberculosis and Homo sapiens. *Immunol Rev.* 2015; 264: 6–24. doi: 10.1111/imr.12264.
- [2] Donoghue ED. Palaeomicrobiology of Tuberculosis. In: Raoult D, Drancourt M, editors. *Paleomicrobiology – Past Human Infections*. Berlin/Heidelberg: Springer-Verlag; 2008. p. 75-97.
- [3] Roberts CA, Buikstra J. *The bioarchaeology of tuberculosis: a global view on a reemerging disease*. Gainesville, Florida: University Press of Florida; 2003.
- [4] Glaziou P, Floyd K, Raviglione MC. Global epidemiology of tuberculosis. *Semin Respir Crit Care Med.* 2018; 39: 271-285. doi: 10.1055/s-0038-1651492.
- [5] Riccardi N, Alagna R, Motta I, Ferrarese M, Castellotti P, Nicolini LA, Diaw MM, Ndiaye M, Cirillo D, Codecasa L, Besozzi G. Towards ending TB: civil community engagement in a rural area of Senegal: results, challenges and future proposal. *Infect Dis (Lond).* 2019; 51: 392-394. doi: 10.1080/23744235.2019.1572920.

- [6] Lönnroth K, Jaramillo E, Williams BG, Dye C, Raviglione M. Drivers of tuberculosis epidemics: the role of risk factors and social determinants. *Soc Sci Med.* 2009; 68: 2240-6. doi: [10.1016/j.socscimed.2009.03.041](https://doi.org/10.1016/j.socscimed.2009.03.041).
- [7] Kaufmann SH. Tuberculosis vaccines: time to think about the next generation. *Semin Immunol* 2013;25172–181. doi: [10.1016/j.smim.2013.04.006](https://doi.org/10.1016/j.smim.2013.04.006).
- [8] Bañuls A-L, Sanou A, Anh NT, Godreuil S. Mycobacterium tuberculosis: ecology and evolution of a human bacterium. *J Med Microbiol.* 2015; 64: 1261-1269. doi: [10.1099/jmm.0.000171](https://doi.org/10.1099/jmm.0.000171).
- [9] Waldron T. *Palaeopathology*. Cambridge: Cambridge University Press; 2009.
- [10] Donoghue ED. Paleomicrobiology of human tuberculosis. *Microbiol Spectr.* 2016; 4. doi: [10.1128/microbiolspec.PoH-0003-2014](https://doi.org/10.1128/microbiolspec.PoH-0003-2014).
- [11] Resnick D, Niwayama G. *Diagnosis of bone and joint disorders*. 4th ed. Philadelphia: Saunders; 2002.
- [12] Aufderheide A, Rodriguez-Martin C. *The Cambridge encyclopaedia of human palaeopathology*. Cambridge: Cambridge University Press; 1998.
- [13] Kappelman J, Alcicek MC, Kazanci N, Schultz M, Ozkul M, Sen S. First Homo erectus from Turkey and implications for migrations into temperate Eurasia. *Am J Phys Anthropol.* 2008; 135: 110-6. <https://doi.org/10.1002/ajpa.20739>.
- [14] Roberts CA, Pfister LA, Mays S. Letter to the editor: was tuberculosis present in Homo erectus in Turkey? *Am J Phys Anthropol.* 2009; 139: 442-4. <https://doi.org/10.1002/ajpa.21056>.
- [15] Rothschild BM, Martin LD, Lev G, Bercovier H, Bar-Gal GK, Greenblatt C, Donoghue H, Spigelman M, Brittain D. Mycobacterium tuberculosis complex DNA from an extinct bison dated 17,000 years before the present. *Clin Infect Dis.* 2001; 33: 305-11. <https://doi.org/10.1086/321886>.
- [16] Baker O, Lee OY, Wu HT, Besra GS, Minnikin DE, Llewellyn G, Williams CM, Maixner F, O’Sullivan N, Zink A, Chamel B, Khawam R, Coqueugniot E, Helmer D, Le Mort F, Perrin P, Gourichon L, Dutailly B, Palfi G, Coqueugniot H, Dutour O. Human tuberculosis predates

domestication in ancient Syria. *Tuberculosis (Edinb)* 2015; 95: S4–S12. doi: 10.1016/j.tube.2015.02.001.

[17] Al-Sarie I, Al-Shiyab A, El-Najjar M. Cases of tuberculosis at 'Ain Ghazal, Jordan. *Paléorient* 1996; 22: 123-128.

[18] Hershkovitz I, Donoghue HD, Minnikin DE, Besra GS, Lee OY-C, Gernaey AM, Galili E, Greenblatt CL, Lemma E, Bar-Gal GK, Spigelman M. Detection and molecular characterization of 9000-year-old *Mycobacterium tuberculosis* from a Neolithic settlement in the eastern Mediterranean. *PLoS One* 2008; 3 (10): e3426. doi: 10.1371/journal.pone.0003426.

[19] Hershkovitz I, Donoghue HD, Minnikin DE, May H, Lee OY, Feldman M, Galili E, Spigelman M, Rothschild BM, Bar-Gal GK. 2015. Tuberculosis origin: the Neolithic scenario. *Tuberculosis (Edinb)* 95 Suppl 1: S122-6. doi: 10.1016/j.tube.2015.02.021.

[20] Nicklisch N, Maixner F, Ganslmeier R, Friederich S, Dresely V, Meller H, Zink A, Alt KW. Rib lesions in skeletons from early neolithic sites in Central Germany: On the trail of tuberculosis at the onset of agriculture. *Am J Phys Anthropol.* 2012; 149: 391-404. doi: 10.1002/ajpa.22137.

[21] Bartels P. Tuberkulose in der jüngeren Steinzeit. *ArchAnthropol.* 1907; 6: 243-255.

[22] Gladykowska-Rzeczycka JJ. Tuberculosis in the past and present in Poland. In: Pálfi G, Dutour O, Deák J, Hutás I, editors. *Tuberculosis: past and present*. Budapest/Szeged: Golden Book Publishers and Tuberculosis Foundation; 1999. p. 561-73.

[23] Masson M, Molnár E, Donoghue HD, Minnikin DE, Lee OY, Besra GS, Bull ID, Pálfi G. 7000-year-old tuberculosis cases from Hungary: osteological and biomolecular evidence. *Tuberculosis (Edinb)* 2015;95 Suppl 1: S13-7. <https://doi.org/10.1016/j.tube.2015.02.007>.

[24] Köhler K, Pálfi G, Molnar E, Zalai-Gaál I, Oszás A, Bánffy E, Kirinó K, Kiss KK, Mende BG. A Late Neolithic Case of Pott's Disease from Hungary. *Int J Osteoarchaeol.* 2012;2 4: 697-703. doi:10.1002/oa.2254.

- [25] Pósa A, Maixner F, Mende BG, Köhler K, Osztás A, Sola C, Dutour O, Masson M, Molnár E, Pálfi G, Zink A. Tuberculosis in Late Neolithic-Early Copper Age human skeletal remains from Hungary. *Tuberculosis (Edinb)* 2015; 95 Suppl.1: S18-22. doi: 10.1016/j.tube.2015.02.011.
- [26] Canci A, Minozzi S, Borgognini Tarli S. New evidence of tuberculousspondylitis from Neolithic Liguria (Italy). *Int J Osteoarchaeol.* 1996; 6: 497–501.
- [27] Formicola V, Milanesi Q, Scarsini C. Evidence of spinal tuberculosis at the beginning of the fourth millennium BC from Arene Candide cave (Liguria, Italy). *Am J Phys Anthropol.* 1987; 72: 1–6. <https://doi.org/10.1002/ajpa.1330720102>.
- [28] Sparacello VS, Roberts CA, Kerudin A, Müller R. A 6500-year-old Middle Neolithic child from Pollera Cave (Liguria, Italy) with probable multifocal osteoarticular tuberculosis. *Int J Paleopath.* 2017; 17: 67-74. doi: 10.1016/j.ijpp.2017.01.004.
- [29] Crubézy E, Ludes B, Poveda JD, Clayton J, Crouau-Roy B, Montagnon D. Identification of Mycobacterium DNA in an Egyptian Pott's disease of 5,400 years old. *C R Acad Sci III.* 1998; 321: 941-51.
- [30] Zink A, Haas CJ, Reischl U, Szeimies U, Nerlich AG. Molecular analysis of skeletal tuberculosis in an ancient Egyptian population. *J Med Microbiol.* 2001; 50: 355–366. doi: 10.1099/0022-1317-50-4-355.
- [31] Nerlich AG, Rohrbach H, Zink A. Paleopathology of ancient Egyptian mummies and skeletons. Investigations on the occurrence and frequency of specific diseases during various time periods in the necropolis of Thebes-West. *Pathologe* 2002; 23: 379– 85.
- [32] Zink AR, Sola C, Reischl U, Grabner W, Rastogi N, Wolf H, Nerlich AG. Characterization of Mycobacterium tuberculosis complex DNAs from Egyptian mummies by spoligotyping. *J Clin Microbiol.* 2003; 41: 359– 67. doi: 10.1128/JCM.41.1.359-367.2003.
- [33] Lalremruata A, Ball M, Bianucci R, Welte B, Nerlich AG, Kun JF, Pusch CM. Molecular identification of falciparum malaria and human tuberculosis co-infections in mummies from the

- Fayum depression (Lower Egypt). *PLoS One* 2013; 8 (4): e60307. <https://doi.org/10.1371/journal.pone.0060307>.
- [34] Okazaki K, Takamuku H, Yonemoto S, Itahashi Y, Lakuhari T, Yoneda M, Chen J. A paleopathological approach to early human adaptation for wet-rice agriculture: the first case of Neolithic spinal tuberculosis at the Yangtze River delta of China. *Int J Paleopathol.* 2019; 24: 236-244. <https://doi.org/10.1016/j.ijpp.2019.01.002>.
- [35] Roberts CA, Buikstra JE. The history of tuberculosis from earliest times to the development of drugs. In: Davies PDO, Gordon SB, Davies G. *Clinical Tuberculosis*. London: CRC Press; 2014.
- [36] Bos KI, Harkins KM, Herbig A, Coscolla M, Weber N, Comas I, Forrest SA, Bryant JM, Harris SR, Schuenemann VJ, Campbell TJ, Majander K, Wilbur AK, Guichon RA, Wolfe Steadman DL, Cook DC, Niemann S, Behr MA, Zumarraga M, Bastida R, Huson D, Nieselt K, Young D, Parkhill J, Buikstra JE, Gagneux S, Stone AC, Krause J. Pre-Columbian mycobacterial genomes reveal seals as a source of New World human tuberculosis. *Nature.* 2014; 514 (7523): 494-7. doi: 10.1038/nature13591.
- [37] Holloway KL, Henneberg RJ, de Barros Lopez M, Henneberg. Evolution of human tuberculosis: a systematic review and meta-analysis of paleopathological evidence. *Homo* 2011; 2011: 402-458. doi: 10.1016/j.jchb.2011.10.001.
- [38] Rivera-Perez JI, Santiago-Rodriguez TM, Toranzos GA. Paleomicrobiology: a snapshot of ancient microbes and approaches to forensic microbiology. *Microbiol Spectr.* 2016; 4 (4). doi: 10.1128/microbiolspec.EMF-0006-2015.
- [39] Barbier M, Wirth T. The evolutionary history, demography, and spread of the *Mycobacterium tuberculosis* complex. 2016. *Microbiol Spectr* 4. doi: 10.1128/microbiolspec.TBTB2-0008-2016.
- [40] Steinbock RT. *Paleopathological Diagnosis and Interpretation: Bone Diseases in Ancient Human Populations*. Springfield, Illinois: Charles C. Thomas; 1976.
- [41] Manchester K. Tuberculosis and leprosy in antiquity: an interpretation. *Med Hist.* 1984; 28: 22-30.

- [42] Clark GA, Kelley MA, Granje JM, Hill CM. The evolution of mycobacterial disease in human populations. *Curr Anthropol.* 1987; 28: 45-62.
- [43] Gutierrez MC, Brisse S, Brosch R, Fabre M, Omaïs B, Marmiesse M, Supply P, Vincent V. Ancient origin and gene mosaicism of the progenitor of *Mycobacterium tuberculosis*. *PLoS Pathog.* 2005; 1: e5. <https://doi.org/10.1371/journal.ppat.0010005>.
- [44] Wirth T, Hildebrand F, Allix-Béguec C, Wölbeling F, Kubica T, Kremer K, van Soolingen D, Rüsche-Gerdes S, Locht C, Brisse S, Meyer A, Supply P, Niemann S. Origin, spread and demography of the *Mycobacterium tuberculosis* complex. *PLoS Pathog.* 2008; 4: e1000160. doi: 10.1371/journal.ppat.1000160.
- [45] Hershberg R, Lipatov M, Small PM, Sheffer H, Niemann S, Homolka S, Roach JC, Kremer K, Petrov DA, Feldman MW, Gagneux S. High functional diversity in *Mycobacterium tuberculosis* driven by genetic drift and human demography. *PloS Biol.* 2008; 6: e311. doi: 10.1371/journal.pbio.0060311.
- [46] Gagneux S, Small PM. Global phylogeography of *Mycobacterium tuberculosis* and implications for tuberculosis product development. *Lancet Infect Dis.* 2007; 7: 328–337. [https://doi.org/10.1016/S1473-3099\(07\)70108-1](https://doi.org/10.1016/S1473-3099(07)70108-1).
- [47] Firdessa R, Berg S, Hailu E, Schelling E, Gumi B, Erenso G, Gadisa E, Kiros T, Habtamu M, Hussein J, Zinsstag J, Robertson BD, Ameni G, Lohan AJ, Loftus B, Comas I, Gagneux S, Tschopp R, Yamuah L, Hewinson G, Gordon SV, Young DB, Aseffa A. Mycobacterial lineages causing pulmonary and extrapulmonary tuberculosis, Ethiopia. *Emerg Infect Dis.* 2013; 19: 460–463. doi: 10.3201/eid1903.120256.
- [48] Brosch R, Gordon SV, Marmiesse M, Brodin P, Buchrieser C, Eiglmeier K, Garnier T, Gutierrez C, Hewinson G, Kremer K, Parsons LM, Pym AS, Samper S, van Soolingen D, Cole ST. A new evolutionary scenario for the *Mycobacterium tuberculosis* complex. *Proc Natl Acad Sci U.S.A.* 2002; 99: 3684–3689. DOI: 10.1073/pnas.052548299.



[49] Gagneux S. Host-pathogen coevolution in human tuberculosis. *Philos Trans R Soc Lond B Biol Sci.* 2012; 367: 850–859. doi: 10.1098/rstb.2011.0316.

[50] Comas I, Coscolla M, Luo T, Borrell S, Holt KE, Kato- Maeda M, Parkhill J, Malla B, Berg S, Thwaites G, Yeboah-Manu D, Bothamley G, Mei J, Wei L, Bentley S, Harris SR, Niemann S, Diel R, Aseffa A, Gao Q, Young D, Gagneux S. Out-of-Africa migration and Neolithic coexpansion of *Mycobacterium tuberculosis* with modern humans. *Nat Genet.* 2013; 45: 1176–1182. doi: 10.1038/ng.2744.