

Washington University School of Medicine

Digital Commons@Becker

2020-Current year OA Pubs

Open Access Publications

7-19-2022

Effect of race on prediction of brain amyloidosis by plasma A β 42/ A β 40, phosphorylated tau, and neurofilament light

Suzanne E Schindler

Rachel L Henson

Yan Li

Benjamin Saef

Krista L Moulder

See next page for additional authors

Follow this and additional works at: https://digitalcommons.wustl.edu/oa_4

Authors

Suzanne E Schindler, Rachel L Henson, Yan Li, Benjamin Saef, Krista L Moulder, David Bradford, Anne M Fagan, Brian A Gordon, Tammie L S Benzinger, Joyce Balls-Berry, Randall J Bateman, Chengjie Xiong, John C Morris, and et al

Effect of Race on Prediction of Brain Amyloidosis by Plasma A β 42/A β 40, Phosphorylated Tau, and Neurofilament Light

Suzanne E. Schindler, MD, PhD, Thomas K. Karikari, PhD, Nicholas J. Ashton, PhD, Rachel L. Henson, MS, Kevin E. Yarasheski, PhD, Tim West, PhD, Mathew R. Meyer, PhD, Kristopher M. Kirmess, PhD, Yan Li, PhD, Benjamin Saef, MS, Krista L. Moulder, PhD, David Bradford, Anne M. Fagan, PhD, Brian A. Gordon, PhD, Tammie L.S. Benzinger, MD, PhD, Joyce Balls-Berry, PhD, Randall J. Bateman, MD, Chengjie Xiong, PhD, Henrik Zetterberg, MD, PhD, Kaj Blennow, MD, PhD, and John C. Morris, MD

Correspondence

Dr. Schindler
schindler.s.e@wustl.edu

Neurology® 2022;99:e245–e257. doi:10.1212/WNL.0000000000200358

Abstract

Background and Objectives

To evaluate whether plasma biomarkers of amyloid (A β 42/A β 40), tau (p-tau181 and p-tau231), and neuroaxonal injury (neurofilament light chain [NfL]) detect brain amyloidosis consistently across racial groups.

Methods

Individuals enrolled in studies of memory and aging who self-identified as African American (AA) were matched 1:1 to self-identified non-Hispanic White (NHW) individuals by age, *APOE* ϵ 4 carrier status, and cognitive status. Each participant underwent blood and CSF collection, and amyloid PET was performed in 103 participants (68%). Plasma A β 42/A β 40 was measured by a high-performance immunoprecipitation–mass spectrometry assay. Plasma p-tau181, p-tau231, and NfL were measured by Simoa immunoassays. CSF A β 42/A β 40 and amyloid PET status were used as primary and secondary reference standards of brain amyloidosis, respectively.

Results

There were 76 matched pairs of AA and NHW participants ($n = 152$ total). For both AA and NHW groups, the median age was 68.4 years, 42% were *APOE* ϵ 4 carriers, and 91% were cognitively normal. AA were less likely than NHW participants to have brain amyloidosis by CSF A β 42/A β 40 (22% vs 43% positive; $p = 0.003$). The receiver operating characteristic area under the curve of CSF A β 42/A β 40 status with the plasma biomarkers was as follows: A β 42/A β 40, 0.86 (95% CI 0.79–0.92); p-tau181, 0.76 (0.68–0.84); p-tau231, 0.69 (0.60–0.78); and NfL, 0.64 (0.55–0.73). In models predicting CSF A β 42/A β 40 status with plasma A β 42/A β 40 that included covariates (age, sex, *APOE* ϵ 4 carrier status, race, and cognitive status), race did not affect the probability of CSF A β 42/A β 40 positivity. In similar models based on plasma p-tau181, p-tau231, or NfL, AA participants had a lower probability of CSF A β 42/A β 40 positivity (odds ratio 0.31 [95% CI 0.13–0.73], 0.30 [0.13–0.71], and 0.27 [0.12–0.64], respectively). Models of amyloid PET status yielded similar findings.

From the Department of Neurology (S.E.S., R.L.H., Y.L., B.S., K.L.M., D.B., A.M.F., J.B.-B., R.J.B., J.C.M.), Knight Alzheimer Disease Research Center (S.E.S., R.L.H., Y.L., B.S., K.L.M., D.B., A.M.F., B.A.G., T.L.S.B., J.B.-B., R.J.B., C.X., J.C.M.), Hope Center for Neurological Disorders (A.M.F.), Mallinckrodt Institute of Radiology (B.A.G., T.L.S.B.), and Division of Biostatistics (C.X.), Washington University School of Medicine, St. Louis, MO; Clinical Neurochemistry Laboratory, Department of Psychiatry and Neurochemistry (T.K.K., N.J.A., H.Z., K.B.), Institute of Neuroscience and Physiology, University of Gothenburg, Mölndal, Sweden; Department of Psychiatry (T.K.K.), University of Pittsburgh, PA; Wallenberg Centre for Molecular and Translational Medicine (N.J.A.), University of Gothenburg, Sweden; Institute of Psychiatry, Psychology and Neuroscience (N.J.A.), Maurice Wohl Institute Clinical Neuroscience Institute, King's College London; NIH Biomedical Research Centre for Mental Health and Biomedical Research Unit for Dementia at South London and Maudsley NHS Foundation (N.J.A.), London, UK; C2N Diagnostics (K.E.Y., T.W., M.R.M., K.M.K.), St. Louis, MO; Department of Neurodegenerative Disease (H.Z.), UCL Institute of Neurology, Queen Square, London; UK Dementia Research Institute at UCL (H.Z.), London, UK; and Hong Kong Center for Neurodegenerative Diseases (H.Z.), China.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by NIH.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

AA = African American; A β = β -amyloid; AD = Alzheimer disease; ADRC = Alzheimer Disease Research Center; AUC = area under the curve; CDR = Clinical Dementia Rating; IQR = interquartile range; LP = lumbar puncture; NfL = neurofilament light chain; NHW = non-Hispanic White; OR = odds ratio; p-tau181 = tau phosphorylated at position 181; PiB = Pittsburgh compound B; ROC = receiver operating characteristic; SUVR = standardized uptake value ratio; t-tau = total tau; UDS = Uniform Data Set.

Discussion

Models predicting brain amyloidosis using a high-performance plasma A β 42/A β 40 assay may provide an accurate and consistent measure of brain amyloidosis across AA and NHW groups, but models based on plasma p-tau181, p-tau231, and NfL may perform inconsistently and could result in disproportionate misdiagnosis of AA individuals.

Biomarkers of Alzheimer disease (AD) brain pathology are used by research studies, clinical trials, and memory clinics for a variety of indications, including to determine whether the etiology of cognitive impairment is likely to be related to AD or another cause. Amyloid PET is a well-established technique to determine whether an individual has significant brain amyloidosis that could be causing or contributing to cognitive impairment; however, amyloid PET is expensive and has limited availability.¹ CSF biomarkers are also highly accurate predictors of brain amyloidosis and are less expensive, but skilled clinicians are required to perform lumbar puncture (LP) procedures, and some individuals perceive LPs as invasive.² Several commercial assays can be used to measure concentrations of CSF β -amyloid (A β) peptide 42 (A β 42), A β 40, total tau (t-tau), and tau phosphorylated at position 181 (p-tau181), and cutoffs consistent with brain amyloidosis have been established.³⁻⁵

Biomarker cutoffs for brain amyloidosis have been defined in cohorts largely comprised of non-Hispanic White (NHW) individuals and then applied to all individuals. However, several studies have found lower levels of CSF t-tau and p-tau181 in African American (AA) individuals as compared with NHW individuals, even after adjusting for factors such as age, sex, APOE ϵ 4 carrier status, and cognitive impairment.⁶⁻⁹ Why AA individuals have lower levels of CSF t-tau and p-tau181 is unknown and could be due to differences in medical comorbidities, biological factors, or social determinants of health.^{8,10,11} Regardless of the underlying reasons, these differences have important implications for the utility of CSF biomarkers. Applying biomarker cutoffs defined in NHW to groups in which the biomarker has not been studied could potentially subject the other groups to additional testing, incorrect medical management, missed opportunities for treatment with AD-specific therapies, and lower enrollment in AD clinical trials.^{9,12} However, it is also highly problematic to adjust the interpretation of medical tests based on race, especially given the heterogeneity represented within racial groups and the dynamic nature of race because it is a social rather than a biological construct.^{9,13,14} It would be preferable to use AD biomarkers that perform accurately and

consistently across racial and ethnic groups. Alternatively, adjusting for the factors that underlie racial differences in AD biomarkers (e.g., medical comorbidities) may be more valid and generalizable across groups.

Over the past 3 years, there has been rapid development of blood-based biomarkers for AD.¹⁵ The PrecivityAD test offered by C2N Diagnostics, which includes highly precise measurement of plasma A β 42/A β 40 and APOE proteotype by mass spectrometry, is now available for clinical use.^{16,17} Multiple plasma p-tau isoforms can also be used as biomarkers of brain amyloidosis, including p-tau181,^{18,19} p-tau217,²⁰⁻²² and p-tau231.²³ Plasma neurofilament light chain (NfL) may also be useful as a nonspecific marker of neuroaxonal injury.²⁴ It is critical to evaluate whether these assays accurately and consistently predict brain amyloidosis across various racial and ethnic groups. In this study, one of the largest cohorts of AA individuals with CSF biomarker and amyloid PET information was used to examine the relationship of these reference measures of brain amyloidosis with the C2N Diagnostics PrecivityAD assay for plasma A β 42/A β 40 as well as Simoa immunoassays for p-tau181, p-tau231, and NfL.

Methods

Participants

This study analyzed samples and data from the Charles F. and Joanne Knight Alzheimer Disease Research Center (ADRC), which includes one of the largest groups of AA individuals in AD research who have undergone CSF collection or amyloid PET. The cohort consists of community-dwelling older adults recruited from the St. Louis area, including participants with and without cognitive impairment, who enrolled in research studies of memory and aging at Washington University in St. Louis. Participants underwent clinical and cognitive assessments using the Uniform Data Set (UDS)²⁵ that includes the Clinical Dementia Rating (CDR)²⁶ and Mini-Mental State Examination.²⁷ The UDS includes the Hollingshead 2 factor index of social position,²⁸ which assigns a social class based on the participant's educational level and the occupation of the head of the participant's household. Presence or absence of

Table 1 Characteristics of the Knight Alzheimer Disease Research Center Matched Cohort

Characteristics	African American participants (n = 76)	Non-Hispanic White participants (n = 76)	p	Adjusted p
Demographics				
Age at CSF collection, y	68.4 (64.9–73.2)	68.4 (64.1–73.1)	NS	
Sex, n (% Female)	44 (58)	39 (51)	NS	
APOE ε4 status, n (% carrier)	32 (42)	32 (42)	NS	
CDR 0/0.5/1 (% >0)	69/4/3 (9)	69/5/2 (9)	NS	
Years of education	16 (12–18)	16 (14–18)	NS	
Hollingshead index	2.0 (2.0–3.5)	2.0 (1.0–3.0)	0.002	
Hypertension, yes/no/not reported (% yes of reported)	51/25/0 (67)	33/40/3 (45)	0.006	
Diabetes, yes/no/not reported (% yes of reported)	21/55/0 (28)	4/69/3 (5)	0.0003	
CSF/plasma to LP interval, y	0.11 (0.05–0.21)	0.08 (0.04–0.23)	NS	
CSF biomarker concentrations				
CSF Aβ42, pg/mL	735 (544–971)	682 (516–883)	NS	NS
CSF Aβ40, pg/mL	9,490 (7,150–11,600)	10,100 (8,880–12,300)	0.07	NS
CSF Aβ42/Aβ40	0.0874 (0.0681–0.0935)	0.0719 (0.0477–0.0870)	0.0003	0.0001
CSF Aβ42/Aβ40 < 0.0673, n (%)	17 (22)	33 (43)	0.006	0.003
CSF total tau, pg/mL	212 (165–287)	290 (217–482)	0.0002	0.002
CSF p-tau181, pg/mL	31 (24.6–41.1)	38.0 (30.4–55.7)	0.002	0.0008
CSF NfL, pg/mL	644 (493–868) ^a	736 (542–973)	0.09	0.08
Plasma biomarker concentrations				
Plasma Aβ42, pg/mL	41.9 (39.3–49.6)	40.9 (37.8–46.3)	0.06	0.03
Plasma Aβ40, pg/mL	409 (380–470)	425 (390–482)	NS	NS
Plasma Aβ42/Aβ40	0.1047 (0.0990–0.1101)	0.0963 (0.0904–0.1028)	<0.0001	<0.0001
Plasma p-tau181, pg/mL	12.3 (10.2–16.2)	14.2 (10.6–19.3)	NS	NS
Plasma p-tau231, pg/mL	8.2 (4.4–11.3)	9.1 (6.6–13.1)	0.09	NS
Plasma NfL, pg/mL	11.1 (7.6–15.5)	11.8 (8.9–16.7)	NS	NS
Amyloid PET				
Amyloid PET Centiloid	2.3 (–1.0–10.1) ^b	10.1 (0.0–33.0) ^c	0.01	0.02
Amyloid PET positive, n (%)	5 (10) ^b	21 (39) ^c	0.0008	0.003

Abbreviations: Aβ = β-amyloid; CDR = Clinical Dementia Rating; LP = lumbar puncture; NfL = neurofilament light; p-tau181 = tau phosphorylated at position 181; p-tau231 = tau phosphorylated at position 231.

Continuous values are presented as median (interquartile range). The significance of differences by self-identified race was evaluated with Wilcoxon rank sum tests for continuous variables and χ^2 or Fisher exact tests for categorical variables. The covariate-adjusted significance of racial differences was evaluated using analysis of covariance models with biomarker concentrations as the outcome measure, race as the predictor variable, and the covariates of age, sex, APOE ε4 carrier status, and cognitive status. Plasma p-tau181 and NfL were transformed with the natural logarithm in covariate-adjusted models.

^a n = 72.

^b n = 49.


^c n = 54.

hypertension or diabetes was noted by the clinician. Race and sex were self-identified.

Participants with CSF biomarker information and adequate aliquots of plasma available for analysis were considered for

inclusion. Each self-identified AA participant was matched 1:1 to a self-identified NHW participant by a computer algorithm. Participants were matched by age at the time of plasma collection (within 2 years), APOE ε4 status (carrier or non-carrier), and cognitive status at the time of plasma collection

Figure 1 Biomarkers by Race



Biomarkers of amyloid (A), tau (B), and neuroaxonal injury (C) are shown by self-identified race. The covariate-adjusted significance of racial differences was evaluated using analysis of covariance models with biomarker concentrations as the outcome measure, race as the predictor variable, and the covariates of sex, age, *APOE* ϵ 4 carrier status, and cognitive status. Plasma tau phosphorylated at position 181 (p-tau181) and neurofilament light chain (NfL) were transformed with the natural logarithm for analysis. Point types denote the following: race: red, African American (AA); black, non-Hispanic White (NHW); cognitive status: open circle, Clinical Dementia Rating (CDR) 0; closed square, CDR >0. A β = β -amyloid.

(cognitively normal [CDR 0] or cognitively impaired [CDR >0]). If >1 NHW participant matched an AA participant, the participant with the closest age was selected.

Standard Protocol Approvals, Registrations, and Patient Consents

Written informed consent was obtained from all participants and their study partners. All procedures were approved by Washington University's Human Research Protection Office.

Genotyping

The *APOE* genotype was determined by genotyping rs7412 and rs429358 with TaqMan genotyping technology.²⁹ Genetic sex determined by sex chromosome-specific analysis was concordant with gender in all individuals in this cohort.

CSF and Plasma Collection and Analysis

CSF and blood samples from each participant were collected at a single session at approximately 8 AM following overnight fasting as previously described.^{5,30} Concentrations of CSF A β 40, A β 42, t-tau, and p-tau181 were measured by chemiluminescent enzyme immunoassay using a fully automated platform (LUMIPULSE G1200; Fujirebio). CSF NfL was measured via commercial ELISA kit (UMAN Diagnostics). Plasma A β 42 and A β 40 were measured in the C2N Diagnostics commercial laboratory with the PrecivityAD immunoprecipitation-mass spectrometry assay.¹⁶ Plasma p-tau181 and p-tau231 were

measured in the Clinical Neurochemistry Laboratory, University of Gothenburg, using in-house Single molecule array (Simoa) assays on an HD-X analyzer (Quanterix), as previously described.^{19,23} Plasma NfL was measured with Quanterix Nf-Light assay kits at Washington University on a HD-X analyzer. All assays were performed by personnel who were blind to participant information.


Amyloid PET

Participants underwent a dynamic scan with either florbetapir (n = 48) or Pittsburgh compound B (PiB; n = 55) in coordination with a structural MRI scan. Regional data from the 30–60 minutes postinjection window for PiB and the 50–70 minutes window for florbetapir were converted to standardized uptake value ratios (SUVRs) using cerebellar gray as a reference and partial volume corrected using a geometric transfer matrix approach based upon the Freesurfer parcellation.³¹ Values from regions where amyloid deposition occurs early in AD were averaged together to represent mean cortical SUVR, which was converted to Centiloid using previously published equations.^{32,33}

Statistical Analysis

The significance of differences by self-identified race were evaluated with Wilcoxon rank sum tests for continuous variables and χ^2 or Fisher exact tests for categorical variables. The covariate-adjusted significance of racial differences were

Figure 2 Relationship of Plasma A β 42/A β 40 With CSF A β 42/A β 40 and Amyloid PET



evaluated using analysis of covariance models with biomarker concentrations as the outcome measure, self-identified race as the predictor variable, and including the covariates of age, sex, *APOE* ϵ 4 carrier status, and cognitive status (cognitively normal [CDR = 0] or cognitively impaired [CDR > 0]). Models used natural logarithm transformed values for CSF and plasma p-tau181 and NfL, which were positively skewed. Models including the interaction between race and *APOE* ϵ 4 carrier status were also evaluated.

CSF A β 42/A β 40 status was chosen as the primary reference standard for brain amyloidosis because all individuals in the study had both CSF and blood collected at the same session, whereas only a subcohort had an amyloid PET scan performed within 2 years of CSF/blood collection. Positive CSF A β 42/A β 40 was defined by a CSF A β 42/A β 40 <0.0673, a cutoff that maximally distinguished amyloid PET status in an overlapping cohort with a receiver operating characteristic area under the curve (ROC AUC) of 0.97.³⁴ Amyloid PET positivity was previously defined as a mean cortical SUVR >1.42 for PiB and >1.19 for florbetapir.^{32,35} Logistic regression models were implemented with CSF A β 42/A β 40 or amyloid PET status as the outcome measure and each plasma biomarker as the predictor variable. Covariate adjusted models included self-identified race, sex, age, *APOE* ϵ 4 carrier status, and cognitive status. Models that in addition included either the interaction between race and *APOE* ϵ 4 carrier status or race and plasma biomarker levels were evaluated. Differences between ROC AUCs were evaluated using the DeLong test.³⁶

Statistical analyses were implemented using SAS 9.4 (SAS Institute Inc.). Plots were created with GraphPad Prism version 9.2.0 (GraphPad Software). All *p* values were from 2-sided tests, and results were deemed statistically significant at *p* < 0.05.

Data Availability

Data are available to qualified investigators upon request to the Knight ADRC (knightadrc.wustl.edu/Research/ResourceRequest.htm).


Results

Participant Characteristics

Based on the inclusion criteria of CSF biomarker information and adequate aliquots of plasma available for analysis, 79 AA and 775 NHW participants were potentially eligible for the study. Each AA participant was matched 1:1 to a NHW participant by age, *APOE* ϵ 4 carrier status and cognitive status. Three AA participants who could not be matched to a NHW participant were not included in the study. The final study cohort included a total of 152 participants (76 AA and 76 matching NHW) who contributed samples that underwent measurement of plasma biomarkers (see Table 1 for cohort characteristics). An amyloid PET scan was performed within 2 years of plasma collection in 49 AA (64%) and 54 NHW (71%) participants (eTable 1, links.lww.com/WNL/B978). All AA participants identified their ethnicity as non-Hispanic.

For both the AA and NHW groups, the median age was 68.4 years, 42% carried at least 1 *APOE* ϵ 4 allele (8% were ϵ 4

Figure 3 Relationship of Plasma p-tau181 With CSF A β 42/A β 40 and Amyloid PET



homozygotes), and 9% were cognitively impaired as defined by a CDR >0. There was no difference in dementia severity by race as measured by the CDR. Both the AA and NHW groups were well-educated (median of 16 years of education), but the AA group had a slightly lower social position than the NHW group as measured by the Hollingshead 2-factor index of social position (median 2.0 [interquartile range (IQR) 2.0–3.5] vs 2.0 [IQR 1.0–3.0], respectively; $p < 0.002$). Because the AA and NHW participants had no significant differences in years of education, this suggests that the median occupational level of the head of household in the AA group was lower (e.g., fewer of the AA participants lived in households headed by executives/major professionals). Compared with NHW individuals, AA individuals were more likely to have hypertension (67% vs 45%; $p = 0.006$) or diabetes (28% vs 5%; $p = 0.0003$).

CSF and Plasma Biomarkers by Race

CSF A β 42 and A β 40 concentrations were not significantly different between the AA and NHW groups (Table 1). However, AA individuals had higher CSF A β 42/A β 40 (median 0.0874 [IQR 0.0681 to 0.0935] vs 0.0719 [0.0477 to 0.0870]; $p < 0.0001$) and lower amyloid PET Centiloid (median 2.3 [IQR -1.0 to 10.1] vs 10.1 [0.0–33.0]; $p = 0.02$), consistent with the AA group having lower average levels of brain amyloidosis compared with the NHW group (Figure 1). In the overall cohort, 22% of the AA and 43% of the NHW groups had brain amyloidosis by CSF A β 42/A β 40 status ($p = 0.003$); in the subcohort with amyloid PET, 10% of AA and 39% of the NHW groups had brain amyloidosis by amyloid PET status ($p = 0.003$). Plasma A β 42 was only slightly higher in the AA group ($p = 0.03$) and plasma A β 40 did not vary by

racial group, but plasma A β 42/A β 40 was markedly higher in the AA group (median 0.1047 [IQR 0.0990–0.1101] vs 0.0963 [0.0904–0.1028]; $p < 0.0001$), again consistent with the AA group having lower average levels of brain amyloidosis compared with the NHW group. CSF total tau and p-tau181 were lower in the AA group than the NHW group ($p = 0.002$ and $p = 0.0008$, respectively), but there were no statistically significant differences in plasma p-tau181 and p-tau231 between racial groups. There was a trend towards lower CSF NfL in AA compared with NHW individuals ($p = 0.08$), but there was no difference in plasma NfL by racial group.

Plasma Biomarkers, CSF A β 42/A β 40 or Amyloid PET Centiloid, and Race

Nonlinear associations between plasma biomarkers and CSF A β 42/A β 40 or amyloid PET Centiloid were examined by Spearman correlations, as depicted in Figures 2 and 3 and eFigures 1–2, links.lww.com/WNL/B978 and summarized in eTable 2. Of the plasma biomarkers, A β 42/A β 40 had the strongest correlations with CSF A β 42/A β 40 ($\rho = 0.52$ [0.39 to 0.63]) and amyloid PET Centiloid (-0.30 [-0.10 to -0.47]) after adjustment for covariates. To examine the relationships between the plasma biomarkers, brain amyloid, and race, biomarker concentrations were modeled as a function of CSF A β 42/A β 40 status and included race, age, sex, APOE ϵ 4 carrier status, and cognitive status as covariates (Table 2). More abnormal (lower) plasma A β 42/A β 40 was associated with NHW race ($p < 0.0001$), male sex ($p < 0.0001$), and positive CSF A β 42/A β 40 status ($p < 0.0001$). In contrast, more abnormal (higher) plasma p-tau181 levels were associated with older age ($p < 0.0001$), positive CSF A β 42/A β 40 status ($p = 0.003$), male

Table 2 Relationship Between Plasma Biomarkers, CSF Aβ42/Aβ40 Status, and Covariates

Parameter	Estimate	SE	p
Plasma Aβ42/Aβ40			
Intercept	0.1052	0.0051	<0.0001
CSF Aβ42/Aβ40 status (positive)	-0.008	0.0013	<0.0001
Race (African American)	0.0060	0.0011	<0.0001
Sex (female)	0.0044	0.0011	<0.0001
Age (y)	-0.00010	0.00007	NS
APOE ε4 status (carrier)	-0.0009	0.0011	NS
Cognitive status (CDR >0)	-0.0010	0.0019	NS
Ln (plasma p-tau181)			
Intercept	1.267	0.311	<0.0001
CSF Aβ42/Aβ40 status (positive)	0.239	0.079	0.003
Race (African American)	-0.044	0.066	NS
Sex (female)	-0.164	0.065	0.01
Age (y)	0.020	0.004	<0.0001
APOE ε4 status (carrier)	0.017	0.068	NS
Cognitive status (CDR >0)	0.278	0.115	0.02
Plasma p-tau231			
Intercept	-1.655	4.474	NS
CSF Aβ42/Aβ40 status (positive)	2.525	1.140	0.03
Race (African American)	-0.970	0.946	NS
Sex (female)	-0.985	0.940	NS
Age (y)	0.160	0.063	0.01
APOE ε4 status (carrier)	-0.190	0.985	NS
Cognitive status (CDR >0)	5.585	1.651	0.0009
Ln (plasma NfL)			
Intercept	-0.710	0.357	0.05
CSF Aβ42/Aβ40 status (positive)	-0.015	0.091	NS
Race (African American)	-0.091	0.075	NS
Sex (female)	-0.052	0.075	NS
Age (y)	0.046	0.005	<0.0001
APOE ε4 status (carrier)	0.092	0.079	NS
Cognitive status (CDR >0)	0.297	0.132	0.03

Abbreviations: Aβ = β-amyloid; CDR = Clinical Dementia Rating; NfL = neurofilament light; p-tau181 = tau phosphorylated at position 181; p-tau231 = tau phosphorylated at position 231.

Analysis of covariate models evaluated the effects of CSF Aβ42/Aβ40 status (positive <0.0673), self-identified race, sex, age, APOE ε4 carrier status, and cognitive status on levels of each plasma biomarker. Plasma p-tau181 and NfL were transformed with the natural logarithm for analysis.

sex ($p = 0.01$), and impaired cognitive status ($p = 0.02$). More abnormal (higher) p-tau231 levels were associated with impaired cognitive status ($p = 0.0009$), older age ($p = 0.01$), and positive CSF Aβ42/Aβ40 status ($p = 0.03$). More abnormal (higher) plasma NfL levels were associated with older age ($p < 0.0001$) and impaired cognitive status ($p = 0.03$). Similar models of plasma biomarker levels including amyloid PET status rather than CSF Aβ42/Aβ40 status yielded similar results except that cognitive status was not a significant predictor in any model (eTables 3–6); few participants with cognitive impairment had amyloid PET data (4 of 103), limiting power to detect differences by cognitive status in these models. Models that included the interaction between race and APOE ε4 carrier status were evaluated, but the interaction was not significant for any model and therefore it was not included in the final analyses.

Correspondence of Plasma Biomarkers With CSF Aβ42/Aβ40 and Amyloid PET Status

Prediction of CSF Aβ42/Aβ40 or amyloid PET status by plasma biomarkers was evaluated by logistic regression analyses, as depicted in Figures 2 and 3 and eFigures 1 and 2, links.lww.com/WNL/B978, shown in eTables 7–11, and summarized in Tables 3 and 4. Models predicting CSF Aβ42/Aβ40 status based on plasma biomarker levels had ROC AUCs as follows: Aβ42/Aβ40, 0.86 (95% CI 0.79–0.92); p-tau181, 0.76 (0.68–0.84); p-tau231, 0.69 (0.60–0.78); and NfL, 0.64 (0.55–0.73). The amyloid probability score, a proprietary modeled value provided by C2N Diagnostics that is based on plasma Aβ42/Aβ40, APOE proteotype and age,¹⁷ had an ROC AUC of 0.89 (0.84–0.95) with CSF Aβ42/Aβ40 status. Comparisons of ROC AUCs showed that plasma Aβ42/Aβ40 had significantly better prediction of CSF Aβ42/Aβ40 status compared with p-tau181, p-tau231, and NfL ($p < 0.05$, 0.004, and <0.0001, respectively; Table 3).

Covariate adjusted models of CSF Aβ42/Aβ40 status incorporating each plasma biomarker and covariates (age, sex, APOE ε4 carrier status, race, and cognitive status) are summarized in Table 4. The model based on plasma Aβ42/Aβ40 had an ROC AUC of 0.90 (0.85–0.96) (eTable 7, links.lww.com/WNL/B978), which was superior to a model of covariates alone (0.82 [0.74–0.89] (eTable 12); $p = 0.006$ for difference in ROC AUCs). In the model of CSF Aβ42/Aβ40 status incorporating plasma Aβ42/Aβ40 and covariates, a higher probability of CSF Aβ42/Aβ40 positivity was associated with APOE ε4 carriers (odds ratio [OR] 5.6 [95% CI 2.0–16]; $p = 0.001$), older age in years (OR 1.12 [1.03–1.21]; $p = 0.007$), and cognitive impairment (OR 9.2 [1.9–46]; $p = 0.007$). Notably, in models incorporating plasma Aβ42/Aβ40 and covariates, race did not significantly affect correspondence with CSF Aβ42/Aβ40 or amyloid PET status.

The covariate-adjusted model for CSF Aβ42/Aβ40 status based on p-tau181 had an ROC AUC of 0.85 (0.79–0.92) (Table 4 and eTable 9, links.lww.com/WNL/B978). In this model, a higher probability of CSF Aβ42/Aβ40 positivity was associated with APOE ε4 carriers (OR 5.7 [2.3–14]; $p = 0.0002$), cognitive

Table 3 CSF A β 42/A β 40 or Amyloid PET Status as Predicted by Plasma A β 42/A β 40 and Covariates

	Unadjusted model			Covariate adjusted model		
	ROC AUC	Biomarker <i>p</i>	Versus plasma A β 42/A β 40 <i>p</i>	ROC AUC	Biomarker <i>p</i>	Versus plasma A β 42/A β 40 <i>p</i>
Prediction of CSF Aβ42/Aβ40 status (n = 152)						
Plasma Aβ42/Aβ40	0.86 (0.79–0.92)	<0.0001	reference	0.90 (0.85–0.96)	<0.0001	reference
Amyloid probability score	0.89 (0.84–0.95)	<0.0001	0.05	0.91 (0.87–0.96)	<0.0001	NS
Ln (plasma p-tau181)	0.76 (0.68–0.84)	<0.0001	<0.05	0.85 (0.79–0.92)	0.007	NS
Plasma p-tau231 (pg/mL)	0.69 (0.60–0.78)	0.0002	0.004	0.85 (0.78–0.91)	0.01	0.07
Ln (plasma NfL)	0.64 (0.55–0.73)	0.008	<0.0001	0.81 (0.74–0.89)	NS	0.005
Covariates alone	NA	NA	NA	0.82 (0.74–0.89)	NA	0.006
Prediction of amyloid PET status (n = 103)						
Plasma Aβ42/Aβ40	0.86 (0.77–0.95)	<0.0001	reference	0.89 (0.82–0.97)	0.0004	reference
Amyloid probability score	0.90 (0.82–0.97)	<0.0001	NS	0.90 (0.84–0.96)	0.0006	NS
Ln (plasma p-tau181)	0.74 (0.63–0.84)	0.002	0.05	0.84 (0.75–0.92)	0.02	NS
Plasma p-tau231 (pg/mL)	0.69 (0.58–0.81)	0.004	0.02	0.84 (0.75–0.92)	0.01	NS
Ln (plasma NfL)	0.55 (0.43–0.67)	NS	<0.0001	0.82 (0.73–0.91)	NS	NS
Covariates alone	NA	NA	NA	0.81 (0.72–0.90)	NA	0.08

Abbreviations: A β = β -amyloid; AUC = area under the receiver operating characteristic curve; CDR = Clinical Dementia Rating; NfL = neurofilament light; p-tau181 = tau phosphorylated at position 181; p-tau231 = tau phosphorylated at position 231; ROC = receiver operating characteristic.

Logistic regression models evaluated prediction of CSF A β 42/A β 40 (positive <0.0673) or amyloid PET status by each plasma biomarker alone (unadjusted models) or plasma biomarkers and the covariates of self-identified race, sex, age, *APOE* ϵ 4 carrier status, and cognitive status (adjusted models). The amyloid probability score is a proprietary modeled value that incorporates plasma A β 42/A β 40, age, and *APOE* proteotype. Plasma p-tau181 and NfL were transformed with the natural logarithm for analysis. For each model, the ROC AUC with 95% CIs is shown. The significance of each biomarker as a predictor in the model (biomarker *p*) and the difference between the ROC AUC for the plasma A β 42/A β 40 model and other models (versus plasma A β 42/A β 40 *p*) is shown.

Table 4 CSF A β 42/A β 40 Status as Predicted by Plasma Biomarkers and Covariates

Parameter	Estimate	SE	<i>p</i>
Plasma Aβ42/Aβ40, ROC AUC 0.90 (0.85–0.96)			
Intercept	13.0	4.7	0.005
Plasma A β 42/A β 40 (pg/mL)	–220	46	<0.0001
Race (African American)	0.058	0.274	NS
Sex (female)	0.843	0.568	NS
Age, y	0.109	0.04	0.007
APOE ϵ 4 status (carrier)	0.865	0.269	0.001
Cognitive status (CDR>0)	1.11	0.41	0.007
Plasma p-tau181, ROC AUC 0.85 (0.79–0.92)			
Intercept	–8.69	2.71	0.001
Ln (plasma p-tau181)	1.53	0.57	0.007
Race (African American)	–0.59	0.22	0.007
Sex (female)	–0.21	0.44	NS
Age, y	0.072	0.035	0.04
APOE ϵ 4 status (carrier)	0.87	0.23	0.0002
Cognitive status (CDR>0)	1.02	0.39	0.009
Plasma p-tau231, ROC AUC 0.85 (0.78–0.91)			
Intercept	–6.95	2.50	0.006
Plasma p-tau231 (pg/mL)	0.098	0.040	0.01
Race (African American)	–0.60	0.22	0.006
Sex (female)	–0.37	0.43	NS
Age, y	0.096	0.034	0.004
APOE ϵ 4 status (carrier)	0.94	0.23	<0.0001
Cognitive status (CDR >0)	1.07	0.38	0.006
Plasma NfL, ROC AUC 0.81 (0.74–0.89)			
Intercept	–6.20	2.41	0.01
Ln (plasma NfL)	–0.097	0.476	NS
Race (African American)	–0.65	0.22	0.003
Sex (female)	–0.50	0.42	NS
Age, y	0.109	0.040	0.007
APOE ϵ 4 status (carrier)	0.89	0.23	<0.0001
Cognitive status (CDR>0)	1.27	0.39	0.001

Abbreviations: A β = β -amyloid; AUC = area under the receiver operating characteristic curve; CDR = Clinical Dementia Rating; NfL = neurofilament light; p-tau181 = tau phosphorylated at position 181; ROC = receiver operating characteristic.

Logistic regression models evaluated prediction of CSF A β 42/A β 40 status (positive <0.0673) by each plasma biomarker and the covariates of self-identified race, sex, age, APOE ϵ 4 carrier status, and cognitive status. Plasma p-tau181 and NfL were transformed with the natural logarithm for analysis. For each model, the ROC AUC with 95% CI is shown.

impairment (OR 7.7 [1.7–36]; *p* = 0.009), and older age in years (OR 1.08 [1.00–1.15]; *p* = 0.04); AA race was associated with a lower probability of positivity (OR 0.31 [0.13–0.73]; *p* = 0.007). Models of CSF A β 42/A β 40 or amyloid PET status based on p-tau231 (eTable 10) or NfL (eTable 11) were also evaluated and are summarized in Table 4.

A model of CSF A β 42/A β 40 status based only on covariates demonstrated that AA race was associated with a lower probability of CSF A β 42/A β 40 positivity (OR 0.27 [0.12–0.64]; *p* = 0.003) (eTable 12, links.lww.com/WNL/B978). AA race significantly decreased the probability of CSF A β 42/A β 40 positivity in models based on plasma p-tau181 (OR 0.31 [0.13–0.73]; *p* = 0.007), p-tau231 (OR 0.30 [0.13–0.71]; *p* = 0.006), or NfL (OR 0.27 [0.12–0.64]; *p* = 0.003) levels. Consistent with these results, AA race decreased the probability of amyloid PET positivity in models including plasma p-tau181 (OR 0.19 [0.06–0.63]; *p* = 0.007), p-tau231 (OR 0.17 [0.05–0.59]; *p* = 0.005), or NfL (OR 0.17 [0.05–0.55]; *p* = 0.003) levels (eTables 9–11, respectively). In contrast, race did not affect the probability of CSF A β 42/A β 40 or amyloid PET positivity associated with plasma A β 42/A β 40 (eTable 7). Models of CSF A β 42/A β 40 status including only cognitively normal individuals (91% of cohort) showed the same major findings as models that included the entire cohort (eTable 13). Models of CSF A β 42/A β 40 status were also evaluated that incorporated either the interaction between race and APOE ϵ 4 carrier status or race and plasma biomarker levels, but neither interaction was significant for any model and therefore the interactions were not included in the final analyses.

Combining Plasma Biomarkers

A model of CSF A β 42/A β 40 status including levels of all plasma biomarkers and covariates had an ROC AUC of 0.92 (0.88–0.96), which was not significantly different from the ROC AUC of the model including A β 42/A β 40 as the only plasma biomarker (eTable 14, links.lww.com/WNL/B978). In the model with all plasma biomarkers, plasma A β 42/A β 40 was the only biomarker that was a significant predictor (*p* < 0.0001): plasma p-tau181, p-tau231, and NfL were not significant predictors of CSF A β 42/A β 40 after adjusting for the effects of plasma A β 42/A β 40 and covariates. In a similar model of amyloid PET status, plasma A β 42/A β 40 and plasma NfL levels were both significant predictors (*p* = 0.0004 and *p* = 0.007, respectively). In models of CSF A β 42/A β 40 or amyloid PET status with all plasma biomarkers and covariates (including plasma A β 42/A β 40), race was not a significant predictor.

Discussion

This study found that the C2N Diagnostics PrecivityAD plasma A β 42/A β 40 assay more accurately classified CSF A β 42/A β 40 or amyloid PET status compared with Simoa-based assays for plasma p-tau181, p-tau231, and NfL in a mostly cognitively normal cohort of matched AA and NHW

research participants. Self-identified race did not affect prediction of CSF A β 42/A β 40 or amyloid PET status by plasma A β 42/A β 40. However, AA individuals had a significantly lower probability of CSF or amyloid PET positivity compared with NHW in models incorporating plasma p-tau181, p-tau231, or NfL levels, suggesting that predictive algorithms for these assays would perform inconsistently across racial groups and that applying cutoffs established in NHW individuals to AA individuals could lead to disproportionate misdiagnosis of AA.

Plasma biomarkers have been almost exclusively studied in non-Hispanic White cohorts, with little data available on the performance of these biomarkers in other groups. A recent study of a multiracial cohort found good performance of plasma p-tau217 in distinguishing clinical, pathologic, and amyloid PET status, but performance of the assay in predicting amyloid PET status across racial groups could not be ascertained because only 40 individuals had amyloid PET data.³⁷ Another study found that plasma p-tau181 and plasma p-tau181/A β 42 were associated with brain amyloidosis and hippocampal atrophy in a Singaporean AD cohort with high burden of cerebrovascular disease, but it did not investigate potential plasma biomarker differences across racial groups.³⁸ Plasma NfL has been studied in a large Latino cohort, but amyloid PET data were only available in a relatively small subset of participants.³⁹ To reduce racial disparities in research and clinical care, it is important to confirm that plasma biomarker assays have accurate and consistent performance in identifying amyloid status across racial and ethnic groups.

Comparing the absolute values of biomarkers corrected for covariates may be misleading in evaluating which biomarkers perform consistently across racial groups. For example, in this study AA individuals had higher average plasma A β 42/A β 40 compared with NHW individuals, but this reflected lower levels of brain amyloidosis in AA individuals and did not affect the probability of CSF A β 42/A β 40 positivity associated with a given plasma A β 42/A β 40 value. In contrast, plasma p-tau181 levels did not vary by race, but AA individuals were less likely to be amyloid positive at a given plasma p-tau181 value. Without a comparison to reference standards, investigators might have concluded that plasma A β 42/A β 40 was more variable across racial groups and that p-tau isoforms were more consistent, when in fact plasma A β 42/A β 40 was accurately detecting differences in brain amyloidosis by racial group. Confirming that plasma biomarker assays have accurate and consistent performance in identifying amyloid status across racial and ethnic groups requires comparison with a reference standard, and not just covariate-adjusted models of absolute levels.

Previous studies have found an inconsistent relationship between amyloid biomarkers and race. One study found that AA individuals had higher measures of amyloid PET⁴⁰; another recent study found the opposite result.¹² Some studies have found no differences in CSF A β 42 levels by racial group,⁶⁻⁸ but the current findings demonstrate that CSF A β 42 alone may not

reveal significant racial differences that are apparent when CSF A β 42/A β 40 is evaluated. The inconsistent relationship between race and amyloid biomarkers could reflect variation in recruitment methods: NHW and AA individuals are often recruited differently (e.g., NHW are more often referred by health care providers and AA individuals are more often referred by community contacts).^{41,42} Recruitment differences could result in racial groups having significantly different comorbidities, social determinants of health, or frequencies of brain amyloidosis. Potential differences in brain amyloidosis by racial group again suggest that comparison of plasma biomarkers with a reference standard, rather than comparison of absolute values, may be more helpful in establishing which plasma biomarker assays are accurate and consistent across racial groups.

One important issue in the fluid biomarker field is that different assays for plasma analytes have widely varying performance. A recent head-to-head comparison of 8 different plasma A β 42/A β 40 assays found ROC AUCs with CSF A β 42/A β 40 status ranging from a maximum of 0.86 for the Washington University assay that is the basis for the C2N assay used in this study down to a minimum of 0.69 for some immunoassays (0.50 is chance alone).⁴³ In another head-to-head comparison study, different p-tau assays yielded somewhat different findings, even for the same p-tau isoform.⁴⁴ The differences in assay performance complicate comparisons of the relationship of different biomarker analytes to factors such as race. For example, it is unclear whether the probability of CSF A β 42/A β 40 or amyloid PET positivity would be affected by race in models incorporating plasma p-tau181, p-tau231, or p-tau217 measured with higher performing assays (e.g., ROC AUC of >0.85 with CSF A β 42/A β 40 or amyloid PET status). Performance of plasma assays may vary markedly in prediction of brain amyloidosis depending on the study cohort. For example, the p-tau181 assay used in the current study performed very well in predicting amyloid PET status in a cohort including both cognitively normal and cognitively impaired individuals (ROC AUC 0.88),¹⁹ but the performance was lower when predicting amyloid PET status in cognitively normal individuals (ROC AUC 0.82).⁴⁵ Overall, use of consistently high-performing assays is needed to make accurate conclusions about comparative associations of biomarkers.

Although this study made use of one of the largest AD research cohorts with CSF and amyloid PET data, there are major limitations in the conclusions. Individuals enrolled in this study were primarily from the greater St. Louis metropolitan area and individuals from other geographic regions may vary in key characteristics such as medical comorbidities or social determinants of health. The very small number of individuals with cognitive impairment (7 of 76 in each group) was not sufficient to allow analysis of the relationships between cognitive impairment, race, and biomarker levels. This study of 76 matched pairs of individuals, in which 6 variables had significant effects, was also not sufficiently powered to evaluate the underlying reasons for the racial differences. The

Hollingshead index of social position demonstrated that AA individuals had a slightly lower social position compared with NHW individuals. However, this measure does not capture the complex social factors that may underlie biomarker differences between the groups. AA individuals had a higher rate of hypertension and diabetes compared with NHW individuals, but the relatively small cohort did not permit a detailed investigation of these effects. For example, only 4 NHW individuals had diabetes, which does not permit analysis of race by diabetes interactions. Although this study is insufficiently powered or does not have the data available to answer many important questions, it does document racial differences in plasma biomarkers that could potentially lead to clinical misdiagnosis, bias clinical trials that use a biomarker cutoff for inclusion,^{12,46} and affect interpretation of biomarkers as a secondary end point. These findings should encourage investigators to evaluate the performance of plasma biomarker assays in diverse cohorts. This report strengthens the justification for the creation of large, diverse cohorts that are adequately powered to evaluate the underlying reasons for racial differences.

It is critical to understand that biomarker differences associated with race likely reflect differences in medical comorbidities, social determinants of health, or the effects of systemic racism rather than inherent biological differences.¹⁰ For example, in this study cohort there were differences in the rates of hypertension and diabetes by racial group, and recent work has demonstrated that major medical comorbidities such as heart and kidney disease may affect plasma biomarker levels.⁴⁷ AD research cohorts have traditionally not collected detailed information about social determinants of health such as economic stability, access to healthy foods, neighborhood safety, and quality of education that may be associated with dementia; the importance of these factors is now gaining greater recognition.⁴⁸ The greater accessibility and acceptance of blood-based AD biomarkers may enable creation of larger cohorts and increased inclusion of groups, such as AA individuals, that have been underrepresented in AD biomarker studies.⁴⁹ Much larger longitudinal studies of diverse cohorts are needed to evaluate the intersection of race, AD biomarkers, cognitive impairment, medical comorbidities, and social determinants of health.⁵⁰ Improved understanding of these complex factors will enable more accurate AD diagnosis and improve patient care for all groups.

Acknowledgment

The authors thank the research volunteers who participated in the studies from which these data were obtained and their families and the Clinical, Fluid Biomarker and Imaging Cores at the Knight Alzheimer Disease Research Center for sample and data collection.

Study Funding

This study was supported by National Institute on Aging grants R01AG070941 (S.E.S.), K23AG053426 (S.E.S.), P30AG066444 (J.C.M.), P01AG003991 (J.C.M.), P01AG026276 (J.C.M.), R01AG067505 (C.X.), and RF1R01AG053550 (C.X.) and the

Cure Alzheimer's Fund (K.L.M.). C2N Diagnostics provided the plasma A β 42/A β 40 assays for this study. T.K.K. was funded by the Alzheimer's Association Research Fellowship (850325), the BrightFocus Foundation (A2020812F), the International Society for Neurochemistry's Career Development Grant, the Swedish Alzheimer Foundation (Alzheimerfonden; AF-930627), the Swedish Brain Foundation (Hjärnfonden; FO2020-0240), the Swedish Dementia Foundation (Demensförbundet), the Swedish Parkinson Foundation (Parkinsonfonden), the Gamla Tjänarinnor Foundation, the Aina (Ann) Wallströms and Mary-Ann Sjöbloms Foundation, the Agneta Prytz-Folkes & Gösta Folkes Foundation (2020-00124), the Gun and Bertil Stohnes Foundation, and the Anna Lisa and Brother Björnsson's Foundation. H.Z. is a Wallenberg Scholar supported by grants from the Swedish Research Council (2018–02532), the European Research Council (681712), Swedish State Support for Clinical Research (ALFGBG-720931), the Alzheimer Drug Discovery Foundation (ADDF) USA (201809–2016862), the AD Strategic Fund and the Alzheimer's Association (ADSF-21-831376-C, ADSF-21-831381-C, and ADSF-21-831377-C), the Olav Thon Foundation, the Erling-Persson Family Foundation, Stiftelsen för Gamla Tjänarinnor, Hjärnfonden, Sweden (FO2019-0228), the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement 860197 (MIRIADE), and the UK Dementia Research Institute at UCL. K.B. is supported by the Swedish Research Council (2017-00915), the Swedish Alzheimer Foundation (AF-742881), Hjärnfonden, Sweden (FO2017-0243), the Swedish state under the agreement between the Swedish government and the County Councils, the ALF-agreement (ALFGBG-715986), and the Alzheimer's Association 2021 Zenith Award (ZEN-21-848,495).

Disclosure

S.E. Schindler has received data on behalf of Washington University from C2N Diagnostics at no cost. T.K. Karikari, N.J. Ashton, and R.L. Henson report no disclosures relevant to the manuscript. K.E. Yarasheski, T. West, M.R. Meyer, and K.M. Kirmess are employees of C2N Diagnostics, which offers the PrecivityAD test described in this article. Y. Li, B. Saef, K.L. Moulder, and D. Bradford report no disclosures relevant to the manuscript. A.M. Fagan has received research funding from Biogen, Centene, Fujirebio, and Roche Diagnostics; is a member of the scientific advisory boards for Roche Diagnostics, Genentech, and Diadem; and consults for DiamiR and Siemens Healthcare Diagnostics Inc. B.A. Gordon reports no disclosures relevant to the manuscript. T.L.S. Benzinger has investigator-initiated research funding from the NIH, the Alzheimer's Association, the Barnes-Jewish Hospital Foundation, and Avid Radiopharmaceuticals (a wholly owned subsidiary of Eli Lilly); participates as a site investigator in clinical trials sponsored by Avid Radiopharmaceuticals, Eli Lilly, Biogen, Eisai, Janssen, and Roche; serves as an unpaid consultant to Eisai and Siemens; and is on the speaker's bureau for Biogen. J. Balls-Berry is a member of the patient advisory board and receives financial support for the Dartmouth University project Implementation of Uterine Fibroid

Option Grid Patient Decision Aids Across Five Organizational Settings (UPFRONT; NCT03985449). R.J. Bateman co-founded C2N Diagnostics, receives income from C2N Diagnostics for serving on the scientific advisory board, and consults for Roche, Genentech, AbbVie, Pfizer, Boehringer-Ingelheim, and Merck. Dr. Bateman and Washington University have equity ownership interest in C2N Diagnostics and receive royalty income based on technology (stable isotope labeling kinetics and blood plasma assay) licensed by Washington University to C2N Diagnostics. Washington University, with Dr. Bateman as coinventor, has submitted the US provisional patent application “Plasma Based Methods for Detecting CNS Amyloid Deposition.” C. Xiong consults for Diadem. H. Zetterberg has served on scientific advisory boards and/or as a consultant for Alector, Eisai, Denali, Roche Diagnostics, Wave, Samumed, Siemens Healthineers, Pinteon Therapeutics, Nervgen, AZTherapies, CogRx, and Red Abbey Labs; has given lectures in symposia sponsored by Cellectric, Fujirebio, Alzecure, and Biogen; and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program. K. Blennow has served as a consultant, on advisory boards, or on data monitoring committees for Abcam, Axon, Biogen, JOMDD/Shimadzu, Julius Clinical, Lilly, MagQu, Novartis, Prothena, Roche Diagnostics, and Siemens Healthineers, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures Incubator Program. J.C. Morris is Chair of the Research Strategy Council of the Cure Alzheimer’s Fund. Go to Neurology.org/N for full disclosures.

Publication History

Received by *Neurology* September 24, 2021. Accepted in final form February 22, 2022. Submitted and externally peer reviewed. The handling editor was Linda Hershey, MD, PhD.

Appendix Authors

Name	Location	Contribution
Suzanne E. Schindler, MD, PhD	Washington University	Design and conceptualization of study, major role in the acquisition of data, analyzed the data, drafted the manuscript for intellectual content
Thomas K. Karikari, PhD	University of Gothenburg	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Nicholas J. Ashton, PhD	University of Gothenburg	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Rachel L. Henson, MS	Washington University	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Kevin E. Yarasheski, PhD	C2N Diagnostics	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Tim West, PhD	C2N Diagnostics	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content

Appendix (continued)

Name	Location	Contribution
Matthew R. Meyer, PhD	C2N Diagnostics	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Kristopher M. Kirmess, PhD	C2N Diagnostics	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Yan Li, PhD	Washington University	Analyzed the data, revised the manuscript for intellectual content
Benjamin Saef, MS	Washington University	Analyzed the data, revised the manuscript for intellectual content
Krista L. Moulder, PhD	Washington University	Interpreted the data, revised the manuscript for intellectual content
David Bradford	Washington University	Interpreted the data, revised the manuscript for intellectual content
Anne M. Fagan, PhD	Washington University	Interpreted the data, revised the manuscript for intellectual content
Brian A. Gordon, PhD	Washington University	Interpreted the data and recommended additional analyses, revised the manuscript for intellectual content
Tammie L.S. Benzinger, MD, PhD	Washington University	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Joyce Balls-Berry, PhD	Washington University	Interpreted the data, revised the manuscript for intellectual content
Randall J. Bateman, MD	Washington University	Interpreted the data and recommended additional analyses, revised the manuscript for intellectual content
Chengjie Xiong, PhD	Washington University	Analyzed the data, revised the manuscript for intellectual content
Henrik Zetterberg, MD, PhD	University of Gothenburg	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
Kaj Blennow, MD, PhD	University of Gothenburg	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content
John C. Morris, MD	Washington University	Major role in the acquisition of data, interpreted the data, revised the manuscript for intellectual content

References

- Johnson KA, Minoshima S, Bohnen NI, et al. Appropriate use criteria for amyloid PET: a report of the Amyloid Imaging Task Force, the Society of Nuclear Medicine and Molecular Imaging, and the Alzheimer’s Association. *Alzheimers Dement*. 2013;9:e1-16.
- Shaw LM, Arias J, Blennow K, et al. Appropriate use criteria for lumbar puncture and cerebrospinal fluid testing in the diagnosis of Alzheimer’s disease. *Alzheimers Dement*. 2018;14:1505-1521.
- Kaplow J, Vandijk M, Gray J, et al. Concordance of Lumipulse cerebrospinal fluid t-tau/Abeta42 ratio with amyloid PET status. *Alzheimers Dement*. 2020;16:144-152.
- Hansson O, Seibyl J, Stomrud E, et al. CSF biomarkers of Alzheimer’s disease concord with amyloid-beta PET and predict clinical progression: a study of fully automated immunoassays in BioFINDER and ADNI cohorts. *Alzheimers Dement*. 2018;14:1470-1481.
- Schindler SE, Gray JD, Gordon BA, et al. Cerebrospinal fluid biomarkers measured by Elecsys assays compared with amyloid imaging. *Alzheimers Dement*. 2018;14:1460-1469.
- Howell JC, Watts KD, Parker MW, et al. Race modifies the relationship between cognition and Alzheimer’s disease cerebrospinal fluid biomarkers. *Alzheimers Res Ther*. 2017;9(1):88.

7. Morris JC, Schindler SE, McCue LM, et al. Assessment of racial disparities in biomarkers for Alzheimer disease. *JAMA Neurol.* 2019;76(3):264-273.
8. Schindler SE, Cruchaga C, Joseph A, et al. African Americans have differences in CSF soluble TREM2 and associated genetic variants. *Neurol Genet.* 2021;7(2):e571.
9. Garrett SL, McDaniel D, Obideen M, et al. Racial disparity in cerebrospinal fluid amyloid and tau biomarkers and associated cutoffs for mild cognitive impairment. *JAMA Netw Open.* 2019;2(12):e1917363.
10. Babulal GM, Quiroz YT, Albeni BC, et al. Perspectives on ethnic and racial disparities in Alzheimer's disease and related dementias: update and areas of immediate need. *Alzheimers Dement.* 2019;15:292-312.
11. Meeker KL, Wisch JK, Hudson D, et al. Socioeconomic status mediates racial differences seen using the AT(N) framework. *Ann Neurol.* 2021;89(2):254-265.
12. Deters KD, Napolioni V, Sperling RA, et al. Amyloid PET imaging in self-identified non-Hispanic black participants of the Anti-Amyloid in Asymptomatic Alzheimer's Disease (A4) study. *Neurology.* 2021;96(11):e1491-e1500.
13. Powe NR. Black kidney function matters: use or misuse of race? *JAMA.* 2020;324:737-738.
14. Vyas DA, Eisenstein LG, Jones DS. Hidden in plain sight: reconsidering the use of race correction in clinical algorithms. *N Engl J Med.* 2020;383(9):874-882.
15. Ashton NJ, Leuzy A, Karikari TK, et al. The validation status of blood biomarkers of amyloid and phospho-tau assessed with the 5-phase development framework for AD biomarkers. *Eur J Nucl Med Mol Imaging.* 2021;48(7):2140-2156.
16. Kirmess KM, Meyer MR, Holubasch MS, et al. The PrecivityAD™ test: accurate and reliable LC-MS/MS assays for quantifying plasma amyloid beta 40 and 42 and apolipoprotein E prototype for the assessment of brain amyloidosis. *Clin Chim Acta.* 2021;519:267-275.
17. West T, Kirmess KM, Meyer MR, et al. A blood-based diagnostic test incorporating plasma Abeta42/40 ratio, ApoE prototype, and age accurately identifies brain amyloid status: findings from a multi cohort validity analysis. *Mol Neurodegener.* 2021;16:30.
18. Moscoso A, Grothe MJ, Ashton NJ, et al. Time course of phosphorylated-tau181 in blood across the Alzheimer's disease spectrum. *Brain.* 2021;144(1):325-339.
19. Karikari TK, Pascoal TA, Ashton NJ, et al. Blood phosphorylated tau 181 as a biomarker for Alzheimer's disease: a diagnostic performance and prediction modelling study using data from four prospective cohorts. *Lancet Neurol.* 2020;19:422-433.
20. Palmqvist S, Janelidze S, Quiroz YT, et al. Discriminative accuracy of plasma phospho-tau217 for Alzheimer disease vs other neurodegenerative disorders. *JAMA.* 2020;324(8):772-781.
21. Barthélemy NR, Horie K, Sato C, Bateman RJ. Blood plasma phosphorylated-tau isoforms track CNS change in Alzheimer's disease. *J Exp Med.* 2020;217(11):217.
22. Thijssen EH, La Joie R, Strom A, et al. Plasma phosphorylated tau 217 and phosphorylated tau 181 as biomarkers in Alzheimer's disease and frontotemporal lobar degeneration: a retrospective diagnostic performance study. *Lancet Neurol.* 2021;20(9):739-752.
23. Ashton NJ, Pascoal TA, Karikari TK, et al. Plasma p-tau231: a new biomarker for incipient Alzheimer's disease pathology. *Acta Neuropathol.* 2021;141(5):709-724.
24. Preische O, Schultz S, Apel A, et al. Serum neurofilament dynamics predicts neurodegeneration and clinical progression in presymptomatic Alzheimer's disease. *Nat Med.* 2019;25:277-283.
25. Morris JC, Weintraub S, Chui HC, et al. The Uniform data Set (UDS): clinical and cognitive variables and descriptive data from alzheimer disease centers. *Alzheimer Dis Assoc Disord.* 2006;20:210-216.
26. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology.* 1993;43(11):2412-2414.
27. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state": a practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12:189-198.
28. Hollingshead Two Factor Index of Social Position, 5th ed. Sage Publications; 1991.
29. Cruchaga C, Kauwe JS, Mayo K, et al. SNPs associated with cerebrospinal fluid phospho-tau levels influence rate of decline in Alzheimer's disease. *PLoS Genet.* 2010;6(9):e1001101.
30. Schindler SE, Bollinger JG, Ovod V, et al. High-precision plasma beta-amyloid 42/40 predicts current and future brain amyloidosis. *Neurology.* 2019;93:e1647-e1659.
31. Su Y, Blazey TM, Snyder AZ, et al. Partial volume correction in quantitative amyloid imaging. *NeuroImage.* 2015;107:55-64.
32. Su Y, Flores S, Wang G, et al. Comparison of Pittsburgh compound B and florbetapir in cross-sectional and longitudinal studies. *Alzheimers Dement.* 2019;11:180-190.
33. Su Y, Flores S, Hornbeck RC, et al. Utilizing the Centiloid scale in cross-sectional and longitudinal PiB PET studies. *Neuroimage Clin.* 2018;19:406-416.
34. Volluz KE, Schindler SE, Henson RL, et al. *Correspondence of CSF Biomarkers Measured by Lumipulse Assays with Amyloid PET. 2021 Alzheimer's Association International Conference;* 2021.
35. Vlassenko AG, McCue L, Jasielec MS, et al. Imaging and cerebrospinal fluid biomarkers in early preclinical Alzheimer disease. *Ann Neurol.* 2016;80(3):379-387.
36. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics.* 1988;44(3):837-845.
37. Brickman AM, Manly JJ, Honig LS, et al. Plasma p-tau181, p-tau217, and other blood-based Alzheimer's disease biomarkers in a multi-ethnic, community study. *Alzheimers Dement.* 2021;17:1353-1364.
38. Chong JR, Ashton NJ, Karikari TK, et al. Plasma P-tau181 to Abeta42 ratio is associated with brain amyloid burden and hippocampal atrophy in an Asian cohort of Alzheimer's disease patients with concomitant cerebrovascular disease. *Alzheimers Dement.* 2021;17(10):1649-1662.
39. O'Bryant S, Petersen M, Hall J, et al. Characterizing plasma NfL in a community-dwelling multi-ethnic cohort: results from the HABLE study. *Alzheimers Dement.* 2022;18(2):240-250.
40. Gottesman RF, Schneider AL, Zhou Y, et al. The ARIC-PET amyloid imaging study: brain amyloid differences by age, race, sex, and APOE. *Neurology.* 2016;87(5):473-480.
41. Raman R, Quiroz YT, Langford O, et al. Disparities by race and ethnicity among adults recruited for a preclinical alzheimer disease trial. *JAMA Netw Open.* 2021;4(7):e2114364.
42. Gleason CE, Norton D, Zuelsdorff M, et al. Association between enrollment factors and incident cognitive impairment in Blacks and Whites: data from the Alzheimer's Disease Center. *Alzheimers Dement.* 2019;15:1533-1545.
43. Janelidze S, Teunissen CE, Zetterberg H, et al. Head-to-head comparison of 8 plasma amyloid-beta 42/40 assays in Alzheimer disease. *JAMA Neurol.* 2021;78(11):1375-1382.
44. Mielke MM, Frank RD, Dage JL, et al. Comparison of plasma phosphorylated tau species with amyloid and tau positron emission tomography, neurodegeneration, vascular pathology, and cognitive outcomes. *JAMA Neurol.* 2021;78(9):1108-1117.
45. Keshavan A, Pannee J, Karikari TK, et al. Population-based blood screening for preclinical Alzheimer's disease in a British birth cohort at age 70. *Brain.* 2021;144(2):434-449.
46. Gottesman RF, Hamilton R. Recruiting diverse populations in clinical trials: how do we overcome selection bias? *Neurology.* 2021;96:509-510.
47. Syrjanen JA, Campbell MR, Algeciras-Schminich A, et al. Associations of amyloid and neurodegeneration plasma biomarkers with comorbidities. *Alzheimers Dement.* Epub 2021 Sep 27.
48. Wilkins CH, Schindler SE, Morris JC. Addressing health disparities among minority populations: why clinical trial recruitment is not enough. *JAMA Neurol.* 2020;77(9):1063-1064.
49. Howell JC, Parker MW, Watts KD, Kollhoff A, Tsvetkova DZ, Hu WT. Research lumbar punctures among African Americans and Caucasians: perception predicts experience. *Front Aging Neurosci.* 2016;8:296.
50. Barnes LL. Alzheimer disease in African American individuals: increased incidence or not enough data? *Nat Rev Neurol.* 2022;18:56-62.

Neurology[®]

Effect of Race on Prediction of Brain Amyloidosis by Plasma A β 42/A β 40, Phosphorylated Tau, and Neurofilament Light

Suzanne E. Schindler, Thomas K. Karikari, Nicholas J. Ashton, et al.
Neurology 2022;99:e245-e257 Published Online before print April 21, 2022
DOI 10.1212/WNL.0000000000200358

This information is current as of April 21, 2022

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/99/3/e245.full
References	This article cites 47 articles, 5 of which you can access for free at: http://n.neurology.org/content/99/3/e245.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Alzheimer's disease http://n.neurology.org/cgi/collection/alzheimers_disease Health disparities http://n.neurology.org/cgi/collection/health_disparities Inclusion, Diversity, Equity, Anti-racism, and Social Justice (IDEAS) http://n.neurology.org/cgi/collection/all_equity_diversity_and_inclusion PET http://n.neurology.org/cgi/collection/pet
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

