

2022

## Psychological outcomes in advanced cancer patients after receiving genomic tumor profiling results

Sabina Vatter

Timothy E. Schlub

Christine E. Napier

Megan C. Best

Nicci Bartley

*See next page for additional authors*

Follow this and additional works at: [https://researchonline.nd.edu.au/ies\\_article](https://researchonline.nd.edu.au/ies_article)

 Part of the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Vatter, S., Schlub, T. E., Napier, C. E., Best, M. C., Bartley, N., Juraskova, I., Meiser, B., Ballinger, M. L., Biesecker, B. B., Goldstein, D., Thomas, D. M., Butow, P., & Members of the PiGeOn Project (2022). Psychological outcomes in advanced cancer patients after receiving genomic tumor profiling results. *Health Psychology, 41* (6), 396-408.

Original article available here:

[10.1037/hea0001181](https://doi.org/10.1037/hea0001181)

This article is posted on ResearchOnline@ND at . For more information, please contact [researchonline@nd.edu.au](mailto:researchonline@nd.edu.au).



---

**Authors**

Sabina Vatter, Timothy E. Schlub, Christine E. Napier, Megan C. Best, Nicci Bartley, Ilona Juraskova, Bettina Meiser, Mandy L. Ballinger, Barbara B. Biesecker, David Goldstein, David M. Thomas, Phyllis Butow, and Members of the PiGeOn Project

© 2022 American Psychological Association.

This article may not exactly replicate the authoritative document published in the APA journal. It is not the copy of record.

Vatter, S., Schlub, T.E., Napier, C.E., Best, M.C., Bartley, N., Juraskova, I., Meiser, B., Ballinger, M.L., Biesecker, B.B., Goldstein, D., Thomas, D.M., Butow, P., & Members of the PiGeOn Project. (2022). Psychological outcomes in advanced cancer patients after receiving genomic tumor profiling results. *Health Psychology, 41*(6), 396-408.

<https://doi.org/10.1037/hea0001181>

This article was accepted for publication in *Health Psychology*, 3 February 2022.

Published version available online at: - <https://doi.org/10.1037/hea0001181>

**Psychological Outcomes in Advanced Cancer Patients After Receiving Genomic Tumour Profiling**

**Results**

Sabina Vatter, PhD<sup>1</sup>, Timothy E. Schlub, PhD,<sup>2</sup> Christine E. Napier, PhD,<sup>3</sup> Megan C. Best, PhD,<sup>1,4</sup> Nicci Bartley, MA<sup>1</sup> Ilona Juraskova, PhD,<sup>5</sup> Bettina Meiser, PhD,<sup>6</sup> Mandy L. Ballinger, PhD,<sup>3,7</sup> Barbara B. Biesecker, PhD,<sup>8</sup> David Goldstein, MD,<sup>9</sup> David M. Thomas, PhD,<sup>3,7</sup> Phyllis Butow, PhD,<sup>1</sup> and Members of the PiGeOn Project

<sup>1</sup> School of Psychology, Faculty of Science, Psycho-Oncology Co-operative Research Group (PoCoG), University of Sydney, Camperdown, NSW 2006, Australia

<sup>2</sup> Sydney School of Public Health, Faculty of Medicine and Health, University of Sydney, NSW 2006, Australia

<sup>3</sup> Cancer Theme, Garvan Institute of Medical Research, Darlinghurst, NSW 2010, Australia

<sup>4</sup> Institute for Ethics and Society, University of Notre Dame Australia, Broadway, NSW 2007, Australia

<sup>5</sup> School of Psychology, Faculty of Science, Centre for Medical Psychology and Evidence-Based Decision-Making (CeMPED), University of Sydney, Camperdown, NSW 2006, Australia

<sup>6</sup> Psychosocial Research Group, Prince of Wales Clinical School, University of New South Wales, Kensington, NSW 2032, Australia

<sup>7</sup> St Vincent's Clinical School, University of New South Wales, Darlinghurst, NSW 2010, Australia

<sup>8</sup> Newborn Screening, Ethics and Disability Studies, RTI International, Washington, DC, USA

<sup>9</sup> Department of Medical Oncology, Prince of Wales Hospital, NSW 2031, Australia

Members of the PiGeOn Project are named authors, plus Ainsley Newson, Kathy Tucker, Richard Vines, Kate Vines, Judy Kirk, Mary-Anne Young, Jacqueline Savard, and Chris Jacobs.

## Supplementary files

Table S1. One-way ANOVA according to actionable via MoST sub-study, actionable via other pathway or non-actionable results for hope (HHI), anxiety (HADS-A), depression (HADS-D) and CGP-specific anxiety (IES) at T2 (n=374)

	Actionable via MoST sub-study		Actionable via different pathway		Non-actionable		F (df)	p-value
	n	M (SD)	n	M (SD)	n	M (SD)		
<b>HHI</b>	79	1.76 (1.21)	139	2.04 (1.23)	124	1.91 (1.22)	1.37 (2)	.254
<b>HADS-A</b>	103	3.09 (1.26)	141	2.94 (1.27)	130	2.82 (1.34)	1.21 (2)	.300
<b>HADS-D</b>	103	3.36 (1.11)	141	3.37 (1.03)	130	3.50 (.97)	.73 (2)	.482
<b>IES</b>	105	2.99 (1.26)	142	2.89 (1.23)	127	2.72 (1.27)	1.46 (2)	.233

Abbreviations: CGP – comprehensive genomic profiling; HADS – Hospital Anxiety and Depression Scale (Anxiety or Depression sub-scale); HHI – Herth Hope Index; IES – Impact of Events Scale.

Table S2. Correlations between hope (HHI), anxiety (HADS-A), depression (HADS-D) and CGP-specific anxiety (IES) at T2

	T2 HHI	T2 HADS-A	T2 HADS-D	T2 IES	Age	Comorbidity Index	T0 Self efficacy	T0 Knowledge	T0 Att. to uncertainty	T0 Perc. Susceptibility	T0 – Perc. Importance of CGP	SES
T2 HADS-A	<b>-.554***</b>											
T2 HADS-D	<b>-.616***</b>	<b>.606***</b>										
T2 IES	<b>-.223***</b>	<b>.356***</b>	<b>.217***</b>									
Age	.135**	<b>-.192***</b>	-.028	-.038								
Comorbidity Index	.072	<b>-.171***</b>	.019	-.015	<b>.613***</b>							
T0 Self-efficacy	<b>.225***</b>	-.098*	-.141**	<b>-.155***</b>	.026	.009						
T0 Knowledge	.018	-.017	-.019	-.030	.028	.047	.026					
T0 Attitude towards uncertainty	.108*	-.005	-.037	-.093*	.018	-.041	<b>.508***</b>	-.071				
T0 Perceived susceptibility	-.148**	.067	.083	-.136**	.020	.019	<b>.154***</b>	-.050	<b>.226***</b>			
T0 Perceived importance of CGP	.038	.066	.022	.098*	.015	-.003	<b>.171***</b>	-.038	<b>.246***</b>	.041		
Socioeconomic status (SES)	.056	-.051	-.035	-.020	-.025	.002	.078*	-.005	.025	.029	-.011	
Time since first cancer diagnosis (months)	-.026	-.024	.012	-.001	.061	.035	.008	.029	-.002	.074*	-.028	.055

Abbreviations: CGP – comprehensive genomic profiling; HADS – Hospital Anxiety and Depression Scale (Anxiety or Depression sub-scale); HHI – Herth Hope Index; IES – Impact of Events Scale; SES – socioeconomic status.

Notes: \*\*\* p < .001, \*\* p < .01, \* p < .05. P-values below the Bonferroni correction (< .00185) highlighted in bold.

Table S3. Multiple hierarchical linear regression models of predictors of hope (HHI), anxiety (HADS-A), depression (HADS-D) and CGP-specific anxiety (IES) at T2 (n=374)

		<b>B</b>	<b>95% CI for B</b>	<b>p-value</b>	<b>R<sup>2</sup></b>	<b>Change in R<sup>2</sup> (p-value)</b>
<b>T2 HHI</b>						
<i>Step 1</i>	T1 HHI	.47	.40 ... .55	<b>.000</b>	.53	.53 (<.001)
<i>Step 2</i>					.67	.14 (<.001)
	T0 self-efficacy	.60	.06 ... 1.13	.031		
	T0 perceived susceptibility	-.02	-.03 ... -.01	<b>.002</b>		
	T2 HADS-A	-.20	-.30 ... -.10	<b>.000</b>		
	T2 HADS-D	-.44	-.56 ... -.32	<b>.000</b>		
<b>T2 HADS-A</b>						
<i>Step 1</i>	T1 HADS-A	.54	.46 ... .61	<b>.000</b>	.56	.56 (<.001)
<i>Step 2</i>	Gender (ref = male)	-.69	-1.20 ... -.19	.007	.67	.11 (<.001)
	Language (ref = only English at home)	.82	.10 ... 1.54	.025		
	T2 HHI	-.11	-.17 ... -.04	.001		
	T2 HADS-D	.29	.20 ... .38	<b>.000</b>		
	T2 IES	.03	.01 ... .05	.008		
<b>T2 HADS-D</b>						
<i>Step 1</i>	T1 HADS-D	.32	.24 ... .40	<b>.000</b>	.39	.39 (<.001)
<i>Step 2</i>	T2 HHI	-.19	-.25 ... -.13	<b>.000</b>	.58	.19 (<.001)
	T2 HADS-A	.27	.20 ... .34	<b>.000</b>		
	Age	.03	.01 ... .05	.002		
	Socioeconomic status	.09	.00 ... .17	.045		
<b>T2 IES</b>						
<i>Step 1</i>	T1 IES	.43	.35 ... .51	<b>.000</b>	.29	.29 (<.001)
<i>Step 2</i>	T0 perceived susceptibility	-.05	-.09 ... -.01	.020	.35	.05 (<.001)
	ECOG (ref = score 1+)	-2.26	-4.34 ... -.17	.034		
	Education (ref = postgrad)	-.96	-1.86 ... -.07	.035		
	T2 HADS-A	.52	.27 ... .78	<b>.000</b>		

Abbreviations: ARIA – Accessibility and Remoteness Index of Australia; B – unstandardized beta; CGP – comprehensive genomic profiling; CI – confidence intervals; ECOG – Eastern Cooperative Oncology Group; HADS – Hospital Anxiety and Depression Scale (Anxiety or Depression sub-scale); HHI – Herth Hope Index; IES – Impact of Events Scale, Tx outcome – treatment (actionable vs non-actionable).

Notes: \*\*\* p < .001, \*\* p < .01. P-values below the Bonferroni correction (< .00185) highlighted in bold.

Table S4. Multinomial logistic regression of predictors of hope (HHI), anxiety (HADS-A), depression (HADS-D) and CGP-specific anxiety (IES) (n=374)

Trajectories	Variables	OR	95% CI	p	OR	95% CI	p	OR	95% CI	p
<b>Herth Hope Index (HHI)</b>		<b>High-High as ref.</b>			<b>High-Low as ref.</b>			<b>Low-High as ref.</b>		
<b>High-High</b>	T0 perceived susceptibility	NA			.95	.93 – .98	.000	NA		
	Comorbidity Index	NA			.51	.33 – .78	.002	NA		
	Previous radiotherapy	NA			3.86	1.05 – >10	.042	NA		
	Previous surgery	NA			NA			2.93	1.27 – 6.77	.012
	Time since diagnosis	NA			NA			.98	.97 – .99	.001
	T1 HADS-A	NA			.99	.49 – .79	.000	.53	.39 – .71	.000
	T1 HADS-D	NA			NA			.02	.00 – .23	.002
<b>High-Low</b>	T0 perceived susceptibility	1.05	1.02 – 1.08	.000	NA			NA		
	Comorbidity Index	1.98	1.28 – 3.05	.003	NA			2.24	1.16 – 4.30	.016
	Previous radiotherapy	.26	.07 – .95	.042	NA			NA		
	Previous surgery	NA			NA			3.52	1.36 – 9.12	.010
	Time since diagnosis	NA			NA			.98	.96 – 1.00	.021
	T1 HADS-A	1.61	1.27 – 2.03	.000	NA			NA		
	T1 HADS-D	NA			NA			NA		
<b>Low-High</b>	T0 perceived susceptibility	NA			NA			NA		
	Comorbidity Index	NA			.45	.23 – .86	.016	NA		
	Previous surgery	.24	.15 – .79	.012	.28	.11 ... 74	.010	NA		
	Time since diagnosis	1.02	1.01 – 1.03	.001	1.02	1.00 – 1.04	.021	NA		
	T1 HADS-A	1.90	1.40 – 2.56	.000	NA			NA		
	T1 HADS-D	NA			NA			NA		
	Medical/science occupation	48.44	4.27 – >10	.002	NA			NA		
<b>Low-Low</b>	T0 perceived susceptibility	1.04	1.01 – 1.06	.002	NA			NA		
	Comorbidity Index	NA			NA			NA		



	Previous radiotherapy	.04	.01 – .24	.001					
	Time since diagnosis						.98	.97 – .1.00 .024	
	T1 HADS-A	1.81	1.24 – 2.31	.000					
	T1 HADS-D	1.40	1.13 – 1.73	.002	1.38	1.07 – 1.79	.014		
	Medical/science occupation	10.6	1.26 – >10	.030					
		1							
<b>HADS-Anxiety (HADS-A)</b>		<b>Low-Low as ref.</b>			<b>High-Low as ref.</b>			<b>Low-High as ref.</b>	
<b>High-High</b>	Age						1.17	1.03 – 1.34 .017	
	SES								
	Comorbidity Index	.55	.31 – .96	.035		n.s.	.20	.07 – .57 .003	
	T1 HHI	.68	.58 – .79	.000					
	T1 HADS-D	1.27	1.05 – 1.54	.013			1.96	1.06 – 3.64 .033	
	T1 IES	1.14	1.08 – 1.21	.000					
	ARIA – major city						101.67	2.53 ... >10 .014	
	ARIA – inner regional						138.62	2.03 ... >10 .022	
	ECOG score	3.64	1.18 – >10	.025					
<b>High-Low</b>	Age						1.17	1.03 – 1.34 .018	
	SES						.52	.31 – .87 .013	
	Comorbidity Index					NA	.23	.08 – .67 .007	
	T1 HHI	.77	.66 – .89	.000					
	T1 HADS-D								
	T1 IES	1.12	1.06 – 1.18	.000					
	ARIA – major city						42.6	1.10 ... >10 .044	
	ECOG score						4		
<b>Low-High</b>	Age	.84	.74 – .96	.008	.85	.75 – .97	.018		
	SES	1.82	1.10 – 3.01	.020	1.93	1.15 – 3.25	.013		
	Comorbidity Index	2.77	1.04 – 7.38	.042	4.29	1.50 – >10	.007	NA	
	T1 HHI								
	T1 HADS-D								
	T1 IES	1.15	1.05 – 1.27	.002					
	ARIA – major city	.01	.00 – .50	.019	.02	.00 – .91	.044		
	ECOG score								

<b>Low-Low</b>	Age							1.19	1.05 – 1.35	.008
	SES							.55	.33 – .91	.020
	Comorbidity Index							.36	.14 – .96	.042
	T1 HHI	NA			1.30	1.13 – 1.51	.000			
	T1 HADS-D									
	T1 IES				.89	.85 – .95	.000	.87	.79 – .95	.002
	ARIA – major city							69.4	2.00 ...	.019
	ECOG score							1	>10	
<b>HADS-Depression (HADS-D)</b>		<b>Low-Low as ref.</b>			<b>High-Low as ref.</b>			<b>Low-High as ref.</b>		
<b>High-High</b>	SES							.50	.29 – .89	.017
	T1 HHI	.62	.45 – .86	.004						
	T1 HADS-A	3.20	1.75 – 5.85	.000				3.42	1.81 – 6.48	.000
<b>High-Low</b>	Relative with cancer				.01	.00 – .24	.007	.02	.00 – .61	.025
	SES	.64	.45 – .91	.013				.50	.33 – .877	.002
	T1 HHI									
	T1 HADS-A	1.88	1.28 – 2.75	.001	NA			2.00	1.31 – 3.07	.001
	Relative with cancer	8.65	1.03 – <10	.047						
<b>Low-High</b>	SES				1.99	1.29 – 3.07	.002			
	T1 HHI	.84	.72 – .98	.025						
	T1 HADS-A				.50	.33 – .77	.001	NA		
	Relative with cancer									
<b>Low-Low</b>	SES				1.56	1.10 – 2.21	.013			
	T1 HHI							.78	.59 – 1.04	.025
	T1 HADS-A	NA			.53	.36 – .78	.001			
<b>Impact of Events Scale (IES)</b>		<b>Low-Low as ref.</b>			<b>High-Low as ref.</b>			<b>Low-High as ref.</b>		
<b>High-High</b>	T0 Perceived susceptibility				.97	.95 – .99	.002			
	T1 HADS-A	1.37	1.17 – 1.60	.000				1.34	1.08 – 1.67	.008
	Speaking non-English at home	7.11	2.23 – <10	.001				8.06	1.11 ... >10	.039

	ECOG score 0				3.45	1.19 – 9.98	.022	
<b>High-Low</b>	T0 Perceived susceptibility				NA			
	T1 HADS-A	1.34	1.14 – 1.57	.000				1.32 1.05 – 1.65 .017
	Speaking non-English at home				NA			
<b>Low-High</b>	ECOG score 0	.26	.11 – .64	.003				.19 .05 – .75 .018
	T0 Perceived susceptibility				NA			
	T1 HADS-A				.76	.61 – .95	.017	
	Speaking non-English at home				NA			
<b>Low-Low</b>	ECOG score 0				5.23	1.33 – <10	.018	
	T0 Perceived susceptibility				NA			
	T1 HADS-A				.75	.64 – .88	.000	
	Speaking non-English at home				NA			
	ECOG score 0				3.81	1.56 – 9.30	.003	

Abbreviations: ARIA – Accessibility and Remoteness Index of Australia; CGP – comprehensive genomic profiling; CI – confidence interval; ECOG – Eastern Cooperative Oncology Group; HADS – Hospital Anxiety and Depression Scale (Anxiety or Depression sub-scale); HHI – Herth Hope Index; IES – Impact of Events Scale; OR – odds ratio; SE – standard error; SES – socioeconomic status.