

7-1-2014

Erratum to: Race, ethnicity, and the duration of untreated psychosis: a systematic review (Soc Psychiatry Psychiatr Epidemiol, DOI 10.1007/s00127-013-0786-8)

Kelly K. Anderson

Centre for Addiction and Mental Health, kelly.anderson@schulich.uwo.ca

Nina Flora

Centre for Addiction and Mental Health

Suzanne Archie

McMaster University

Craig Morgan

King's College London

Kwame McKenzie

Centre for Addiction and Mental Health

Follow this and additional works at: <https://ir.lib.uwo.ca/paedpub>

Citation of this paper:

Anderson, Kelly K.; Flora, Nina; Archie, Suzanne; Morgan, Craig; and McKenzie, Kwame, "Erratum to: Race, ethnicity, and the duration of untreated psychosis: a systematic review (Soc Psychiatry Psychiatr Epidemiol, DOI 10.1007/s00127-013-0786-8)" (2014). *Paediatrics Publications*. 629.

<https://ir.lib.uwo.ca/paedpub/629>

Erratum to: Race, ethnicity, and the duration of untreated psychosis: a systematic review

Kelly K. Anderson · Nina Flora · Suzanne Archie ·
Craig Morgan · Kwame McKenzie

Published online: 7 December 2013
© Springer-Verlag Berlin Heidelberg 2013

Erratum to: Soc Psychiatry Psychiatr Epidemiol
DOI 10.1007/s00127-013-0786-8

Table 3 was incorrectly formatted during the production process. The corrected version of Table 3 is shown below:

The online version of the original article can be found under
doi:[10.1007/s00127-013-0786-8](https://doi.org/10.1007/s00127-013-0786-8).

K. K. Anderson (✉) · N. Flora · K. McKenzie
Social and Epidemiological Research, Centre for Addiction and
Mental Health (CAMH), 455 Spadina Avenue, Suite 300,
Toronto, ON M5S 2G8, Canada
e-mail: kelly.anderson@camh.ca

S. Archie
Department of Psychiatry and Behavioural Neurosciences,
McMaster University, 25 Charlton Avenue East, Suite 703,
Hamilton, ON L8N 1Y2, Canada

C. Morgan
Section of Social Psychiatry, Institute of Psychiatry, King's
College London, De Crespigny Park, London SE5 8AF, UK

K. McKenzie
Department of Psychiatry, University of Toronto, 455 Spadina
Avenue, Suite 300, Toronto, ON M5S 2G8, Canada

Table 3 Findings on duration of untreated psychosis and racial/ethnic group for studies included in the systematic review ($n = 10$)

Study	Start point for DUP	End point for DUP	DUP tool	Race/ethnicity measurement	Race/ethnicity categories	DUP (weeks)		
						<i>n</i>	Median	IQR
Canada								
Archie et al. [27]	Onset of psychosis	Earlier of date of initiation of antipsychotic medication for at least 1 month, or date of entry to EI services	CORS	Self-report	White ^b Black ^b Asian ^b Other ^b	118 30 25 20	22 31 29 20	3–52 3–57 7–63 3–66
van der Ven et al. [21]	Onset of psychotic symptoms ^a	Initiation of antipsychotic treatment ^a	CORS	Staff assigned based on place of origin	White non-migrants ^b African and Caribbean ^b Asian ^b Central/South America ^b Middle East/North African ^b European/North American ^b	142 36 25 17	15 17 22 11	6–52 5–34 10–56 3–75 8–66
England								
Brunet et al. [31]	One symptom from the positive scale of the PANSS ≥ 4 , or a cluster of symptoms totaling ≥ 7 , for longer than 2 weeks	Antipsychotic treatment at recommended dosage levels for 1 month, or leading to symptom reduction	Pathways Encounter Form	Not described	White Black South Asian Mixed	21 3 26 4	N/A N/A	N/A
Drake [28]	Onset of delusions and hallucinations	Not described	Ad-Hoc Algorithm	Self-report ^b	White ^b African-Caribbean ^b Other ^b	216 19 13	12 9 10	5–36 6–28 6–40
Ghali et al. [20]	One symptom from the positive scale of the PANSS ≥ 4	(1) Initiation of regular antipsychotic treatment with adherence of at least 75 % for 1 month (used in meta-analysis) (2) Date of referral to early intervention services	NOS	Staff assigned	White-British White-other Black-British Black-Caribbean Black-African South Asian	134 84 126 21 110	16 10 14 8 8	49 28 26 34 22
Morgan et al. [15]	Presence of positive psychotic symptoms for at least 1 week	Contact with secondary mental health services	PPHS	Self-report staff assigned	White-British ^b African-Caribbean ^b Black-African ^b	60 217 129 68	9 8 12 8	23 2–39 4–77 1–21

Table 3 continued

Study	Start point for DUP	End point for DUP	DUP tool	Race/ethnicity measurement	Race/ethnicity categories	DUP (weeks)		
						<i>n</i>	IQR	
New Zealand								
Turner et al. [30]	One positive scale symptom on the PANSS ≥ 4	Initiation of antipsychotic treatment	Not described	Not described	Non-Aboriginal ^b Aboriginal (Maori) ^b	135 24	4 9	1–17 2–35
Singapore								
Pek et al. [26]	Onset of delusions, hallucinations, thought disorder, or disorganized behavior	Establishment of definitive diagnosis and treatment	Not described	Not described	Chinese Malay Indian Others	245 64 18 7	N/A	N/A
USA								
Compton et al. [32]	Onset of positive psychotic symptoms	First hospital admission	CORS SOS	Not described	White ^b Black ^b	21 243	20 31	3–119 7–119
Haas et al. [29]	Onset of psychotic symptoms	First administration of antipsychotic medication	Not described	Not described	White Black Asian Hispanic Other	68 24 4 4 3	N/A	N/A

N/A Data not available, *DUP* duration of untreated psychosis, *IQR* interquartile range, *EI* early intervention, *CORS* circumstances of onset or relapse schedule, *PANSS* Positive and Negative Syndrome Scale, *NOS* Nottingham Onset Schedule, *PPHS* Psychiatric and Personal History Schedule, *SOS* Symptom Onset in Schizophrenia Inventory

^a As per prior paper [39]

^b Data obtained from corresponding author