

On the Hodge conjecture for quasi-smooth intersections in toric varieties

UGO BRUZZO[‡] AND WILLIAM D. MONTOYA[¶]

[‡] SISSA (Scuola Internazionale Superiore di Studi Avanzati),
Via Bonomea 265, 34136 Trieste, Italy;

Departamento de Matemática, Universidad Federal da Paraíba,
Campus I, Jardim Universitário, 58051-900, João Pessoa, PB, Brazil;

INFN (Istituto Nazionale di Fisica Nucleare), Sezione di Trieste;

IGAP (Institute for Geometry and Physics), Trieste;

Arnold-Regge Center for Algebra, Geometry
and Theoretical Physics, Torino;

[¶] Instituto de Matemática, Estatística e Computação Científica,
Universidade Estadual de Campinas, Rua Sérgio Buarque de Holanda 651,
13083-859, Campinas, SP, Brazil

Abstract

We establish the Hodge conjecture for some subvarieties of a class of toric varieties. First we study quasi-smooth intersections in a projective simplicial toric variety, which is a suitable notion to generalize smooth complete intersection subvarieties in the toric environment, and in particular quasi-smooth hypersurfaces. We show that under appropriate conditions, the Hodge Conjecture holds for a very general quasi-smooth intersection subvariety, generalizing the work on quasi-smooth hypersurfaces of the first author and Grassi in [3]. We also show that the Hodge Conjecture holds asymptotically for suitable quasi-smooth hypersurface in the Noether-Lefschetz locus, where “asymptotically” means that the degree of the hypersurface is big enough, under the assumption that the ambient variety $\mathbb{P}_{\Sigma}^{2k+1}$ has Picard group \mathbb{Z} . This extends to a class of toric varieties Otwinowska’s result in [18].

Date: Revised May 25, 2021

2010 Mathematics Subject Classification: 14C22, 14J70, 14M25

Keywords: Noether-Lefschetz theory, Hodge Conjecture, toric varieties

Email: bruzzo@sisssa.it, montoya@unicamp.br

1 Introduction

A projective simplicial toric variety \mathbb{P}_Σ^d satisfies the Hodge Conjecture, i.e., every cohomology class in $H^{p,p}(\mathbb{P}_\Sigma^d, \mathbb{Q})$ is a linear combination of algebraic cycles. On the one hand, by the Lefschetz hyperplane theorem, the Hodge conjecture holds true for every hypersurface and $p < \frac{d-1}{2}$ and by the hard Lefschetz theorem also for $p > \frac{d-1}{2}$. Moreover, by Theorem 1.1 in [3], when $p = \frac{d-1}{2}$, $d = 2k + 1$ and \mathbb{P}_Σ^{2k+1} is an Oda variety with an ample class β such that $k\beta - \beta_0$ is nef, where β_0 is the anticanonical class, the Hodge conjecture with rational coefficients holds for a very general hypersurface in the linear system $|\beta|$.

The notion of Oda varieties was introduced in [2]. Let us recall that the Cox ring of a toric variety \mathbb{P}_Σ is graded over the class group $\text{Cl}(\mathbb{P}_\Sigma)$, and that one has an injection $\text{Pic}(\mathbb{P}_\Sigma) \rightarrow \text{Cl}(\mathbb{P}_\Sigma)$.

Definition 1.1. *Let \mathbb{P}_Σ be a toric variety with Cox ring S . \mathbb{P}_Σ is said to be an Oda variety if the multiplication morphism $S_{\alpha_1} \otimes S_{\alpha_2} \rightarrow S_{\alpha_1 + \alpha_2}$ is surjective whenever the classes α_1 and α_2 in $\text{Pic}(\mathbb{P}_\Sigma)$ are ample and nef, respectively.*

In [17] Mavlyutov proved a Lefschetz type theorem for quasi-smooth intersection subvarieties, and moreover using the ‘‘Cayley trick’’ he related the cohomology of a quasi-smooth subvariety $X = X_{f_1} \cap \dots \cap X_{f_s} \subset \mathbb{P}_\Sigma^d$ to the cohomology of a quasi-smooth hypersurface $Y \subset \mathbb{P}_\Sigma^{d+s-1}$. This allows us to prove a Noether-Lefschetz type theorem, namely:

Theorem 2.5. *Let \mathbb{P}_Σ^d be an Oda projective simplicial toric variety. For a very general quasi-smooth intersection subvariety X cut off by f_1, \dots, f_s such that $d + s = 2(\ell + 1)$ and*

$$\sum_{i=1}^s \deg(f_i) - \beta_0$$

is nef, one has

$$H^{\ell+1-s, \ell+1-s}(X, \mathbb{Q}) = i^* \left(H^{\ell+1-s, \ell+1-s}(\mathbb{P}_\Sigma^d, \mathbb{Q}) \right).$$

From this one obtains the following result about the Hodge conjecture for quasi-smooth intersections.

Corollary 2.7. *If \mathbb{P}_Σ^d is an Oda projective simplicial toric variety, the Hodge Conjecture holds for a very general quasi-smooth intersection subvariety X cut off by f_1, \dots, f_s such that $d + s$ is even and $\sum_{i=1}^s \deg(f_i) - \beta_0$ is nef.*

Let T be the open subset of $|\beta|$ corresponding to quasi-smooth hypersurfaces, and let $\mathcal{H}^{2k} = R^{2k}\pi_*\mathbb{C} \otimes_{\mathbb{C}} \mathcal{O}_T$ be the Hodge bundle on T ; here $\pi: \mathcal{X} \rightarrow T$ is the tautological family

on T , and $d = 2k + 1$. We restrict \mathcal{H}^{2k} to a contractible open subset $U \subset T$. The bundle \mathcal{H}^{2k} has a Hodge decomposition

$$\mathcal{H}^{2k} = \bigoplus_{p+q=2k} \mathcal{H}^{p,q}$$

but this is not holomorphic. On the other hand, the bundles that make up the Hodge filtration

$$F^p \mathcal{H}^{2k} = \bigoplus_{p=0}^{2k} \mathcal{H}^{2k-p,p}$$

are holomorphic; to see this one can use the *period map* (which in particular we write for $p = k$)

$$\mathcal{P}^{k,2k}: U \rightarrow \text{Grass}(b_k, H^{2k}(X_{u_0}, \mathbb{C}))$$

where $b_k = \dim F^k H^{2k}(X_{u_0}, \mathbb{C})$ for a fixed point $u_0 \in U$; this map sends $f \in U$ to the subspace $F^k H^{2k}(X_f, \mathbb{C}) \subset H^{2k}(X_f, \mathbb{C}) = H^{2k}(X_{u_0}, \mathbb{C})$. This map is holomorphic (see [15] and [5, Prop. 3.4]). But, by the very definition of the period map (see also [19], Section 10.2.1 for the smooth case)

$$F^k \mathcal{H}^{2k} \simeq (\mathcal{P}^{k,2k})^* \mathcal{U}_k,$$

where \mathcal{U}_k is the tautological bundle on the Grassmannian $\text{Grass}(b_k, H^{2k}(X_{u_0}, \mathbb{C}))$, so that the bundles $F^k \mathcal{H}^{2k}$ are indeed holomorphic.

Pushing ahead the ideas developed in [5] and [4], let λ_f be a nonzero class in the primitive cohomology $H^{k,k}(X_f, \mathbb{Q})/H^{k,k}(\mathbb{P}_{\Sigma}^{2k+1}, \mathbb{Q})$, and let U be a contractible open subset of T around f , so that $\mathcal{H}_{|U}^{2k}$ is constant. Moreover, let $\lambda \in \mathcal{H}^{2k}(U)$ be the section defined by λ_f and let $\bar{\lambda}$ be its image in $(\mathcal{H}^{2k}/F^k \mathcal{H}^{2k})(U)$. One has

Proposition 1.2. *The local Noether-Lefschetz loci can be defined as*

$$N_{\lambda,U}^{k,\beta} := \{G \in U \mid \bar{\lambda}_G = 0\}$$

where $\beta = \deg(f)$.

The following result is Theorem 1.2 in [4].

Theorem. *Let $\mathbb{P}_{\Sigma}^{2k+1}$ be an Oda variety with an ample class β such that $k\beta - \beta_0 = n\eta$, where β_0 is the anticanonical class, η is a primitive ample class, and $n \in \mathbb{N}$. Let*

$$m_{\beta} = \max\{i \in \mathbb{N} \mid i\eta \leq \beta\}. \quad (1)$$

For every positive ϵ there is a positive δ such that for every $m \geq \max(\frac{1}{\delta}, m_{\beta})$ and $d \in [1, m\delta]$, and every nontrivial Hodge class $\lambda \in F^k \mathcal{H}^{2k}(U)$ such that

$$\text{codim } N_{\lambda,U}^{k,\beta} \leq d \frac{m_{\beta}^k}{k!},$$

for every $f \in N_{\lambda, U}^{k, \beta}$, there exists a k -dimensional variety $V \subset X_f$ with $\deg V \leq (1 + \epsilon)d$. Here $\deg V$ is taken with respect to the ample divisor η , i.e.,

$$\deg V = [V] \cdot \eta^k.$$

Based on this, in this paper we obtain the following result.

Theorem 4.2. *Under the same hypotheses of the previous theorem, assume also that $\text{Pic}(\mathbb{P}_{\Sigma}^{2k+1}) = \mathbb{Z}$. Then, if $V \subset X_f$ is a nonempty quasi-smooth intersection subvariety of $\mathbb{P}_{\Sigma}^{2k+1}$ for some $f \in N_{\lambda, U}^{k, \beta}$, there exists $c \in \mathbb{Q}^*$ such that $\lambda_f = c\lambda_V$, where λ_V is the class of V in $H_{\text{prim}}^{k, k}(X_f, \mathbb{Q})$.*

In other words, λ_f is algebraic.

In his paper [11] A. Dan proves a form of our Theorem 4.2 for smooth hypersurfaces in odd-dimensional projective spaces \mathbb{P}^{2k+1} which is not asymptotic. So our result is more general in two ways, as we consider *quasi-smooth intersections* in *toric varieties* with $h^{k, k} = 1$ (for instance, weighted or fake projective spaces); however, our result is asymptotic.

In Section 3 we give an extension of the notion of Gorenstein ideal to Cox rings; this may have some interest on its own.

Acknowledgements. We thank Paolo Aluffi for useful discussions, and Antonella Grassi for developing with the first author the foundations on which this work is based. We are very thankful to the referee for her/his very careful reading, and the many suggestions and remarks which allowed us to greatly improve the presentation of this paper. The first author's research is partly supported by PRIN "Geometry of algebraic varieties" and GNSAGA-INdAM. The second author acknowledges support from FAPESP postdoctoral grant No. 2019/23499-7.

2 Very general quasi-smooth intersections

Let f_1, \dots, f_s be weighted homogeneous polynomials in the Cox ring $S = \mathbb{C}[x_1, \dots, x_n]$ of \mathbb{P}_{Σ}^d . Their zero locus $V(f_1, \dots, f_s)$ defines a closed subvariety $X \subset \mathbb{P}_{\Sigma}^d$. Let $U(\Sigma) = \mathbb{A}^n - Z(\Sigma)$, where $Z(\Sigma)$ is the irrelevant locus, i.e., $Z(\Sigma) = \text{Spec } B$, where B is the irrelevant ideal.

Definition 2.1. [17] *X is a codimension s quasi-smooth intersection if $V(f_1, \dots, f_s) \cap U(\Sigma)$ is either empty or a smooth intersection subvariety of codimension s in $U(\Sigma)$.*

This notion generalizes that of smooth complete intersection in a projective space. For $s = 1$ it reduces to the notion of *quasi-smooth hypersurface*, see Def. 3.1 in [1]. If we regard \mathbb{P}_Σ^d as an orbifold, then a hypersurface X is quasi-smooth when it is a sub-orbifold of \mathbb{P}_Σ^d ; heuristically, “ X has only singularities coming from the ambient variety.”

We also have a Lefschetz type theorem in this context.

Proposition 2.2 ([17] Proposition 1.4). *Let $X \subset \mathbb{P}_\Sigma^d$ be a closed subset, defined by homogeneous polynomials $f_1, \dots, f_s \in B$. Then the natural map $i^* : H^i(\mathbb{P}_\Sigma^d) \rightarrow H^i(X)$ is an isomorphism for $i < d - s$ and an injection for $i = d - s$. In particular, this is true if the hypersurfaces cut by the polynomials f_i are ample.*

Hence if $p \neq \frac{d-s}{2}$ every cohomology class in $H^{p,p}(X)$ is a linear combination of algebraic cycles. So let us see what happens when $p = \frac{d-s}{2}$. The idea is to relate the Hodge structure of a quasi-smooth intersection variety $X = X_{f_1} \cap \dots \cap X_{f_s}$ in \mathbb{P}_Σ^d with the Hodge structure of a quasi-smooth hypersurface Y in a toric variety $\mathbb{P}_{X,\Sigma}^{d+s-1}$ whose fan depends on X and Σ .

Proposition 2.3. *Let $X = X_{f_1} \cap \dots \cap X_{f_s}$ be quasi-smooth intersection subvariety in \mathbb{P}_Σ^d cut off by homogeneous polynomials $f_1 \dots f_s$. There exists a projective simplicial toric variety $\mathbb{P}_{X,\Sigma}^{d+s-1}$ and a quasi-smooth hypersurface $Y \subset \mathbb{P}_{X,\Sigma}^{d+s-1}$ such that for $p \neq \frac{d+s-1}{2}, \frac{d+s-3}{2}$*

$$H_{\text{prim}}^{p-1, d+s-1-p}(Y) \simeq H_{\text{prim}}^{p-s, d-p}(X).$$

Proof. One constructs $\mathbb{P}_{X,\Sigma}^{d+s-1}$ via the so-called “Cayley trick”. Let L_1, \dots, L_s be the line bundles associated to the quasi-smooth hypersurfaces X_1, \dots, X_s , and so let $\mathbb{P}(E)$ be the projective bundle of $E = L_1 \oplus \dots \oplus L_s$. It turns out that $\mathbb{P}(E)$ is a $d + s - 1$ - dimensional projective simplicial toric variety whose Cox ring is

$$\mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_s]$$

where $S = \mathbb{C}[x_1, \dots, x_n]$ is the Cox ring of \mathbb{P}_Σ^d . The hypersurface Y is cut off by the polynomial $F = y_1 f_1 + \dots + y_s f_s$ and is quasi-smooth by Lemma 2.2 in [17]. Moreover, combining Theorem 10.13 in [1] and Theorem 3.6 in [17], we have that

$$H_{\text{prim}}^{p-1, d+s-1-p}(Y) \simeq R(F)_{(d+s-p)\beta - \beta_0} \simeq H_{\text{prim}}^{p-s, d-p}(X)$$

for $p \neq \frac{d+s-1}{2}, \frac{d+s-3}{2}$ as desired. \square

Here $R(F)$ is the Jacobian ring of Y , i.e., the quotient of the Cox ring

$$R(F) = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_s] / J(F),$$

where $J(F)$ is the ideal generated by the derivatives of F , see [1].

Remark 2.4. With the same notation of Proposition 2.3, note that we have a well defined map

$$\begin{aligned}\phi: |\beta_1| \times \cdots \times |\beta_s| &\rightarrow |\beta| \\ (f_1, \dots, f_s) &\mapsto f_1 y_1 + \cdots + f_s y_s.\end{aligned}$$

Moreover, by the Noether-Lefschetz theorem NL_β is a countable union of closed sets $\cup_i C_i$ and hence $\cup \phi^{-1}(C_i)$ is too. \triangle

We have a Noether-Lefschetz type theorem, namely,

Theorem 2.5. *Let \mathbb{P}_Σ^d be an Oda projective simplicial toric variety. Then for a very general quasi-smooth intersection subvariety X cut off by f_1, \dots, f_s such that $d + s = 2(k + 1)$ and $\sum_{i=1}^s \deg(f_i) - \beta_0$ is nef, one has that*

$$H^{k+1-s, k+1-s}(X, \mathbb{Q}) = i^* (H^{k+1-s, k+1-s}(\mathbb{P}_\Sigma^d, \mathbb{Q}))$$

So we get a natural generalization of the Noether-Lefschetz loci.

Definition 2.6. *The Noether-Lefschetz locus $NL_{\beta_1, \dots, \beta_s}$ of quasi-smooth intersection varieties is the locus of s -tuples (f_1, \dots, f_s) such that $X = X_{f_1} \cap \dots \cap X_{f_s}$ is quasi-smooth intersection with $f_i \in |\beta_i|$ and $H^{k+1-s, k+1-s}(X, \mathbb{Q}) \neq i^* (H^{k+1-s, k+1-s}(\mathbb{P}_\Sigma^d, \mathbb{Q}))$.*

Now we consider the Hodge conjecture for very general quasi-smooth intersection subvarieties in \mathbb{P}_Σ^d .

Corollary 2.7. *If \mathbb{P}_Σ^d is a Oda projective simplicial toric variety, the Hodge Conjecture holds for a very general quasi-smooth intersection subvariety X cut off by f_1, \dots, f_s such that $d + s = 2(k + 1)$ and $\sum_{i=1}^s \deg(f_i) - \beta_0$ is nef.*

Proof. First note that by Theorem 4.1 in [13] the projective simplicial toric variety \mathbb{P}_Σ^{2k+1} is Oda and since X is very general the quasi-smooth hypersurface Y is very general as well. So applying the Noether-Lefschetz theorem one has that $h_{\text{prim}}^{k, k}(Y) = 0 = h_{\text{prim}}^{k+1-s, k+1-s}(X)$ or equivalently every $(k + 1 - s, k + 1 - s)$ cohomology class is a linear combination of algebraic cycles. \square

3 Cox-Gorenstein ideals

We shall need a partial generalization of Macaulay's theorem (see e.g. Thm. 6.19 in [20] for the classical theorem). This generalization is basically contained in the work of Cox and Cattani-Cox-Dickenstein [9, 7].

Let S be the Cox ring of a complete simplicial toric variety \mathbb{P}_Σ . This is graded over the effective classes in the class group $\text{Cl}(\mathbb{P}_\Sigma)$ and [8]

$$S^\alpha \simeq H^0(\mathbb{P}_\Sigma, \mathcal{O}_{\mathbb{P}_\Sigma}(\alpha)).$$

As $\mathcal{O}_{\mathbb{P}_\Sigma}(\alpha)$ is coherent and \mathbb{P}_Σ is complete, each S^α is finite-dimensional over \mathbb{C} ; in particular, $S^0 \simeq \mathbb{C}$.

Lemma 3.1. *For every effective $N \in \text{Cl}(\mathbb{P}_\Sigma)$, the set of classes $\alpha \in \text{Cl}(\mathbb{P}_\Sigma)$ such that $N - \alpha$ is effective is finite.*

Proof. Since the torsion submodule of $\text{Cl}(\mathbb{P}_\Sigma)$ is finite, we may assume that $\text{Cl}(\mathbb{P}_\Sigma)$ is free. Then the exact sequence

$$0 \rightarrow M \rightarrow \text{Div}_{\mathbb{T}}(\mathbb{P}_\Sigma) \rightarrow \text{Cl}(\mathbb{P}_\Sigma) \rightarrow 0$$

splits, and we may identify $\text{Cl}(\mathbb{P}_\Sigma)$ with a free subgroup of $\text{Div}_{\mathbb{T}}(\mathbb{P}_\Sigma)$, generated by a subset $\{D_1, \dots, D_r\}$ of \mathbb{T} -invariant divisors. A class in $\text{Cl}(\mathbb{P}_\Sigma)$ is effective if and only its coefficients on this basis are nonnegative, whence the claim follows. \square

We shall give a definition of *Cox-Gorenstein ideal* of the Cox rings which generalizes to toric varieties the definition given by Otwinowska in [18] for projective spaces. Let $B \subset S$ be the irrelevant ideal, and for a graded ideal $I \subset B$, denote by $V_{\mathbb{T}}(I)$ the corresponding closed subscheme of \mathbb{P}_Σ .

Definition 3.2. *A graded ideal I of S contained in B is said to be a Cox-Gorenstein ideal of socle degree $N \in \text{Cl}(\mathbb{P}_\Sigma)$ if*

1. *there exists a \mathbb{C} -linear form $\Lambda \in (S^N)^\vee$ such that for all $\alpha \in \text{Cl}(\mathbb{P}_\Sigma)$*

$$I^\alpha = \{f \in S^\alpha \mid \Lambda(fg) = 0 \text{ for all } g \in S^{N-\alpha}\}; \tag{2}$$

2. $V_{\mathbb{T}}(I) = \emptyset$.

Remark 3.3. Cox-Gorenstein ideals need not be Artinian. Property 2 in this definition replaces that condition. \triangle

Proposition 3.4. *Let $R = S/I$. If I is Cox-Gorenstein then*

1. $\dim_{\mathbb{C}} R^N = 1$;

2. the natural bilinear morphism

$$R^\alpha \times R^{N-\alpha} \rightarrow R^N \simeq \mathbb{C} \quad (3)$$

is nondegenerate whenever α and $N - \alpha$ are effective.

Proof. 1. From eq. (2) we see that the sequence

$$0 \rightarrow I^N \rightarrow S^N \xrightarrow{\Lambda} \mathbb{C} \rightarrow 0$$

is exact.

2. Define $\Phi: R^\alpha \times R^{N-\alpha} \rightarrow \mathbb{C}$ as $\Phi(x, y) = \Lambda(\bar{x}\bar{y})$, where \bar{x}, \bar{y} are pre-images of x, y in S . One easily checks that this is well defined and that via the isomorphism $R^N \simeq k$ it coincides with the pairing (3). Now if $x \in R^\alpha$ and $\Phi(x, y) = 0$ for all $y \in R^{N-\alpha}$ then $\Lambda(\bar{x}\bar{y}) = 0$ for all $\bar{y} \in S^{N-\alpha}$ so that $\bar{x} \in I^\alpha$, i.e., $x = 0$. \square

Let f_0, \dots, f_d be homogeneous polynomials, $f_i \in S^{\alpha_i}$, where $d = \dim \mathbb{P}_\Sigma$ and each α_i is ample, and let $N = \sum_i \alpha_i - \beta_0$, where β_0 is the anticanonical class of \mathbb{P}_Σ . Assume that the f_i have no common zeroes in \mathbb{P}_Σ , i.e., $V_{\mathbb{T}}(I) = \emptyset$ if $I = (f_0, \dots, f_d)$.

In [1, 9, 7] it is shown that for each $G \in S^N$ one can define a meromorphic d -form ξ_G on \mathbb{P}_Σ by letting

$$\xi_G = \frac{G \Omega}{f_0 \cdots f_d}$$

where Ω is a Euler form on \mathbb{P}_Σ . The form ξ_G determines a class in $H^d(\mathbb{P}_\Sigma, \omega)$, where ω is the canonical sheaf of \mathbb{P}_Σ (the sheaf of Zariski d -forms on \mathbb{P}_Σ), and in turn the trace morphism $\text{Tr}_{\mathbb{P}_\Sigma}: H^d(\mathbb{P}_\Sigma, \omega) \rightarrow \mathbb{C}$ associates a complex number to G , so we can define $\Lambda \in (S^N)^\vee$ as

$$\Lambda(G) = \text{Tr}_{\mathbb{P}_\Sigma}([\xi_G]) \in \mathbb{C}. \quad (4)$$

Finally, we can prove a toric version of Macaulay's theorem.

Theorem 3.5. *The linear map defined in Eq. (4) satisfies the condition in Definition 3.2. Therefore, the ideal $I = (f_0, \dots, f_d)$ is a Cox-Gorenstein ideal of socle degree N .*

Proof. By Theorem 6 in [7] the map Λ establishes an isomorphism $R^N \simeq \mathbb{C}$. Hence, if $f \in S^\alpha$ is such that $\Lambda(fg) = 0$ for all $g \in S^{N-\alpha}$, then $fg \in I^N$, which implies $f \in I^\alpha$. On the other hand, it is clear that $\Lambda(fg) = 0$ if $f \in I^\alpha$ and $g \in S^{N-\alpha}$. \square

Another example is given in terms of *toric Jacobian ideals*. For every ray $\rho \in \Sigma(1)$ we shall denote by v_ρ its rational generator, and by x_ρ the corresponding variable in the Cox ring. Recall that d is the dimension of the toric variety \mathbb{P}_Σ , while we denote by $r = \#\Sigma(1)$ the number of rays. Given $f \in S^\beta$ one defines its *toric Jacobian ideal* as

$$J_0(f) = \left(x_{\rho_1} \frac{\partial f}{\partial x_{\rho_1}}, \dots, x_{\rho_r} \frac{\partial f}{\partial x_{\rho_r}} \right).$$

We recall from [1] the definition of nondegenerate hypersurface and some properties (Def. 4.13 and Prop. 4.15).

Definition 3.6. *Let $f \in S(\Sigma)^\beta$, with β an ample Cartier class. The associated hypersurface X_f is nondegenerate if for all $\sigma \in \Sigma$ the affine hypersurface $X_f \cap O(\sigma)$ is a smooth codimension one subvariety of the orbit $O(\sigma)$ of the action of the torus \mathbb{T}^d .*

Proposition 3.7. *1. Every nondegenerate hypersurface is quasi-smooth.
2. If f is generic then X_f is nondegenerate.*

The following is part of Prop. 5.3 in [9], with some changes in the terminology.

Proposition 3.8. *Let $f \in S(\Sigma)^\beta$, and let $\{\rho_1, \dots, \rho_d\} \subset \Sigma(1)$ be such that $v_{\rho_1}, \dots, v_{\rho_d}$ are linearly independent.*

1. *The toric Jacobian ideal of f coincides with the ideal*

$$\left(f, x_{\rho_1} \frac{\partial f}{\partial x_{\rho_1}}, \dots, x_{\rho_d} \frac{\partial f}{\partial x_{\rho_d}} \right).$$

2. *The following conditions are equivalent:*

- (a) *f is nondegenerate;*
- (b) *the polynomials $x_{\rho_i} \frac{\partial f}{\partial x_{\rho_i}}$, $i = 1, \dots, r$, do not vanish simultaneously on X_f ;*
- (c) *the polynomials f and $x_{\rho_i} \frac{\partial f}{\partial x_{\rho_i}}$, $i = 1, \dots, d$, do not vanish simultaneously on X_f .*

3. *If moreover β is ample and f is nondegenerate, then $J_0(f)$ is a Cox-Gorenstein ideal of socle degree $N = (d+1)\beta - \beta_0$, where β_0 is the anticanonical class of \mathbb{P}_Σ^d .*

4 Asymptotic Hodge conjecture

In this section we prove Theorem 4.2. Let us recall part of the notation and assumptions of [4]. Let $\mathbb{P}_{\Sigma}^{2k+1}$ be an Oda variety with an ample Cartier class β such that $k\beta - \beta_0 = n\eta$, where β_0 is the anticanonical class, η is a primitive ample class and $n \in \mathbb{N}$.

We need to define a pre-order in the group

$$N^1(\mathbb{P}_{\Sigma}^{2k+1}) = \text{Pic}(\mathbb{P}_{\Sigma}^{2k+1}) \otimes \mathbb{Q}/\text{numerical equivalence},$$

by letting $\alpha < \alpha'$ if $\alpha' - \alpha$ is an effective class.

Let $X_f \in |\beta|$ be a quasi-smooth hypersurface in the Noether-Lefschetz locus associated to a nontrivial Hodge class $\lambda \in F^k \mathcal{H}^{2k}(U)$. Again, its degree is computed by intersecting with the ample class η , i.e., $\deg X_f = [X_f] \cdot \eta$. Let r be number of rays of Σ , so that $r \geq 2(k+1)$. Assuming that n is big enough, it follows from Proposition 4.7 or Theorem 6.1 in [4] that there exists a k -dimensional subvariety V of X_f satisfying the following conditions:

- $\deg V \leq 2\delta m_{\beta}$ with $0 < \delta < \frac{1}{4(r-(k+1))}$ (the number m_{β} was defined in Eq. (1));
- the graded ideals I_V and

$$E = \left\{ g \in S^{\bullet} \mid \sum_{i=1}^b \lambda_i \int_{\text{Tub } \gamma_i} \frac{gh\Omega_0}{f^{k+1}} = 0 \text{ for all } h \in S^{N-\bullet} \right\}, \quad (5)$$

coincide in degree less than or equal to $(m_{\beta} - 2 - (r-j) \deg V) \eta$ for some j , with $0 < j < r$. Here $\text{Tub}(-)$ is the adjoint of the residue map, and $N = (k+1)\beta - \beta_0$ is the socle degree of the *Cox-Gorenstein ideal* E , while

$$\lambda_f = \left(\sum_{i=1}^b \lambda_i \gamma_i \right)^{pd}$$

is the Poincaré dual of some rational combination of the homology cycles γ_i generating $H_{2k}(X_f, \mathbb{Q})$. Moreover, via the isomorphism $T_f U \simeq S^{\beta}$, the degree β summand E^{β} of E is identified with the tangent space $T_f N_{\lambda, U}^{k, \beta}$ to the Noether-Lefschetz locus, so that E^{β} contains the degree β part $J(f)^{\beta}$ the Jacobian ideal of f .

Lemma 4.1. *The toric Jacobian ideal $J_0(f)$ is contained in E .*

Proof. $J_0(f) \subset J(f)$, so that $J_0(f)^{\beta} \subset J(f)^{\beta} \subset E^{\beta}$, and since $J_0(f)$ is generated in degree β , one has $J_0(f) \subset E$. \square

We denote by λ_V the class of V in $H_{\text{prim}}^{k,k}(X_f, \mathbb{Q})$. In the following theorem we assume that $\text{Pic}(\mathbb{P}_{\Sigma}^{2k+1}) = 1$, i.e., that $\mathbb{P}_{\Sigma}^{2k+1}$ is a (possibly fake) weighted projective space [6, 14] (cf. [10] Exer. 5.1.13). This implies that $h^{p,p}(\mathbb{P}_{\Sigma}^{2k+1}) = 1$ for all p .

Theorem 4.2. *If V is a smooth intersection subvariety, there exists $c \in \mathbb{Q}^*$ such that $\lambda_f = c\lambda_V$.*

Proof. We divide the proof in three steps.

Step I: $\lambda_V \neq 0$. For clarity, for every cohomology class of a subvariety we denote in the cohomology of which ambient variety we consider it (so we write $[V]_{X_f}$ and $[V]_{\mathbb{P}_{\Sigma}^{2k+1}}$). Since $V \subset X_f$ is a regular embedding we have

$$\begin{aligned} [V]_{X_f}^2 &= \int_V c_k(N_{V/X_f}) = \int_V \left[c(N_{V/\mathbb{P}_{\Sigma}^{2k+1}}) / c(N_{X_f/\mathbb{P}_{\Sigma}^{2k+1}|_V}) \right]_k \\ &= \int_{\mathbb{P}_{\Sigma}^{2k+1}} [V]_{\mathbb{P}_{\Sigma}^{2k+1}} \cup \Xi_k \end{aligned} \quad (6)$$

where Ξ_k is the component in $H^{k,k}(\mathbb{P}_{\Sigma}^{2k+1})$ of

$$\Xi = \frac{\prod_i (1 + A_i)}{1 + [X_f]_{\mathbb{P}_{\Sigma}^{2k+1}}};$$

here A_1, \dots, A_{k+1} are the classes in $\text{Cl}(\mathbb{P}_{\Sigma}^{2k+1})$ of the hypersurfaces that cut the smooth intersection subvariety V . The claim is proved by contradiction: if $[V]_{X_f}$ is the restriction of a class in $H^{k,k}(\mathbb{P}_{\Sigma}^{2k+1})$, i.e.,

$$[V]_{X_f} = b\eta|_{X_f}^k$$

for some b , then comparing this with (6) we obtain

$$\deg V = m_k \deg X_f, \quad (7)$$

where m_k is defined by $\Xi_k = m_k \eta^k$. But (7) cannot be true when $\deg X_f$ is big enough.

Step II. Let E_{alg} and E be the Cox-Gorenstein ideals associated to λ_V and λ_f , respectively, as in equation (5). To prove the theorem it is enough to show that $E = E_{\text{alg}}$. Note that $I_V + J_0(f)$ is contained in E and E_{alg} . Moreover, since $V \subset X_f$, and f is quasi-smooth, there exist $K_1, \dots, K_{k+1} \in B$ such that $f = A_1 K_1 + \dots + A_{k+1} K_{k+1}$ and $(A_1, \dots, A_{k+1}, K_1, \dots, K_{k+1})$ is a Cox-Gorenstein ideal with socle degree N ; this will follow from the next step, which concludes the proof.

Step III. It is enough to show that every Cox-Gorenstein ideal \mathcal{I} of socle degree N containing $I_V + J_0(f)$ also contains $(A_1, \dots, A_{k+1}, K_1, \dots, K_{k+1})$. By assumption

$$\left(A_j, j \in \{1, \dots, k+1\}, \sum_{j=1}^{k+1} x_i \frac{\partial A_j}{\partial x_i} K_j, i \in 1, \dots, r \right) \subset \mathcal{I}.$$

Let us see that $K_j \in \mathcal{I}$ for every $j \in \{1, \dots, k+1\}$. Let $M \in \text{Mat}(r \times (k+1))$ be the matrix $[x_i \frac{\partial A_j}{\partial x_i}]$ and K the column $(K_j)_{j \in \{1, \dots, k+1\}}$. Let $I \subset \{1, \dots, r\}$ with cardinality $k+1$ and let M_I be the matrix obtained extracting the $i \in I$ -arrows of M . We have that $\sum_{j=1}^{k+1} x_i \frac{\partial A_j}{\partial x_i} K_j = (MK)_i = (M_I K)_i$; multiplying by the adjoint of M_I we get that $\det(M_I)K_j \in \mathcal{I}$ for all $j \in \{1, \dots, k+1\}$. On one hand the ideal $(\mathcal{I} : K_j)$ contains the ideal

$$\mathcal{J} = I_V + \langle \det M_I \mid I \subset \{1, \dots, r\}, \#I = k+1 \rangle.$$

Since V is a smooth complete intersection subvariety, it follows that \mathcal{J} is base point free, and therefore it contains a complete intersection Cox-Gorenstein ideal \mathcal{J}' by the toric Macaulay theorem, Theorem 3.5. Since \mathcal{J} is generated in degree less than or equal to $(\deg V)\eta$, we can take \mathcal{J}' with the same property. It follows that

$$\text{soc}(\mathcal{J}') \leq 2(k+1)(\deg V)\eta - \beta_0 \leq 2rm_\beta\delta\eta - \beta_0.$$

On the other hand if $K_j \notin \mathcal{I}$ then $(\mathcal{I} : K_j)$ contains a Cox-Gorenstein ideal with socle degree

$$N - \deg K_j \geq N - \beta = k\beta - \beta_0;$$

then comparing the above two inequalities and keeping in mind that $r \geq 2(k+1)$, we get

$$\delta \geq \frac{1}{2r} \geq \frac{1}{4(r - (k+1))},$$

which is absurd. □

References

- [1] D. A. Batyrev, V. V. Cox, *On the Hodge structure of projective hypersurfaces in toric varieties*, Duke Math. J. **75** (1994) 293–338. [5](#), [8](#), [9](#)
- [2] U. Bruzzo, A. Grassi, *The Noether-Lefschetz locus of surfaces in toric threefolds*, Commun. Contemp. Math. No. 5 (2017) 1750070 (20 pages). [2](#)
- [3] —, *On the Hodge conjecture for hypersurfaces in toric varieties*, Commun. Anal. Geom. **28** (2020) 1773–1786. [1](#), [2](#)
- [4] U. Bruzzo, W.D. Montoya, *An asymptotic description of the Noether-Lefschetz components in toric varieties*, arXiv:1905.01570 [math.AG]. [3](#), [10](#)

- [5] —, *Codimension bounds for the Noether-Lefschetz components for toric varieties*, arXiv:2001.01960 [math.AG]. To appear in Eur. J. Math. [3](#)
- [6] W. Buczyńska, *Fake weighted projective spaces*, English translation of *Toryczne przestrzenie rzutowe*, Magister thesis, 2002. arXiv:0805.1211 [math.AG]. [11](#)
- [7] E. Cattani, D. Cox, A. Dickenstein, *Residues in toric varieties*, Compositio Math. **108** (1997) 35–76. [6](#), [8](#)
- [8] D. A. Cox, *The homogeneous coordinate ring of a toric variety*, **4** (1995) 17–50. [7](#)
- [9] —, *Toric residues*, Ark. Mat. **75** (1996) 73–96. [6](#), [8](#), [9](#)
- [10] D. Cox, J. Little, H. Schenck, *Toric varieties*, Graduate Studies in Mathematics, 124. American Mathematical Society, Providence, RI, 2011. xxiv+841 pp. [11](#)
- [11] A. Dan, *Noether-Lefschetz locus and a special case of the variational Hodge conjecture: using elementary techniques*, in “Analytic and algebraic geometry,” Hindustan Book Agency, New Delhi 2017, pp. 107–115. [4](#)
- [12] I. Dolgachev, *Weighted projective varieties*, in Lecture Notes in Math. **956**, Springer-Verlag, Berlin (1982), pp. 34–71.
- [13] A. Ikeda, *Subvarieties of generic hypersurfaces in a non-singular projective toric variety*, Math. Z. **263** (2009) 923–937. [6](#)
- [14] T. Kawasaki, *Cohomology of twisted projective spaces and lens complexes*, Math. Ann. **206** (1973) 243–248. [11](#)
- [15] K. F. Liu, X. B. Zhuang, *Deformations of complex orbifolds and the period maps*, Sci. China Math. **63** (2020) 83–100. [3](#)
- [16] T. Maeno, J. Watanabe, *Lefschetz elements of Artinian Gorenstein algebras and Hessians of homogeneous polynomials*, Illinois J. Math. **53** (2009) 591–603.
- [17] A. Mavlyutov, *Cohomology of Complete Intersections In Toric Varieties*, Pacific J. Math. **191** (1999) 133–144. [2](#), [4](#), [5](#)
- [18] A. Otwinowska, *Composantes de petite codimension du lieu de Noether-Lefschetz: un argument asymptotique en faveur de la conjecture de Hodge*, J. Alg. Geom. **12** (2003) 307–320. [1](#), [7](#)
- [19] C. Voisin, *Hodge Theory and Complex Algebraic Geometry I*, Cambridge University Press, Cambridge, UK (2002). [3](#)
- [20] —, *Hodge Theory and Complex Algebraic Geometry II*, Cambridge University Press, Cambridge, UK (2003). [6](#)