Aalborg Universitet



Inbreeding constrains host adaptive capacity and microbiome diversity

Ørsted, Michael; Yashiro, Erika; Hoffmann, Ary Anthony; Kristensen, Torsten Nygård

Creative Commons License CC BY 4.0

Publication date: 2022

Document Version Publisher's PDF, also known as Version of record

Link to publication from Aalborg University

Citation for published version (APA):

Ørsted, M., Yashiro, E., Hoffmann, A. A., & Kristensen, T. N. (2022). *Inbreeding constrains host adaptive capacity and microbiome diversity*. Abstract from Evolution and Population genetics in Denmark, Copenhagen, Denmark. http://popgen.dk/epic/#session7

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Inbreeding constrains adaptive capacity and microbiome diversity

Michael Ørsted¹, Erika Yashiro¹, Ary A. Hoffmann^{1,2}, and Torsten N. Kristensen¹

¹ Department of Chemistry and Bioscience, Section for Bioscience and Engineering, Aalborg University ² School of Biosciences, University of Melbourne, Australia.

The consequences of a reduction in population size commonly experienced by threatened species are normally interpreted in terms of loss of genetic variation, increased inbreeding, and associated inbreeding depression and reduced adaptive potential. However, large-scale empirical studies testing consequences of population genetic bottlenecks on genome-wide variation and evolutionary adaptation are surprisingly scarce. Also, it is becoming increasingly clear that microbial symbionts may play a so-far underappreciated role in conservation biology/genetics, as fitness effects of host population bottlenecks might also be mediated through the host microbiome diversity. Here we present results from an evolution experiment with 120 lines of Drosophila melanogaster having experienced inbreeding caused by low population size for a variable number of generations. Genetic variation in inbred lines and in outbred control lines was assessed by genotyping-by-sequencing (GBS). All lines were subsequently reared on a novel stressful medium for 10 generations during which egg-to-adult viability, body mass, productivity, and extinctions were scored. We found strong positive correlations between levels of genetic variation and evolutionary response in all investigated traits. We also found that lines with lower genetic diversity were at greater risk of extinction. Secondly, we amplified the bacterial 16S rRNA gene in a subset of the lines and found that host population bottlenecks constrained microbiome richness and diversity, with interacting effects of host genetic variation and microbial diversity on host fitness. Combined, this work presents convincing support for long-standing evolutionary theory, while providing novel insights about how population bottlenecks reduce hologenomic variation (combined host and microbial genetic variation). Thus, while the current biodiversity crisis typically focus on population sizes and genetic variation of eukaryotes, an additional focal point should be the microbial diversity carried by these hosts, which in turn may influence host fitness and adaptability and ultimately the persistence of populations.