

University of Groningen

Response to: Thoughts on Tissue Stromal Vascular Fraction for Early Scar Healing

van Dongen, Joris A; Boxtel, Joeri V; Uguten, Mustafa; Brouwer, Linda A; Vermeulen, Karin M; Melenhorst, Wynand B; Niessen, Frank B; Harmsen, Martin C; Stevens, Hieronymus P; van der Lei, B

Published in:
Aesthetic Surgery Journal

DOI:
[10.1093/asj/sjac098](https://doi.org/10.1093/asj/sjac098)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van Dongen, J. A., Boxtel, J. V., Uguten, M., Brouwer, L. A., Vermeulen, K. M., Melenhorst, W. B., Niessen, F. B., Harmsen, M. C., Stevens, H. P., & van der Lei, B. (2022). Response to: Thoughts on Tissue Stromal Vascular Fraction for Early Scar Healing. *Aesthetic Surgery Journal*, 42(8), NP560–NP561. <https://doi.org/10.1093/asj/sjac098>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Response to: Thoughts on Tissue Stromal Vascular Fraction for Early Scar Healing

Joris A. van Dongen, MD, PhD[✉]; Joeri van Boxtel, MD[✉];
Mustafa Uguten, MD; Linda A. Brouwer, BAS; Karin M. Vermeulen,
PhD; Wynand B. Melenhorst, MD, PhD; Frank B. Niessen, MD, PhD;
Martin C. Harmsen, PhD[✉]; Hieronymus P. Stevens, MD, PhD; and
Berend van der Lei, MD, PhD

Aesthetic Surgery Journal
2022, 1–2

© The Author(s) 2022. Published
by Oxford University Press on behalf
of The Aesthetic Society. All rights
reserved. For permissions, please
e-mail: journals.permissions@oup.com
<https://doi.org/10.1093/asj/sjac098>
www.aestheticsurgeryjournal.com

OXFORD
UNIVERSITY PRESS

Editorial Decision date: April 11, 2022; online publish-ahead-of-print April 23, 2022.

We would like to thank Sun and Zhao for their response and comments¹ on our randomized clinical trial studying the effect of tissue-stromal vascular fraction (tSVF) on early scar healing.² Their first comment is that the histologic evidence for an extracellular matrix in tSVF that functions as a scaffold is lacking. However, we feel there is enough evidence presented: the presence of an extracellular matrix in tSVF—isolated by the fractionation of adipose tissue (FAT) procedure—has clearly been demonstrated by Masson's trichrome staining in this study. In a previous study, we were able to decellularize tSVF (obtained by means of the FAT procedure) and to isolate extracellular matrix, which was entirely free of cells.³ This decellularized extracellular matrix was then digested with pepsin and turned into a self-assembled hydrogel. Multiple experiments subsequently demonstrated the capability of such an extracellular matrix–derived hydrogel to bind and release paracrine factors from adipose-derived stromal cells (ASCs) in a controlled fashion over time.³ These results show that adipose extracellular matrix functions as a scaffold.

The second comment is that mechanical fractionation of adipose tissue will reduce the viability of ASCs, a claim that is based on the authors being unable to culture ASCs derived from tSVF by an explant technique. Their findings, however, in our opinion do not justify this conclusion because enzymatic isolation of tSVF derived from mechanical fractionation still yields large numbers of ASCs. In contrast to enzymatic isolation, mechanical isolation preserves all cell-cell connections, including connection to

the extracellular matrix and small vessels.⁴ Therefore, in tSVF, ASCs are still attached around vessels as pericytes and/or vascular precursor cell types.⁵ Hence, using an explant technique to extract ASCs from tSVF is significantly more difficult than just culturing ASCs present in a single cell suspension of cellular SVF. One might even speculate that the viability of ASCs after mechanical isolation is even better than after enzymatic isolation because

Dr van Dongen is a plastic surgery resident, Department of Plastic Surgery, University Medical Center Utrecht, Utrecht, the Netherlands. Dr van Boxtel is a plastic surgery resident, Department of Plastic Surgery, Catharina Hospital Eindhoven, Eindhoven, the Netherlands. Dr Uguten is a plastic surgery resident, Department of Plastic Surgery, Haga Hospital, the Hague, the Netherlands. Ms Brouwer is a technician and Dr Harmsen is a professor of cardiovascular regenerative medicine, Department of Pathology and Medical Biology, University of Groningen and University Medical Center of Groningen, Groningen, the Netherlands. Dr Vermeulen is an epidemiologist, Department of Epidemiology, University of Groningen and University Medical Center of Groningen, Groningen, the Netherlands. Dr Melenhorst is a plastic surgeon, Department of Plastic Surgery, Diaconessenhuis Utrecht, Utrecht, the Netherlands. Dr Niessen is a plastic surgeon in private practice in Amsterdam, the Netherlands. Dr Stevens is a plastic surgeon in private practice in Rotterdam, the Netherlands. Dr van der Lei is a professor of aesthetic surgery, Department of Plastic Surgery, University of Groningen and University Medical Center of Groningen, Groningen, the Netherlands.

Corresponding Author:

Dr Berend van der Lei, University Medical Center of Groningen, Hanzplein 1, 9713 GZ Groningen, the Netherlands.
E-mail: info@berendvanderlei.nl; Twitter: [@professorvanderlei](https://twitter.com/professorvanderlei)

ASCs remain in their natural niche. Recently, a cleaved caspase-3 staining of freshly isolated tSVF showed no cell death in the first 24 hours after mechanical isolation in our laboratory (unpublished data). A disadvantage of mechanical vs enzymatic isolation of SVF is that mechanical isolation might be more variable because of performance variability among clinicians. Therefore, not all isolated tSVF will be free of adipocytes. Yet, not all adipocytes stain positive for perilipin A in tSVF, suggesting the presence of “ghost” adipocytes, which are disrupted cells with their shape still intact.²

Our prospective study has clearly demonstrated an accelerated early scar healing by a single injection of tSVF in the wound after mammoplasty at 6 months postsurgery. This accelerated wound healing effect, however, was no longer detectable at 12 months postsurgery. This observation might indicate that a single injection of tSVF is not that effective for accelerated wound healing in the long term. Multiple injections with tSVF are probably necessary to obtain long-term results. In situations with disturbed wound healing, such as chronic diabetic wounds, perianal fistulas, etc, however, a single injection of tSVF appears to be more effective.⁶ In these situations, healing inflammation plays a central role.

Disclosures

The authors declared no potential conflicts of interest with respect to the research, authorship, and publication of this article. Dr van der Lei is an international editor for *Aesthetic Surgery Journal*.

Funding

The authors received no financial support for the research, authorship, and publication of this article.

REFERENCES

1. Zhao H, Sun Z. Thoughts on tissue stromal vascular fraction for early scar healing. *Aesthet Surg J*. In press.
2. Van Dongen JA, Van Boxtel J, Uguten M, et al. Tissue stromal vascular fraction improves early scar healing: a prospective randomized multicenter clinical trial. *Aesthet Surg J*. 2021:sjab431. doi: [10.1093/asj/sjab431](https://doi.org/10.1093/asj/sjab431). [Epub ahead of print]
3. Van Dongen JA, Getova V, Brouwer LA, et al. Adipose tissue-derived extracellular matrix hydrogels as a release platform for secreted paracrine factors. *J Tissue Eng Regen Med*. 2019;13(6):973-985. doi: [10.1002/term.2843](https://doi.org/10.1002/term.2843)
4. van Dongen JA, Tuin AJ, Spiekman M, Jansma J, van der Lei B, Harmsen MC. Comparison of intraoperative procedures for isolation of clinical grade stromal vascular fraction for regenerative purposes: a systematic review. *J Tissue Eng Regen Med*. 2018;12(1):e261-e274. doi: [10.1002/term.2407](https://doi.org/10.1002/term.2407)
5. Corselli M, Chen CW, Sun B, et al. The tunica adventitia of human arteries and veins as a source of mesenchymal stem cells. *Stem Cells Dev*. 2021;21(8):1299-1308. doi: [10.1089/scd.2011.0200](https://doi.org/10.1089/scd.2011.0200)
6. Schouten WR, Arkenbosch JH, van der Woude CJ et al. Efficacy and safety of autologous adipose-derived stromal vascular fraction enriched with platelet-rich plasma in flap repair of transsphincteric cryptoglandular fistulas. *Tech Coloproctol*. 2021;25(12):1301-1309. doi: [10.1007/s10151-021-02524-6](https://doi.org/10.1007/s10151-021-02524-6)