

University of Groningen

Severe acute hepatitis and acute liver failure of unknown origin in children

Contributors Survey; de Kleine, Ruben H.; Lexmond, Willem S.; Buescher, Gustav; Sturm, Ekkehard; Kelly, Deidre; Lohse, Ansgar W.; Lenz, Dominic; Jorgensen, Marianne Horby

Published in:
Eurosurveillance

DOI:
[10.2807/1560-7917.ES.2022.27.19.2200369](https://doi.org/10.2807/1560-7917.ES.2022.27.19.2200369)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Contributors Survey, de Kleine, R. H., Lexmond, W. S., Buescher, G., Sturm, E., Kelly, D., Lohse, A. W., Lenz, D., & Jorgensen, M. H. (2022). Severe acute hepatitis and acute liver failure of unknown origin in children: a questionnaire-based study within 34 paediatric liver centres in 22 European countries and Israel, April 2022. *Eurosurveillance*, 27(19). <https://doi.org/10.2807/1560-7917.ES.2022.27.19.2200369>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Severe acute hepatitis and acute liver failure of unknown origin in children: a questionnaire-based study within 34 paediatric liver centres in 22 European countries and Israel, April 2022

Ruben H de Kleine^{1,2}, Willem S Lexmond^{2,3}, Gustav Buescher^{2,4}, Ekkehard Sturm^{2,5}, Deidre Kelly^{2,6}, Ansgar W Lohse^{2,4}, Dominic Lenz^{2,7*}, Marianne Hørby Jørgensen^{2,8*}, on behalf of the contributors to the survey⁹

1. Department of Surgery, Division of Hepatobiliary Surgery and Liver Transplantation, University of Groningen, University Medical Centre Groningen, Groningen, The Netherlands
2. European Reference Network for hepatological diseases (ERN RARE-LIVER)
3. Department of Paediatrics, Division of Gastroenterology and Hepatology, Beatrix Children's Hospital, University of Groningen, Groningen, The Netherlands
4. Department of Internal Medicine, University Medical Centre Hamburg-Eppendorf, Hamburg, Germany
5. Paediatric Gastroenterology/ Hepatology, University Hospital Tuebingen, Tuebingen, Germany
6. University of Birmingham, and The Liver Unit, Birmingham Women & Children's Hospital, Birmingham, UK
7. Division of Neuropaediatrics and Paediatric Metabolic Medicine, Centre for Paediatric and Adolescent Medicine, University Hospital Heidelberg, Heidelberg, Germany
8. Department of Paediatric and Adolescent Medicine, Rigshospitalet, Copenhagen, Denmark
9. The contributors to the survey are acknowledged at the end of the article

* These authors contributed equally to this work and share last authorship.

Correspondence: Ruben de Kleine (R.de.kleine@umcg.nl)

Citation style for this article:

de Kleine Ruben H, Lexmond Willem S, Buescher Gustav, Sturm Ekkehard, Kelly Deidre, Lohse Ansgar W, Lenz Dominic, Hørby Jørgensen Marianne, on behalf of the contributors to the survey. Severe acute hepatitis and acute liver failure of unknown origin in children: a questionnaire-based study within 34 paediatric liver centres in 22 European countries and Israel, April 2022. Euro Surveill. 2022;27(19):pii=2200369. <https://doi.org/10.2807/1560-7917.ES.2022.27.19.2200369>

Article submitted on 29 Apr 2022 / accepted on 12 May 2022 / published on 12 May 2022

To detect potential concern about severe acute hepatitis in children, we conducted a survey among 50 ERN RARE-LIVER centres. By 26 April 2022, 34 centres, including 25 transplant centres, reported an estimated median of 3–5, 0–2 and 3–5 cases in 2021, 2020 and 2019 and a mean of 2 (range: 0–8) cases between January and April 2022 (mean in 10 large liver transplant centres: 3). Twelve centres reported suspicion of an increase, but no rise.

Following a report by the United Kingdom (UK) on 5 April 2022 on the occurrence of cases of severe acute hepatitis in children aged 16 years or under, the World Health Organization (WHO) raised concerns about the possibility of an epidemic [1,2]. By 21 April, 169 possible or confirmed cases were reported fulfilling the WHO case definition [3]. The cause of the hepatitis is unknown but a link to a virus infection has been suggested due to the epidemiological pattern of cases [4,5]. The hepatitis can progress to paediatric acute liver failure (pALF) necessitating urgent liver transplantation to avoid multi-organ failure [6]. We intended to assess whether a rise in incidence of severe acute hepatitis or pALF could be observed between 1 January and 26 April 2022 in comparison to previous years, within the European Reference Network on Hepatological Diseases (ERN RARE-LIVER) [7].

Questionnaire survey

The web-based questionnaire survey available from the Supplementary material was sent to 50 (associate) member centres of the ERN RARE-LIVER involved in the treatment of children with hepatitis. Using the European Union (EU) Survey tool, the survey started on 15 April and was open for responses until 26 April. Data retrieved were managed from the secure EU Survey tool. Contributors were asked to register their name and email before entering anonymous data from their own case-defined patients at their centre. After the upload, a summary of their data was sent back for verification via email.

The survey questionnaire was divided into two parts with the first part related to the incidence and treatment of paediatric hepatitis and pALF patients fulfilling the definition of the Paediatric Acute Liver Failure Study Group (Box), in the 3 preceding years [8]. Participants were asked to provide an estimate with the options 0–2, 3–5, 6–10, 11–20 patients.

Questions in the second part of the survey focussed on patients seen in 2022 in line with the WHO definition of 15 April 2022 [1]. Responses captured anonymous details on all children admitted with severe hepatitis or pALF (Box) and data on demographics (age, sex,

Box

Case definition of severe non-A-E hepatitis and definition of paediatric acute liver failure

Severe non-A-E hepatitis

- Rise in aspartate-aminotransferase (AST) or alanine-aminotransferase (ALT) of more than 500 units per litre (U/L).
- Age \leq 16 years.
- Since 1 January 2022.
- No A-E hepatitis virus detected.

Paediatric acute liver failure

- Evidence of acute hepatic injury (elevated ALT).
- Spontaneous INR \geq 2.0.
- Or INR \geq 1.5 with signs of hepatic encephalopathy.

INR: international normalised ratio.

aetiology), clinical course (cause identified, possible cause, indetermined) and outcome (survival with native liver, liver transplantation indicated, overall survival) were entered.

This rapid communication was written with STROBE criteria in consideration [9].

Participating centres and responses

The survey was completed by 34 centres from 22 countries of which 31 are (associate) members of the ERN RARE-LIVER. This network is dedicated to the improvement of care for rare liver diseases by collaboration through innovation, research and exchange of data for all involved patients [7]. The location of participating centres is mapped in the Figure.

All centres treat children with hepatitis, while 25 of the responding 34 centres also perform paediatric liver transplantation. Except for contributing centres in Switzerland and Israel, all participants were from the EU. No data from the UK were entered in the query.

Incidence of severe acute hepatitis or pALF

Of the 34 participating centres, 22 reported no suspicion of an increase of children with severe hepatitis between January and April 2022, with 10 centres reporting no new paediatric hepatitis patients in this period. All centres reported seeing patients with hepatitis in all previous 3 years with further details listed in Table 1.

In the study period of January to April 2022, a mean of 2 (range: 0–8) cases was detected. Twelve centres reported a suspected increase but documented no rise in numbers. The 11 large transplant centres with 16 or more paediatric liver transplantations per year, reported a mean of 2.5 cases (range: 0–5) referred for transplantation in the 3.8 months of 2022, while in 2019, 2020 and 2021 on average 4.9 (range: 0–10),

3.7 (range: 0–10) and 4.9 (range: 0–10) cases per year were reported, respectively.

Details of children with severe hepatitis and paediatric acute liver failure in 2022

We received detailed information on 65 children with severe hepatitis ($n = 59$) or pALF ($n = 33$, 4 missing data, INR \geq 2.0) treated at 19 different centres. One was excluded since they were over 16 years of age. Four of 64 children did not fulfil the criteria for severe hepatitis but were included since they presented with the more severe condition of pALF. Therefore, some groups show overlap in numbers.

An extensive diagnostic work-up was documented in all children (Table 2). In 11 children the cause for the clinical condition was identified, while the cause remained indeterminate for 27 children (26 severe hepatitis and 10 pALF, with one child in the pALF group not fulfilling the hepatitis criteria). The centres reported a possible cause of the severe hepatitis or pALF episode without final certainty in 26 children (24 severe hepatitis; 17 pALF, with two children in the pALF group not fulfilling the hepatitis criteria).

A previously known cause for severe acute hepatitis in children, such as an underlying medical condition, was present in 16 of 64 children, comprising congenital defects ($n = 4$), immune deficiency ($n = 1$), metabolic disorders ($n = 3$) as well a wide range of other disease like a structural heart defect, bone marrow disease, immune therapy for malignancy or structural defects otherwise unclassified. Different viruses were detected in 13 children with adenovirus being present in four (Table 2). Five of 64 children (aged 7 years or older) were vaccinated against coronavirus disease (COVID-19). These five received the vaccination at least 4 weeks before the onset of symptoms of hepatitis with no report of adverse effects or complaints in between. No link between the vaccination and the liver disease was suspected by any of the contributors.

In total four children received a liver transplant, three died before an organ was available. One child with Wilson's disease (cause identified) died after transplantation (Table 3).

Discussion

We present results from a European survey among specialised liver centres across Europe and in Israel to provide further insight for the recently reported increase in severe acute hepatitis cases in children. Compared with the average of cases in each of the full previous years 2019–21, there was no absolute increase of cases within stated criteria in the study period, based on the data from the contributing centres. However, the data for 2022 comprise only the first 3.8 months of the year and should be considered preliminary. Extrapolation of these data into the future is tempting but should be done with caution.

FIGURE

Location of the centres participating in the survey (n = 34)



TABLE 1

Centres' reported estimated numbers of children with severe hepatitis or paediatric acute liver failure, European Reference Network Rare Liver survey, 1 January 2019–26 April 2022 (n = 34)

Number of children with pALF [8] per centre	2019	2020	2021	2022 ^a
	Number of centres	Number of centres	Number of centres	Number of centres (exact number of cases)
0–2	16	18	5	11 (0) 6 (1) 7 (2)
3–5	10	9	9	3 (3) 2 (4) 3 (5)
6–10	8	7	8	1 (8)
11–20	0	0	2	0

pALF: paediatric acute liver failure.

^a Data cover only 3.8 months and comprises pALF and acute severe hepatitis.

TABLE 2

Characteristics of children with severe hepatitis or paediatric acute liver failure by cause, ERN RARE-LIVER survey, 1 January–26 April 2022 (n = 64)

Baseline characteristics	All children	Cause identified	Possible cause identified	No cause identified
	(mean, range) n = 64	(mean, range) n = 11	(mean, range) n = 27	(mean, range) n = 26
Sex female vs male	28 vs 35 (1 missing)	5 vs 6	17 vs 9 (1 missing)	13 vs 13
Age	7.7 years (28 days–16 years)	5.5 years (28 days–16 years)	5.8 years (66 days–16 years) (2 missing)	5.8 years (56 days–16 years)
Admission to clinic	64	11	27	26
Laboratory findings				
ALT (U/L) ^a	2,871 (90–16,686)	1,115 (90–6,350)	3,640 (100–16,686)	2,736 (515–9,000)
Bilirubin (µmol/L) ^a	100 (0–470)	106 (1–534)	107 (0–470)	88 (4–468)
INR ^a	2.98 (0.89–16.70)	3.18 (1.13–7.00)	3.49 (1.00–16.70)	2.29 (0.96–6.70)
NH ₃	104 (26–472)	95 (45–215)	105 (26–266)	108 (27–472)
Viruses detected	13	2 (enterovirus, EBV)	9 patients, multiple viruses (4 adenovirus, 4 SARS-CoV-2, 1 rotavirus, 1 influenza A virus, 2 EBV) ^b	2 (rotavirus, influenza A virus)
Active SARS-CoV-2 infection at admission (n, %)	3 (5%)	0	3 (11%)	0
SARS-CoV-2 vaccinated (n, %)	5 (3 missing data) (8%)	0 (2 missing data)	4 (1 missing data)	1

EBV: Epstein-Barr virus; INR: international normalised ratio; NH₃: ammonia; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2.

^a Highest value selected.

^b Co-infection in one patient with rotavirus, adenovirus and SARS-CoV-2. Another patient with EBV and adenovirus.

TABLE 3

Outcome of children with severe acute hepatitis or paediatric acute liver failure by cause identified ERN RARE-LIVER survey, 1 January–26 April 2022 (n = 64)

Outcome as at 26 April	All children (mean, range) n = 64 (4 missing data)	Cause identified n = 11	Possible cause identified (mean, range) n = 27 (2 missing data)	No cause identified n = 26 (2 missing data)
Ongoing hepatitis	9	0	5	4
Survival with native liver	43	8	17	18
LTX (n)	4	1	1	2
Deceased before/after LTX (n)	3/1	2/1	1/0	0/0

LTX: liver transplantation.

While existing data on the hepatitis increase in children the UK and the United States suggest a role for adenovirus infections [1,10], we did not detect adenoviruses in the majority of patients, nor another uniform viral infection. Interestingly, severe hepatitis with abdominal symptoms suggestive of a gastrointestinal virus, was reported in 1923 [11] following the 1918 influenza pandemic. It was then considered to be related to susceptibility to viruses people had not been exposed to during social containment. This parallelism supports a possible role for viral pathogens in the current situation. The exact pathomechanism causing hepatitis remains largely unknown, however, an interaction between the immature or naïve immune system and the liver might play an important role. Based on the portal vein blood flow, directly coming from the intestine, and the described clinical presentation of abdominal pain, illness and vomiting [1], such a trigger might be coming from an infection with one of the enteric viruses, such as adenovirus type 41 and 42.

Our study is limited by the format of a self-reported questionnaire which comes with its own form of reporting bias. Sixteen of the 50 approached centres did not respond. This may have led to an overestimation of the actual numbers within the network since the mean number was calculated only from responders. Absolute numbers of paediatric severe hepatitis remain low and numbers of pALF have been even lower in 2022 and previously. Therefore, formal conclusions are difficult to make. Our survey does not give additional information to prompt extra caution or specific protective measures at this very moment. This variation of hepatitis and pALF occurrence needs further investigation and could benefit from a pan-European collaboration to research and biobank material for modern viral detection techniques in order to identify the underlying cause. Awareness among paediatric hepatologists about a possible rise in hepatitis is high and this should be a good basis for establishing a reliable platform for cutting edge research.

Conclusion

Currently, there is no clear overall increase in the occurrence of severe hepatitis or pALF in children in Europe, when we compare the reported data of previous years to the preliminary data from the study period of 3.8 months in 2022. By 26 April, only a minority of the centres within the ERN RARE-LIVER appear to be affected by a possible rise in hepatitis cases. No increase in virus-derived hepatitis could be detected at this point in time within the whole group of participating centres. However, a close monitoring will be necessary in order to achieve early signal detection of this clinical condition.

Acknowledgements

We acknowledge the help of M. Faber, Gastrointestinal Infections, Zoonosis and Tropical Infections (Division 35), Department for Infectious Disease Epidemiology, Robert Koch Institute, Berlin, Germany for his help and expertise in the field of epidemiology.

Contributors to the survey

Panayiota Protopapa, Paediatric Liver GI and Nutrition Centre Makarios Hospital, Nicosia, Cyprus

Eyal Shteyer, Shaare Zedek Medical Center, Jerusalem, Israel

Raffaele Iorio, University of Naples Federico II, Napoli, Italy

Ieva Pukite, Children's Clinical University Hospital Riga, Latvia, Latvia

Ruta Kucinskiene, Lithuanian University of Health Sciences Hospital, Kaunas, Lithuania

Anne-Marie Grima, Mater Dei Hospital, Msida, Malta

Bart Koot, Amsterdam University Medical Centre, Amsterdam, the Netherlands

Tudor Lucian Pop, 2nd Pediatric Clinic of the University of Medicine and Pharmacy Cluj-Napoca, Romania

Ekkehard Sturm, University Hospital Tuebingen, Tuebingen, Germany

Ruth De Bruyne, Ghent University Hospital, Ghent, Belgium

Stephennne, Cliniques Universitaires Saint Luc Bruxelles, Belgium

Jurica Vukovic, Department of Pediatrics, University Hospital Centre Zagreb, Croatia

Katarina Mitrova, Department of Paediatrics, University Hospital Motol, Prague, Czech Republic

Helene Kvistgaard, Aarhus University Hospital, Aarhus, Denmark

Marianne Hørby Jørgensen, Rigshospitalet, Copenhagen, Denmark

Laura Merras-Salmio, Helsinki University Hospital, New Children's Hospital, Finland

Geraldine Héry, Bicêtre Hospital, Paris, France

Dominique Debray, Necker Hospital, Paris, France

Mathias Ruiz, Hospital Femme Mere Enfant du CHU de Lyon, Lyon, France

Dominic Lenz, University Hospital Heidelberg, Heidelberg, Germany

Sebastian Schulz-Jürgensen, University Hospital Hamburg-Eppendorf, Hamburg, Germany

Eberhard Lurz, von Hauner Children's Hospital, LMU Munich, Germany

Virag Hives, First Department of Paediatrics, Semmelweis University, Budapest, Hungary

Mara Cananzi, University Hospital of Padova, Padova, Italy

Lorenzo D'Antiga, Hospital Papa Giovanni XXIII, Bergamo, Italy

Angelo di Giorgio, Hospital Papa Giovanni XXIII, Bergamo, Italy

Michele Pinon, AOU Città della Salute e della Scienza di Torino, Italy

Willem Lexmond, University Medical Centre Groningen, Groningen, The Netherlands

Diana Kamińska, Children's Memorial Health Institute Warsaw, Poland

Isabel Gonçalves Costa, Hospitais da Universidade de Coimbra, CHUC, Portugal

Jernej Breclj, University Children's Hospital Ljubljana, Slovenia

Gema Muñoz Bartolo, Hospital Infantil La Paz, Madrid, Spain

Jesús Quintero, Hospital Universitari Vall d'Hebron, Barcelona, Spain

Björn Fischler, Karolinska University Hospital, Stockholm, Sweden

Valerie McLin, University Hospitals Geneva, Geneva, Switzerland

Ethical statement

For conducting this study an institutional review board approval was obtained at the University Medical Centre Groningen, Groningen, The Netherlands (Reference M22.296493).

Conflict of interest

None declared.

Authors' contributions

R de Kleine conceptualised the study, designed the query, analysed the data and wrote the paper. W Lexmond designed the query, analysed the data, wrote the paper and designed the Figure. G Buescher managed the query EU tool, implemented the query, processed and analysed the data and wrote the paper. E. Sturm conceptualised the study, analysed the data and edited the paper. A Lohse and D Kelly conceptualised the study, supervised and edited the paper. D Lenz and M Hørby Jørgensen conceptualised the study, coordinated communications and edited the paper.

References

1. World Health Organization (WHO). Acute hepatitis of unknown aetiology – the United Kingdom of Great Britain and Northern Ireland. Geneva: WHO; 15 Apr 2022. Available from: <https://www.who.int/emergencies/disease-outbreak-news/item/acute-hepatitis-of-unknown-aetiology---the-united-kingdom-of-great-britain-and-northern-ireland>
2. European Centre for Disease Prevention and Control (ECDC). Increase in severe acute hepatitis cases of unknown aetiology in children. Stockholm: ECDC; 28 Apr 2022. Available from: <https://www.ecdc.europa.eu/sites/default/files/documents/RRR-20220420-218.pdf>
3. World Health Organization (WHO). Multi-Country – Acute, severe hepatitis of unknown origin in children. Geneva: WHO; 23 Apr 2022. Available from: <https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON376>
4. Christie B. Scientists search for cause of hepatitis cases in children. *BMJ*. 2022;377:0982. <https://doi.org/10.1136/bmj.0982> PMID: 35422430
5. United Kingdom Health Security Agency (UKHSA). Increase in acute hepatitis cases of unknown aetiology in children. London: UKHSA; 8 Apr 2022. Available from: <https://www.gov.uk/government/publications/hepatitis-increase-in-acute-cases-of-unknown-aetiology-in-children/increase-in-acute-hepatitis-cases-of-unknown-aetiology-in-children>
6. Squire s RH Jr, Shneider BL, Bucuvalas J, Alonso E, Sokol RJ, Narkewicz MR, et al. Acute liver failure in children: the first 348 patients in the pediatric acute liver failure study group. *J Pediatr*. 2006;148(5):652-8. PMID: 16737880
7. European Reference Network (ERN). Hepatological Diseases (ERN RARE-LIVER). What is ERN RARE-LIVER? [Accessed: 10 May 2022]. Available from: <https://rare-liver.eu/about/what-is-ern-rare-liver>
8. Squires RH Jr, Shneider BL, Bucuvalas J, Alonso E, Sokol RJ, Narkewicz MR, et al. Acute liver failure in children: the first 348 patients in the pediatric acute liver failure study group. *J Pediatr*. 2006;148(5):652-8. <https://doi.org/10.1016/j.jpeds.2005.12.051> PMID: 16737880
9. Vandembroucke JP, von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al. , STROBE Initiative. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *PLoS Med*. 2007;4(10):e297. <https://doi.org/10.1371/journal.pmed.0040297> PMID: 17941715
10. Mücke MM, Zeuzem S. The recent outbreak of acute severe hepatitis in children of unknown origin – what is known so far. *J Hepatol*. 2022;So168-8278(22)00271-9. <https://doi.org/10.1016/j.jhep.2022.05.001>

11. Williams H. Epidemic jaundice in New York State, 1921-1922. JAMA. 1923;80(8):532-4. <https://doi.org/10.1001/jama.1923.02640350014007>

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2022.