

**IV Meeting on Astrophysical Spectroscopy -
A&M DATA - Atmosphere**

May 30 to June 2, 2022, Fruška Gora, Serbia

**BOOK OF ABSTRACTS AND
CONTRIBUTED PAPERS**

**Edited by Vladimir A. Srećković, Milan S. Dimitrijević,
Nikola Veselinović and Nikola Cvetanović**

A&M DATA



Belgrade 2022

Scientific Committee

Milan S. Dimitrijević, **Co-Chairman**
Vladimir A. Srećković, **Co-Chairman**
Nebil Ben Nessib, Saudi Arabia
Nikolay Bezuglov, Russia
Magdalena Christova, Bulgaria
Nikola Cvetanović, Serbia
Stevica Djurović, Serbia
Saša Dujko, Serbia
Rafik Hamdi, Tunisia
Darko Jevremović, Serbia
Bratislav Marinković, Serbia
Zoran Mijić, Serbia
Aleksandra Nina, Serbia
Bratislav Obradović, Serbia
Luka Č. Popović, Serbia
Branko Predojević, Republic of Srpska
Maja Rabasović, Serbia
Sylvie Sahal-Bréchet, France

Local Organizing Committee

Vladimir A. Srećković (Chair), Institute of Physics Belgrade
Nikola Veselinović, Institute of Physics Belgrade
Lazar Gavanski, Faculty of Sciences – University of Novi Sad
Nataša Simić, Faculty of Sciences – University of Novi Sad
Veljko Vujčić, Astronomical Observatory, Belgrade
Radomir Banjanac, Institute of Physics Belgrade
Aleksandra Kolarski, Institute of Physics Belgrade
Milan S. Dimitrijević, Astronomical Observatory, Belgrade

Organizers:

Institute of Physics Belgrade, Serbia,
Astronomical Observatory Belgrade, Serbia and
Faculty of Sciences – University of Novi Sad, Serbia

Text arrangement by computer: Tanja Milovanov

ISBN

Published and copyright © by Institute of Physics Belgrade, Pregrevica 118,
11080 Belgrade Serbia

Financially supported by the Ministry of Education, Science and Technological
Development of Serbia

Production: Skripta Internacional, Mike Alasa 54, Beograd in 50 copies

SCIENTIFIC RATIONALE

Spectroscopy is a powerful tool for the analysis of radiation from different plasmas in astronomy, laboratory, fusion research, atmospheric research and industry. Effective theoretical analysis, synthesis and modelling of stellar spectra as well as the spectra from other plasma sources, depends on atomic data and their sources. In particular, for the modelling of stellar atmospheres and opacity calculations a large amount of atomic data is needed, since we do not know *a priori* the chemical composition of a stellar atmosphere. Consequently, the development of databases with atomic data and astroinformatics is important for stellar spectroscopy.

The Conference is planned as an opportunity to consider above mentioned aspects of spectroscopic research on plenary sessions and then to work on the special mini-projects, which will result in common papers to be published in international astronomical journals after the Conference.

Venue

Fruška Gora (Ceptor, Andrevlje), Serbia

CONTENTS

Zlatko Majlinger, Milan S. Dimitrijević and Vladimir A. Srećković: <i>Curve fitting method of Stark width determination – example of H I line in G191-B2B spectrum</i>	9
Slavica Malinović-Milićević: <i>Clear-sky spectral UV radiation modeling</i> (Invited lecture)	10
Željka D. Nikitović and Zoran M. Raspopović: <i>Rate coefficients of He⁺ ions in CF₄ gas</i>	16
Milan S. Dimitrijević, Magdalena D. Christova, Nenad Milovanović and Sylvie Sahal-Bréchet: <i>Stark broadening of Zn II spectral lines in stellar atmospheres</i> (Invited lecture)	17
Maja S. Rabasovic, Vladimir Sreckovic, Bratislav P. Marinkovic and Dragutin Sevic: <i>Analysis of laser induced plasma plume in atmosphere: artificial neural network approach</i> (Invited lecture)	19
Maja Kuzmanoski, Luka Ilić and Slobodan Ničković: <i>Spatial variability of mineral dust single scattering albedo based on DREAM model</i> (Invited lecture)	21
Aleksandra Kolarski and Vladimir Srećković: <i>Lower Ionosphere perturbations due to Solar X-ray flares simultaneously monitored on two VLF signals with close GCPs</i>	23
Aleksandra Kolarski: <i>Monitoring effects of low intensity X-ray Solar flares from 23/24 Solar cycle minimum on VLF signals recorded in Belgrade</i> (Progress report)	24
Aleksandra Kolarski: <i>Numerical simulations of subionospheric VLF propagation under influence of moderate Solar X-ray flare events</i> (Invited lecture)	25
Aleksandra Nina, Vladimir Čadež, Luka Č. Popović and Milan Radovanović: <i>Periodic variations of ionospheric Wait's parameters caused by changes in the intensity of incoming solar hydrogen Ly radiation</i>	32
Oleg R. Odalović: <i>GNSS signals as a tool for detection of the influence of solar radiation of terrestrial ionosphere</i> (Invited lecture)	37
Dušan S. Petković, Oleg R. Odalović and Aleksandra Nina: <i>Influence of the periodic changes in the incoming solar hydrogen Ly-α radiation intensity on the total electron content in the ionospheric D-region</i>	39

Z. Mijatović, S. Djurović, T. Gajo and I. Savić: <i>Approximation of the shape of hydrogen H_b spectral line with Voigt profiles</i>	44
Mihailo Savić, Nikola Veselinović, Aleksandar Dragić, Dimitrije Maletić, Dejan Joković, Vladimir Udovičić, Radomir Banjanac and David Knežević: <i>The study of atmospheric effects on cosmic ray muons in the Low Background Laboratory for Nuclear Physics at the Institute of Physics Belgrade</i> (Invited lecture)	48
Bratislav P. Marinković, Stefan Ivanović and Zoran Mijić: <i>Data analysis on Serbian participation in COST Actions: Celebrating 50 years of research networks</i> (Progress report)	49
Vladimir Srećković, Ljubinko Ignjatović, Milan Dimitrijević, Nikolai Bezuglov and Andrey Klyucharev: <i>Rate coefficients and cross-sections for some collisional processes involving Rydberg atoms</i>	58
Vladimir Srećković, Veljko Vujčić, Aleksandra Kolarski, Jelena Barović and Ognyan Kounchev: <i>Low ionosphere modeling: new data and models</i>	59
Veljko Vujčić and Vladimir A. Srećković: <i>A&M Data: Processing and Modeling in Real Time</i>	60
Vladimir Srećković, Jelena Barović and Gordana Jovanović: <i>Ionosphere modeling and intense radiation</i>	62
Vladimir A. Srećković and Sanja Tošić: <i>Collisional and radiative processes involving some small molecules: A&M data</i>	63
Lamia Abu El Maati, Mahmoud Ahmad, I. S. Mahmoud, Sahar G. Tawfik, Najah Alwadii, Nabil Ben Nessib and Milan S. Dimitrijević: <i>Atomic structure and transition parameters of the V XVIII carbon-like ion</i> (Invited lecture)	64
Violeta M. Petrović, Sanja D. Tošić, Hristina Delibašić Marković and Ivan D. Petrović: <i>Investigation of Laser Induced Breakdown Threshold</i> (Invited lecture)	66
Ljiljana M. Brajović and Miodrag Malović: <i>Gravity satellite missions measurement data for atmospheric density estimation</i> (Invited lecture)	68
Jelena B. Maljković, Jelena Vukalović, Zoran Pešić, F. Blanco, G. García and Bratislav P. Marinković: <i>Absolute differential cross section for elastic electron scattering from halothane molecule at 50eV</i> (Invited lecture)	70

Zoran R. Mijić and Bratislav P. Marinković: <i>Statistics of Management Committee Members from Serbia in COST Actions</i> (Progress report)	75
Saša Dujko, Danko Bošnjaković, Ilija Simonović and Zoran Lj. Petrović: <i>Collisional and transport processes of low-energy positrons in molecular hydrogen</i> (Invited lecture)	82
N. M. Sakan, I. Traparić, V. A. Srećković and M. Ivković: <i>The usage of perception, feed and deep feed forward artificial neural networks on the European South Observatory data as a test model for determining stellar spectral type</i>	84
Nikola Cvetanović and Bratislav M. Obradović: <i>Stark broadening method of Hydrogen Balmer beta for low-density atmospheric pressure discharges</i>	86
Darko Jevremovic: <i>History of Serbian involvement in LSST</i> (Invited lecture)	88
Zoran Simić and Nenad Sakan: <i>On the Stark broadening parameters of Ir II spectral lines</i>	89
Zoran Mijić, Maja Kuzmanoski, Luka Ilić, Aleksander Kovačević and Darko Vasiljević: <i>Review of atmospheric aerosol optical properties profiling and lidar station activities in Serbia</i> (Invited lecture)	90
PROGRAMME	99
AUTHORS' INDEX	103

Clear-sky spectral UV radiation modeling

Slavica Malinović-Milićević

*Geographical Institute “Jovan Cvijić”, Serbian Academy of Sciences and Arts,
Djure Jakšića 9, 11000 Belgrade, Serbia
E-mail: s.malinovic-milicevic@gi.sanu.ac.rs*

Abstract

The parametric model NEOPLANTA was used to study the effects of various atmospheric conditions on terrestrial spectral UV irradiance. The capability of the model to correctly reproduce processes in the atmosphere is tested by changing input parameters such as ozone content, solar zenith angle, amount and type of aerosols, and altitude. The results of the model testing on input parameter change were satisfactory. As is expected, the amount of UV radiation reaching the surface is the most dependable on ozone layer thickness and solar zenith angle. Erythmal irradiance is more dependent on changes in the amount of ozone in the atmosphere than spectral UV irradiance, showing a significantly greater impact of UV-B radiation on human skin than radiation in the UV-A part of the spectrum.

Introduction

Although ultraviolet (UV) radiation (UVR) is a small part of the solar spectrum, its effect on the living world is very important because it is highly energetic and biologically very active. UVR is divided into the UV-C range (100–280 nm), UV-B range (280–315 nm), and UV-A range (315–400 nm). Surface UVR is predominantly within the UV-A range, while the UV-B wavelengths are largely filtered out by stratospheric ozone and make less than 5% of surface UVR. The UV-C radiation does not reach the earth's surface because it is completely absorbed in the upper atmosphere by ozone and oxygen. Besides ozone content, the ambient UVR depends on many other variables such as the position of the Sun, cloud cover, atmospheric aerosols type and content, surface albedo, and altitude. Processes are very complex since factors that control attenuation vary as a function of time and location.

The detection of the large depletion of stratospheric ozone in the last three decades of the 20th century (Molina and Rowland, 1974; Kaurola et al., 2000; etc) initiated increased public and scientific interest in the state of stratospheric ozone levels and variability of UV radiation and led to the establishment of the Montreal Protocol on Substances that Deplete the Ozone Layer in 1987. However, although mitigation activities over the past three decades significantly limited the production

of ozone-depleting substances and stabilized the ozone levels, UV radiation levels are still high (EEAP, 2019).

Although the ground-based UV monitoring network constantly growing since its formation in the 1980s, the measuring sites are still sparsely located, while UV data are, in most cases, limited to the period up to 25 years. Most of the stations are equipped with broadband meters (75% in Europe) that measures irradiance over a certain wavelength range and convert it into UV index, while the spectral measurement is even rarer (Schmalwieser, 2017). Due to lack of the long-term measurements different modeling techniques based on radiative transfer, and empirical method has been established (Diffey, 1977; Ruggaber et al., 1994, Mayer and Kylling, 2005, Lindfors and Vuilleumier, 2005; Rieder *et al.*, 2008; Paulescu *et al.*, 2010; etc). Radiative transfer models use available atmospheric parameters and estimate the UV radiation considering the scattering and absorption processes that take place in the radiation path through the atmosphere. In general, those models are more accurate than empirical, although their accuracy depends on the availability and reliability of input parameters.

This paper aims to improve our understanding of variations in the surface UVR using the parametric model NEOPLANTA which was developed at the University of Novi Sad (Malinović, 2003) as the first original model in Serbia.

Analysis and results

Description of the model

The numerical model NEOPLANTA calculates the intensity of clear sky direct and diffuse solar UVR, the corresponding erythemal UVR, and the UV index on the horizontal surface. The calculation of direct and diffuse UVR was performed with a wavelength resolution of 1 nm in the range of 280-400 nm. The model simulates the effects of the absorption of the UVR by O₃, SO₂, and NO₂, and absorption and scattering by aerosol and air molecules in the atmosphere. It calculates the spectral irradiance for a given solar zenith angle (SZA), but there is also the possibility of calculating the UV index for the whole day at intervals of half an hour from sunrise to sunset. It is designed so that it can be used in any location in the world at any time.

In the model, the atmosphere is divided into parallel layers (maximum 40), and it is assumed that each layer is a homogeneous medium with constant meteorological parameters. The vertical resolution of the model is 1 km for altitudes less than 25 km and 5 km for altitudes between 25 km and 100 km, while the thickness of the last layer varies depending on the altitude. The values of the meteorological elements in each layer are determined using vertical profiles. The model contains data on the vertical profile of meteorological parameters of the standard atmosphere (Kurucz et al., 1984), but it is possible to use measured or modeled values. The influence of the surface on the intensity of UVR was calculated using the spectral albedo values for nine different types of surfaces.

The required input parameters are the local geographic coordinates and time or SZA, altitude, spectral albedo, and the total amount of gases. The model includes its vertical gas profiles (Ruggaber et al. 1994) and extinction cross-sections (Burrows et al. 1999; Bogumil et al. 2000), extraterrestrial solar irradiance shifted to terrestrial wavelength (Koepke et al. 1998), aerosol optical properties for 10 different aerosol types (Hess et al. 1998), and spectral albedo for nine different ground surface types (Ruggaber et al. 1994). Output data are spectral direct, diffuse, and global irradiance divided into the UV-A and UV-B part of the spectrum, erythemal irradiance, UV index (UVI), spectral optical depth, and spectral transmittance for each atmospheric component. All outputs are computed at the lower boundary of each layer.

The UV irradiance is calculated as the sum of the direct and the diffuse components. Calculation of the direct part of UVR is carried out by Beer-Lambert law. The direct irradiance $E_{dir}(\lambda)$ at wavelength λ received at ground level by unit area is given by:

$$E_{dir}(\lambda) = E_0(\lambda)T(\lambda), \quad (1)$$

where $E_0(\lambda)$ is the extraterrestrial irradiance corrected for the actual Sun-Earth distance and $T(\lambda)$ is the total transmittance that includes O_3 , NO_2 , SO_2 , aerosol, and air transmittances. Each transmittance is calculated using optical depth $\tau(\lambda)$ is the product of extinction coefficient $\beta(\lambda)$ and ray path through the atmosphere s :

$$T(\lambda) = \exp[-\tau(\lambda)] = \exp[-\beta(\lambda)s]. \quad (2)$$

The extinction coefficient of UV radiation, $\beta(\lambda)$, is calculated by the product of the cross-sectional area, $\sigma(\lambda)$, and layer particle concentration, N .

The diffuse UVR, $E_{dif}(\lambda)$, is divided into three components: (i) the Rayleigh scattering component, $E_{ray}(\lambda)$ (ii) the aerosol scattering component, $E_{aer}(\lambda)$ and (iii) the component that accounts for multiple reflections of irradiance between the ground and the air, $E_{rf}(\lambda)$. Calculation of $E_{dif}(\lambda)$ is made using an improved set of the equation described by Bird and Riordan (1986) and Blattner (1983) that were modified to include the wavelength range calculated by the NEOPLANTA model. The downward fraction is calculated from the same transmittance functions used to determine the direct UV irradiance. Calculation procedure of $E_{dif}(\lambda)$ is described in more detail in Malinović (2003) and Malinovic et al. (2006).

Evidence of an increasing number of skin cancer cases (IARC, 1992) has made it necessary to inform the public about the risk of solar radiation. Therefore, to take into account the potential erythemal effects of UVR on human skin calculated UV spectrum (in $Wm^{-2}nm^{-1}$) was multiplied by the erythema spectral weighting function ($s_{er}(\lambda)$) adopted by the CIE (CIE, 2019) thus providing the erythemal irradiance ($E_{er}(\lambda)$, in Wm^{-2}):

$$E_{er}(\lambda) = \int E(\lambda)s_{er}(\lambda)d\lambda, \quad (3)$$

where $E(\lambda)$ is spectral UV irradiance ($E_{dir}(\lambda) + E_{dif}(\lambda)$). $E_{er}(\lambda)$ assesses the potential of UV radiation to induce erythema (sunburn) and it is usually expressed as UVI ($UVI = E_{er} \times 40$). The erythemal irradiance accumulated over time gives the erythemal radiant exposure (H_{er} , in Jm^{-2}) (CIE, 2019).

Sensitivity of the model to input atmospheric parameters

Similar to measurements, models are subject to uncertainties that can be divided into two groups, numerical errors and uncertainties due to input parameters. Numerical errors can be avoided by programming and testing, while the uncertainties due to input parameters can be examined by sensitivity analysis and comparison with measured data.




Fig. 1. Dependence of spectral UV irradiance on (a) ozone thickness and (b) air pollution levels; (c) dependence of UVI on altitude, and (d) dependence of share of diffuse UV irradiance on SZA.

All the input parameters of the model have an impact on the intensity of UVR reaching the surface, but not to the same extent. The capability of the model to correctly reproduce processes in the atmosphere is tested by changing input parameters. The tests have shown that the model can properly simulate changes in the intensity of UVR as well as the ratio of direct and diffuse UVR with the change in input parameters. The simulations qualitatively agree with the existing knowledge: (i) the intensity of UVR decreases with increasing the thickness of the ozone layer (Fig. 1a), the amount of aerosol (Fig. 1b), altitude, and the increase in

SZA (Fig. 1c), (ii) the contribution of diffuse UVR increases with increasing SZA (Fig. 1 d), the amount of aerosols, the presence of aerosols with a higher amount of water-soluble particles, air humidity, and more reflecting surface. Elevation of the surface above sea level has a small effect, but it becomes considerable in mountainous areas. Simulations have shown that ozone absorbs the largest amount of radiation while SO_2 and NO_2 have little impact, except in heavily polluted areas. It has been shown that aerosols can also greatly affect the amount and composition of UVR on the surface, while the effect on the surface reflection is small, except in the case of snow-covered areas. It has also been shown that the contribution of diffuse radiation in clear-sky UVR reaching the surface is large and that its amount depends mostly on SZA.

While extraterrestrial radiation $E_0(\lambda)$ is almost constant over time in UV-B and UV-A range $T(\lambda)$ transmittance is highly dependent on the composition of the Earth's atmosphere and is therefore very variable over time. However, as Fig. 1a shows, while $E_0(\lambda)$ is highly variable on the wavelength scale, the $T(\lambda)$ forms a relatively smooth curve. The processes in the Earth's atmosphere, mainly ozone absorption and Rayleigh scattering, vary smoothly with wavelength, define the transmittance $T(\lambda)$ and form the UV-B cutoff. The resulting spectrum $E(\lambda)$ in Fig. 2b is the product of graphs in Fig. 1a. Erythemal irradiance $E_{er}(\lambda)$ has a maximum in the wavelength range of 300 to 320 nm. $E_{er}(\lambda)$ and therefore it is more dependent on changes in the amount of ozone in the atmosphere than $E(\lambda)$, showing a significantly greater impact of UV-B radiation on human skin than radiation in the UV-A part of the spectrum (Fig. 2c).




Fig.2. (a) Extraterrestrial UV spectrum $E_0(\lambda)$, and transmittance of the atmosphere $T(\lambda)$, (b) measured and modeled spectral UV irradiance $E(\lambda)$, and measured and modeled erythemal UV irradiance at 12:10 CET, 1 July 2021, Uccle, Belgium.

Conclusions

The parametric model NEOPLANTA was used to study the effects of various atmospheric conditions on terrestrial spectral UV irradiance. The capability of the model to correctly reproduce processes in the atmosphere is tested by changing input parameters such as ozone content, SZA, amount and type of aerosols, and altitude. Tests showed that the intensity of UVR decreases with increasing the thickness of the ozone layer, the amount of aerosol, altitude, and the increase in

SZA. It has been also confirmed that the contribution of diffuse UVR increases with increasing SZA, the amount of aerosols, the presence of aerosols with a higher amount of water-soluble particles, air humidity, and more reflecting surface. Elevation of the surface above sea level has a small effect, but it becomes considerable in mountainous areas. Erythema irradiance is more dependent on changes in the amount of ozone in the atmosphere than spectral UVR, showing a significantly greater impact of UV-B radiation on human skin than radiation in the UV-A range. The results are in accordance with the existing knowledge.

References

- Bird E.R., Riordan C., 1986, *J. Clim. Appl. Meteorol.* 25, 87–97.
- Blattner W., 1983, Radiation Research Associates, Fort Worth, TX, 104 pp.
- Bogumil K., Orphal J., Burrows J.P., 2000, Proc ERS- ENVISTAT Symp., Gothenburg, Sweden, ESA–ESTEC, pp 11
- Burrows J.P., Richter A., Dehn A., Deters B et al., 1999, *J. Quant. Spectrosc. Radiat. Transf.* 61,509–517.
- CIE, 2019, Erythema Reference Action Spectrum and Standard Erythema Dose; ISO/CIE 17166:2019 (E), CIE Standard. CIE Publications, Wien, Austria.
- Diffey B.L., 1977, *Phys. Med. Biol.*, 22, 309–316.
- EEAP, 2019, Nairobi: Environmental Effects Assessment Panel, United Nations Environment Programme (UNEP) 390 pp.
- Hess M., Koepke P., Schult I., 1998, *Bull Am Meteorol Soc* 79, 831–844
- IARC, 1992, *Risks Hum.*, 1992, 55, 1–316.
- Kaurola J., Taalas P., Koskela T., Borkowski J., Josefsson W. 2000, *J. Geophys. Res.*, 105, 20813–20820.
- Koepke P., Basis A., Balis D., Buchwitz M., et al., 1998, *Photochem. Photobiol.*, 67,657–662
- Kurucz R.L., Furenlid I., Brault J., Testerman L., 1984, National Solar Observatory Atlas No. 1
- Lindfors A., Vuilleumier, L., 2005, *J. Geophys. Res.* 110, D02104.
- Malinovic S., Mihailovic D.T., Kapor D., Mijatovic Z., Arsenic I.D., 2006, *J. Appl. Meteorol. Climatol.* 45, 1171–1177.
- Malinović, S., 2003, M.S. thesis, University of Novi Sad, 103 pp.
- Mayer B., Kylling A., 2005, *Atmos. Chem. Phys.*, 5, 1855–1877.
- Molina M.J., Rowland F.S., 1974, *Nature*, 249, 810–812.
- Paulescu M., Stefu N., Tulcan-Paulescu E., Calinoiu D., et al., 2010, *Atmos. Res.* 96, 141–148.
- Rieder H.E., Holawe F., Simic S., Blumthaler M., et al., 2008, *Atmos. Chem. Phys.* 8, 6309–6323.
- Ruggaber A., Dlug R., Nakajima T., 1994, *J. Atmos. Chem.*, 18,171–210.
- Schmalwieser A.W., Gröbner J., Blumthaler M., Klotz B., et al., 2017, *Photochem. Photobiol. Sci.*, 16(9), 1349–1370.