

University of Groningen

How applicable is geospatial analysis in maternal and neonatal health in sub-Saharan Africa?

Alemu, Sisay Mulugeta; Tura, Abera Kenay; Gurgel do Amaral, Gabriel S; Moughalian, Catherine; Weitkamp, Gerd; Stekelenburg, Jelle; Biesma, Regien

Published in:
Journal of global health

DOI:
[10.7189/jogh.12.04066](https://doi.org/10.7189/jogh.12.04066)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Alemu, S. M., Tura, A. K., Gurgel do Amaral, G. S., Moughalian, C., Weitkamp, G., Stekelenburg, J., & Biesma, R. (2022). How applicable is geospatial analysis in maternal and neonatal health in sub-Saharan Africa? A systematic review. *Journal of global health, 12*, [04055]. <https://doi.org/10.7189/jogh.12.04066>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

How applicable is geospatial analysis in maternal and neonatal health in sub-Saharan Africa? A systematic review

Sisay Mulugeta Alemu¹,
Abera Kenay Tura^{2,3}, Gabriel S
Gurgel do Amaral⁴, Catherine
Moughalian¹, Gerd Weitkamp⁵,
Jelle Stekelenburg^{1,6}, Regien
Biesma¹

¹Global Health Unit, Department of Health Sciences, University Medical Center Groningen, Groningen, the Netherlands

²School of Nursing and Midwifery, College of Health and Medical Sciences, Haramaya University, Harar, Ethiopia

³Department of Obstetrics and Gynaecology, University Medical Centre Groningen, University of Groningen, Groningen, the Netherlands

⁴Department of Health Sciences, Community and Occupational Medicine, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

⁵Department of Cultural Geography, Faculty of Spatial Sciences, University of Groningen, Groningen, the Netherlands

⁶Department Obstetrics & Gynaecology, Leeuwarden Medical Centre, Leeuwarden, the Netherlands

Background Sub-Saharan Africa (SSA) has the world's highest maternal and neonatal morbidity and mortality and has shown the slowest progress in reducing them. In addition, there is substantial inequality in terms of maternal and neonatal morbidity and mortality in the region. Geospatial studies can help prioritize scarce resources by pinpointing priority areas for implementation. This systematic review was conducted to explore the application of geospatial analysis to maternal and neonatal morbidity and mortality in SSA.

Methods A systematic search of PubMed, SCOPUS, EMBASE, and Web of Science databases was performed. All observational and qualitative studies that reported on maternal or neonatal health outcomes were included if they used a spatial analysis technique and were conducted in a SSA country. After removing duplicates, two reviewers independently reviewed each study's abstract and full text for inclusion. Furthermore, the quality of the studies was assessed using the Joanna Briggs Institute (JBI) critical appraisal checklists. Finally, due to the heterogeneity of studies, narrative synthesis was used to summarize the main findings, and Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline was strictly followed to report the review results. A total of 56 studies were included in the review.

Results We found that geospatial analysis was used to identify inequalities in maternal and neonatal morbidity, mortality, and health care utilization and to identify gaps in the availability and geographic accessibility of maternal health facilities. In addition, we identified a few studies that used geospatial analysis for modelling intervention areas. We also detected challenges and shortcomings, such as unrealistic assumptions used by geospatial models and a shortage of reliable, up-to-date, small-scale georeferenced data.

Conclusions The use of geospatial analysis for maternal and neonatal health in SSA is still limited, and more detailed spatial data are required to exploit the potential of geospatial technologies fully.

Correspondence to:

Sisay Mulugeta Alemu
Global Health Unit, Department of Health Sciences
University Medical Center Groningen
AD, Groningen
The Netherlands
s.m.alemu@umcg.nl

Sub-Saharan Africa (SSA) records the highest maternal and neonatal morbidity and mortality in the world [1]. Although SSA is home to only 13% of the world's population, 66% of maternal deaths worldwide occur in this region [2,3], with substantial inequality between and within countries in the region [3,4]. In addition, the region has shown the slowest progress in reducing maternal and neonatal morbidity and mortality [5].

Despite the global consensus to reduce maternal and neonatal morbidity and mortality [6,7], and despite these deaths being largely preventable [8], there are many challenges in SSA that slow progress. Low coverage, poor quality, and inequities in the provision of emergency obstetric and newborn care (EmONC) remain a challenge in many SSA countries [9]. In low- and middle-income countries (LMICs), and in particular, in the region of SSA, prioritizing interventions and resource allocation to areas where maternal and neonatal morbidity and mortality are most likely to happen, is necessary to ensure that the available resources are optimally used [10].

Geospatial analysis is one of the tools for improving decision-making by identifying geographical inequality and priority areas for implementation [11]. With the growing availability of spatial and/or temporal data at higher resolutions [12], the geographical coverage of maternal and neonatal health care can now be monitored with increasing accuracy. Geospatial analysis allows one to identify geographical inequalities in maternal and neonatal health by locating high and low morbidity and mortality areas. It can also help to determine the possible reasons for such disparities [13]. As a result, findings from geospatial analysis can help target intervention programs to areas where maternal and neonatal deaths are most likely to occur. Geospatial analysis can also suggest where to locate resources, such as ambulances and maternity waiting homes, to ensure timely access to EmONC facilities [14]. Other applications include identifying gaps in health facilities' availability and geographic accessibility, demand analysis, pinpointing ideal or priority locations for building new hospitals, health care data mapping and communication, and health program evaluation. Despite numerous applications of geospatial analysis techniques, its application for maternal and newborn health planning and priority setting in SSA was particularly rare. However, it is quickly gaining recognition, and more research outputs are being published. Therefore, there is a need to review the application of geospatial analysis in SSA. The aim of this review is, therefore, to explore the application of geospatial analysis in maternal and neonatal morbidity and mortality in SSA.

METHODS

Protocol registration

The review protocol was registered in the prospective registry of systematic reviews (PROSPERO) with registration ID CRD42021224930. We reported the review following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline [15] (Appendix S1 in the [Online Supplementary Document](#)).

Data source and search strategy

A systematic search of relevant publications was made in PubMed, SCOPUS, EMBASE, and Web of Science from inception until January 05, 2021. A comprehensive search strategy was developed for PubMed and adapted for the rest of the databases in consultation with a medical information specialist. The core search terms and phrases were 'maternal health,' 'neonatal health,' 'spatial analysis,' and 'sub-Saharan Africa'. The last search was done on January 05, 2021. In addition, bibliographies of identified articles and grey literature were hand-searched (Appendix S2 in the [Online Supplementary Document](#)).

Eligibility criteria

All observational (ecological, cross-sectional, case-control, cohort, survey, and surveillance reports) and qualitative studies were included. In contrast, commentaries, case reports, case series, anonymous reports, conference abstracts, letters, protocols, systematic reviews, meta-analyses, and editorials were excluded. Articles were included if they fulfil the following criteria: 1) contain data on maternal or neonatal health outcomes (ie, morbidity, mortality, or health care utilization during pregnancy, childbirth, or postpartum), 2) used a spatial analysis technique, and 3) conducted in SSA countries. There was no restriction on language or time of publication of the study. However, studies that only describe the geographical distribution of health outcomes without spatial analysis (eg, comparing maternal mortality rates between different provinces of a country) were excluded. We also excluded one study because the full text was not accessible, and we failed to get a response from the corresponding authors.

Screening and selection

Screening and selection of the studies were made using the Covidence, a web-based platform for systematic reviews [16]. At first, all studies obtained from the databases were exported to Covidence. After duplicates were removed, the title and abstract of each study were screened independently by two reviewers for inclusion. Then, two reviewers independently screened the full text of all potentially important articles. Any disagree-

ment between reviewers was resolved by discussion. The agreement between reviewers during the screening was 'good' (Cohen's kappa=0.63) [17]. The screening and selection process of the reviewed articles was illustrated using a PRISMA flow diagram (Figure 1).

Quality assessment/ risk of bias and data extraction

The quality of studies that meet the eligibility criteria was assessed using the Joanna Briggs Institute (JBI) critical appraisal checklists. JBI comprises checklists that look at the methodological quality of a study to determine the extent to which it has addressed the possibility of bias in its design, conduct, and analysis [18]. Only studies with 'good' quality were selected for data extraction and analysis. For all included studies, data about the title, year of publication, journal of publication, aim or research questions, focus (ie, maternal, neonatal, or both), the scope of the study (regional, national, subnational), country, outcome variables, study design, data sources, study period, study participants/population, sample size, spatial analysis technique used, funding sources, findings, and the conclusion were extracted. For one-third of the studies, data extraction and quality assessment were done by two individuals independently. The agreement between the two individuals was assessed to be "good." Subsequently, one reviewer did a quality assessment and extracted the remaining studies to save time.

Data synthesis and reporting

We used narrative synthesis and a table to summarize the main findings. Meta-analysis was not possible since the included studies were heterogeneous in terms of population, methods, and outcomes. PRISMA guideline [15] was strictly followed to report the review results.




Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of literature screening and selection process.

RESULTS

Characteristics of included studies

From a total of 11 499 studies returned from the search, 56 studies that fulfilled the inclusion criteria were included in the narrative synthesis (Figure 1, Appendix S3 in the [Online Supplementary Document](#)). The majority (57%) of geospatial studies were national-level studies, while only 9% were regional studies. More than a quarter (30%) of the studies were done in Ethiopia. More than three-fourth (79%) of the studies were on maternal health, followed by neonatal health (12%) and both (9%). Nine in ten (91%) of the studies were cross-sectional, and a half (54%) used Demographic and Health Survey (DHS) data. Global spatial autocorrelation (29%), spatial scan statistics (27%), and different travel time/accessibility measuring techniques (23%) were the most commonly used spatial analysis methods (Table 1 and Figure 2).

Application of geospatial techniques in Sub Saharan Africa

Geospatial analysis was applied for a) identifying inequalities in maternal and neonatal morbidity and mortality, and b) health care utilization; c) identifying gaps in availability and geographic accessibility of maternal health facilities; d) highlighting the effect of availability and accessibility on maternal health care utilization; and e) modelling intervention areas.

Identifying inequalities in maternal and neonatal morbidity and mortality

Three studies [19–21] applied geospatial analysis to describe the distribution of maternal and pregnan-

Table 1. Descriptive summary of the characteristics of geospatial studies in maternal and neonatal morbidity and mortality

		NUMBER	PERCENT			NUMBER	PERCENT
Study period	Before 2000	1	1.8%	Study design	Cross-sectional	51	91.1%
	2000 to 2009	9	16.1%		Cohort	3	5.4%
	2010 to 2014	22	39.3%		Case-control	1	1.8%
	2015 to 2021	24	42.9%		Others	1	1.8%
	Total	56	100%		Total	56	100%
Scope	National	32	57.1%	Analysis method	Global spatial autocorrelation	16	28.6%
	Sub-national	19	33.9%		Spatial scan statistics	15	26.8%
	Regional	5	8.9%		Travel time/accessibility modelling	13	23.2%
	Total	56	100%		Different logistic regression models	11	19.6%
Focus	Maternal	44	78.6%		Hot spot analysis	10	17.9%
	Neonatal	7	12.5%		Spatial interpolation	7	12.5%
	Both	5	8.9%		Different Bayesian models	7	12.5%
	Total	56	100%		Exploratory spatial analysis	6	10.7%
Country	Ethiopia	17	30.4%		Local spatial autocorrelation	3	5.4%
	Ghana	9	16.1%		Incremental spatial autocorrelation	2	3.6%
	Nigeria	6	10.7%		Geographic Weighted Regression	2	3.6%
	Kenya	5	8.9%		Zero-inflated negative binomial model	2	3.6%
	Mozambique	4	7.1%		Anselin Local Moran's I	2	3.6%
	Tanzania	3	5.4%		Linear regression	2	3.6%
	Zambia	2	3.6%	Others	16	28.6%	
	Malawi	2	3.6%	Total†	114	204%	
	Uganda	1	1.8%	Data source	DHS survey	30	53.6%
	South Africa	1	1.8%		Health facility surveys	10	17.9%
	Senegal	1	1.8%		Secondary analysis of RCT	4	7.1%
	Regional *	5	8.9%		Secondary analysis of cohort data	3	5.4%
	Total	56	100%		HDSS	3	5.4%
	Outcome	Skilled birth attendance, ANC, and PNC	32		57.1%	HMIS	3
Access to health care		12	21.4%		WorldPop database	3	5.4%
Neonatal mortality		4	7.1%		Census	3	5.4%
Pregnancy-related mortality		3	5.4%		Primary data collection	3	5.4%
Perinatal mortality		2	3.6%		Road network data	2	3.6%
Abortion		2	3.6%	Others	13	23.2%	
Stillbirth		1	1.8%	Total‡	77	138%	
Total		56	100%				

ANC – antenatal care, PNC – postnatal care, DHS – demographic health survey, RCT – randomized control trial, HMIS – health management information systems, HDSS – health and demographic surveillance system

*Out of the five regional studies, one study in East (ie, Ethiopia, Kenya, Malawi, Mozambique, Tanzania, Zambia, and Zimbabwe) and West Africa (Benin, Burkina Faso, Ghana, Guinea, Mali, Niger, and Nigeria), one study contained 49 and another 48 countries, and the remaining two studies in Mali, Guinea and Liberia, Benin, Niger, and Nigeria.

†The total percentage is more than 100 because some studies used more than one analysis technique.

‡The total percentage is more than 100 because some studies used more than one data source.

cy-related mortality. First, a national study in Zambia [19] found significant differences in the country's lifetime risk of pregnancy-related deaths. This study showed a lower lifetime risk of pregnancy-related death clusters in Lusaka, Muchinga, Copperbelt, Northwestern, and Southern provinces. In contrast, clusters of higher lifetime risk were found in Western and Luapula provinces. Similarly, a study in two rural districts in southern Tanzania [20] showed geographical differences in maternal mortality even within a relatively small geographical area. Conversely, a study in a small geographic area on the west side of South Africa [21] with approximately 11 000 household inhabitants found no evidence of spatial clustering of maternal mortality in the study area.

Seven studies [13,22-27] reported the geographical distribution of stillbirth, perinatal, and neonatal mortality at regional, national, and subnational levels. A large study in all 49 SSA countries indicated no spatial clustering of neonatal mortality between countries [22]. On the other hand, a study in seven East African and seven West African countries found higher neonatal mortality in Ethiopia, Kenya, and Tanzania compared to the rest of the countries [23]. Studies in Malawi [24], Nigeria [25], and Ethiopia [26,27] also found inequalities in perinatal deaths within the countries. For instance, relatively lower perinatal deaths in Ethiopia were found in Addis Ababa, some parts of Tigray, Gambella, Benishangul-Gumuz, and Southern Nations Nationalities




Figure 2. Number of national and subnational geospatial studies conducted on maternal and neonatal health outcomes in each sub-Saharan Africa (SSA) country.

and Peoples Region, whereas significantly higher perinatal deaths were recorded in the Somali region, South Amhara, North Oromia, and Southern Afar [26,27]. For example, in one study, women within a specific hotspot area in the Somali region of Ethiopia had a 22.5 times higher risk of experiencing stillbirth than women outside the hotspot area [27].

Two studies [28,29] which were conducted in Ghana and Ethiopia on abortion, showed a significant variation in abortion rates in those countries [28,29]. In Ghana, there was a distinct north-south difference in the prevalence of abortion, with higher rates in the southern part of the country [28]. Similarly, in Ethiopia, significant spatial variation, where higher abortion was recorded in the northern Tigray region, border areas of Oromia, Amhara, and Southern Nations Nationalities and Peoples Region, was observed [29].

Identifying inequalities in maternal health care utilization

Twenty-seven studies investigated inequalities in maternal health care utilization [30-56]. All reported significant disparities in maternal health care utilization at regional, national, or subnational levels. For example, a regional study among five neighbouring West African countries showed that utilization was significantly higher in some regions of Nigeria and Benin, while the lowest utilization was reported among women residing in some areas of Niger and Mali [30].

Inequality in maternal health care utilization was also identified at a national level. For instance, in Ethiopia, a relatively higher utilization was found in Addis Ababa [31-37], Dire Dawa [31,34-37], Tigray [31,33-39], and Harari [31,33,35,36]. A relatively low utilization was found in regions of Somali [31,33,34,36,37,39], Afar [31,33-35,37], Oromia [32,33,35,36,39], most of Amhara [33-35,37,40], Gambela [32,33,37,39], Benishangul Gumuz [34,35], and parts of Southern Nations Nationalities and Peoples Region [32,34-36,39,40]. Similarly, a significant north-south divide in the likelihood of maternal health care utilization was reported in Nigeria. Most Northern states have a significantly lower maternal health services utilization compared with the southern states [41-45], except for a few southern states (such as part of Cross River [43], Bayelsa [42,43], Imo and Anambra state [45]). The north-south division in the utilization of maternal health care was also evident in Ghana, with women from the north being more likely to utilize maternal health care services [46-49]. However, one study in Ghana reported a significant difference between the southern and northern regions only for skilled birth attendance but not for antenatal care utilization [50]. A study in Uganda also reported marked geographic variation in maternal health care utilization across the country [51].

Most geospatial studies also demonstrated variation in maternal health care services utilization at a subnational level. For example, studies in Ethiopia [52,53], Kenya [54], and Mozambique [55] found subnational clustering of maternal health care utilization at the different administrative levels. Another study that focused on selected districts in Ethiopia's four most populous regions (Tigray, Oromia, Amhara, and Southern Nations Nationalities and Peoples Region) found evidence of geographic clustering in the utilization of skilled birth attendance but not for antenatal care [56].

Identifying gaps in availability and geographic accessibility of health facilities

A regional study from 48 countries [57] reported that all countries meet the geographical accessibility target of 80% of all women located within 2h of the nearest comprehensive emergency obstetric and newborn care (C-EmONC) at the national level except Eritrea (71%). This is with the assumption that all hospitals in SSA have a C-EmONC capacity. On the other hand, a national study in Ghana estimated only 55% of women in Ghana reach their nearest C-EmONC facility within two hours. Furthermore, 29% of Ghanaian women reside

more than four hours from the C-EmONC facility [58]. However, substantial variations were also observed regarding geographic accessibility at the subnational level. The same study that reported all SSA countries meet the geographical accessibility target [57] also observed a substantial variability at subnational levels, with the majority (56%) of the countries having at least one administration unit not meeting the target at administrative level-1 (the largest subnational administrative unit of a country) and 74% at administrative level-2 (the largest subnational administrative unit of a country). Malawi, for example, has 82% of the country's administrative level-2 not meeting the target. Similarly, in Ethiopia, although the national average distance to EmONC facilities was 12.8 km, it varies from as long as 27.1 km in the Somali region to as short as a kilometre in bigger cities of Addis Ababa and Dire Dawa [32,38]. Similar phenomena were observed in Ghana, where 82% of women can access C-EmONC facilities within two hours in the metropolitan region of Greater Accra while only less than a quarter in the two predominantly rural Northern and Western regions [58]. Studies in Kenya, Malawi, Tanzania, Mali, Guinea, and Liberia also reported substantial disparities in access within the country and even at the subnational level [54,59-62].

Looking at the geographic accessibility during inter health facility referral, the median distance between health facilities that can(not) perform caesareans section was 32km (ie, 41-minute driving time) in Senegal [63]. On the contrary, in two regions of Ethiopia (Tigray and Amhara), the average time between the two facilities was as high as two hours [64].

Regarding availability, according to a regional study, all SSA countries meet the World Health Organization (WHO) availability of one C-EmONC facility per 500 000 population targets at the national level, except for Ethiopia (which only completes 77% of the target) and Senegal (96% of the target). In addition, there was an average of two hospitals per 500 000 people across SSA. However, at the subnational level, 58% of countries have at least one administrative unit not meeting this target at administrative level-1 (the largest subnational administrative unit of a country) and nearly all (95%) at administrative level-2 (the largest subnational administrative unit of a country) [57]. Also, in the Upper West region of Ghana, there was approximately one C-EmONC facility per 170 000 populations, thus exceeding the availability target by a factor of three. However, only one-third of women in that region can access these facilities within two hours. Similarly, in the Northern Region of Ghana, where one C-EmONC facility was present per 300 000 population, only a quarter can access a C-EmONC facility within two hours of travel [58].

Highlighting the effect of availability and geographic accessibility on maternal health care utilization

Most studies examining the effect of distance on health care utilization found that the further a woman lives from a health facility, the less likely she is to receive necessary maternal health care [38,53,65-67]. Some studies, however, found a threshold distance beyond which maternal health service utilization ceased to decline with additional distance [67,68]. In contrast, studies in Ghana and western Kenya did not find any correlation between distance to a health facility and maternal health care utilization [46,54].

Also, in two of the reviewed studies, the number of facilities around where a woman lives was strongly associated with maternal health care utilization [55,59]. Another study revealed that the effect of health service availability on maternal health care utilization was much stronger during childbirth than during pregnancy [65].

In addition to the availability and accessibility of health facilities, the quality of the service at the facility strongly influenced maternal health care utilization. Studies in different countries of SSA confirmed that higher service quality at health facilities increases the odds of maternal health service utilization [38,46,51]. However, two studies in different parts of Ghana failed to find such an association [65,68].

Modelling interventions

So far, only two studies [64,69] have used geospatial analysis to identify areas that can benefit most from specific interventions. Both studies, one of which is in Mozambique and the other in Ethiopia, modelled the impact of two interventions (ie, improving transportation and communication and upgrading strategically selected facilities) on the population's access to facilities.

In Mozambique, upgrading 37 geographically strategic health facilities to perform caesarean delivery would result in an additional 4% of the population (equivalent to 968 846 people) reaching a higher health facility within 2 hours. For the remaining population, access would be improved even though it would still take more than 2 hours to get to a higher health facility. Furthermore, when the above intervention was combined with making ambulances/transportation and communication in facilities available where none exist, it reduced mean inter-facility referral travel time from 2.5 hours to 1.8 hours [69].

Similar interventions were modelled to decrease the distance between a higher functioning facility that routinely conducts caesarean deliveries with the rest of the facilities in two regions of Ethiopia. Upgrading only seven selected health facilities improved the population served by first-level facilities located within the 2-hour transfer time to higher facilities from 70% to 80% (ie, translated into serving an additional 3.1 million people). Also, providing each facility with its transportation and communication if one or both were missing would result in 83% of the overall population being served by first-level facilities located within the 2-hour transfer time to higher facilities. Combining the above two interventions would improve the number of women and infants served by a first-level facility located within 2 hours of a facility providing lifesaving obstetric surgery from 70% to 90% (ie, translated into serving an additional 8.6 million populations) [64].

DISCUSSION

To our knowledge, this is the first systematic review to map all studies that have applied geospatial analysis for maternal and neonatal morbidity and mortality in SSA. We found that geospatial analysis was commonly used to identify inequalities in maternal and neonatal morbidity, mortality, and health care utilization and identify gaps in the availability and geographic accessibility of maternal health facilities. In addition, we found few studies that used geospatial analysis for modelling intervention areas.

Of the several applications of geospatial analysis, the most used application was the identification of areas with significantly higher and lower maternal and neonatal morbidity, mortality, and health care utilization (such as antenatal care, skilled birth attendance, and postnatal care). The majority of the studies reported significant disparities in maternal and neonatal morbidity, mortality, and health care utilization at regional, national, or subnational levels. For instance, in Ethiopia, maternal and neonatal morbidity, mortality, and health care utilization were better in urban and semi-urban areas such as Addis Ababa, Dire Dawa, Harari, and Tigray; whereas, in Nigeria and Ghana, most studies reported a significant north-south divide. The findings of those studies could be used as input for prioritizing certain areas for resource allocation during national health planning. In addition, some studies pinpointed small areas sub-nationally with disproportional morbidity, mortality, and health care utilization which can be valuable input for subnational or local health planning according to the extent of the study. However, the availability of reliable, up-to-date, small-scale georeferenced data are still limited. This might require primary georeferenced data collection, which can be costly. Most of the reviewed studies use DHS as a data source. The DHS Program is United States Agency for International Development (USAID) funded nationally representative survey on fertility, family planning, maternal and child health, gender, HIV/AIDS, malaria, and nutrition for over 90 developing countries [70]. The finding that most of the reviewed studies use DHS as a data source is comparable with the finding of other reviews conducted on different study populations [71] and study areas [72]. Despite the large sample size and availability in regular five-year intervals, DHS only contains data aggregated at the survey cluster level [70]. Availability of household or individual level georeferenced data could further enable a more accurate prediction of the inequality with further resolution. In addition, most of the reviewed studies were done in a few countries, leaving parts of SSA not covered at all or only investigated as part of a few regional studies. All the national and subnational geospatial studies on maternal and neonatal health were done in 11 out of the 49 SSA countries. More than a quarter of those national and subnational studies were conducted in one country, Ethiopia. On the other hand, the number of geospatial studies done after 2010 is more than five times the number of studies done between 2000 and 2009, indicating better progress recently.


The other identified application of geospatial studies in maternal and neonatal health was the identification of gaps in maternal health care availability and geographic accessibility. We saw that though overall, women and neonates' geographic access to and availability of health facilities in SSA is low, this varies significantly between different countries in the region and between areas within countries. We also observed how the national summary estimates of availability and geographic accessibility, which are usually the ones used for monitoring purposes by countries, may obscure marked subnational variation. Such findings have two major implications. First, it illustrates that using national summary estimates of availability and geographic accessibility singlehandedly for monitoring purposes could be misleading. Hence, national-level availability indicators, such as the WHO's "at least one is C-EmONC for every 500 000 populations" indicator [73] and geographical accessibility targets such as Global Surgery's "women should be able to reach a lifesaving blood transfusion or caesarean sections within a maximum of two hours of travel" indicator [74], should always be used jointly with the performance of similar indicators sub-nationally. Second, such a finding also indicates which areas in the country are performing better than others regarding the availability and geographic accessibility of maternal health facilities. Such knowledge is essential for planning where to focus scarce re-

sources for improving the availability and geographic accessibility of maternal health facilities. However, we also identified several limitations on geospatial studies on maternal facilities' availability and geographic accessibility. First, most geospatial models used to estimate the travel time to a health facility assumes patients to walk to the nearest road and then continue the journey driving. Yet, walking is the predominant form of transportation in rural Africa due to the lack of infrastructure and motorized transport services [75]. Indeed, a study in Siaya County of Kenya showed that 53 to 87 percent of women walked to the nearest maternal health care facility [61,76]. In addition, the travel speed assumptions used by geospatial models for mechanized transport were generous. For instance, for most studies, travel time was calculated by assuming a driving speed of 100, 50, and 30 km/h for primary, secondary, and tertiary roads, respectively, while a walking speed of 5 km/h was set to areas without a road. Studies conducted in Rwanda [77] and Sierra Leone [78] checked the agreement between travel time reported by the patient and calculated by the geospatial model with the above travel speed assumptions. In both studies, patient-reported travel time was longer than estimated by the geospatial model. As a result, those estimations usually underrate the actual time it takes a patient to reach a health facility. However, when conservative estimates of 50 km/h to primary roads, 20 km/h to secondary roads, and 5 km/h to tertiary roads while walking speed of 1.5-2.5 km/h for all other areas were used, the travel time estimated by the geospatial model was closer to the patient-reported time [78]. Moreover, most geospatial studies ignored seasonal variation when estimating geographic access to health facilities. The speed assumed for studies applies to the dry season. Yet, spatial access to health facilities generally decreases during the wet season for all modes of transport due to the increase in travel times imposed by precipitation and flooding. For instance, in southern Mozambique, at the peak of the dry season, nearly half of women lived within a two-hour travel time to the nearest C-EmONC facility that offers lifesaving maternal care. The population of women dropped twenty-seven percent during the rainy season because of weather-induced increases in travel in travel time. What is more, thirteen of the 417 neighbourhoods were isolated entirely from health facilities at some point [57]. Furthermore, studies about the accessibility of inter-facility referrals were usually estimations of the direct travel times, which did not consider the availability of transportation and communication delays in facilities. For instance, in a remote region in western Tanzania, the estimated median direct travel times between pairs of a facility with and without a caesarean section (ie, not considering the availability of transportation and communication delays) was around 26-minute. However, less than two percent of the referring facilities had an ambulance at the facility. To access the remaining facilities, the patient has to either find a means of transportation herself (53%) or has to wait for an ambulance to come from other facilities such as the district health office (45%) [79]. Also, 75% do not have a mobile phone, a landline (phone/radio receiver), or both. Therefore, median direct travel times between pairs of facilities with and without caesarean section becomes 107 minutes [79].

Modelling and predictions based on the existing spatial data are one application of geospatial techniques. However, our review revealed that this application was not commonly used in SSA. The few studies in this regard quantified the potential effect of new interventions such as improving interfacility transportation and communication, and upgrading strategically selected facilities to enhance access to health facilities [64,69]. In addition to quantifying the potential effect of an intervention on a population, modelling and prediction geospatial studies can be used to indicate what percentage of the population would benefit from a specific intervention, such as building a new health facility or road construction in a specific area. It can also indicate geographic areas that would benefit most from a selected intervention. Such application of geospatial techniques is specifically vital in resource-limited areas like SSA to help policymakers, and health planners make evidence-based decisions. However, more geospatial modelling studies are needed to learn about the specific advantages and limitations of the method.

This review has several strengths and limitations. It is the first study to review the application of geospatial techniques in maternal and neonatal health in SSA. Additionally, in consultation with a medical information specialist, we developed a broader search strategy to minimize the chance of missing relevant studies. Hence, we screened a total of 8184 articles for eligibility. Furthermore, there was no language or time restriction in the inclusion of articles. However, our study also has limitations that should be considered during the interpretation of the results. First, studies included in this review used different geospatial analysis methods; hence they are not exactly comparable to each other. Additionally, we only included articles indexed in PubMed, SCOPUS, EMBASE, and Web of Science databases; hence, there is a possibility that some potentially relevant studies indexed elsewhere may have been missed. We could not manage two databases we planned to search, which were African Journals Online (AJOL) and WHO Global Health Library. WHO Global Health Library was not accessible, while AJOL was difficult to search using Boolean operators. Finally, we could not do a meta-analysis due to the heterogeneity of the studies.

In conclusion, the use of geospatial analysis for maternal and neonatal health in SSA is still limited, and more detailed spatial data are required to exploit the potential of geospatial technologies fully. The available studies used geospatial analysis to identify inequalities in maternal and neonatal morbidity, mortality, and health care utilization and identify gaps in maternal health facilities' availability and geographic accessibility. In addition, we identified a few studies that used geospatial analysis for modelling intervention areas. The findings of those studies could be used as input for planning where to focus the scarce resources by prioritizing areas where maternal and neonatal morbidity and mortality are most likely to occur. On the other hand, we also identified challenges and shortcomings of the reviewed studies. First, the availability of reliable, up-to-date, small-scale georeferenced data are still a challenge. Similarly, women's access to health facilities should be perceived as more complex than the availability of a health facility and the calculated distance to a health facility.



Acknowledgments: We thank Tesfa Habtewold and Melisa Castañeda for their assistance with screening and Karin Sijtsma for her guidance in preparing the search strategy.

Ethical approval: No ethical approval was required for this study

Funding: The authors acknowledge funding from University Medical Center Groningen (UMCG) Ph.D. grant for S.M.A.

Authorship contributions: Conception or design of the work, SMA, AKT, GW, JS and RB; data collection, SMA.; data analysis and interpretation, SMA, AKT, GG, CM, GW, JS and RB; drafting the article, SMA; critical revision of the article, AKT, GG, CM, GW, JS and RB. All authors have read and approved the manuscript prior to submission.

Disclosure of interest: The authors completed the ICMJE Disclosure of Interest F (available upon request from the corresponding author) and disclose no relevant interests.

Additional material

Online Supplementary Document

REFERENCES

- 1 Maternal and newborn health. Available: <https://www.unicef.org/health/maternal-and-newborn-health>. Accessed: 21 September 2021.
- 2 The African Academy of Sciences. From minding the gap to closing the gap: Science to transform maternal and newborn survival and stillbirths in sub-Saharan Africa in the Sustainable Development Goals era. The African Academy of Sciences: Nairobi, Kenya; 2018.
- 3 World Health Organization WB. United Nations Population Fund & United Nations Children's Fund (UNICEF). Trends in maternal mortality: 1990-2015: estimates from WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division. World Health Organization: Geneva; 2015.
- 4 Maternal Health. Available: <https://www.afro.who.int/health-topics/maternal-health>. Accessed: 29 April 2019.
- 5 The Partnership for Maternal Newborn and Child Health. Opportunities for Africa's newborns: Practical data, policy and programmatic support for newborn care in Africa. WHO on behalf of The Partnership for Maternal Newborn and Child Health: Geneva; 2006.
- 6 UN General Assembly. United Nations Millennium Declaration, Resolution Adopted by the General Assembly. General Assembly: New York; 2000.
- 7 United Nations. The Sustainable Development Goals Report 2019. United Nations: New York; 2019.
- 8 United Nations Population Fund. Costing the Three Transformative Results. United Nations: New York; 2020.
- 9 Hofmeyr GJ, Haws RA, Bergström S, Lee AC, Okong P, Darmstadt GL, et al. Obstetric care in low-resource settings: what, who, and how to overcome challenges to scale up? *Int J Gynaecol Obstet.* 2009;107:S21-S44. Medline:19815204 doi:10.1016/j.ijgo.2009.07.017
- 10 Ruhago GM, Ngalesoni FN, Norheim OF. Addressing inequity to achieve the maternal and child health millennium development goals: looking beyond averages. *BMC Public Health.* 2012;12:1119. Medline:23270489 doi:10.1186/1471-2458-12-1119
- 11 Colston JM, Burgert CR. Using Geospatial Analysis to Inform Decision Making In Targeting Health Facility-Based Programs. MEASURE Evaluation: Chapel Hill; 2014.
- 12 Tatem AJ. WorldPop, open data for spatial demography. *Sci Data.* 2017;4:170004. Medline:28140397 doi:10.1038/sdata.2017.4
- 13 Banda M, Kazembe L, Lewycka S, King C, Phiri T, Masache G, et al. Spatial modelling of perinatal mortality in Mchinji, Malawi. *Spat Spatiotemporal Epidemiol.* 2016;16:50-8. Medline:26919755 doi:10.1016/j.sste.2015.11.004
- 14 Sabde Y, De Costa A, Diwan V. A spatial analysis to study access to emergency obstetric transport services under the public private "Janani Express Yojana" program in two districts of Madhya Pradesh, India. *Reprod Health.* 2014;11:57. Medline:25048795 doi:10.1186/1742-4755-11-57
- 15 Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6:e1000097. Medline:19621072 doi:10.1371/journal.pmed.1000097
- 16 Babineau J. Product review: covidence (systematic review software). *J Can Health Libr Assoc.* 2014;35:68-71. doi:10.5596/c14-016

- 17 Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med*. 2005;37:360-3. Medline:15883903
- 18 Institute JB. Joanna Briggs Institute reviewers' manual: 2014 edition. Australia: The Joanna Briggs Institute. 2014.
- 19 Banda R, Sandøy IF, Fylkesnes K, Janssen F. Lifetime risk of pregnancy-related death among Zambian women: district-level estimates from the 2010 census. *J Popul Res (Canberra)*. 2016;33:263-81. doi:10.1007/s12546-016-9172-1
- 20 Manyeh AK, Nathan R, Nelson G. Maternal mortality in ifakara health and demographic surveillance system: Spatial patterns, trends and risk factors, 2006 - 2010. *PLoS One*. 2018;13:e0205370. Medline:30346950 doi:10.1371/journal.pone.0205370
- 21 Tlou B, Sartorius B, Tanser F. Space-time patterns in maternal and mother mortality in a rural South African population with high HIV prevalence (2000-2014): Results from a population-based cohort. *BMC Public Health*. 2017;17:543. Medline:28578674 doi:10.1186/s12889-017-4463-9
- 22 Kayode GA, Grobbee DE, Amoakoh-Coleman M, Ansah E, Uthman OA, Klipstein-Grobusch K. Variation in neonatal mortality and its relation to country characteristics in sub-Saharan Africa: An ecological study. *BMJ Glob Health*. 2017;2:e000209. Medline:29104766 doi:10.1136/bmjgh-2016-000209
- 23 Grady SC, Frake AN, Zhang Q, Bene M, Jordan DR, Vertalka J, et al. Neonatal mortality in East Africa and West Africa: A geographic analysis of district-level demographic and health survey data. *Geospat Health*. 2017;12:501. Medline:28555482 doi:10.4081/gh.2017.501
- 24 Kazembe LN, Mpeketula PMG. Quantifying spatial disparities in neonatal mortality using a structured additive regression model. *PLoS One*. 2010;5:e11180. Medline:20567519 doi:10.1371/journal.pone.0011180
- 25 Adebayo SB, Fahrmeir L, Klasen S. Analyzing infant mortality with geospatial categorical regression models: A case study for Nigeria. *Econ Hum Biol*. 2004;2:229-44. Medline:15464004 doi:10.1016/j.ehb.2004.04.004
- 26 Yadeta TA, Mengistu B, Gobena T, Regassa LD. Spatial pattern of perinatal mortality and its determinants in Ethiopia: Data from Ethiopian Demographic and Health Survey 2016. *PLoS One*. 2020;15:e0242499. Medline:33227021 doi:10.1371/journal.pone.0242499
- 27 Tesema GA, Gezie LD, Nigatu SG. Spatial distribution of stillbirth and associated factors in Ethiopia: A spatial and multilevel analysis. *BMJ Open*. 2020;10:e034562. Medline:33115888 doi:10.1136/bmjopen-2019-034562
- 28 Owoo NS, Lambon-Quayefio MP, Onuoha N. Abortion experience and self-efficacy: Exploring socioeconomic profiles of GHANAIAN women. *Reprod Health*. 2019;16:117. Medline:31349789 doi:10.1186/s12978-019-0775-9
- 29 Tesema GA, Mekonnen TH, Teshale AB. Spatial distribution and determinants of abortion among reproductive age women in Ethiopia, evidence from Ethiopian demographic and health survey 2016 data: Spatial and mixed-effect analysis. *PLoS One*. 2020;15:e0235382. Medline:32598398 doi:10.1371/journal.pone.0235382
- 30 Gayawan E, Omolofe OT. Analyzing Spatial Distribution of Antenatal Care Utilization in West Africa Using a Geo-additive Zero-Inflated Count Model. *Spat Demogr*. 2016;4:245-62. doi:10.1007/s40980-016-0027-3
- 31 Yeneneh A, Alemu K, Dadi AF, Alamirrew A. Spatial distribution of antenatal care utilization and associated factors in Ethiopia: Evidence from Ethiopian demographic health surveys. *BMC Pregnancy Childbirth*. 2018;18:242. Medline:29914403 doi:10.1186/s12884-018-1874-2
- 32 Tegegne TK, Chojenta C, Getachew T, Smith R, Loxton D. Antenatal care use in Ethiopia: A spatial and multilevel analysis. *BMC Pregnancy Childbirth*. 2019;19:399. Medline:31675918 doi:10.1186/s12884-019-2550-x
- 33 Tessema ZT, Anmut Y. Spatial distribution and determinants of an optimal ANC visit among pregnant women in Ethiopia: Further analysis of 2016 Ethiopian demographic health survey. *BMC Pregnancy Childbirth*. 2020;20:137. Medline:32131759 doi:10.1186/s12884-020-2795-4
- 34 Tessema ZT, Akalu TY. Spatial Pattern and Associated Factors of ANC Visits in Ethiopia: Spatial and Multilevel Modeling of Ethiopian Demographic Health Survey Data. *Adv Prev Med*. 2020;2020:4676591. Medline:32922999 doi:10.1155/2020/4676591
- 35 Muluneh AG, Anmut Y, Ayele TA. Spatial clustering and determinants of home birth after at least one antenatal care visit in Ethiopia: Ethiopian demographic and health survey 2016 perspective. *BMC Pregnancy Childbirth*. 2020;20:97. Medline:32046677 doi:10.1186/s12884-020-2793-6
- 36 Teshale AB, Alem AZ, Yeshaw Y, Kebede SA, Liyew AM, Tesema GA, et al. Exploring spatial variations and factors associated with skilled birth attendant delivery in Ethiopia: Geographically weighted regression and multilevel analysis. *BMC Public Health*. 2020;20:1444. Medline:32977789 doi:10.1186/s12889-020-09550-3
- 37 Tesema GA, Mekonnen TH, Teshale AB. Individual and community-level determinants, and spatial distribution of institutional delivery in Ethiopia, 2016: Spatial and multilevel analysis. *PLoS One*. 2020;15: e0242242. Medline:33180845 doi:10.1371/journal.pone.0242242
- 38 Tegegne TK, Chojenta C, Getachew T, Smith R, Loxton D. Giving birth in Ethiopia: a spatial and multilevel analysis to determine availability and factors associated with healthcare facility births. *BJOG*. 2020;127:1537-46. Medline:32339407 doi:10.1111/1471-0528.16275
- 39 Sisay MM, Geremew TT, Demlie YW, Alem AT, Beyene DK, Melak MF, et al. Spatial patterns and determinants of postnatal care use in Ethiopia: Findings from the 2016 demographic and health survey. *BMJ Open*. 2019;9: e025066. Medline:31189672 doi:10.1136/bmjopen-2018-025066
- 40 Tessema ZT, Tiruneh SA. Spatio-temporal distribution and associated factors of home delivery in Ethiopia. Further multilevel and spatial analysis of Ethiopian demographic and health surveys 2005-2016. *BMC Pregnancy Childbirth*. 2020;20:342. Medline:32493302 doi:10.1186/s12884-020-02986-w
- 41 Gayawan E. A poisson regression model to examine spatial patterns in antenatal care utilisation in Nigeria. *Popul Space Place*. 2014;20:485-97. doi:10.1002/psp.1775
- 42 Gayawan E. Spatial analysis of choice of place of delivery in Nigeria. *Sex Reprod Healthc*. 2014;5:59-67. Medline:24814440 doi:10.1016/j.srhc.2014.01.004

- 43 Wong KLM, Radovich E, Owolabi OO, Campbell OMR, Brady OJ, Lynch CA, et al. Why not? Understanding the spatio clustering of private facility-based delivery and financial reasons for homebirths in Nigeria. *BMC Health Serv Res.* 2018;18:397. doi:10.1186/s12913-018-3225-4
- 44 Adedokun ST, Uthman OA. Women who have not utilized health Service for Delivery in Nigeria: Who are they and where do they live? *BMC Pregnancy Childbirth.* 2019;19:93. Medline:30866841 doi:10.1186/s12884-019-2242-6
- 45 Ononokpono DN, Gayawan E, Adedini SA. Regional variations in the use of postnatal care in Nigeria: a spatial analysis. *Women Health.* 2020;60:440-55. Medline:31328689 doi:10.1080/03630242.2019.1643816
- 46 Dotse-Gborgbortsi W, Tatem AJ, Alegana V, Utazi CE, Ruktanonchai CW, Wright J. Spatial inequalities in skilled attendance at birth in Ghana: a multilevel analysis integrating health facility databases with household survey data. *Trop Med Int Health.* 2020;25:1044-54. Medline:32632981 doi:10.1111/tmi.13460
- 47 Johnson FA, Padmadas SS, Brown JJ. On the spatial inequalities of institutional versus home births in Ghana: A multilevel analysis. *J Community Health.* 2009;34:64-72. Medline:18830808 doi:10.1007/s10900-008-9120-x
- 48 Dankwah E, Feng C, Kirychuck S, Zeng W, Lepnum R, Farag M. Assessing the contextual effect of community in the utilization of postnatal care services in Ghana. *BMC Health Serv Res.* 2021;21:40. Medline:33413362 doi:10.1186/s12913-020-06028-1
- 49 Bosomprah S, Dotse-Gborgbortsi W, Aboagye P, Matthews Z. Use of a spatial scan statistic to identify clusters of births occurring outside Ghanaian health facilities for targeted intervention. *Int J Gynaecol Obstet.* 2016;135:221-4. Medline:27527530 doi:10.1016/j.ijgo.2016.04.016
- 50 Iyanda AE, Oppong JR, Hamilton P, Tiwari C. Using GIS to detect cluster and spatial disparity in maternal health indicators: a need for social health interventions. *Soc Work Public Health.* 2018;33:449-66. Medline:30426852 doi:10.1080/19371918.2018.1543628
- 51 Sprague DA, Jeffery C, Crossland N, House T, Roberts GO, Vargas W, et al. Assessing delivery practices of mothers over time and over space in Uganda, 2003-2012. *Emerg Themes Epidemiol.* 2016;13:9. Medline:27307784 doi:10.1186/s12982-016-0049-8
- 52 Kurji J, Talbot B, Bulcha G, Bedru KH, Morankar S, Gebretsadik LA, et al. Uncovering spatial variation in maternal healthcare service use at subnational level in Jimma Zone, Ethiopia. *BMC Health Serv Res.* 2020;20:703. Medline:32736622 doi:10.1186/s12913-020-05572-0
- 53 Nigatu AM, Gelaye KA, Degefe DT, Birhanu AY. Spatial variations of women's home delivery after antenatal care visits at lay Gayint District, Northwest Ethiopia. *BMC Public Health.* 2019;19:677. Medline:31159775 doi:10.1186/s12889-019-7050-4
- 54 Prudhomme O'Meara W, Platt A, Naanyu V, Cole D, Ndege S. Spatial autocorrelation in uptake of antenatal care and relationship to individual, household and village-level factors: Results from a community-based survey of pregnant women in six districts in western Kenya. *Int J Health Geogr.* 2013;12:55. Medline:24314170 doi:10.1186/1476-072X-12-55
- 55 Agadjanian V, Yao J, Hayford SR. Place, time and experience: Barriers to universalization of institutional child delivery in rural Mozambique. *Int Perspect Sex Reprod Health.* 2016;42:21-31. Medline:28770025 doi:10.1363/42e0116
- 56 Defar A, Okwaraji YB, Tigabu Z, Persson LÅ, Alemu K. Geographic differences in maternal and child health care utilization in four Ethiopian regions; A cross-sectional study. *Int J Equity Health.* 2019;18:173. Medline:31718658 doi:10.1186/s12939-019-1079-y
- 57 Wigley AS, Tejedor-Garavito N, Alegana V, Carioli A, Ruktanonchai CW, Pezzulo C, et al. Measuring the availability and geographical accessibility of maternal health services across sub-Saharan Africa. *BMC Med.* 2020;18:237. Medline:32895051 doi:10.1186/s12916-020-01707-6
- 58 Gething PW, Johnson FA, Frempong-Ainguah F, Nyarko P, Baschieri A, Aboagye P, et al. Geographical access to care at birth in Ghana: A barrier to safe motherhood. *BMC Public Health.* 2012;12:991. Medline:23158554 doi:10.1186/1471-2458-12-991
- 59 Gao X, Kelley DW. Understanding how distance to facility and quality of care affect maternal health service utilization in Kenya and haiti: A comparative geographic information system study. *Geospat Health.* 2019;14:92-102. Medline:31099519 doi:10.4081/gh.2019.690
- 60 Chen YN, Schmitz MM, Serbanescu F, Dynes MM, Maro G, Kramer MR. Geographic Access Modeling of Emergency Obstetric and Neonatal Care in Kigoma Region, Tanzania: Transportation Schemes and Programmatic Implications. *Glob Health Sci Pract.* 2017;5:430-45. Medline:28839113 doi:10.9745/GHSP-D-17-00110
- 61 Ouko JJO, Gachari MK, Sichangi AW, Alegana V. Geographic information system-based evaluation of spatial accessibility to maternal health facilities in Siaya County, Kenya. *Geogr Res.* 2019;57:286-98. doi:10.1111/1745-5871.12339
- 62 Ononokpono DN, Baffour B, Richardson A. Mapping maternal healthcare access in selected west african countries. *Etude Popul Afr.* 2020;34:5082-105.
- 63 Cavallaro FL, Benova L, Dioukhane E, Wong K, Sheppard P, Faye A, et al. What the percentage of births in facilities does not measure: readiness for emergency obstetric care and referral in Senegal. *BMJ Glob Health.* 2020;5: e001915. Medline:32201621 doi:10.1136/bmjgh-2019-001915
- 64 Bailey PE, Keyes EB, Parker C, Abdullah M, Kebede H, Freedman L. Using a GIS to model interventions to strengthen the emergency referral system for maternal and newborn health in Ethiopia. *Int J Gynaecol Obstet.* 2011;115:300-9. Medline:21982854 doi:10.1016/j.ijgo.2011.09.004
- 65 Masters SH, Burstein R, Amofah G, Aboagye P, Kumar S, Hanlon M. Travel time to maternity care and its effect on utilization in rural Ghana: A multilevel analysis. *Soc Sci Med.* 2013;93:147-54. Medline:23906132 doi:10.1016/j.socscimed.2013.06.012
- 66 Nesbitt RC, Gabrysch S, Laub A, Soremekun S, Manu A, Kirkwood BR, et al. Methods to measure potential spatial access to delivery care in low- and middle-income countries: A case study in rural Ghana. *Int J Health Geogr.* 2014;13:25. Medline:24964931 doi:10.1186/1476-072X-13-25
- 67 Mwaliko E, Downing R, O'Meara W, Chelagat D, Obala A, Downing T, et al. "not too far to walk": The influence of distance on place of delivery in a western Kenya health demographic surveillance system. *BMC Health Serv Res.* 2014;14:212. Medline:24884489 doi:10.1186/1472-6963-14-212

- 68 Dotse-Gborgbortsi W, Dwomoh D, Alegana V, Hill A, Tatem AJ, Wright J. The influence of distance and quality on utilisation of birthing services at health facilities in Eastern Region, Ghana. *BMJ Glob Health*. 2020;4:e002020. Medline:32154031 doi:10.1136/bmjgh-2019-002020
- 69 Keyes EB, Parker C, Zissette S, Bailey PE, Augusto O. Geographic access to emergency obstetric services: A model incorporating patient bypassing using data from Mozambique. *BMJ Glob Health*. 2019;4:e000772. Medline:31321090 doi:10.1136/bmjgh-2018-000772
- 70 Rutstein SO, Rojas G. *Guide to DHS statistics*. Calverton, MD: ORC Macro. 2006;38.
- 71 Manda S, Haushona N, Bergquist R. A Scoping Review of Spatial Analysis Approaches Using Health Survey Data in Sub-Saharan Africa. *Int J Environ Res Public Health*. 2020;17:3070. Medline:32354095 doi:10.3390/ijerph17093070
- 72 Ferreira LZ, Blumenberg C, Utazi CE, Nilsen K, Hartwig FP, Tatem AJ, et al. Geospatial estimation of reproductive, maternal, newborn and child health indicators: a systematic review of methodological aspects of studies based on household surveys. *Int J Health Geogr*. 2020;19:41. Medline:33050935 doi:10.1186/s12942-020-00239-9
- 73 Bailey P, Lobis S, Maine D, Fortney JA. *Monitoring emergency obstetric care: a handbook*. World Health Organization; 2009.
- 74 Meara JG, Leather AJ, Hagander L, Alkire BC, Alonso N, Ameh EA, et al. Global Surgery 2030: evidence and solutions for achieving health, welfare, and economic development. *Lancet*. 2015;386:569-624. Medline:25924834 doi:10.1016/S0140-6736(15)60160-X
- 75 United Nations Economic Commission for Africa. Sixth Session of the Committee on Food Security and Sustainable Development (CFSSD-6)/Regional Implementation Meeting (RIM) for CSD-18: Africa Review Report on Transport. United Nations Economic Commission for Africa: Addis Ababa, Ethiopia; 2009.
- 76 Kohler PK, Akullian A, Okanda J, Otieno G, Kinuthia J, Voss J, et al. Distance to HIV and Antenatal Care: A Geospatial Analysis in Siaya County, Kenya. *J Assoc Nurses AIDS Care*. 2019;30:548-55. Medline:30694879 doi:10.1097/JNC.000000000000050
- 77 Rudolfson N, Gruendl M, Nkurunziza T, Kateera F, Sonderman K, Nihiwacu E, et al. Validating the Global Surgery Geographical Accessibility Indicator: Differences in Modeled Versus Patient-Reported Travel Times. *World J Surg*. 2020;44:2123-30. Medline:32274536 doi:10.1007/s00268-020-05480-8
- 78 van Duinen AJ, Adde HA, Fredin O, Holmer H, Hagander L, Koroma AP, et al. Travel time and perinatal mortality after emergency caesarean sections: An evaluation of the 2-hour proximity indicator in Sierra Leone. *BMJ Glob Health*. 2020;5:e003943. Medline:33355267 doi:10.1136/bmjgh-2020-003943
- 79 Schmitz MM, Serbanescu F, Arnott GE, Dynes M, Chaote P, Msuya AA, et al. Referral transit time between sending and first-line receiving health facilities: A geographical analysis in Tanzania. *BMJ Glob Health*. 2020;4:e001568. Medline:31478017 doi:10.1136/bmjgh-2019-001568