

University of Groningen

Hand function in patients with distal radius fractures after home-based kinaesthetic motor imagery training

Broekstra, Dieuwke C; Mouton, Leonora J; van der Sluis, Corry K; IJpma, Frank F A; Stenekes, Martin W

Published in:
Journal of Hand Surgery-British and European Volume

DOI:
[10.1177/17531934221075945](https://doi.org/10.1177/17531934221075945)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Broekstra, D. C., Mouton, L. J., van der Sluis, C. K., IJpma, F. F. A., & Stenekes, M. W. (2022). Hand function in patients with distal radius fractures after home-based kinaesthetic motor imagery training. *Journal of Hand Surgery-British and European Volume*, 47(2). <https://doi.org/10.1177/17531934221075945>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Hand function in patients with distal radius fractures after home-based kinaesthetic motor imagery training

Dear Editor,

The effect of limb immobilization on the brain after a distal radial fracture (DRF) might be partly responsible for postinjury functional impairment. Conversely, motor imagery (MI) might diminish neural changes due to immobilization (Meugnot et al., 2015). This (non-blinded) randomized-controlled trial assessed whether women with a DRF who performed kinaesthetic MI training during cast immobilization had better short-term functional outcomes than controls receiving cast immobilization only.

For this study, conducted between 2011 and 2020 (ethically approved METc2011/102), eligible patients were women (45–75 years) with a DRF treated with closed reduction and cast immobilization, without pre-existent upper extremity disorders and with sufficient MI ability (score on Vividness of Motor Imagery Questionnaire ≤ 72 points). They were randomly allocated to the MI or control group. We did not inform the controls about MI. A sample size calculation yielded a total sample size of 52 participants using the F-test family (repeated measures analysis of variance [RM ANOVA], between-within interaction): effect size 0.2, alpha 5%, power 80%.

One week after cast application (T0), participants in the MI group received verbal and written instructions about the MI protocol, while controls only received general information. Cast immobilization was continued for 3–5 weeks (median 4.0, Table 1). All participants were contacted by phone once a week to monitor their recovery and compliance to the MI training. The first measurements (T1) took place directly after cast removal, and the second (T2) 2 weeks later.

All participants received finger mobilization instructions. Only MI participants were instructed to perform kinaesthetic MI at home four times each day, between T0–T1. MI consisted of 10 repetitions of

imagining: (1) making a fist and slowly knocking on the table making exaggerated movements, mimicking flexion-extension; (2) a horizontal line with two dots at each end drawn on the tabletop, and that you touch the dots with the thumb and little finger, mimicking radial-ulnar deviation; (3) turning a key in a keyhole, mimicking pronation-supination. This set was repeated three times. Participants registered the number of times they performed MI training.

The Patient-Rated Wrist Evaluation (PRWE) was the primary outcome variable. Secondary outcomes were strength, range of motion (ROM) and pain. We measured strengths using a digital Jamar and pinch grip dynamometer (H500 Hand Kit, Biometrics Ltd, Newport, UK). These measurements included grip force, three-jaw chuck pinch, key pinch and two-point pinch. The averages of three measurements were used. Active flexion, extension, radial deviation, ulnar deviation, pronation and supination were measured once using a digital goniometer (R500 Range of Motion Kit, Biometrics Ltd, Newport, UK). Strength and ROM were measured in the non-affected hand as reference. Finally, pain intensity and relief were measured using visual analogue scales (VAS).

Differences between the two groups on PRWE scores, Δ strength and Δ ROM (unaffected-affected) were analysed by (robust) linear regression, with the variables mentioned above as outcome and group as a predictor (intention-to-treat). The analyses were adjusted as the MI group had their dominant hands affected more often. Mann-Whitney *U*-tests were applied to determine differences in VAS pain and relief scores.

Forty-three out of 66 patients giving consent were included: 24 in the MI and 19 in the control group. The main reasons for exclusion were surgery and drop-out before the study started. Study groups did not differ regarding patient and fracture characteristics, except that more dominant hands were affected in the MI group (75% versus 53%, Table 1). Descriptive statistics for grip strength and ROM at T1 and T2 (raw scores instead of change scores) are presented in Online Table S1.

Table 1. Comparison of the MI and control group on baseline characteristics (T0), PRWE, VAS, ROM, grip and pinch strength at T1 and T2.

	T0				T1					T2				
	MI	N	Control	N	MI	N	Control	N	p-value	MI	N	Control	N	p-value
Age, years (mean, SD)	58.7 (7.5)	24	58.8 (6.7)	19										
Immobilization time, weeks (median, IQR)	4.0 (4.0–4.0)	24	4.0 (4.0–4.0)	19										
Time T1–T2, days (median, IQR)	14.0 (14.0–19.3)	24	14.0 (14.0–23.0)	19										
Fracture type (n, %)		24		19										
Extra-articular	18 (75%)		16 (84%)											
Intra-articular	6 (25%)		3 (16%)											
AO classification	—	24		19										
23A	16 (67%)		14 (74%)											
23B	2 (8%)		2 (11%)											
23C	6 (25%)		3 (16%)											
Dominant hand affected (n, %)	18 (75%)	24	10 (53%)	19										
Hand dominance (n, %)		24		19										
Right	20 (83%)		17 (89%)											
Left	3 (13%)		2 (11%)											
Bimanual	1 (4%)		0 (0%)											
MI adherence ^a , % (median, IQR)	92 (85–95)		NA											
PRWE score (mean, SD)	48.0 (23.2)	20	44.3 (19.4)	15	0.450	35.3 (20.7)	20	32.4 (19.2)	19	0.917				
VAS Pain score ^b (median, IQR)	4.0 (2.0–18.0)	23	10.0 (2.5–23.5)	19	0.477	8.5 (3.3–15.3)	22	9.0 (3.0–13.5)	19	0.990				
VAS Pain relief score ^b (median, IQR)	78 (49.0–90.0)	23	69.0 (48.0–90.5)	19	0.742	81.5 (37.0–86.3)	22	67.0 (35.0–81.0)	19	0.346				
Δ ROM ^c , °(mean, SD)														
Extension	26 (17)	21	35 (14)	19	0.082	14 (13)	22	18 (16)	18	0.377				
Flexion	36 (21)	21	39 (12)	19	0.740	22 (15)	22	16 (22)	19	0.627				
Pronation	12 (14)	21	8 (20)	18	0.500	3 (13)	21	4 (12)	19	0.688				
Supination	26 (26)	21	24 (22)	18	0.885	21 (23)	21	16 (22)	19	0.469				
Radial deviation	7 (8)	21	6 (7)	19	0.332	4 (7)	22	3 (5)	19	0.289				
Ulnar deviation	13 (8)	21	9 (6)	19	0.077	8 (8)	22	9 (6)	19	0.519				
Δ Strength ^c , kg (mean, SD)														
Power grip	15.2 (5.7)	21	15.6 (3.6)	16	0.685	10.5 (5.8)	22	11.5 (5.0)	19	0.346				
Key pinch	2.6 (1.2)	21	2.9 (1.3)	18	0.390	1.7 (1.2)	22	2.2 (1.5)	19	0.163				
Three-jaw pinch	3.6 (1.7)	21	3.4 (1.2)	17	0.897	2.2 (1.4)	22	2.4 (1.5)	19	0.428				
Two-point pinch	2.2 (1.0)	21	1.9 (0.9)	17	0.492	1.0 (1.2)	22	1.4 (1.1)	19	0.085				

MI: motor imagery; N: number; SD: standard deviation; IQR: interquartile range; PRWE: Patient-Rated Wrist Evaluation; VAS: visual analogue scale; AO: Arbeitsgemeinschaft für Osteosynthesefragen; ROM: range of motion.

^aAdherence was calculated as the actual number of MI sessions, divided by the number of training sessions that should have been done according to protocol.

^bTested with the Mann–Whitney *U*-test.

^cFor range of motion and strength, the differences between the unaffected and affected side are reported; a lower score representing better outcome.

No significant differences between groups were found in any outcome variables at T1 and T2 (Table 1), which indicates that MI did not influence functional outcomes or pain within 2 weeks after immobilization.

A limitation is the limited number of participants caused by our institution's change in policy to shift focus toward complex care. The absence of statistically significant effects does not necessarily mean that no effects of MI exist. However, the 95% confidence intervals of the regression coefficient of the PRWE did not contain the minimal clinically important change reported in similar populations (McCreary et al., 2020). So, we are 95% confident that in the population from which we recruited our sample, the difference between the groups is so small that it is not meaningful to the patient. Another limitation lies in the moment of inclusion. Due to the acute nature of the injury, patients could only be included 1 week after the immobilization started. Since cortical reorganization occurs within several days (Meugnot et al., 2015), the most considerable cortical changes might already have taken place before inclusion, limiting the effectiveness of the MI training. Finally, there is no consensus about the best MI protocol. Different protocols (Online Table S2) might be responsible for contrasting findings reported in the literature (Dilek et al., 2018; Korbus and Schott, 2020).


Declaration of conflicting interests The authors declare no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding The authors received no financial support for the research, authorship, and/or publication of this article.

Ethical approval This study and amendments were reviewed and approved by the Institutional Review Board (METc2011/102), and conforms to the Declaration of Helsinki.

Informed consent All participants gave written informed consent.


Trial registration number This study has been registered on August 13, 2013 in Clinicaltrials.gov, number NCT01921062.

ORCID iD Diewuke C Broekstra  <https://orcid.org/0000-0002-7134-7007>

Supplemental material Supplemental material for this article is available online.

References

- Dilek B, Ayhan C, Yagci G, Yakut Y. Effectiveness of the graded motor imagery to improve hand function in patients with distal radius fracture: a randomised controlled trial. *J Hand Ther.* 2018, 31: 2–9.e1.
- Korbus H, Schott N. Does mental practice or mirror therapy help prevent functional loss after distal radius fracture? A randomised controlled trial. *J Hand Ther.* 2020, 24: S0894-1130(20)30207–6.
- McCreary DL, Sandberg BC, Bohn DC, Parikh HR, Cunningham BP. Interpreting patient-reported outcome results: is one minimum clinically important difference really enough? *Hand (NY).* 2020, 15: 360–4.
- Meugnot A, Agbangla NF, Almecija Y, Toussaint L. Motor imagery practice may compensate for the slowdown of sensorimotor processes induced by short-term upper-limb immobilisation. *Psychol Res.* 2015, 79: 489–99.

Diewuke C. Broekstra^{1,*} , **Leonora J. Mouton²,**
Corry K. van der Sluis³, **Frank F. A. IJpma⁴** and
Martin W. Steneker¹

¹*Department of Plastic Surgery, University Medical Center Groningen, Groningen, The Netherlands*

²*Department of Human Movement Sciences, University Medical Center Groningen, Groningen, The Netherlands*

³*Department of Rehabilitation Medicine, University Medical Center Groningen, Groningen, The Netherlands*

⁴*Department of Trauma Surgery, University Medical Center Groningen, Groningen, The Netherlands*

*Corresponding author: d.c.broekstra@umcg.nl

Twitter: @DCBroekstra

© The Author(s) 2022

Article reuse guidelines:

sagepub.com/journals-permissions

doi: 10.1177/17531934221075945 available online at <http://jhs.sagepub.com>