

University of Groningen

The influence of the dietary exposome on oxidative stress in pregnancy complications

Prins, Jelmer R; Schoots, Mirthe H; Wessels, Jule I; Campmans-Kuijpers, Marjo J E; Navis, Gerjan J; van Goor, Harry; Robertson, Sarah A; van der Beek, Eline M; Sobrevia, Luis; Gordijn, Sanne J

Published in:
Molecular Aspects of Medicine

DOI:
[10.1016/j.mam.2022.101098](https://doi.org/10.1016/j.mam.2022.101098)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Prins, J. R., Schoots, M. H., Wessels, J. I., Campmans-Kuijpers, M. J. E., Navis, G. J., van Goor, H., Robertson, S. A., van der Beek, E. M., Sobrevia, L., & Gordijn, S. J. (2022). The influence of the dietary exposome on oxidative stress in pregnancy complications. *Molecular Aspects of Medicine*, 87, [101098]. <https://doi.org/10.1016/j.mam.2022.101098>

Copyright


Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



The influence of the dietary exposome on oxidative stress in pregnancy complications

Jelmer R. Prins^{a,*}, Mirthe H. Schoots^{b,1}, Jule I. Wessels^a, Marjo J.E. Campmans-Kuijpers^c, Gerjan J. Navis^d, Harry van Goor^b, Sarah A. Robertson^e, Eline M. van der Beek^{f,g}, Luis Sobrevia^{b,h,i,j,k,1}, Sanne J. Gordijn^a

^a Department of Obstetrics and Gynaecology, University Medical Center Groningen (UMCG), University of Groningen, Groningen, the Netherlands

^b Department of Pathology and Medical Biology, Division of Pathology, UMCG, University of Groningen, Groningen, the Netherlands

^c Department of Gastroenterology and Hepatology, UMCG, University of Groningen, Groningen, the Netherlands

^d Department of Medicine, Division of Nephrology, UMCG, University of Groningen, Groningen, the Netherlands

^e Robinson Research Institute and Adelaide Medical School, University of Adelaide, Adelaide, Australia

^f Department of Pediatrics, UMCG, University of Groningen, Groningen, the Netherlands

^g Nestlé Institute of Health Sciences, Nestlé Research, Lausanne, Switzerland

^h Cellular and Molecular Physiology Laboratory (CMPL), Department of Obstetrics, Division of Obstetrics and Gynaecology, School of Medicine, Faculty of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile

ⁱ Department of Physiology, Faculty of Pharmacy, Universidad de Sevilla, Seville, Spain

^j Medical School (Faculty of Medicine), Sao Paulo State University (UNESP), Brazil

^k University of Queensland Centre for Clinical Research (UQCCR), Faculty of Medicine and Biomedical Sciences, University of Queensland, Queensland, Australia

¹ Tecnológico de Monterrey, Eutra, The Institute for Obesity Research (IOR), School of Medicine and Health Sciences, Monterrey, Nuevo León, Mexico

ARTICLE INFO

Keywords:

Exposome
Pregnancy
Oxidative stress
Inflammation
Diet
Placenta

ABSTRACT

Pregnancy complications including fetal growth restriction, preeclampsia, and preterm birth, as well as gestational diabetes, affect one in every four to five pregnancies. Accumulating evidence indicates that increased production of reactive oxygen species accompanies these complications. Given that reactive oxygen species are cell stress-inducing agents, they may have a causal role in disease pathophysiology, although the exact mechanisms by which they contribute to pregnancy complications are not completely understood. Since many environmental and lifestyle factors and exposures are known to modulate reactive oxygen species production, the exposome of pregnant women could contribute to increased generation of reactive oxygen species. The objective of this review is to provide a comprehensive overview of the endogenous and exogenous exposome factors that regulate reactive species in healthy and complicated pregnancies. We also provide a description of dietary interventions aimed at the reduction of reactive species in order to attenuate adverse pregnancy outcome. Dietary interventions in general hold minimal risk in pregnancy and could therefore be considered a promising therapeutic approach.

1. Introduction

The perinatal period is of cardinal importance for the lifelong health of every individual. Suboptimal fetal circumstances encompass risk factors for developing chronic disease later in life. Diabetes, hypertension, stroke and coronary artery disease, for instance, have all been linked to suboptimal perinatal outcomes including placental malfunction (Hoffman et al., 2021). Preeclampsia, fetal growth restriction

(FGR), and preterm labour, common manifestations of the “great obstetrical syndromes”, have common aetiologies (Brosens et al., 2011). These conditions are each strongly linked to the growth and functional development of the placenta, and exhibit a long subclinical phase, becoming evident only when pregnancy progresses to the stage that the compensatory mechanisms can no longer be sustained. This subclinical phase provides a window of opportunity for potential interventions to protect against the later onset of overt symptoms.

* Corresponding author.

E-mail address: j.r.prins@umcg.nl (J.R. Prins).

¹ These authors contributed equally.

The term “great obstetrical syndromes” (GOS) was first used to describe pregnancy-related disorders with a placental component as part of their aetiology. GOS refers to preterm labour and premature rupture of membranes, preeclampsia, spontaneous pregnancy loss, stillbirth, and abnormal fetal growth (Brosens et al., 2011). GOS is steadily rising worldwide and around 15% of all pregnancies are complicated by GOS (Mastrolia and Cetin, 2020), most of them with a high recurrence risk. The overarching concept is that these aetiologies arise from events during fetal development that impact the maternal-fetal exchange of nutrients, oxygen, waste products, and toxins; initiating subclinical pathology that progresses to clinical manifestation over the course of pregnancy. These events include exposure to endogenous metabolites along with exogenous nutrients, as the consequence of different exposome factors. The exposome is defined as the sum of all internal (endogenous) and external (exogenous) non-genetic factors that influence human health throughout life, including perinatal life (Wild, 2005, 2012). These factors can be subdivided into three domains: (1) an internal domain (*endo*-exposome), for example hormones, inflammation and oxidative stress; (2) a specific external domain (*ecto*-exposome), for example infectious agents, diet and lifestyle; and (3) a general external domain that includes education, socio-economic status and mental burden (Wild, 2012). During pregnancy the developing fetus is exposed to different *endo*- and *ecto*-exposome factors, most notably including oxidative stress, nutrition, and inflammation (Robinson and Vrijheid, 2015; Valero et al., 2021). A particular example of a complication of pregnancy in which the exposome is involved is gestational diabetes mellitus (GDM). GDM is a metabolic disease in which the combination of maternal predisposition and placental factors results in progressively impaired glucose tolerance that becomes evident from the end of the second trimester to the third trimester of pregnancy (American Diabetes, 2020; Gabbay-Benziv and Baschat, 2015; Yamamoto et al., 2018). GDM is a pathology where different *endo*- and *ecto*-exposome factors disturb the maternal-fetal interaction (mainly through effects on the micro- and macrovascular endothelial function) resulting in negative consequences for the health of both the mother and the fetus with long-lasting post-natal sequelae (Valero et al., 2021).

The pathogenesis underpinning each of the aforementioned pregnancy complications remains elusive, although a variety of predisposing factors, including a pre-pregnancy maternal body mass index (BMI) of at least 30 kg/m² and genetic predisposition, are recognized (Brosens et al., 2011). Both inflammation and excessive oxidative stress—imbalance between the production of reactive species, such as reactive oxygen species (ROS) and reactive nitrogen species (RNS), and the antioxidant defence system—are strongly associated with the ‘placental syndrome’ that underpins the great obstetrical syndromes (Mannaerts et al., 2018; Schoots et al., 2018). Placentation that is insufficiently deep results in maternal vascular malperfusion (Khong et al., 2016), secondary to increased oxidative stress associated with impaired maternal immune regulation. On the other hand, suboptimal placental invasion can itself lead to an earlier and less organized initial onset of blood flow in the placental intervillous spaces, also resulting in excessive oxidative stress (Redman, 2014; Redman et al., 2022). Oxidative stress can thus have a significant role in both types of placental insufficiency due to abnormal remodelling of maternal vasculature and the resulting endothelial dysfunction (Aouache et al., 2018; Roberts, 2014; Schoots et al., 2018).

Oxidative stress and inflammation are closely related as inflammatory reactions cause oxidative stress and oxidative stress perpetuates an inflammatory response (Zhang, 2010). Inflammatory triggers that cause a shift from immune tolerance towards immune effector activation, and thus a harmful immune response, may result in reactions that can lead to increased oxidative stress and a higher risk of adverse pregnancy outcomes (Nadeau-Vallée et al., 2016).

The exact role of oxidative stress and inflammatory processes in both the onset and the progression of pregnancy complications remains unclear. However, the involvement of these processes in the pathology of

GOS also provide promising therapeutic options (Robertson et al., 2019; Schoots et al., 2018). Many *ecto*-exposome factors are known to influence oxidative stress pathways and inflammation, including dietary intake and (cessation of) smoking (Tenório et al., 2019). Options for pharmaceutical interventions during pregnancy are limited, as pharmaceutical companies are often reluctant to test medication in pregnant women. Dietary interventions carry only little risk in pregnancy, making dietary substrates a promising future therapeutic approach in pregnancy (Amati et al., 2019). In investigating the effect of dietary substrates on pregnancy complications, a wide variety of dietary treatment options have been reported, ranging from changes in complete dietary patterns to single nutrients or mineral supplementations. It is also known that the immune response can be modulated by dietary intake (Childs et al., 2019). As such, nutrition can be considered a strong candidate for a modifiable *ecto*-exposome factor that can be targeted in order to reduce oxidative stress and inflammation in pregnancy.

This narrative review summarizes current understanding of oxidative stress in pregnancy related to placental dysfunction and the potential for attenuating oxidative stress caused by *endo*- and *ecto*-exposome factors through dietary interventions.

2. Exposome-induced oxidative stress and inflammation in pregnancy

2.1. Oxidative stress in pregnancy

Oxidative stress itself should be regarded as an endogenous factor, belonging to the *endo*-exposome oxidative stress (Wild, 2012). In broad terms, pregnancy is a physiological state of prolonged, mildly elevated oxidative stress due to the high oxygen metabolic need for fetal growth and development (Wu et al., 2016). This mildly elevated oxidative stress is the result of physiological inflammatory phenomena of pregnancy that are required for the angiogenic processes needed to expand the placental vascular bed (Wu et al., 2016).

Common oxygen-derived ROS and RNS products include superoxide anion (O₂^{•-}), hydrogen peroxide (H₂O₂), and hydroxyl radical (•HO), nitric oxide (•NO), and peroxynitrite (ONOO⁻). Under physiological conditions, ROS and RNS play fundamental roles in cellular metabolism, cell signalling cascades, and the expression of genes. Increased metabolic activity of mitochondria in the placenta and over-generation of ROS induce oxidative stress during pregnancy (Aouache et al., 2018). ROS levels are negatively affected by several antioxidant enzymes that can be divided into two categories, namely, the enzymatic antioxidants superoxide dismutase (SOD), catalase (CAT), peroxiredoxin (Prx), and glutathione peroxidase (GPx), and the non-enzymatic antioxidants vitamin C, vitamin E, and glutathione (GSH). If this antioxidant capacity is not enough to scavenge the increased amount of ROS, oxidative stress leads to oxidative damage characterized by DNA- and protein damage and lipid peroxidation (Aouache et al., 2018) (Fig. 1).

ROS and RNS are generated continuously from various sources, and in healthy pregnancy they are eliminated by the fetoplacental unit producing abundant corresponding antioxidants (Wu et al., 2016). Under physiological conditions, cytotrophoblasts and villous stromal cells synthesize new antioxidants when exposed to ROS. Assuming that increased oxidative stress is of critical importance in the development of placental pathology, antioxidant therapy might be considered as a preventive therapeutic option. For example, several clinical studies have been conducted to investigate efficacy of strategies to improve the antioxidant capacity of pregnant women in an effort to reduce the risk of GDM (Chan et al., 2021). This has been recently reviewed showing that dietary supplementation of vitamin D and myoinositol reduced the risk of GDM (Chan et al., 2021).

Oxidative stress is measured in maternal blood and placental tissue by various indices. In maternal blood, malondialdehyde (MDA), the breakdown product of fatty acid oxidation, is an indicator of lipid peroxidation and of oxidative damage. MDA normally binds to




Fig. 1. Increased oxidative stress and its consequences in pregnancy. Physiological oxidative stress is seen in pregnancy due to perturbation in the delicate balance between the availability and activity of antioxidants and pro-oxidants. In this physiological process, the normal metabolism in pregnancy generates several pro-oxidants, mainly reactive oxygen (O₂^{•-}, H₂O₂, •HO) or nitrogen (•NO, ONOO⁻) species. The biological production and activity of these species is regulated by antioxidant mechanisms, including enzymes (SOD, CAT, Prx, GPx) and micronutrients (vitamin C, vitamin E, GSH). The equilibrated state of oxidative stress allows normal uterine vascular adaptation and placental development, securing a healthy pregnancy. When the metabolic state of a pregnant woman is altered, the pro-oxidant generation and biological activity surpass the protective action of antioxidants. This phenomenon may be further changed due to reduced generation and biological activity of antioxidants. The result of this condition is dysregulated vascularization (*Abnormal vascularization*), resulting in diseases of pregnancy such as preeclampsia, gestational diabetes mellitus, and fetal growth restriction. O₂^{•-} superoxide anion, H₂O₂ hydrogen peroxide, •HO hydroxyl radical, •NO nitric oxide, ONOO⁻ peroxynitrite, SOD superoxide dismutase, CAT catalase, Prx peroxiredoxin, GPx glutathione peroxidase, GSH glutathione.

thiobarbituric acid-reactive substances (TBARS) formed as a by-product of lipid peroxidation and is elevated in preeclampsia. In addition to the previously mentioned antioxidant enzymes, total antioxidant capacity and gasotransmitters are also used as indices. The antioxidant capacity in plasma can be measured by the Ferric Reducing Ability of Plasma (FRAP) as a reflection of the oxidative stress status (Benzie and Strain, 1996). One of the best markers is free thiols, which have a strong predictive capacity. Plasma-free thiols have recently been shown to be significantly decreased in preeclampsia reflecting oxidative stress (Bos et al., 2019; Schoots et al., 2021). Therefore, a required level of thiols might be required to protect the mother and fetus from this hypertensive disorder. Other components of the *endo-exposome* are involved in the modulation of angiogenesis in the fetoplacental unit by an equilibrium between pro- and anti-angiogenic factors. Fetal or placental pro-angiogenic factors that are detectable in the maternal blood include placental growth factor (PlGF) and heme oxygenase-1 (HO-1). Anti-angiogenic factors include soluble fms-like tyrosine kinase 1 (sFlt-1) and endoglin (ENG). Moreover, plasma markers of endothelial activation and placental dysfunction are the *endo-exposome* factors plasminogen-activator inhibitor 1 (PAI-1) and 2 (PAI-2). Finally, an increase in inflammation (*endo-exposome*) could contribute to oxidative stress, as pro-inflammatory cytokines are produced in response to ROS and subsequently stimulate further ROS production by target cells (Burton and Jauniaux, 2011).

Oxidative stress can be aggravated by a variety of *ecto-exposome* factors including environmental pollutants, lifestyle factors such as tobacco smoking, and the amount of dietary fat consumption (Pizzino et al., 2017). Also, heavy metals consumed as environmental contaminants, such as arsenic, mercury, copper, and lead, are associated with higher oxidative stress increasing the risk of preeclampsia and preterm birth (Stone et al., 2021) and GDM (Gómez-Roig et al., 2021; Valero et al., 2021). Thus, adequate regulation of the inflammatory mediators and the *ecto-exposome* factors involved in triggering this state is essential for pregnancy success, and excessive ROS can perturb this.

2.2. Placental pathology associated with oxidative stress

The placental lesion most often related to oxidative stress and inflammation and the concurrent clinical conditions is maternal

vascular malperfusion (MVM) (Fig. 2). This lesion is characterized, among other things, by placental hypoplasia (low weight, maldevelopment), infarction, accelerated villous maturation, and decidual arteriopathy (Khong et al., 2016). Due to impaired spiral artery remodelling, the oxygen tension in the placenta rises disproportionately, leading to a rise in ROS and eventually to oxidative stress (Schoots et al., 2018). Another common placental lesion, chronic villitis of unknown aetiology, is often seen in (late) FGR (Fig. 3C and D). Due to a chronic lymphocytic CD8⁺ T cell infiltration of maternal origin the villi are damaged, thus reducing the amount of functional placental tissue available to participate in the exchange of oxygen and nutrients to the fetus (Khong et al., 2016). In GDM, the placental parenchyma shows, in varying degree, delayed maturation with decreased number of vasculosyncytial membranes (Fig. 3A and B) (Huynh et al., 2015); thereby reducing the capacity of the placenta to deliver enough nutrients and oxygen to the fetus and remove the necessary waste products. As a compensatory mechanism, the number of villous capillaries increases, known as chorangiogenesis (Carrasco-Wong et al., 2020) (Fig. 3). However, the diffusion distance remains increased due to villous stromal proliferation. The increase in villous stromal cells and in villous capillaries results from dysregulation of the glucose and oxygen metabolic pathways, although the exact mechanisms remains unclear (Jauniaux and Burton, 2006).

2.3. Inflammation in pregnancy

Inflammatory responses are also considered *endo-exposome* factors. The immune system plays a fundamental role in pregnancy. The principal role of the immune system is to protect against potentially dangerous microbial infections and tumor cells. In pregnancy the maternal immune system, however, faces a major challenge. The embryo and gestational tissues arising from the fusion of the male and female gametes express both paternal and maternal major histocompatibility (MHC) antigens. Therefore, the fetus is semi-allogeneic to the maternal immune system. The maternal immune system must remain tolerant to the semi-allogeneic embryo and gestational tissue on the one hand and facilitate normal immune competence to protect against pathogens on the other hand. Adaptation of the maternal immune system during early pregnancy is therefore required in order to create an environment in which both immune tolerance and immune




Fig. 2. Histology of placental lesions in preeclampsia. Representative light microscope photographs of Hematoxylin/Eosin-stained placental samples. The excessive oxidative stress and inflammation in preeclampsia results in structural abnormalities in the placental vasculature leading to malperfusion. The most characteristics structural abnormalities include (A, B) placental infarction (*), (C) accelerated villous maturation with increased syncytial knotting (arrowheads), (D) decidual arteriopathy with necrosis of the vessel wall (*), and (E) presence and influx of foamy macrophages (arrowheads).




Fig. 3. Histology of placental lesions in gestational diabetes mellitus and fetal growth restriction. Representative light microscope photographs of Hematoxylin/Eosin-stained placental samples. The excessive oxidative stress and inflammation in gestational diabetes mellitus and fetal growth restriction result in structural abnormalities in the placental vasculature leading to altered perfusion and fetal growth. Gestational diabetes mellitus is accompanied by delayed maturation of the parenchyma where there is an increase in villous capillaries (A, B). Placentas from pregnancies with fetal growth restriction show with chronic villitis (E-F) presenting lymphocytic infiltration (*) in the villi (C). Infiltration of T-cells is identified by positive CD3 staining (D).

protection can coexist (Munoz-Suano et al., 2011; Poole and Claman, 2004).

The maternal immune response adapts to accommodate the fetus in multifactorial ways, and an aberration in any aspect can potentially lead to deterioration of fetal health. When such a deterioration occurs, this could be interpreted as derailment of the *endo*-exposome. To prevent a harmful maternal immune response towards the fetus, some parts of the immune response are evaded, while other parts are directed to induce antigen-specific tolerance (Munoz-Suano et al., 2011). Both maternal and fetal factors cooperate to create an immunological ‘fetus friendly’ environment. The role and balance of different leukocyte lineages in gestational tissues are modified to secure a healthy pregnancy outcome. This includes restricting access and modifying the phenotypes of macrophages, dendritic cells, natural killer (NK) cells, and lymphocytes in various compartments of the uterus and placental membranes (Munoz-Suano et al., 2011; Robertson et al., 2019; van der Zwan et al., 2020). This restricted access is modulated by the production of several

endogenous factors including immunoregulatory cytokines, chemokines, hormones and prostaglandins from both uterus and placental tissue.

Macrophages (especially the more anti-inflammatory subsets), dendritic cells, and regulatory T-cells (Treg-cells) that mediate immune tolerance also influence vascular dynamics through their potent anti-inflammatory capacity. Treg-cells are central players in maintaining immune tolerance (Guerin et al., 2009). These cells are powerful suppressors of CD4⁺ Th1/Th17 cells, inhibit generation of CD8⁺ cytotoxic T cells and NK cells, and interact with macrophages and dendritic cells for suppressing inflammation and maintaining immune quiescence. Uterine NK cells promote structural changes in the decidual vasculature for placental invasion and development (Faas and de Vos, 2017). The fetus, and its placental tissue in particular, contribute by bypassing or inactivating various maternal immune cells, through regulation of MHC antigen expression, excretion of complement regulatory proteins, and displaying other protective proteins. The fact that trophoblast has

modified expression of classical HLA class I (with absence of HLA-A and HLA-B, and attenuated HLA-C) is considered one of the key protective mechanisms from the fetal side. These classical HLA antigens are polymorphic, and the paternally inherited antigens are therefore recognized as foreign. Nonclassical Ib antigens (HLA-E, HLA-F, HLA-G) are expressed instead. These Ib antigens are relative non-polymorphic and are not recognized as foreign by the maternal immune system (Poole and Claman, 2004).

For GDM it is now becoming clear that inflammatory processes play a significant role in the disease pathophysiology. Underlying mechanisms include a shift away from a pro-tolerance immune environment due to diminishment of protective mechanisms or dominating exogenous or endogenous pro-inflammatory agents. For example, deficient signalling to A_{2B} adenosine receptors activation results in a lower capacity to reduce the generation of O₂^{•-} and ONOO⁻, thereby increasing oxidative stress. This phenomenon also leads to insufficient protection against inflammation due to reduced generation of the anti-inflammatory cytokines IL-10 and adiponectin, but increased generation of the pro-inflammatory cytokines TNF and IL-6 in GDM and gestational diabetes (Cabalín et al., 2019; Cornejo et al., 2021). In preeclampsia, failure of trophoblast cells to invade and remodel the maternal spiral arteries results in exaggeration of a systemic inflammatory process. Immuno-competent cells are activated and hypoxic conditions, excessive inflammation and/or oxidative stress occurs resulting in increased trophoblast failure (Chiarello et al., 2020).

3. Dietary interventions to prevent pregnancy complications

Therapeutic approaches to modulate excessive oxidative stress and the associated adverse inflammatory reactions in pregnancy complications could benefit pregnant women and their fetuses. However, due to

potential teratogenic effects, pharmaceutical interventions during pregnancy are scarcely studied and thus have limited approvals for community use. Recently, prevention of pregnancy disorders through dietary intake has received more attention as a tractable and potentially effective intervention.

Nutritional interventions in patients with different forms of pregnancy complication and different outcome measures have been examined. One of these investigations is the Dietary Approaches to Stop Hypertension (DASH) (Asemi et al., 2013, 2014). This diet was originally designed to normalize blood pressure in patients with hypertension. The DASH dietary pattern is low in fat, with a focus on reduced saturated fats compared with a high-fat diet. The rationale for the study was that the DASH diet would be expected to have a beneficial effect on pregnancy outcomes and reduces macrosomia in fetuses of mothers with GDM. Unfortunately, analyses of the data are hampered by the many differences between these two diets besides the fat content.

Furthermore, it is reported that dietary intervention in women at high risk for GDM may reduce inflammation and the risk of GDM. A reduction and improvement in carbohydrate quality rather than a restriction in the high-fat content in the diet plays a major role. The habitual diet plays an important role in the improvement that can be expected from dietary adaptation as seen in women with GDM (García-Patterson et al., 2019; Yamamoto et al., 2018). Therefore, dietary intake has been considered as a potential treatment option (Fig. 4). Below we summarize data of the most promising nutritional interventions, especially in relation to GOS.

3.1. L-arginine

The cationic amino acid L-arginine is considered as the biological precursor for [•]NO, a potent endothelium derived vasodilator (Moncada




Fig. 4. Dietary and other interventions to prevent pregnancy complications. Among the several intervention strategies implemented to avoid pregnancy complications, diet and supplementation with micronutrients are described. Diet interventions with a selection of healthy foods result in reducing insulin resistance and increasing the antioxidant capacity in pregnant women. Supplementation with essential trace elements such as Selenium reduced the level of prooxidant molecules. A generalized mechanism to improve placental function after diet or Selenium is a higher generation of nitric oxide (NO) by NO synthase (NOS) favoured by higher uptake of its substrate L-arginine (L-Arg). This pathway seems favoured compared with the use of L-Arg to the synthesis of polyamines and proteins. Supplementation of patients with the micronutrients vitamins E (Vit E), C (Vit C), and D (Vit D) diminishes the formation of lipid hydroperoxides (LOOH) from peroxy radicals (LOO[•]) when the anion superoxide (O₂^{•-}) levels are higher in membrane lipids. This phenomenon results from the action of Vit E reducing (red line) the formation of LOOH, causing oxidation of Vit E (oxVit E). Vit E is restored from oxVit E involving Vit C. These mechanisms result in a lower membrane lipids oxidation. Vit E, Vit C and Vit D function restore endothelial dysfunction in the vasculature and other tissues. Garlic flavonoids and Omega-3 fatty acids also restored placental function involving the mother-to-fetus transplacental transfer of icosapentaenoic acid (EPA), docosahexaenoic acid (DHA), and alpha-linolenic acid (ALA). DHA is reported to accumulate

in the placenta in diseases showing reduced or deficient membrane transport activity for this type of Omega-3. A healthy diet, physiological levels of Selenium, vitamins and Omega-3 fatty acids result in reducing oxidative stress and inflammation in pregnancy. This phenomenon reduces (↓) the risk of developing gestational diabetes mellitus with a less more evident effect (↔) in preeclampsia. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

et al., 1988). Increased synthesis of \bullet NO is associated with normal vascular adaptation under physiological conditions, while decreased \bullet NO synthesis has been reported in preeclampsia (Rytlewski et al., 2005). It is proposed that uptake of L-arginine is a phenomenon linked to generation of \bullet NO via the endothelial NO synthase (eNOS) in the fetoplacental endothelium (Ramírez et al., 2018). This phenomenon seems to be restricted to the human cationic amino acid transporters 1 (hCAT-1) since the Michaelis-Menten constant (Km) for its activity was in the range of this plasma membrane transporter isoform, i.e., apparent Km 50–350 μ mol/L (Mann et al., 2003). Also, the potential link between L-arginine transport and \bullet NO synthesis may involve a very high-affinity transport mechanism (Km \sim 2 μ mol/L) describing system y+L activity (Mann et al., 2003; Ramírez et al., 2018). Thus, the availability of L-arginine for its uptake via hCAT1 and system y+L into nearby eNOS localized in the vascular endothelium may be critical for the vascular adaptive regulatory mechanisms opposing vasoconstriction in preeclampsia (Vadillo-Ortega et al., 2011).

During pregnancy, an L-arginine rich diet as a source of substrate for the \bullet NO synthesis and consequent vasodilation shows promise in protecting against preeclampsia. Although L-arginine results in vasodilation through \bullet NO, the influence of L-arginine administration on blood pressure is disputed and the mechanism underlying its antihypertensive action is unclear. While studies of high doses for short periods have not demonstrated effects on blood pressure, prolonged low doses dietary supplementation of L-arginine (3 g/day) for 3 weeks has shown to decrease blood pressure through increased synthesis and bioavailability of \bullet NO in women with preeclampsia (Rytlewski et al., 2005). Although a recent meta-analysis suggests that intervention with L-arginine may reduce FGR and result in increased birth weight, no reduction in the occurrence of preeclampsia in human pregnancies could be confirmed (Terstappen et al., 2020). Another study showed however that in high-risk pregnancies, daily intake of medical food bars containing L-arginine plus antioxidant vitamins significantly reduced the incidence of preeclampsia compared to placebo (Vadillo-Ortega et al., 2011). The protective effects may be greatest when onset of the supplement is before 24 weeks of gestation (Terstappen et al., 2020; Vadillo-Ortega et al., 2011).

In early-onset preeclampsia, the human placental microvascular endothelial cells (hPMECs) show reduced capacity of \bullet NO synthesis compared with hPMECs from late-onset preeclampsia (Escudero et al., 2009). The proposed mechanism behind these differences in the capacity of NO synthesis in hPMECs involves an elevated extracellular adenosine concentration as reported in human umbilical blood in preeclamptic pregnancies. Adenosine activates A_{2A}/A_{2B} adenosine receptors that induce the expression of VEGF which, via activation of VEGFR1/2, increases the activity of the inducible NOS form (iNOS) expressed in hPMECs. In early-onset, and in a minor degree in late-onset preeclampsia, the released VEGF in response to adenosine binds to sFlt-1 restricting VEGF biological actions. It is worth noting that L-arginine is taken up by the endothelium via plasma membrane mechanisms with a maximal transport capacity –defined as the ratio between the maximal velocity of transport over the affinity of membrane transporters for this amino acid (Mann et al., 2003)– could be rate-limiting for the \bullet NO synthesis via eNOS (McCabe et al., 2000). The transport activity via hCAT-1 isoform and hCAT-2B, a splice variant of hCAT-2A isoform, is the main mechanism leading to uptake of L-arginine (apparent Km 50–350 μ mol/L) and is concentrative leading to \sim 10-fold intracellular accumulation of L-arginine compared with its plasma concentration (\sim 100 μ mol/L). Since the affinity of eNOS for L-arginine is \sim 2–3 μ mol/L, it may be a limiting mechanism for the synthesis of NO in preeclampsia. Therefore, supplementing women with preeclampsia with this amino acid may not necessarily restore the reduced \bullet NO synthesis by the fetoplacental endothelium.

3.2. Fatty acids

Omega-3 (N-3) fatty acids possess antioxidant and anti-inflammatory properties and play a role in promoting placental vasculature (Jones et al., 2014). The precise effect on lowering oxidative stress depends on the dosage, background diet and experimental conditions (Shoji et al., 2009). The three types of omega-3s include eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA), and alpha-linolenic acid (ALA). The long chain EPA and DHA are found in fatty fish, while ALA is found in oils from plant origin, such as flaxseed oil and chia seed. Interestingly, DHA supplementation in women with GDM, despite their pre-pregnancy BMI, resulted in an increased DHA level in the mother but not in the fetal circulation (Gázquez et al., 2021). This observation has been partially explained by a reduced placental transport of DHA likely through the major facilitator super family domain containing 2a (Mfsd2a) transporter (Prieto-Sánchez et al., 2017) resembling mechanisms operating in the blood-brain barrier (Nguyen et al., 2014). The reported accumulation in the placenta in GDM pregnancies may reflect reduced transport to the fetus (Larqué et al., 2014), since offspring born after GDM pregnancies have been reported to have lower DHA status at birth.

Thus, maternal supplementation with DHA in pregnancy may be useful to counteract GDM-associated oxidative stress and inflammatory response, while having minor or no direct effect on the fetus, and a likely positive indirect effect.

Omega-3 fatty acids might act better in combination with vitamin E because of its sensitivity to oxidation and strong synergistic activity on related markers. In a co-administration trial in women with GDM, administration of 1000 mg n-3 fatty acids from flaxseed oil plus 400 IU vitamin E per day resulted in improved oxidative stress biomarkers (TAC, NO, and MDA) (Jamilian et al., 2017). There was no improvement of the inflammation biomarker ‘high-sensitive C-reactive-protein’ and other pregnancy outcomes (Jamilian et al., 2017).

The short chain fatty acid alpha-linolenic acid (ALA) is known as a universal oxidant because of its hydrophobic and hydrophilic characteristics. Administration of ALA improved markers of glucose metabolism and some markers of liver function, and increased adiponectin (A), A/L ratio and A/H ratio and decrease of L/A values (Asfalalah et al., 2019a, 2019b). Thus, although omega-3 fatty acids and especially ALAs seem to have beneficial effects on metabolic profiles, biomarkers of oxidative stress, inflammation, the effects on adverse pregnancy outcomes are less imminent. This phenomenon may in part be explained by the fact that an increase in the n6 linoleic acid will impact the production of LCPUFAs, e.g., suppressed DHA and EPA production favouring a higher ARA status. Interestingly, the modern Western diet contains an imbalance in the omega-6/-3 ratio due to lower omega-3 intake and higher omega-6 consumption (Simopoulos, 2002).

3.3. Vitamin B₁₂, C, D and E

Vitamins and minerals have a substantial positive influence on the health of women and their fetuses. Adequate maternal micronutrient status is critical during pregnancy, and an inadequate micronutrient status associates with several pregnancy complications (Allen, 2005; Gomes et al., 2020). Antioxidant vitamin supplementation is considered a therapeutic or even preventive option for reducing inflammation and oxidative stress in placenta pathology.

Vitamin B₁₂ (cobalamins), member of the vitamin B complex, is a cofactor for several enzymes involved in carbohydrates, fatty acids and amino acids metabolism, maintaining fetal growth and maternal metabolic state. In addition to these functions, vitamin B₁₂ is also a scavenger of ROS, particularly O₂^{•-} and vitamin B₁₂ deficiency increases the risk of GDM (Krishnaveni et al., 2009; Lai et al., 2018; Offringa et al., 2021; Radzicka et al., 2019). Patients with GDM are increasingly treated with the oral hypoglycaemic drug metformin. Since metformin restricts the absorption of vitamin B₁₂ (Aroda et al., 2016), metformin-treated women with GDM may present lower levels of vitamin B₁₂ (Gatford

et al., 2013). However, metformin does not alter the active form of vitamin B₁₂, holotranscobalamin, suggesting that a reduced absorption of vitamin B₁₂ may not alter the level of its active form. Interestingly, no effects of vitamin B₁₂ on the development of the classical GOS have been reported in the literature.

Vitamin C and E are often supplemented in combination because *in vivo* and *in vitro* studies have shown that vitamin C and E act in synergy. Vitamin E is considered crucial in interrupting the cycle of lipid peroxidation, while vitamin C enhances the capacity of vitamin E to preclude lipid peroxidation by renewing oxidized vitamin E to its reduced form. Women at low risk of preeclampsia show no further risk reduction after supplementation of vitamin C solely (Kiondo et al., 2014), or after vitamin C plus E supplementation (Roberts et al., 2010; Rumbold et al., 2015). A trend of lower incidence of pregnancy induced hypertension was shown after 100 mg tocotrienol-rich-fraction of palm oil vitamin E per day (Mahdy et al., 2013). However, in women with essential hypertension, there was no reduction in risk of superimposed preeclampsia or aggravation of hypertension after supplementation (Kalpdev et al., 2011). These findings were similar to previous reports showing that supplementation did not reduce the incidence of preeclampsia, irrespective of risk at enrolment or early onset (Poston et al., 2006). In a group of pregnant women with type 1 diabetes mellitus, i.e. at high risk for developing preeclampsia, the DABIT trial also failed to demonstrate reduction of the risk of preeclampsia and gestational hypertension after supplementation with vitamin E combined with vitamin C (McCance et al., 2010). Similarly, another study assessed among high- and low-risk women whether vitamin C plus E supplementation reduces the risk of developing gestational hypertension (Xu et al., 2010). This study concluded that supplementation did not reduce the risk of preeclampsia, and unexpectedly found that supplementation with vitamins C and E increased the risk of fetal loss or perinatal death.

The lack of beneficial effects of vitamins C and E supplementation in some studies could be explained by the fact that these women may have had an adequate antioxidant status at study initiation. There is some evidence that in women with low antioxidant status, such as in developing countries and low socioeconomic status, early administration of enriched milk containing several minerals and vitamins starting in first trimester, including 200 mg vitamin C and 400 mg vitamin E, reduced the incidence of preeclampsia (Wibowo et al., 2012). The effects on birthweights are even more diverse, with some studies finding lower birth weights in the groups receiving antioxidants or high dose vitamin E (Boskovic et al., 2005; Poston et al., 2006). Furthermore, it might also well be that too high concentrations of antioxidants may affect the normal physiology of reactive species.

Regarding preterm birth, there are conflicting reports of the effects of vitamin C and E supplementation. As ascorbic acid is one of several factors associated with preterm premature membrane rupture (PPROM, <37 weeks of gestation), it is thought to have a potential preventive effect. Vitamin C is involved in collagen synthesis and may maintain the integrity of the chorioamniotic membranes. In fact, reduced vitamin C levels are associated with spontaneous preterm birth. Contradictory results of supplementation are found regarding the risk of premature rupture of membranes (PROM). Supplementation of 100 mg vitamin C per day has been shown to reduce the incidence of PROM in healthy pregnant women (Casanueva et al., 2005). Whereas other trials found higher rates of PROM and PPRM (Steyn et al., 2003; Xu et al., 2010).

Besides vitamin C and E and their combination, vitamin D has also been studied extensively in pregnant women. Vitamin D is shown to have an immunomodulatory effect, and may play a role in maintaining immune tolerance in pregnancy (Schröder-Heurich et al., 2020). Studies on the effects of supplementation with vitamin D on GOS show inconsistent findings (Haugen et al., 2009; O'Callaghan and Kiely, 2018). Some studies report no influence of administration of vitamin D on the prevalence of preeclampsia (Mirzakhani et al., 2016; Sablok et al., 2015), whilst others describe a positive effect on the prevalence of preeclampsia (Ali et al., 2019; Fogacci et al., 2020). There are

contradictory reports on the effect of birth weight and the prevalence of preterm labor (Ali et al., 2019; Fogacci et al., 2020; Harreiter et al., 2019; Sablok et al., 2015). Both Fogacci et al. (2020) and Ali et al. (2019) found a decrease in the prevalence of FGR. It is reported that supplementation with vitamin D in 2487 women reduced the risk of preeclampsia, FGR and preterm birth and was dependent on the doses of vitamin D but independent of supplementing with vitamin D alone or together with calcium (Fogacci et al., 2020). Although these findings are difficult to reconcile, it remains possible that a beneficial effect of vitamin D supplementation to prevent or alleviate the signs and symptoms of preeclampsia may apply in certain high risk populations (Ali et al., 2019; Fogacci et al., 2020; Sablok et al., 2015).

More extensively the effects of vitamin D on women with GDM have been described. Women with GDM showed low vitamin D levels compared with the level of this micronutrient in normal pregnancies (>20 ng/L, measured as serum 25-hydroxyvitamin D, 25(OH)D) (Farant et al., 2009; Gunasegaran et al., 2021; Liu et al., 2020; Palacios et al., 2019; Zhang et al., 2008). Vitamin D supplementation varied between 200 and 1000 IU/day (5–25 mg/day) to reduce the risk of developing GDM. Vitamin D in combination with calcium or in combination with omega-3 fatty acids improve the glucose and lipid metabolism of patients with GDM (Gunasegaran et al., 2021; Huang et al., 2021). However, supplementation of vitamin D is an approach that has been not conclusive in women with GDM (Ali et al., 2019; Palacios et al., 2019; Sablok et al., 2015). A unique intramuscular administration of a high dose of vitamin D (300,000 IU) has been shown to improve the status of its metabolic product 25(OH)D₃ in women with GDM (Hosseinzadeh-Shamsi-Anar et al., 2012). Recently, co-administration of vitamin D in combination with zinc, magnesium and calcium in women with GDM showed a significant reduction in serum high-sensitivity C-reactive protein and plasma malondialdehyde concentrations, as well as a significant increase in total antioxidant capacity levels compared to placebo (Jamilian et al., 2019).

Even when vitamin D levels are improved, there is a question of how this approach relates to the duration of the beneficial effect in these patients and their fetuses and newborns. The biological actions of vitamin D include activation of vitamin D receptors in the fetoplacental endothelium (Molinari et al., 2011; Uberti et al., 2014; Wu et al., 2021) protecting against oxidative stress and endothelial dysfunctional metabolism, including a reduced mitochondrial oxidative phosphorylation (Uberti et al., 2014). Thus, lower level of vitamin D and altered activation of vitamin D receptor in GDM may increase O₂^{•-} which scavenges NO limiting most biological actions of this gasotransmitter (Mahdi et al., 2021). HUVECs from GDM show increased eNOS activity, thus, a reduced eNOS activation due to deficient vitamin D activation of vitamin D receptors and the increased NO scavenging by O₂^{•-} may not be enough to limit the NO-mediated inhibition of cytochrome c oxidase seen in GDM (Sobrevia et al., 2020).

3.4. Selenium

Similar to vitamins, minerals with antioxidant function are also associated with pregnancy outcomes and have been used in several trials. A good example of this is selenium. Selenium (Se) is an essential trace element that functions via incorporation in protein forming selenoproteins. These proteins have both antioxidant and anti-inflammatory effects. Se possesses anti-inflammatory activity due to its regulatory effect on expression of pro-inflammatory genes, including *selenoprotein S* gene and the release of pro-inflammatory cytokines.

Selenium is mostly obtained from dietary intake of nuts, cereals, meat, mushrooms, fish, and eggs. The dose of intake remains subject of discussion; one study recommends dosage of 10 up to 56 µg/day while others recommend the intake of 25–50 µg/day, irrespective of diet. The recommended allowance is 55 µg Se/day for healthy adult non-pregnant women; however, it is likely that substantially higher levels are required during pregnancy. Higher blood level of Se associates with improved

markers of oxidative stress, inflammation, insulin metabolism and obstetrical outcomes. The concentration of Se varies in a wide range in both pregnant women and non-pregnant women, due to variable global distribution and ubiquitous dietary sources. The blood concentration of Se reflects the daily intake and higher blood concentration of D-glucose correlates with lower Se blood levels in pregnant women with GDM (Hawkes et al., 2004), a phenomenon likely due to the potential state of insulin resistance in these women.

Supplementation with Se decreases oxidative stress and alleviates inflammation in pregnant women. Selenium reduces inflammation by inhibiting NF- κ B, toll-like receptors and 38 kDa mitogen-activated protein kinases (p38^{mapk}) pathways in several cell types including HUVECs (Zheng et al., 2008). Oxidative stress is also reversed by Se due to its capacity to activate D-glucose uptake and metabolism thus reducing the state of hyperglycaemia seen in patients with GDM (Kar-amali et al., 2020). In a trial performed in Iranian women from Arak (Asemi et al., 2015; Jamilian et al., 2018; Razavi et al., 2016), administration of Se did not change the maternal HOMA β -cell function, lipid profile, plasma NO level, or total antioxidant capacity. Although Se supplementation resulted in beneficial effects on glucose metabolism and oxidative stress and inflammation markers in pregnant women, it did not reduce the incidence of PE, and no similar results were described for FGR and preterm birth. These results should be considered with caution however, since they refer to a small sample size and women from the Arak region have higher selenium concentrations than women from other regions of Iran.

Serum levels of at least 100 μ g Se/L are needed for maintaining adequate Se mediated antioxidant capacity. Because both plasma levels below 45 μ g/L and decreased GPx activity are associated with advanced pregnancy outcomes, supplementation should be considered for those having suboptimal blood Se levels. However, the underlying mechanism of the role of Se and the Se-dependent enzymes stays unclear. Moreover, the dosage, timing and duration of Se supplementation during pregnancy is still subject of debate.

3.5. Iron

Pregnant women are at risk of developing iron deficiency anemia. Therefore, the World Health Organization (WHO) recommends iron supplementation early in pregnancy (World Health Organization (WHO), 2016). Anemic pregnant women have a higher incidence of preterm birth and offspring with low birth weight (Casanueva and Viteri, 2003). Korkmaz et al. (2014) found in a randomized double-blind controlled trial that iron supplementation during first trimester is associated with higher levels of oxidative stress. No correlation with preeclampsia was found (Korkmaz et al., 2014). On the other hand, Siddiqui et al. (2011) found an increased risk of developing preeclampsia in women with pre-existent high serum iron levels (Siddiqui et al., 2011). Also, in a study in which iron supplementation before 16 weeks gestational age was started, an increased risk of developing hypertension after 20 weeks gestational age was apparent (Jirakittidul et al., 2018). Iron influences the production of ROS in the placenta contributing to oxidative stress (Casanueva and Viteri, 2003) via the formation of hydroxide (OH⁻) and hydroxyl radical (*HO) (Mannaerts et al., 2018). It is interesting that both a low haemoglobin (Hb) status as well as a high Hb status are associated with pregnancy complications (Aly et al., 2016; Casanueva and Viteri, 2003; Yong et al., 2022). Increase in iron (both pre-existent plasma ferritin levels and supplementary iron) is also associated with a higher risk of developing GDM (Asadi et al., 2019; Mannaerts et al., 2018; Zhang et al., 2021). So, although iron supplementation could be beneficial in anemic pregnant women, iron levels need to be monitored to prevent overload and secondary complications like GDM.

4. Discussion and conclusion

Oxidative stress and inflammation are involved in the pathophysiology of various pregnancy complications collectively described as “the Great Obstetrical Syndromes”. Nutrient and dietary interventions can play a role in favourably attenuating the processes of oxidative stress and inflammation by configuring a more protective exposome. It is known that unhealthy diets characterized by energy dense but not nutrient dense foods result in more unfavourable outcomes attributable to inflammation and oxidative stress. However, the beneficial effects of various dietary interventions on pregnancy outcomes through modulation of inflammatory processes and oxidative stress is still controversial. Research on this topic is challenging for several reasons:

- (1) A causative relation between adverse outcomes, dietary intake, oxidative stress and inflammation is hard to prove, largely because of the complex heterogeneity of many risk factors that include not only *ecto*-exposome and *endo*-exposome, but also inter-related and independent factors like BMI and smoking, as well as socio-economic status and ethnicity.
- (2) The onset of supplementation during pregnancy might simply be too late to have a beneficial effect. Most of the tested interventions are initiated in the second trimester of pregnancy while most likely the underlying causal events giving rise to placental dysfunction occur prior to that time. Therefore, introduction of antioxidant or vitamin supplementation before or around conception might be more effective, since a woman's health and nutrition status before pregnancy is crucial for improving pregnancy health (Hanson et al., 2015).
- (3) The interventions have been conducted in a wide variety of study populations (baseline characteristics), mostly in studies with a small number of participants and different outcome measurements. There were differences in nutritional state, race, social economic status, background, food patterns, culture, start, duration, dosage and route of administration of the interventions. Due to differences in study design and objectives, the outcomes cannot be compared directly.
- (4) The susceptibility of the developing fetus is an important consideration, so once potential interventions are identified they must be evaluated for their safety and any effects on the development of the embryo and fetus.
- (5) Dietary intake exists of many constituents that interact together. It is not to be expected that changing one single component for a limited period of time will lead to immediately measurable effect.
- (6) Systematic report bias can occur through non-objective self-reported food questionnaires and interventions can be confounded by this bias.

In other diseases, for example inflammatory bowel disease (IBD), preventive dietary interventions are thought to be useful (Campmans-Kuijpers and Dijkstra, 2021). Some of the underlying oxidative stress and immunological inflammatory mechanisms are comparable between IBD and pregnancy complications. Therefore, the proven effective treatment of diet in IBD might be translated to candidate dietary treatments for pregnancy complications (Campmans-Kuijpers and Dijkstra, 2021). In contrast, the desired effect in IBD patients (less disease activity) is different to that in pregnancy (less FGR and hypertension), which could influence patient motivation. In the future, diets with focus on changes in multiple nutritional aspects could be developed as a candidate intervention, considering the potency as well as the complexity of nutrition. Since a balanced diet consisting of healthy foods is by definition not harmful, it might be beneficial if it also solves nutritional deficiencies. Because pharmacotherapeutic treatment is restricted in pregnancy, pregnant woman reasonably would be expected to benefit from healthy nutritional approaches, regardless of impact on GOS outcomes.

A limitation of the current review is that a considerable proportion of the included studies were published by only a limited number of research groups. Most trials were of small sample size, short duration and often local dietary recommendations and/or supplementation guidelines were not known, nor patient compliance reported. Most randomized controlled trials of supplementation during pregnancy to reduce pregnancy complications reported controversial or negative results, despite the expected protective associations between dietary antioxidant and pregnancy outcomes demonstrated in observational studies.

In summary, the multiple exogenous and endogenous factors involved in placental and fetal development and growth are crucial elements of the exposome to which the mother and fetus are exposed (Kupper and Huppertz, 2021; Valero et al., 2021). Dietary interventions envisage reaching levels of the modulatory factors involved in suppressing oxidative stress to a physiological level, in order to promote positive pregnancy outcomes in women at high risk for pregnancy complications. In this phenomenon, the activation of the *endo*-exposome as consequence of dietary interventions might be essential to explain the inflammation and redox regulatory mechanisms involved in achieving a healthy pregnancy. Evaluation of dietary interventions will need to account for the many other factors that predispose to GOS. For example, increased maternal age is associated with increased pregnancy complications like preeclampsia and GDM (Attali and Yogev, 2021), and an interaction between age and nutritional effects may exist.

In conclusion, nutrition is considered key to maternal optimal health before and during pregnancy to optimize birth outcomes. Inflammation and oxidative stress have been implicated in the most common pregnancy complications like GOS, and interest in diet as a possible positive factor has grown. Now is the time for further assessment of anti-inflammatory and fully-fledged nutrition in pregnancy with attention on the timing, duration and contents of the dietary intervention, with the benefit of a reduced risk of potential adverse effects compared to pharmacotherapeutic interventions.

Declaration of competing interest

The authors declare no competing interests.

Acknowledgements

EvdB is an employee of Nestlé Research, part of Société des Produits de Nestlé, Switzerland. LS thanks the support from the Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) [grant number 1190316], Chile, and International Sabbaticals (University Medical Center Groningen, University of Groningen, The Netherlands) from the Vicerectorate of Academic Affairs, Academic Development Office of the Pontificia Universidad Católica de Chile.

Funding sources had no role in the study design, in the collection, analysis and interpretation of data, in the writing of the report, and in the decision to submit the article for publication.

References

- Ali, A.M., Alobaid, A., Malhis, T.N., Khattab, A.F., 2019. Effect of vitamin D3 supplementation in pregnancy on risk of pre-eclampsia - randomized controlled trial. *Clin. Nutr.* 38, 557–563. <https://doi.org/10.1016/j.clnu.2018.02.023>.
- Allen, L.H., 2005. Multiple micronutrients in pregnancy and lactation: an overview. *Am. J. Clin. Nutr.* 81, 1206S–1212S. <https://doi.org/10.1093/ajcn/81.5.1206>.
- Aly, S., Fayed, H., Ahmed, S., Abdella, A., Tamam, A.-A., Mohammed, N., 2016. Effects of oral iron (ferrous versus ferric) supplementation on oxidative stress and antioxidant status in pregnant women with iron deficiency: controlled trial. *Egypt. J. Haematol.* 41, 31. <https://doi.org/10.4103/1110-1067.186392>.
- Amati, F., Hassounah, S., Swaka, A., 2019. The impact of Mediterranean dietary patterns during pregnancy on maternal and offspring health. *Nutrients* 11, 1098. <https://doi.org/10.3390/nu11051098>.
- American Diabetes, A., 2020. 2. Classification and diagnosis of diabetes: standards of medical care in diabetes-2020. *Diabetes Care* 43, S14–S31. <https://doi.org/10.2337/dc20-S002>.
- Aouache, R., Biquard, L., Vaiman, D., Miralles, F., 2018. Oxidative stress in preeclampsia and placental diseases. *Int. J. Mol. Sci.* 19, 1496. <https://doi.org/10.3390/ijms19051496>.
- Aroda, V.R., Edelstein, S.L., Goldberg, R.B., Knowler, W.C., Marcovina, S.M., Orchard, T. J., Bray, G.A., Schade, D.S., Temprosa, M.G., White, N.H., Crandall, J.P., 2016. Long-term metformin use and vitamin B12 deficiency in the diabetes prevention program outcomes study. *J. Clin. Endocrinol. Metab.* 101, 1754–1761. <https://doi.org/10.1210/jc.2015-3754>.
- Asadi, N., Vafaei, H., Kasraeian, M., Yoosofi, S., Faraji, A., Abbasi, L., 2019. Effects of prophylactic iron supplementation on outcome of nonanemic pregnant women. *J. Chin. Med. Assoc.* 82, 840–844. <https://doi.org/10.1097/JCMA.000000000000184>.
- Asemi, Z., Jamilian, M., Mesdaghinia, E., Esmailzadeh, A., 2015. Effects of selenium supplementation on glucose homeostasis, inflammation, and oxidative stress in gestational diabetes: randomized, double-blind, placebo-controlled trial. *Nutrition* 31, 1235–1242. <https://doi.org/10.1016/j.nut.2015.04.014>.
- Asemi, Z., Samimi, M., Tabassi, Z., Esmailzadeh, A., 2014. The effect of DASH diet on pregnancy outcomes in gestational diabetes: a randomized controlled clinical trial. *Eur. J. Clin. Nutr.* 68, 490–495. <https://doi.org/10.1038/ejcn.2013.296>.
- Asemi, Z., Samimi, M., Tabassi, Z., Sabihi, S., Esmailzadeh, A., 2013. A randomized controlled clinical trial investigating the effect of DASH diet on insulin resistance, inflammation, and oxidative stress in gestational diabetes. *Nutrition* 29, 619–624. <https://doi.org/10.1016/j.nut.2012.11.020>.
- Aslalah, H., Jamilian, M., Khosrowbeygi, A., 2019a. Elevation of the adiponectin/leptin ratio in women with gestational diabetes mellitus after supplementation with alpha-lipoic acid. *Gynecol. Endocrinol.* 35, 271–275. <https://doi.org/10.1080/09513590.2018.1519795>.
- Aslalah, H., Jamilian, M., Rafiei, F., Khosrowbeygi, A., 2019b. Reduction in maternal serum values of glucose and gamma-glutamyltransferase after supplementation with alpha-lipoic acid in women with gestational diabetes mellitus. *J. Obstet. Gynaecol. Res.* 45, 313–317. <https://doi.org/10.1111/jog.13842>.
- Attali, E., Yogev, Y., 2021. The impact of advanced maternal age on pregnancy outcome. *Best Pract. Res. Clin. Obstet. Gynaecol.* 70, 2–9. <https://doi.org/10.1016/j.bpobgyn.2020.06.006>.
- Benzie, I.F.F., Strain, J.J., 1996. The ferric reducing ability of plasma (FRAP) as a measure of “antioxidant power”: the FRAP assay. *Anal. Biochem.* 239, 70–76. <https://doi.org/10.1006/abio.1996.0292>.
- Bos, M., Schoots, M.H., Fernandez, B.O., Lelinska, M.M., Lau, L.C., Eikmans, M., Goor, H. V., Gordijn, S.J., Pasch, A., Feelisch, M., van der Hoorn, M.-L.P., 2019. Reactive species interactome alterations in oocyte donation pregnancies in the absence and presence of pre-eclampsia. *Int. J. Mol. Sci.* 20 <https://doi.org/10.3390/ijms20051150>.
- Boskovic, R., Gargaun, L., Oren, D., Djulus, J., Koren, G., 2005. Pregnancy outcome following high doses of Vitamin E supplementation. *Reprod. Toxicol.* 20, 85–88. <https://doi.org/10.1016/j.reprotox.2005.01.003>.
- Brosens, I., Pijnenborg, R., Vercruyse, L., Romero, R., 2011. The “Great Obstetrical Syndromes” are associated with disorders of deep placentation. *Am. J. Obstet. Gynecol.* 204 <https://doi.org/10.1016/j.ajog.2010.08.009>.
- Burton, G.J., Jauniaux, E., 2011. Oxidative stress. *Best Pract. Res. Clin. Obstet. Gynaecol.* 25, 287–299. <https://doi.org/10.1016/j.bpobgyn.2010.10.016>.
- Cabalín, C., Villalobos-Labra, R., Toledo, F., Sobrevia, L., 2019. Involvement of A2B adenosine receptors as anti-inflammatory in gestational diabetes. *Mol. Aspect. Med.* 66, 31–39. <https://doi.org/10.1016/j.mam.2019.01.001>.
- Campmans-Kuijpers, M.J.E., Dijkstra, G., 2021. Food and food groups in inflammatory bowel disease (IBD): the design of the groningen anti-inflammatory diet (GrAID). *Nutrients* 13, 1067. <https://doi.org/10.3390/nu13041067>.
- Carrasco-Wong, I., Moller, A., Giachini, F.R., Lima, V.V., Toledo, F., Stojanova, J., Sobrevia, L., San Martín, S., 2020. Placental structure in gestational diabetes mellitus. *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1866, 165535 <https://doi.org/10.1016/j.bbadis.2019.165535>.
- Casanueva, E., Ripoll, C., Tolentino, M., Morales, R.M., Pfeffer, F., Vilchis, P., Vellido-Ortega, F., 2005. Vitamin C supplementation to prevent premature rupture of the chorioamniotic membranes: a randomized trial. *Am. J. Clin. Nutr.* 81, 859–863. <https://doi.org/10.1093/ajcn/81.4.859>.
- Casanueva, E., Viteri, F.E., 2003. Iron and oxidative stress in pregnancy. *J. Nutr.* 133, 1700S–1708S. <https://doi.org/10.1093/jn/133.5.1700S>.
- Chan, K.Y., Wong, M.M.H., Pang, S.S.H., Lo, K.K.H., 2021. Dietary supplementation for gestational diabetes prevention and management: a meta-analysis of randomized controlled trials. *Arch. Gynecol. Obstet.* 303, 1381–1391. <https://doi.org/10.1007/s00404-021-06023-9>.
- Chiarello, D.I., Abad, C., Rojas, D., Toledo, F., Vázquez, C.M., Mate, A., Sobrevia, L., Marín, R., 2020. Oxidative stress: normal pregnancy versus preeclampsia. *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1866, 165354 <https://doi.org/10.1016/j.bbadis.2018.12.005>.
- Childs, C., Calder, P., Miles, E., 2019. Diet and immune function. *Nutrients* 11, 1933. <https://doi.org/10.3390/nu11081933>.
- Cornejo, M., Fuentes, G., Valero, P., Vega, S., Grismaldo, A., Toledo, F., Pardo, F., Moore-Carrasco, R., Subiabre, M., Casanella, P., Faas, M.M., Goor, H., Sobrevia, L., 2021. Gestational diabetes and foetoplacental vascular dysfunction. *Acta Physiol.* 232 <https://doi.org/10.1111/apha.13671>.
- Escudero, C., Puebla, C., Westermeier, F., Sobrevia, L., 2009. Potential cell signalling mechanisms involved in differential placental angiogenesis in mild and severe preeclampsia. *Curr. Vasc. Pharmacol.* 7, 475–485. <https://doi.org/10.2174/157016109789043865>.
- Faas, M.M., de Vos, P., 2017. Uterine NK cells and macrophages in pregnancy. *Placenta* 56, 44–52. <https://doi.org/10.1016/j.placenta.2017.03.001>.

- Farrant, H.J.W., Krishnaveni, G.V., Hill, J.C., Boucher, B.J., Fisher, D.J., Noonan, K., Osmond, C., Veena, S.R., Fall, C.H.D., 2009. Vitamin D insufficiency is common in Indian mothers but is not associated with gestational diabetes or variation in newborn size. *Eur. J. Clin. Nutr.* 63, 646–652. <https://doi.org/10.1038/ejcn.2008.14>.
- Fogacci, S., Fogacci, F., Banach, M., Michos, E.D., Hernandez, A.V., Lip, G.Y.H., Blaha, M. J., Toth, P.P., Borghi, C., Cicero, A.F.G., 2020. Vitamin D supplementation and incident preeclampsia: a systematic review and meta-analysis of randomized clinical trials. *Clin. Nutr.* 39, 1742–1752. <https://doi.org/10.1016/j.clnu.2019.08.015>.
- Gabbay-Benziv, R., Baschat, A.A., 2015. Gestational diabetes as one of the “great obstetrical syndromes” – the maternal, placental, and fetal dialog. *Best Pract. Res. Clin. Obstet. Gynaecol.* 29, 150–155. <https://doi.org/10.1016/j.bpobgyn.2014.04.025>.
- García-Patterson, A., Balsells, M., Yamamoto, J.M., Kellett, J.E., Solà, I., Gich, I., van der Beek, E.M., Hadar, E., Castañeda-Gutiérrez, E., Heinonen, S., Hod, M., Laitinen, K., Olsen, S.F., Poston, L., Rueda, R., Rust, P., van Lieshout, L., Schelk, B., Murphy, H. R., Corcoy, R., 2019. Usual dietary treatment of gestational diabetes mellitus assessed after control diet in randomized controlled trials: subanalysis of a systematic review and meta-analysis. *Acta Diabetol.* 56, 237–240. <https://doi.org/10.1007/s00592-018-1238-4>.
- Gatford, K.L., Houda, C.M., Lu, Z.X., Coat, S., Baghurst, P.A., Owens, J.A., Sikaris, K., Rowan, J.A., Hague, W.M., 2013. Vitamin B₁₂ and homocysteine status during pregnancy in the metformin in gestational diabetes trial: responses to maternal metformin compared with insulin treatment. *Diabetes Obes. Metabol.* 15, 660–667. <https://doi.org/10.1111/dom.12080>.
- Gázquez, A., Giménez-Bañón, M., Prieto-Sánchez, M., Martínez-Graciá, C., Suárez, C., Santaella-Pascual, M., Galdo-Castañeira, L., Ballesteros-Meseguer, C., Vioque, J., Martínez-Villanueva, M., Avilés-Plaza, F., Noguera-Velasco, J., Morales, E., García-Marcos, L., Larqué, E., 2021. Self-reported DHA supplementation during pregnancy and its association with obesity or gestational diabetes in relation to DHA concentration in cord and maternal plasma: results from NELA, a prospective mother-offspring cohort. *Nutrients* 13, 843. <https://doi.org/10.3390/nu13030843>.
- Gomes, F., Bourassa, M.W., Adu-Afaruwah, S., Ajello, C., Bhutta, Z.A., Black, R., Catarino, E., Chowdhury, R., Dalmiya, N., Dwarkanath, P., Engle-Stone, R., Gernand, A.D., Goudet, S., Hodinott, J., Kästel, P., Manger, M.S., McDonald, C.M., Mehta, S., Moore, S.E., Neufeld, L.M., Osendarp, S., Ramachandran, P., Rasmussen, K.M., Stewart, C., Sudfeld, C., West, K., Bergeron, G., 2020. Setting research priorities on multiple micronutrient supplementation in pregnancy. *Ann. N. Y. Acad. Sci.* 1465, 76–88. <https://doi.org/10.1111/nyas.14267>.
- Gómez-Roig, M.D., Pascal, R., Cahuana, M.J., García-Algar, O., Sebastiani, G., Andreu-Fernández, V., Martínez, L., Rodríguez, G., Iglesia, I., Ortiz-Arrabal, O., Mesa, M.D., Cabero, M.J., Guerra, L., Llurba, E., Domínguez, C., Zanini, M.J., Foraster, M., Larqué, E., Cabañas, F., Lopez-Azorín, M., Pérez, A., Loureiro, B., Pallás-Alonso, C.R., Escuder-Vieco, D., Vento, M., 2021. Environmental exposure during pregnancy: influence on prenatal development and early life: a comprehensive review. *Fetal Diagn. Ther.* 48, 245–257. <https://doi.org/10.1159/000514884>.
- Guerin, L.R., Prins, J.R., Robertson, S.A., 2009. Regulatory T-cells and immune tolerance in pregnancy: a new target for infertility treatment? *Hum. Reprod. Update* 15, 517–535. <https://doi.org/10.1093/humupd/dmp004>.
- Gunasegaran, P., Tahmina, S., Daniel, M., Nanda, S.K., 2021. Role of vitamin D-calcium supplementation on metabolic profile and oxidative stress in gestational diabetes mellitus: a randomized controlled trial. *J. Obstet. Gynaecol. Res.* 47, 1016–1022. <https://doi.org/10.1111/jog.14629>.
- Hanson, M.A., Bardsley, A., De-Regil, L.M., Moore, S.E., Oken, E., Poston, L., Ma, R.C., McAuliffe, F.M., Maleta, K., Purandare, C.N., Yajnik, C.S., Rushwan, H., Morris, J.L., 2015. The international federation of gynecology and obstetrics (FIGO) recommendations on adolescent, preconception, and maternal nutrition: “think nutrition first”[#]. *Int. J. Gynecol. Obstet.* 131, S213–S253. [https://doi.org/10.1016/S0020-7292\(15\)30034-5](https://doi.org/10.1016/S0020-7292(15)30034-5).
- Harreiter, J., Desoye, G., van Poppel, M.N.M., Kautzky-Willer, A., Dunne, F., Corcoy, R., Devlieger, R., Simmonds, D., Adelantado, J.M., Damm, P., Mathiesen, E.R., Jensen, D. M., Anderson, L.L.T., Lapolla, A., Dalfra, M.G., Bertolotto, A., Wender-Ozegowska, E., Zawiejska, A., Hill, D.J., Snoek, F.J., 2019. The effects of lifestyle and/or vitamin D supplementation interventions on pregnancy outcomes: what have we learned from the DALI studies? *Curr. Diabetes Rep.* 19, 162. <https://doi.org/10.1007/s11892-019-1282-7>.
- Haugen, M., Brantsæter, A.L., Trogstad, L., Alexander, J., Roth, C., Magnus, P., Meltzer, H.M., 2009. Vitamin D supplementation and reduced risk of preeclampsia in nulliparous women. *Epidemiology* 20, 720–726. <https://doi.org/10.1097/EDE.0b013e3181a70f08>.
- Hawkes, W.C., Alkan, Z., Lang, K., King, J.C., 2004. Plasma selenium decrease during pregnancy is associated with glucose intolerance. *Biol. Trace Elem. Res.* 100, 19–30. <https://doi.org/10.1385/BTER:100:1:019>.
- Hoffman, D.J., Powell, T.L., Barrett, E.S., Hardy, D.B., 2021. Developmental origins of metabolic diseases. *Physiol. Rev.* 101, 739–795. <https://doi.org/10.1152/physrev.00002.2020>.
- Hosseinzadeh-Shamsi-Anar, M., Mozaffari-Khosravi, H., Salami, M.A., Hadinedoushan, H., Mozayan, M.R., 2012. The efficacy and safety of a high dose of vitamin d in mothers with gestational diabetes mellitus: a randomized controlled clinical trial. *Iran. J. Med. Sci.* 37, 159–165.
- Huang, S., Fu, J., Zhao, R., Wang, B., Zhang, M., Li, L., Shi, C., 2021. The effect of combined supplementation with vitamin D and omega-3 fatty acids on blood glucose and blood lipid levels in patients with gestational diabetes. *Ann. Palliat. Med.* 10, 5652–5658. <https://doi.org/10.21037/apm-21-118>.
- Huynh, J., Dawson, D., Roberts, D., Bentley-Lewis, R., 2015. A systematic review of placental pathology in maternal diabetes mellitus. *Placenta* 36, 101–114. <https://doi.org/10.1016/j.placenta.2014.11.021>.
- Jamilian, M., Hashemi Dizaji, S., Bahmani, F., Taghizadeh, M., Memarzadeh, M.R., Karamali, M., Akbari, M., Asemi, Z., 2017. A randomized controlled clinical trial investigating the effects of omega-3 fatty acids and vitamin E Co-supplementation on biomarkers of oxidative stress, inflammation and pregnancy outcomes in gestational diabetes. *Can. J. Diabetes* 41, 143–149. <https://doi.org/10.1016/j.cjcd.2016.09.004>.
- Jamilian, M., Mirhosseini, N., Eslahi, M., Bahmani, F., Shokrpour, M., Chamani, M., Asemi, Z., 2019. The effects of magnesium-zinc-calcium-vitamin D co-supplementation on biomarkers of inflammation, oxidative stress and pregnancy outcomes in gestational diabetes. *BMC Pregnancy Childbirth* 19, 107. <https://doi.org/10.1186/s12884-019-2258-y>.
- Jamilian, M., Samimi, M., Afshar Ebrahimi, F., Aghadavod, E., Mohammadbeigi, R., Rahimi, M., Asemi, Z., 2018. Effects of selenium supplementation on gene expression levels of inflammatory cytokines and vascular endothelial growth factor in patients with gestational diabetes. *Biol. Trace Elem. Res.* 181, 199–206. <https://doi.org/10.1007/s12011-017-1045-8>.
- Jauniaux, E., Burton, G.J., 2006. Villous histomorphometry and placental bed biopsy investigation in type I diabetic pregnancies. *Placenta* 27, 468–474. <https://doi.org/10.1016/j.placenta.2005.04.010>.
- Jirakitidul, P., Sirichotiyakul, S., Ruengorn, C., Techatraksa, K., Wiriyastrivaj, B., 2018. Effect of iron supplementation during early pregnancy on the development of gestational hypertension and pre-eclampsia. *Arch. Gynecol. Obstet.* 298, 545–550. <https://doi.org/10.1007/s00404-018-4821-6>.
- Jones, M.L., Mark, P.J., Waddell, B.J., 2014. Maternal dietary omega-3 fatty acids and placental function. *Reproduction* 147, R143–R152. <https://doi.org/10.1530/REP-13-0376>.
- Kalpdev, A., Saha, S.C., Dhawan, V., 2011. Vitamin C and E supplementation does not reduce the risk of superimposed PE in pregnancy. *Hypertens. Pregnancy* 30, 447–456. <https://doi.org/10.3109/10641955.2010.507840>.
- Karamali, M., Dastyar, F., Badakhsh, M.H., Aghadavod, E., Amirani, E., Asemi, Z., 2020. The effects of selenium supplementation on gene expression related to insulin and lipid metabolism, and pregnancy outcomes in patients with gestational diabetes mellitus: a randomized, double-blind, placebo-controlled trial. *Biol. Trace Elem. Res.* 195, 1–8. <https://doi.org/10.1007/s12011-019-01818-z>.
- Khong, T.Y., Mooney, E.E., Ariel, I., Balmus, N.C.M., Boyd, T.K., Brundler, M.-A., Derricott, H., Evans, M.J., Faye-Petersen, O.M., Gillan, J.E., Heazell, A.E.P., Heller, D.S., Jacques, S.M., Keating, S., Kelehan, P., Maes, A., McKay, E.M., Morgan, T.K., Nikkels, P.G.J., Parks, W.T., Redline, R.W., Scheimberg, I., Schoots, M. H., Sebire, N.J., Timmer, A., Turowski, G., van der Voorn, J.P., van Lijnschoten, I., Gordijn, S.J., 2016. Sampling and definitions of placental lesions: amsterdam placental workshop group consensus statement. *Arch. Pathol. Lab. Med.* 140, 698–713. <https://doi.org/10.5858/arpa.2015-0225-CC>.
- Kiondo, P., Wamuyu-Maina, G., Wandabwa, J., Bimenya, G.S., Tumwesigye, N.M., Okong, P., 2014. The effects of vitamin C supplementation on pre-eclampsia in Mulago Hospital, Kampala, Uganda: a randomized placebo controlled clinical trial. *BMC Pregnancy Childbirth* 14, 283. <https://doi.org/10.1186/1471-2393-14-283>.
- Korkmaz, V., Ozkaya, E., Seven, B.Y., Duzguner, S., Karlı, M.F., Kucukozkan, T., 2014. Comparison of oxidative stress in pregnancies with and without first trimester iron supplement: a randomized double-blind controlled trial. *J. Matern. Neonatal Med.* 27, 1535–1538. <https://doi.org/10.3109/14767058.2013.863869>.
- Krishnaveni, G.V., Hill, J.C., Veena, S.R., Bhat, D.S., Wills, A.K., Karat, C.L.S., Yajnik, C. S., Fall, C.H.D., 2009. Low plasma vitamin B12 in pregnancy is associated with gestational ‘diabesity’ and later diabetes. *Diabetologia* 52, 2350–2358. <https://doi.org/10.1007/s00125-009-1499-0>.
- Kupper, N., Huppertz, B., 2021. The endogenous exosome of the pregnant mother: placental extracellular vesicles and their effect on the maternal system. *Mol. Aspect. Med.* 100955. <https://doi.org/10.1016/j.jam.2021.100955>.
- Lai, J.S., Pang, W.W., Cai, S., Lee, Y.S., Chan, J.K.Y., Shek, L.P.C., Yap, F.K.P., Tan, K.H., Godfrey, K.M., van Dam, R.M., Chong, Y.S., Chong, M.F.F., 2018. High folate and low vitamin B12 status during pregnancy is associated with gestational diabetes mellitus. *Clin. Nutr.* 37, 940–947. <https://doi.org/10.1016/j.clnu.2017.03.022>.
- Larqué, E., Pagán, A., Prieto, M.T., Blanco, J.E., Gil-Sánchez, A., Zornoza-Moreno, M., Ruiz-Palacios, M., Gázquez, A., Demmelair, H., Parrilla, J.J., Koletzko, B., 2014. Placental fatty acid transfer: a key factor in fetal growth. *Ann. Nutr. Metab.* 64, 247–253. <https://doi.org/10.1159/000365028>.
- Liu, Z., Meng, T., Liu, J., Xu, X., Luo, S., Jin, C., Han, N., Wang, H.-J., 2020. The individual and joint effects of maternal 25(OH)D deficiency and gestational diabetes on infant birth size. *Nutr. Metabol. Cardiovasc. Dis.* 30, 2398–2405. <https://doi.org/10.1016/j.numecd.2020.07.046>.
- Mahdi, A., Cortese-Krott, M.M., Kelm, M., Li, N., Pernow, J., 2021. Novel perspectives on redox signaling in red blood cells and platelets in cardiovascular disease. *Free Radic. Biol. Med.* 168, 95–109. <https://doi.org/10.1016/j.freeradbiomed.2021.03.020>.
- Mahdy, Z.A., Siraj, H.H., Khaza’ai, H., Mutalib, M.S.A., Azwar, M.H., Wahab, M.A., Dali, A.Z.H.M., Jaafar, R., Ismail, N.A.M., Jamil, M.A., Adeeb, N., 2013. Does palm oil vitamin E reduce the risk of pregnancy induced hypertension? *Acta Medica (Hradec Kral. Czech Republic)* 56, 104–109. <https://doi.org/10.14712/18059694.2014.17>.
- Mann, G.E., Yudilevich, D.L., Sobrevia, L., 2003. Regulation of amino acid and glucose transporters in endothelial and smooth muscle cells. *Physiol. Rev.* 83, 183–252. <https://doi.org/10.1152/physrev.00022.2002>.
- Mannaerts, D., Faes, E., Cos, P., Briede, J.J., Gyselaers, W., Cornette, J., Gorbanev, Y., Bogaerts, A., Spaanderma, M., Van Craenenbroeck, E., Jacquemyn, Y., 2018. Oxidative stress in healthy pregnancy and preeclampsia is linked to chronic

- inflammation, iron status and vascular function. *PLoS One* 13, e0202919. <https://doi.org/10.1371/journal.pone.0202919>.
- Mastrolia, S.A., Cetin, I., 2020. The “great obstetrical syndromes.” https://doi.org/10.1007/978-3-030-14782-2_21, 411–430.
- McCabe, T.J., Fulton, D., Roman, L.J., Sessa, W.C., 2000. Enhanced electron flux and reduced calmodulin dissociation may explain “calcium-independent” eNOS activation by phosphorylation. *J. Biol. Chem.* 275, 6123–6128. <https://doi.org/10.1074/jbc.275.9.6123>.
- McCance, D.R., Holmes, V.A., Maresh, M.J., Patterson, C.C., Walker, J.D., Pearson, D.W., Young, I.S., 2010. Vitamins C and E for prevention of pre-eclampsia in women with type 1 diabetes (DAPIT): a randomised placebo-controlled trial. *Lancet* 376, 259–266. [https://doi.org/10.1016/S0140-6736\(10\)60630-7](https://doi.org/10.1016/S0140-6736(10)60630-7).
- Mirzakhani, H., Litonjua, A.A., McElrath, T.F., O'Connor, G., Lee-Parritz, A., Iverson, R., Macones, G., Strunk, R.C., Bacharier, L.B., Zeiger, R., Hollis, B.W., Handy, D.E., Sharma, A., Laranjo, N., Carey, V., Qiu, W., Santolini, M., Liu, S., Chhabra, D., Enquobahrie, D.A., Williams, M.A., Loscalzo, J., Weiss, S.T., 2016. Early pregnancy vitamin D status and risk of preeclampsia. *J. Clin. Invest.* 126, 4702–4715. <https://doi.org/10.1172/JCI89031>.
- Molinari, C., Uberti, F., Grossini, E., Vacca, G., Carda, S., Invernizzi, M., Cisari, C., 2011. 1 α ,25-Dihydroxycholecalciferol induces nitric oxide production in cultured endothelial cells. *Cell. Physiol. Biochem.* 27, 661–668. <https://doi.org/10.1159/000330075>.
- Moncada, S., Palmer, R.M., Higgs, E.A., 1988. The discovery of nitric oxide as the endogenous nitrovasodilator. *Hypertension* 12, 365–372. <https://doi.org/10.1161/01.HYP.12.4.365>.
- Munoz-Suano, A., Hamilton, A.B., Betz, A.G., 2011. Gimme shelter: the immune system during pregnancy. *Immunol. Rev.* 241, 20–38. <https://doi.org/10.1111/j.1600-065X.2011.01002.x>.
- Nadeau-Vallée, M., Obari, D., Palacios, J., Brien, M.-È., Duval, C., Chemtob, S., Girard, S., 2016. Sterile inflammation and pregnancy complications: a review. *Reproduction* 152, R277–R292. <https://doi.org/10.1530/REP-16-0453>.
- Nguyen, L.N., Ma, D., Shui, G., Wong, P., Cazenave-Gassiot, A., Zhang, X., Wenk, M.R., Goh, E.L.K., Silver, D.L., 2014. Mfsd2a is a transporter for the essential omega-3 fatty acid docosahexaenoic acid. *Nature* 509, 503–506. <https://doi.org/10.1038/nature13241>.
- O'Callaghan, K., Kiely, M., 2018. Systematic review of vitamin D and hypertensive disorders of pregnancy. *Nutrients* 10, 294. <https://doi.org/10.3390/nu10030294>.
- Offringa, A.K., Bourgonje, A.R., Schrier, M.S., Deth, R.C., van Goor, H., 2021. Clinical implications of vitamin B12 as redox-active cofactor. *Trends Mol. Med.* 27, 931–934. <https://doi.org/10.1016/j.molmed.2021.07.002>.
- Palacios, C., Trak-Fellermeier, M.A., Martínez, R.X., Lopez-Perez, L., Lips, P., Salisi, J.A., John, J.C., Peña-Rosas, J.P., 2019. Regimens of vitamin D supplementation for women during pregnancy. *Cochrane Database Syst. Rev.* <https://doi.org/10.1002/14651858.CD013446>.
- Pizzino, G., Irrera, N., Cucinotta, M., Pallio, G., Mannino, F., Arcoraci, V., Squadrone, F., Altavilla, D., Bitto, A., 2017. Oxidative stress: harms and benefits for human health. *Oxid. Med. Cell. Longev.* 1–13. <https://doi.org/10.1155/2017/8416763>, 2017.
- Poole, J.A., Claman, H.N., 2004. Immunology of pregnancy: implications for the mother. *Clin. Rev. Allergy Immunol.* 26, 161–170. <https://doi.org/10.1385/CRIAI:26:3:161>.
- Poston, L., Briley, A., Seed, P., Kelly, F., Shenan, A., 2006. Vitamin C and vitamin E in pregnant women at risk for pre-eclampsia (VIP trial): randomised placebo-controlled trial. *Lancet* 367, 1145–1154. [https://doi.org/10.1016/S0140-6736\(06\)68433-X](https://doi.org/10.1016/S0140-6736(06)68433-X).
- Prieto-Sánchez, M.T., Ruiz-Palacios, M., Blanco-Carnero, J.E., Pagan, A., Hellmuth, C., Uhl, O., Peissner, W., Ruiz-Alcaraz, A.J., Parrilla, J.J., Koletzko, B., Larqué, E., 2017. Placental MFSD2a transporter is related to decreased DHA in cord blood of women with treated gestational diabetes. *Clin. Nutr.* 36, 513–521. <https://doi.org/10.1016/j.clnu.2016.01.014>.
- Radzicka, S., Ziolkowska, K., Zaborowski, M.P., Brazert, J., Pietryga, M., 2019. Serum homocysteine and vitamin B12 levels in women with gestational diabetes mellitus. *Ginekol. Pol.* 90, 381–387. <https://doi.org/10.5603/GP.2019.0066>.
- Ramírez, M.A., Morales, J., Cornejo, M., Blanco, E.H., Mancilla-Sierpe, E., Toledo, F., Beltrán, A.R., Sobrevia, L., 2018. Intracellular acidification reduces L-arginine transport via system y⁺L but not via system y⁺/CATs and nitric oxide synthase activity in human umbilical vein endothelial cells. *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1864, 1192–1202. <https://doi.org/10.1016/j.bbadis.2018.01.032>.
- Razavi, M., Jamilian, M., Kashan, Z.F., Heidari, Z., Mohseni, M., Ghandi, Y., Bagherian, T., Asemi, Z., 2016. Selenium supplementation and the effects on reproductive outcomes, biomarkers of inflammation, and oxidative stress in women with polycystic ovary syndrome. *Horm. Metab. Res.* 48, 185–190. <https://doi.org/10.1055/s-0035-1559604>.
- Redman, C., 2014. The six stages of pre-eclampsia. *Pregnancy Hypertens.* 4, 246. <https://doi.org/10.1016/j.preghy.2014.04.020>.
- Redman, C.W.G., Staff, A.C., Roberts, J.M., 2022. Syncytiotrophoblast stress in preeclampsia: the convergence point for multiple pathways. *Am. J. Obstet. Gynecol.* 226, S907–S927. <https://doi.org/10.1016/j.ajog.2020.09.047>.
- Roberts, J.M., 2014. Pathophysiology of ischemic placental disease. *Semin. Perinatol.* 38, 139–145. <https://doi.org/10.1053/j.semperi.2014.03.005>.
- Roberts, J.M., Myatt, L., Spong, C.Y., Thom, E.A., Hauth, J.C., Leveno, K.J., Pearson, G. D., Wapner, R.J., Varner, M.W., Thorp, J.M., Mercer, B.M., Peaceman, A.M., Ramin, S.M., Carpenter, M.W., Samuels, P., Sciscione, A., Harper, M., Smith, W.J., Saade, G., Sorokin, Y., Anderson, G.B., 2010. Vitamins C and E to prevent complications of pregnancy-associated hypertension. *N. Engl. J. Med.* 362, 1282–1291. <https://doi.org/10.1056/NEJMoa0908056>.
- Robertson, S.A., Green, E.S., Care, A.S., Moldenhauer, L.M., Prins, J.R., Hull, M.L., Barry, S.C., Dekker, G., 2019. Therapeutic potential of regulatory T cells in preeclampsia—opportunities and challenges. *Front. Immunol.* 10 <https://doi.org/10.3389/fimmu.2019.00478>.
- Robinson, O., Vrijheid, M., 2015. The pregnancy exposome. *Curr. Environ. Heal. Rep.* 2, 204–213. <https://doi.org/10.1007/s40572-015-0043-2>.
- Rumbold, A., Ota, E., Nagata, C., Shahrook, S., Crowther, C.A., 2015. Vitamin C supplementation in pregnancy. *Cochrane Database Syst. Rev.* <https://doi.org/10.1002/14651858.CD004072.pub3>.
- Rytlewski, K., Olszanecki, R., Korbut, R., Zdebski, Z., 2005. Effects of prolonged oral supplementation with L-arginine on blood pressure and nitric oxide synthesis in preeclampsia. *Eur. J. Clin. Invest.* 35, 32–37. <https://doi.org/10.1111/j.1365-2362.2005.01445.x>.
- Sablok, A., Batra, Aruna, Thariani, K., Batra, Achla, Bharti, R., Aggarwal, A.R., Kabi, B.C., Chellani, H., 2015. Supplementation of vitamin D in pregnancy and its correlation with foeto-maternal outcome. *Clin. Endocrinol.* 83, 536–541. <https://doi.org/10.1111/cen.12751>.
- Schoots, M.H., Bourgonje, M.F., Bourgonje, A.R., Prins, J.R., van Hoorn, E.G.M., Abdulle, A.E., Muller Kobold, A.C., van der Heide, M., Hillebrands, J.-L., van Goor, H., Gordijn, S.J., 2021. Oxidative stress biomarkers in fetal growth restriction with and without preeclampsia. *Placenta* 115, 87–96. <https://doi.org/10.1016/j.placenta.2021.09.013>.
- Schoots, M.H., Gordijn, S.J., Scherjon, S.A., van Goor, H., Hillebrands, J.-L., 2018. Oxidative stress in placental pathology. *Placenta* 69, 153–161. <https://doi.org/10.1016/j.placenta.2018.03.003>.
- Schröder-Heurich, B., Springer, C.J.P., von Versen-Höyneck, F., 2020. Vitamin D effects on the immune system from periconception through pregnancy. *Nutrients* 12, 1432. <https://doi.org/10.3390/nu12051432>.
- Shoji, H., Franke, C., Demmelmaier, H., Koletzko, B., 2009. Effect of docosahexaenoic acid on oxidative stress in placental trophoblast cells. *Early Hum. Dev.* 85, 433–437. <https://doi.org/10.1016/j.earlhumdev.2009.02.003>.
- Siddiqui, I.A., Jaleel, A., Kadri, H.M.F. Al, Saeed, W. Al, Tamimi, W., 2011. Iron status parameters in preeclamptic women. *Arch. Gynecol. Obstet.* 284, 587–591. <https://doi.org/10.1007/s00404-010-1728-2>.
- Simopoulos, A., 2002. The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed. Pharmacother.* 56, 365–379. [https://doi.org/10.1016/S0753-3322\(02\)00253-6](https://doi.org/10.1016/S0753-3322(02)00253-6).
- Sobrevia, L., Valero, P., Grismaldo, A., Villalobos-Labra, R., Pardo, F., Subiabre, M., Armstrong, G., Toledo, F., Vega, S., Cornejo, M., Fuentes, G., Marín, R., 2020. Mitochondrial dysfunction in the fetoplacental unit in gestational diabetes mellitus. *Biochim. Biophys. Acta (BBA) - Mol. Basis Dis.* 1866, 165948 <https://doi.org/10.1016/j.bbadis.2020.165948>.
- Steyn, P.S., Odendaal, H.J., Schoeman, J., Stander, C., Fanie, N., Grové, D., 2003. A randomised, double-blind placebo-controlled trial of ascorbic acid supplementation for the prevention of preterm labour. *J. Obstet. Gynaecol. (Lahore)* 23, 150–155. <https://doi.org/10.1080/014436103000074673>.
- Stone, J., Suttrave, P., Gascoigne, E., Givens, M.B., Fry, R.C., Manuck, T.A., 2021. Exposure to toxic metals and per- and polyfluoroalkyl substances and the risk of preeclampsia and preterm birth in the United States: a review. *Am. J. Obstet. Gynecol. MFM* 3, 100308. <https://doi.org/10.1016/j.ajogmf.2021.100308>.
- Tenório, M.B., Ferreira, R.C., Moura, F.A., Bueno, N.B., de Oliveira, A.C.M., Goulart, M. O.F., 2019. Cross-talk between oxidative stress and inflammation in preeclampsia. *Oxid. Med. Cell. Longev.* 1–26. <https://doi.org/10.1155/2019/8238727>, 2019.
- Terstappen, F., Tol, A.J.C., Gremmels, H., Wever, K.E., Paauw, N.D., Joles, J.A., M van der Beek, E., Lely, A.T., 2020. Prenatal amino acid supplementation to improve fetal growth: a systematic review and meta-analysis. *Nutrients* 12, 2535. <https://doi.org/10.3390/nu12092535>.
- Uberti, F., Lattuada, D., Morsanuto, V., Nava, U., Bolis, G., Vacca, G., Squarzanti, D.F., Cisari, C., Molinari, C., 2014. Vitamin D protects human endothelial cells from oxidative stress through the autophagic and survival pathways. *J. Clin. Endocrinol. Metab.* 99, 1367–1374. <https://doi.org/10.1210/jc.2013-2103>.
- Vadillo-Ortega, F., Perichart-Perera, O., Espino, S., Avila-Vergara, M.A., Ibarra, I., Ahued, R., Godines, M., Parry, S., Macones, G., Strauss, J.F., 2011. Effect of supplementation during pregnancy with L-arginine and antioxidant vitamins in medical food on pre-eclampsia in high risk population: randomised controlled trial. *BMJ* 342. <https://doi.org/10.1136/bmj.d2901> d2901–d2901.
- Valero, P., Fuentes, G., Cornejo, M., Vega, S., Grismaldo, A., Pardo, F., García-Rivas, G., Hillebrands, J.-L., Faas, M.M., Casanello, P., van der Beek, E.M., van Goor, H., Sobrevia, L., 2021. Exposome and foetoplacental vascular dysfunction in gestational diabetes mellitus. *Mol. Aspect. Med.* 101019 <https://doi.org/10.1016/j.mam.2021.101019>.
- van der Zwan, A., van Unen, V., Beyrend, G., Laban, S., van der Keur, C., Kapsenberg, H. J.M., Höllt, T., Chuva de Sousa Lopes, S.M., van der Hoorn, M.-L.P., Koning, F., Claas, F.H.J., Eikmans, M., Heidt, S., 2020. Visualizing dynamic changes at the maternal-fetal interface throughout human pregnancy by mass cytometry. *Front. Immunol.* 11 <https://doi.org/10.3389/fimmu.2020.571300>.
- Wibowo, N., Purwosunu, Y., Sekizawa, A., Farina, A., Idriansyah, L., Fitriana, I., 2012. Antioxidant supplementation in pregnant women with low antioxidant status. *J. Obstet. Gynaecol. Res.* 38, 1152–1161. <https://doi.org/10.1111/j.1447-0756.2012.01855.x>.
- Wild, C.P., 2012. The exposome: from concept to utility. *Int. J. Epidemiol.* 41, 24–32. <https://doi.org/10.1093/ije/dyr236>.
- Wild, C.P., 2005. Complementing the genome with an “exposome”: the outstanding challenge of environmental exposure measurement in molecular epidemiology. *Cancer Epidemiol. Biomarkers Prev.* 14, 1847–1850. <https://doi.org/10.1158/1055-9965.EPI-05-0456>.
- World Health Organization (WHO), 2016. WHO Recommendations on Antenatal Care for a Positive Pregnancy Experience.

- Wu, F., Tian, F.-J., Lin, Y., Xu, W.-M., 2016. Oxidative stress: placenta function and dysfunction. *Am. J. Reprod. Immunol.* 76, 258–271. <https://doi.org/10.1111/aji.12454>.
- Wu, M., Wu, Y., Xu, K., Lin, L., 2021. Protective effects of 1,25 dihydroxyvitamin D₃ against high-glucose-induced damage in human umbilical vein endothelial cells involve activation of Nrf2 antioxidant signaling. *J. Vasc. Res.* 58, 267–276. <https://doi.org/10.1159/000515512>.
- Xu, H., Perez-Cuevas, R., Xiong, X., Reyes, H., Roy, C., Julien, P., Smith, G., von Dadelszen, P., Leduc, L., Audibert, F., Moutquin, J.-M., Piedboeuf, B., Shatenstein, B., Parra-Cabrera, S., Choquette, P., Winsor, S., Wood, S., Benjamin, A., Walker, M., Helewa, M., Dubé, J., Tawagi, G., Seaward, G., Ohlsson, A., Magee, L.A., Olatunbosun, F., Gratton, R., Shear, R., Demianczuk, N., Collet, J.-P., Wei, S., Fraser, W.D., 2010. An international trial of antioxidants in the prevention of preeclampsia (INTAPP). *Am. J. Obstet. Gynecol.* 202 <https://doi.org/10.1016/j.ajog.2010.01.050>, 239.e1-239.e10.
- Yamamoto, J.M., Kellett, J.E., Balsells, M., García-Patterson, A., Hadar, E., Solà, I., Gich, I., van der Beek, E.M., Castañeda-Gutiérrez, E., Heinonen, S., Hod, M., Laitinen, K., Olsen, S.F., Poston, L., Rueda, R., Rust, P., van Lieshout, L., Schelkde, B., Murphy, H.R., Corcoy, R., 2018. Gestational diabetes mellitus and diet: a systematic review and meta-analysis of randomized controlled trials examining the impact of modified dietary interventions on maternal glucose control and neonatal birth weight. *Diabetes Care* 41, 1346–1361. <https://doi.org/10.2337/dc18-0102>.
- Yong, H.Y., Mohd Shariff, Z., Mohd Yusof, B.N., Rejali, Z., Tee, Y.Y.S., Bindels, J., van der Beek, E.M., 2022. Early pregnancy hemoglobin is associated with the risk of gestational diabetes mellitus: a retrospective cohort study. *Br. J. Nutr.* 1–8. <https://doi.org/10.1017/S000711452100502X>.
- Zhang, C., Qiu, C., Hu, F.B., David, R.M., van Dam, R.M., Bralley, A., Williams, M.A., 2008. Maternal plasma 25-hydroxyvitamin D concentrations and the risk for gestational diabetes mellitus. *PLoS One* 3, e3753. <https://doi.org/10.1371/journal.pone.0003753>.
- Zhang, K., 2010. Integration of ER stress, oxidative stress and the inflammatory response in health and disease. *Int. J. Clin. Exp. Med.* 3, 33–40.
- Zhang, X., Wu, M., Zhong, C., Huang, L., Zhang, Y., Chen, R., Zhou, X., Xu, S., Li, Q., Cui, W., Wang, X., Chen, X., Lin, L., Zhang, G., Xiong, G., Sun, G., Yang, X., Hao, L., Jin, Z., Yang, N., 2021. Association between maternal plasma ferritin concentration, iron supplement use, and the risk of gestational diabetes: a prospective cohort study. *Am. J. Clin. Nutr.* 114, 1100–1106. <https://doi.org/10.1093/ajcn/nqab162>.
- Zheng, H.-T., Zhou, L.-N., Huang, C.-J., Hua, X., Jian, R., Su, B.-H., Fang, F., 2008. Selenium inhibits high glucose- and high insulin-induced adhesion molecule expression in vascular endothelial cells. *Arch. Med. Res.* 39, 373–379. <https://doi.org/10.1016/j.arcmed.2007.12.007>.

Jelmer R Prins is an obstetrician-scientist and works at the University Medical Center Groningen. His research focuses on immune-mediated pregnancy complications. He has extensive experience in analysis of immune parameters in pregnancy, including characterization of blood, placental and uterine lymphocytes, and is involved in several translational pregnancy studies within the UMCG.

Mirthe H Schoots is a pathologist and works at the University Medical Center Groningen. Her regions of interest are fetal and placental pathology, neuropathology and autopsy-

related pathology. Her current research focuses on fetal growth restriction, placental pathology and placental oxidative stress.

Jule I Wessels is a physician with an interest in obstetrics and dietary interventions.

Marjo J E Campmans-Kuijpers is a post-doctoral researcher, clinical epidemiologist and research dietician with a special focus on diet in IBD. She has developed the Groningen anti-inflammatory diet (GrAID) for IBD patients, which is currently being investigated in the Vita-GrAID trial.

Gerjan J Navis is a nephrologist and professor in nephrology and diet. Her research focusses on diet and kidney diseases.

Harry van Goor is Professor in Experimental Renal Pathology with a PhD in Medical Sciences. His current research focusses on the systemic Redox status as moveable biomarker in various disease conditions. He also works on gasotransmitters like hydrogen sulfide with the aim to use these as therapeutical options in experimental and clinical conditions.

Sarah A Robertson is Professor of Reproductive Immunology at the Robinson Research Institute, University of Adelaide, and a Fellow of the Australian Academy of Science and Australian Academy for Health and Medical Sciences. Her research focuses on the biological mechanisms of immune tolerance in conception and pregnancy.

Eline M van der Beek obtained her doctoral degree at the University of Utrecht in the Netherlands. She did a postdoc at Wageningen University, after which she stayed as an Assistant Professor before moving to the food industry. She is currently appointed as Professor in Nutritional Programming at the Pediatric Department of the University Medical Centre in Groningen, the Netherlands, next to her role as Head of the Nestlé Institute of Health Sciences in Lausanne, Switzerland. Her research focuses on the contribution of maternal health and nutrition status in offspring growth and development.

Luis Sobrevia is a BSc in biological sciences holding an MSc in Physiological Sciences from the Universidad de Concepción (Chile), a PhD in Physiology and Medical Sciences and postdoctoral training in vascular physiology King's College London from University of London (UK). His research line regards human vascular endothelial dysfunction in diseases of pregnancy involving cell signalling through adenosine receptors and insulin receptors and the role of membrane transport systems in this phenomenon.

Sanne J Gordijn is an obstetrician. She is involved in standardisation in the defining FGR and structuring research protocol by development of a minimum reporting set and a core outcome set. She is cofounder and project leader of a large multicentre RCT: DRIGITAT considering SGA and the role of Doppler indices. She started a large worldwide collaboration of pathologists that resulted in an important, widely acknowledged placental definition paper and recently an authoritative textbook of placental disorders. Dr Gordijn is involved in placental translational research to understand and influence the pathophysiological background of common obstetric conditions, mainly focussed on FGR.