

3-2-2020

## Comparing Self-Reported Medication Adherence Measures With Hair Antiretroviral Concentration Among People Living With HIV in Guangxi, China

Quan Zhang

Xiaoming Li Ph.D.

University of South Carolina, xiaoming@mailbox.sc.edu

Shan Qiao Ph.D.

University of South Carolina, shanqiao@mailbox.sc.edu

For more of this author's additional works at: <https://scholarcommons.sc.edu/>

[sph\\_health\\_promotion\\_education\\_behavior\\_facpub](#)

Yuejiao Zhou

 Part of the [Public Health Education and Promotion Commons](#)


---

### Publication Info

Published in *AIDS Research and Therapy*, Volume 17, Issue 8, 2020.

© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

This Article is brought to you by the Health Promotion, Education, and Behavior at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact [digres@mailbox.sc.edu](mailto:digres@mailbox.sc.edu).



# The role of mental health on the relationship between food insecurity and immunologic outcome among people living with HIV in Guangxi, China

Cheng Chen, Xueying Yang, Chengbo Zeng, Xiaoming Li, Shan Qiao & Yuejiao Zhou

To cite this article: Cheng Chen, Xueying Yang, Chengbo Zeng, Xiaoming Li, Shan Qiao & Yuejiao Zhou (2020) The role of mental health on the relationship between food insecurity and immunologic outcome among people living with HIV in Guangxi, China, Health Psychology and Behavioral Medicine, 8:1, 636-648, DOI: [10.1080/21642850.2020.1854762](https://doi.org/10.1080/21642850.2020.1854762)

To link to this article: <https://doi.org/10.1080/21642850.2020.1854762>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



[View supplementary material](#)



Published online: 08 Dec 2020.



[Submit your article to this journal](#)



Article views: 512








[View related articles](#)



[View Crossmark data](#)

# The role of mental health on the relationship between food insecurity and immunologic outcome among people living with HIV in Guangxi, China

Cheng Chen <sup>a</sup>, Xueying Yang <sup>b,c</sup>, Chengbo Zeng <sup>b,c</sup>, Xiaoming Li <sup>b,c</sup>, Shan Qiao <sup>b,c</sup> and Yuejiao Zhou<sup>d</sup>

<sup>a</sup>School of Public Health, School of Medicine, Shanghai Jiao Tong University, Shanghai, People's Republic of China; <sup>b</sup>Department of Health Promotion, Education, and Behavior, University of South Carolina Arnold School of Public Health, Columbia, SC, USA; <sup>c</sup>South Carolina SmartState Center for Healthcare Quality (CHQ), University of South Carolina Arnold School of Public Health, Columbia, SC, USA; <sup>d</sup>Guangxi Center for Disease Control and Prevention, Nanning, People's Republic of China

## ABSTRACT

**Background:** Previous studies showed that food insecurity could adversely affect clinical outcomes of people living with HIV (PLWH). The mental health pathways of such effects are suggested in existing literature, but empirical data are limited in resource-limited settings.

**Methods:** This cross-sectional study aims to explore the role of depressive symptoms and anxiety on the association between food insecurity and CD4 counts among a sample of 2,987 PLWH in Guangxi, China. Path analysis was used to examine a hypothetical model and delta z test was used to assess the indirect effects of food insecurity on CD4 counts through depressive symptoms and anxiety.

**Results:** The prevalence of food insecurity in this sample was 25.3%, and the median CD4 counts were 318 cells/mm<sup>3</sup>. In correlation analyses, food insecurity was not directly associated with LogCD4 but was associated with depressive symptoms and anxiety. Path analysis indicated a significant indirect effect from food insecurity to LogCD4 through depressive symptoms, but not anxiety.

**Conclusion:** Improving mental health among PLWH with food insecurity may be beneficial for treatment outcomes. Besides intervening food insecurity, an intervention targeting depressive symptoms could improve the immunologic outcomes of PLWH.

## ARTICLE HISTORY

Received 31 August 2020  
Accepted 19 November 2020


## KEYWORDS

Mental health; Food insecurity; Immunologic outcome; CD4; HIV/AIDS

## Introduction

According to Chinese Centers for Disease Control and Prevention (CDC), the number of people living with HIV (PLWH) has reached 1.25 million in 2018 (Chinese Centers for

**CONTACT** Xueying Yang  [xueyyang@mailbox.sc.edu](mailto:xueyyang@mailbox.sc.edu)  Department of Health Promotion, Education, and Behavior, Arnold School of Public Health, University of South Carolina, 915 Greene Street Suite 408, Columbia, SC, USA

 Supplemental data for this article can be accessed at <https://doi.org/10.1080/21642850.2020.1854762>

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Disease Prevention and Control, 2018). With the worldwide availability of antiretroviral therapy (ART), the mortality of PLWH has been decreased in recent years (Herbst et al., 2009; Hogg et al., 1997; Palella et al., 1998), resulting in the longer life expectancy (Antiretroviral Therapy Cohort Collaboration, 2017). However, according to a 2016 meta-analysis, only around 40% of PLWH were on ART and 36% of those on ART achieved viral suppression in China (Levi et al., 2016). While effective ART can significantly improve the immune recovery of the majority of PLWH, there is still a certain proportion of PLWH who remain at relatively low CD4 levels (Kaufmann et al., 2003).

CD4 cells (also known as CD4+ T cells) are white blood cells that fight infection. CD4 count is an indicator of immune function in PLWH and one of the key determinants for the need of opportunistic infection prophylaxis. CD4 counts are obtained from bloodwork as part of laboratory monitoring for HIV infection. As an immunological outcome, CD4 count is a critical indicator for classifying clinical stages of HIV infection, evaluating treatment efficacy, and changing the medication when necessary (Bentwich, 2005). Lower CD4 counts are reported to be associated with an increased risk of opportunistic infection (Lewden et al., 2007). Yet the immunological outcome was not satisfactory in China, with the median CD4 count ranging from only 118 to 182 cells/mm<sup>3</sup> (Huang et al., 2015; Ruan et al., 2010; Zhang et al., 2009). Therefore, it is important to explore potential factors that impede immunological recovery among PLWH. Factors associated with lower CD4 count in previous studies included older age (Kaufmann et al., 2003), treatment interruption, non-adherence to ART (Gross, Bilker, Friedman, & Strom, 2001; Kaufmann et al., 2011), and food insecurity (Aibibula et al., 2016).

Food insecurity, defined as a limited or uncertain ability to acquire acceptable foods in socially acceptable ways, or limited or uncertain availability of nutritionally adequate and safe foods (Anderson, 1990), has been found to be associated with multiple health problems of PLWH, such as unprotected sex, suboptimal treatment adherence (Weiser et al., 2009b), incomplete viral suppression (McMahon, Wanke, Elliott, Skinner, & Tang, 1999), and greater mortality (Weiser et al., 2009a). Food insecurity is common among PLWH in resource-poor settings, ranging from 24% to 81% (Mamlin et al., 2009; Palar et al., 2015; Wang et al., 2011).

The underlying mechanisms about the negative effect of food insecurity on HIV outcomes can be explained in several perspectives. In pharmacokinetic perspective, food intake is recommended for processing, absorption, and optimal clinical benefits for certain ART regimens. For example, pharmacokinetic studies showed as much as a 30% increase in the plasma concentration of darunavir (Prezista) when taken with food (Gribble et al., 2000) and administration of atazanavir (Reyataz) with a light meal increases drug plasma concentration by 70% (Gribble et al., 2000). Prescribing ART that requires food to those who are food insecure could therefore diminish the efficacy of ART even when patients are adherent (Kalichman et al., 2015). In behavioral perspective, food insecurity can interfere with medication adherence by disrupting daily routines, impairing memory and attention, impeding adherence strategies, and reducing motivation (Frega, Duffy, Rawat, & Grede, 2010; Singer, Weiser, & McCoy, 2015;). In addition, in the face of limited resources, demands for food may compete with resources needed to procure medicines. The need to make tradeoffs between paying for food and other medical care (including transport costs to clinic) leading to poor retention in

medical care (Young, Wheeler, McCoy, & Weiser, 2014), which would influence the treatment outcomes (e.g. CD4 counts).

A 2016 literature review (Aibibula et al., 2016) revealed that only 8 studies investigated the association between food insecurity and immunological outcomes (e.g. CD4 counts). According to this systematic review, PLWH with food insecurity had 1.32 times higher odds of having lower CD4 counts compared to PLWH without food insecurity. All 8 studies except one were conducted in the U.S. Thus, there is a dearth of such data in China, such as Guangxi. According to China National Health Accounts Report, Guangxi has a heavy burden on medical expense in 2008 (China National Health Economics, 2005). Guangxi's poverty rate is 17.1%, compared to the whole China's 8.3% (Deng, 2012), and the average monthly salary of Guangxi is 1087 yuan, only half of the national average (Gong, 2011). However, there is a paucity of data investigating food insecurity among PLWH in Guangxi.

Given the negative impact of food insecurity on HIV treatment outcomes (e.g. CD4 counts), understanding the underlying mechanisms could potentially inform the future interventions to improve HIV treatment outcomes. Previous literature suggested that inadequate dietary intake can lead to mental health problems (McIntyre, Williams, Lavorato, & Patten, 2013), which in turn leads to compromised immunity among PLWH, especially decrease in CD4 counts (Ickovics et al., 2001; Kemeny et al., 1994; Leserman, 2008; Patterson et al., 1996). Among all the mental health problems, psychiatric problems such as depressive and anxiety symptoms have been reported to be highly prevalent among PLWH (with the prevalence of 36% and 16%, respectively) (Atkinson et al., 2009; Bing et al., 2001; Kemeny et al., 1994;). Moreover, some studies (Burack et al., 1993; Leserman et al., 1997) demonstrated a faster decline in CD4 counts in PLWH with depressive symptom than those PLWH without depressive symptoms. Since food insecurity can interplay with depressive and anxiety symptoms as suggested in the existing literature (Laraia, Siega-Riz, Gundersen, & Dole, 2006; Vozoris & Tarasuk, 2003; Whitaker, Phillips, & Orzol, 2006), it is plausible that these two mental health problems may play a role in the relationship between food insecurity and CD4 counts. Hence, the current study aims to examine the relationship between food insecurity and CD4 counts and to explore the effects of depressive and anxiety symptoms on such relationship in Guangxi, China. We hypothesized that (1) food insecurity is negatively associated with CD4 counts; and (2) food insecurity can affect CD4 counts through depressive symptoms and anxiety.

## Materials and methods

### Study site and participants

Data were derived from a cross-sectional survey conducted in Guangxi, China from October 2012 to August 2013. Details of the survey were described previously (Yang et al., 2019). Briefly, 2 cities and 10 rural counties that had the largest cumulative number of reported HIV/AIDS cases were selected as the survey sites. Staff from local CDC and healthcare workers from local health centers were selected and trained to facilitate the data collection. Individuals who were aged  $\geq 18$  years old and with a confirmed HIV diagnosis were eligible to participate. The aforementioned staff conducted face-to-

face interviews in private rooms after obtained informed consent from the participants. The latest CD4 counts were retrieved from the participants' medical records with appropriate informed consent. With an approximate 10% refusal rate, 3002 PLWH were recruited and 2987 completed questionnaires. The study protocol was approved by the Institutional Review Boards at the Wayne State University in the United States and Guangxi CDC in China.

## Measures

### Sociodemographic characteristics

The sociodemographic characteristics included gender, age, ethnicity (Han/others), marital status (single/separated, married/cohabitated, divorced/widowed), religion, education attainment (illiteracy/primary school, middle school, college), employment status (no job, part-time, full-time), place of household registration, and monthly household

**Table 1.** Socio-demographic characteristics.

Variables	N (%)	LogCD4	
		Mean $\pm$ SD	P
<b>Total</b>	2987 (100%)	-	-
<b>LogCD4</b>	2855 (95.6%)	5.57 $\pm$ 0.93	
<b>Age (years)</b>			<b>&lt;.001</b>
$\leq 30$	404 (13.5)	5.66 $\pm$ 1.04	
30-50	1864 (62.4)	5.58 $\pm$ 0.94	
$> 50$	719 (24.1)	5.47 $\pm$ 0.81	
<b>Duration of Diagnosis (Mean <math>\pm</math> SD, years)</b>		3.63 $\pm$ 2.44	<b>&lt;.001</b>
$\leq 5$	2230 (74.7)	5.49 $\pm$ 0.97	
6~10	710 (23.8)	5.78 $\pm$ 0.75	
$> 11$	47 (1.5)	5.80 $\pm$ 0.71	
<b>Gender</b>			<b>&lt;.001</b>
Male	1876 (62.8)	5.48 $\pm$ 0.96	
Female	1111 (37.2)	5.71 $\pm$ 0.84	
<b>Ethnicity</b>			0.817
Han	2109 (70.7)	5.56 $\pm$ 0.94	
Others	873 (29.3)	5.58 $\pm$ 0.90	
<b>Levels of Education</b>			<b>0.018</b>
Less than primary school	901 (30.3)	5.52 $\pm$ 0.91	
Middle school or above	2075 (69.7)	5.58 $\pm$ 0.93	
<b>Employment status</b>			<b>0.001</b>
No job	800 (26.9)	5.45 $\pm$ 1.09	
Part-time	992 (33.4)	5.57 $\pm$ 0.87	
Full-time	1182 (39.7)	5.64 $\pm$ 0.83	
<b>Household's monthly income (RMB)</b>			0.595
$< 2000$	2442 (82.6)	5.57 $\pm$ 0.92	
$\geq 2000$	516 (17.4)	5.56 $\pm$ 0.94	
<b>Marital status</b>			<b>0.017</b>
Single/Separated	386 (13.2)	5.43 $\pm$ 1.10	
Married/cohabitated	2011 (68.9)	5.59 $\pm$ 0.92	
Divorced/Widowed	520 (17.8)	5.55 $\pm$ 0.81	
<b>Household registration</b>			0.167
The local city	2783(93.4)	5.57 $\pm$ 0.91	
Other cities in Guangxi	168 (5.6)	5.39 $\pm$ 1.17	
Other provinces	29 (1.0)	5.79 $\pm$ 0.69	
<b>Whether on ART</b>			<b>&lt;.001</b>
Yes	2146 (72.1)	5.53 $\pm$ 0.93	
No	829 (27.9)	5.63 $\pm$ 1.05	

Note: \* The number of the participants is not equal to the total sample size of some variables due to missing data.

Bold values indicate the significant variables with p-value  $< 0.05$ .

income (in Chinese currency, RMB). Participants were also asked to provide HIV-related information, including duration since HIV diagnosis and their HIV treatment status. The log transformed CD4 count (LogCD4) was used as the dependent outcome since its distribution was highly skewed (Table 1).

### **Food insecurity**

Food insecurity was defined by one question: 'In the past year, does your household have enough food to meet the needs of the whole family? (Yes/no)'. Participants who responded 'No' were categorized as experienced food insecurity, while participants who answered 'Yes' were categorized as not experienced food insecurity.

### **Mental health**

**Depression:** The 10-item Center for Epidemiologic Studies Depression Scale (CESD-10) (Andresen, Malmgren, Carter, & Patrick, 1994), a shortened version from the standard 20-item scale, was used to measure depression (Radloff, 1977). The CESD-10 consists of questions about how many depressive symptoms a person has experienced in the past week (item samples: 'I can't concentrate on what I'm doing', 'I'm full of hope for the future', 'I don't sleep well'). Each item has four choices from 'never or seldom' (Chinese Centers for Disease Prevention and Control, 2018) to 'often' (Herbst et al., 2009), resulting in a total score ranging from 10 to 40, with a higher total score representing a higher level of depressive symptoms. The Cronbach alpha for the CESD-10 was 0.79 in this study after reversing certain items.

**Anxiety:** The Brief Anxiety Scale (Zung, 1971) was used to measure anxiety. The scale consists of 20 questions about the feelings a person has experienced in the past week (item samples: 'I feel easily upset or panic', 'I have a headache and a sore neck', 'My face was hot and red'). (Response categories: 1 = never or seldom, 4 = often). A summary score was created by adding up the score of each item, with a higher total score representing a higher level of anxiety symptoms. After reversing certain items, this scale exhibited good internal reliability (Cronbach alpha = 0.90).

### **Data analysis**

First, descriptive statistics were reported on socio-demographic characteristics (e.g. age, gender, marital status). Mean and standard deviation (SD) were used to describe continuous variables, and frequencies were used for categorical variables. Bivariate analyses were carried out using t tests or ANOVA to examine the relationships between socio-demographic characteristics and logCD4. Second, bivariate correlation analyses were performed to examine the associations among food insecurity, depressive symptom, anxiety, and logCD4. Third, path model was conducted to examine the associations among depressive symptoms, anxiety, logCD4, and food insecurity while adjusting for covariates that were significantly associated with logCD4 in bivariate analyses. Depressive symptoms, anxiety, and logCD4 were standardized before conducting path analysis. Delta z tests were used to examine the roles of depressive symptoms and anxiety between food insecurity and logCD4. Both unstandardized (see Appendix) and standardized path



coefficients were reported in this study. Descriptive statistics and correlation analyses were performed using SPSS software version 22 (SPSS Inc., Chicago, IL). Path analysis was performed using Mplus version 7.0 (Muthen & Muthen, Los Angeles, CA).

## Results

### Sociodemographic characteristics

As shown in Table 1, among a total of 2987 participants, the average age was 42.46 years old (SD = 12.84); more than half of them (62.8%, 1876/2987) were male. The majority of the participants were of Han ethnicity (70.7%, 2109/2987). Around 70% of participants were married or in a cohabiting relationship (68.9%, 2011/2987), and had received middle school or above education (69.7%, 2075/2987). Most participants (82.6%, 2442/2987) had a household's monthly income less than 2000 RMB. In terms of the treatment, the majority of them (72.1%) were on ART. The mean scores of CESD-10 scale and anxiety scale were 19.5 (SD = 4.8) and 31.1 (SD = 8.9), respectively.

### Association between background characteristics and outcome

The prevalence of food insecurity among PLWH in this study was 25.3%. The median CD4 count was 318 cells/mm<sup>3</sup> (interquartile range: 195–459 cells/mm<sup>3</sup>), with the mean and SD of logCD4 of 5.57 (SD = 0.93). Table 1 shows the associations between socio-demographic characteristics and logCD4. PLWH who were female ( $p < .001$ ), younger ( $p < .001$ ), having a full time job ( $p = 0.001$ ), being married/cohabited with boyfriend or girlfriend ( $p = 0.017$ ), and having a household registration of other provinces ( $p = 0.167$ ) were more likely to have a higher logCD4 than their counterparts.

### Correlations among depressive symptom, anxiety, logCD4 and food insecurity

Table 2 shows correlations among depressive symptom, anxiety, logCD4, and food insecurity. Results of correlation analyses indicated that both depressive symptom and anxiety were positively associated with food insecurity, and their Spearman correlation coefficients were 0.242 ( $p < .001$ ) and 0.257 ( $p < .001$ ), respectively. Both depressive symptom and anxiety were negatively associated with logCD4, and their Spearman correlation coefficients were  $-0.092$  ( $p < .001$ ) and  $-0.080$  ( $p < .001$ ), respectively. Food insecurity was not significantly associated with logCD4 (correlation coefficient = 0.003,  $p = 0.89$ ).

**Table 2.** Correlation matrix.

	1	2	3	4
1. Depression	1.000			
2. Anxiety	0.735***	1.000		
3. LogCD4	-0.092***	-0.080***	1.000	
4. Food insecurity	0.242***	0.257***	0.003	1.000

Notes: \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ .



### Path analysis

While adjusting for covariates that were significantly associated with CD4 counts in bivariate analyses, path model revealed that food insecurity was not significantly associated with logCD4 but was significantly related to both depressive symptom (std.β = 0.229,  $p < .001$ ) and anxiety (std.β = 0.247,  $p < .001$ ). Depression was negatively associated with logCD4 (std.β = -0.062,  $p = 0.041$ ), while the relationship between logCD4 and anxiety was not statistically significant (std.β = -0.004,  $p = 0.897$ ) (Table 3).

### Direct and indirect effects of path model

Direct pathway from food insecurity to logCD4 was not significant (std.β = 0.004,  $p = 0.837$ ), but indirect pathway from food insecurity to logCD4 through depressive symptom was significant (std.β = -0.014, delta  $z = -2.01$ ,  $p = 0.045$ ) (Table 4). Of note, such indirect pathway through anxiety was not statistically significant (std.β = -0.001, delta  $z = -0.13$ ,  $p = 0.897$ ). This result indicated that food insecurity may affect CD4 count indirectly through depressive symptom but not anxiety (Figure 1). Unstandardized path coefficients were shown in the Appendix.

### Discussion

To the best of our knowledge, this is one of the first studies to investigate the relationship between food insecurity and CD4 counts among PLWH in China. The finding underscored food insecurity issue among PLWH in China. Although the prevalence of food insecurity among PLWH in the current study is lower than the rate in other cities of China (e.g. 37.2% in Chengdu) or other countries (e.g. 63% in the US) (Hu et al., 2011; McMahon et al., 1999), the results addressed its role in compromising the immunological recovery of HIV patients in China. Addressing food insecurity should become an integral part of HIV care.

Although no direct effect between food insecurity and CD4 counts was found, results from the current study revealed the indirect association between the two variables through depressive symptom after controlling for potential confounders. Beside intervening food insecurity among PLWH, the government needs to pay more attention to this impoverished population with mental health problems in order to improve their treatment outcomes. These findings highlighted the importance of addressing food insecurity as part of comprehensive HIV care in order to improve HIV-related treatment outcomes.

**Table 3.** Path coefficients.

Paths*	Std.β	95% C.I.	S.E.	p-value
Food insecurity → Depression	0.229	0.191~0.268	0.020	<b>&lt;.001</b>
Food insecurity → Anxiety	0.247	0.207~0.283	0.019	<b>&lt;.001</b>
Food insecurity → LogCD4	0.004	-0.036~0.042	0.020	0.837
Depression → LogCD4	-0.062	-0.123~-0.007	0.041	<b>0.041</b>
Anxiety → LogCD4	-0.004	-0.063~0.054	0.030	0.897

\*Adjusted for significant socio-demographic characteristics in bivariate analyses: age, gender, levels of education, employment status, marital status, whether on ART, and duration of diagnosis.

Bold value indicate the significant associations with p-value <0.05.

**Table 4.** Mediation analysis.


Effect*	Std.β	95% C.I.	S.E.	p-value
Total effect	-0.011	-0.049~0.024	0.019	0.550
Indirect effect				<b>0.002</b>
Depression	-0.014	-0.028~-0.001	0.007	<b>0.045</b>
Anxiety	-0.001	-0.016~0.014	0.008	0.897
Direct effect	0.004	-0.036~0.042	0.020	0.837

\*Adjusted for significant socio-demographic characteristics in bivariate analyses: age, gender, levels of education, employment status, marital status, whether on ART, and duration of diagnosis.

Bold value indicate the significant associations with p-value <0.05.

It's worth noting that the insignificant association between food insecurity and CD4 counts was not congruent with previous results from existing studies which suggested a negative effect of food insecurity on CD4 counts (Kalichman et al., 2010 Palar et al., 2015; Weiser et al., 2014;). The study settings may partially account for such inconsistency. Compared with China, some previous studies were conducted in countries where the starvation problem is more acute, such as Africa (Benzekri et al., 2017 Weiser et al., 2014;). HIV patients there can be greatly affected by their nutritious status. In addition, different measures of food insecurity could also account for such inconsistency. For example, previous studies measured food insecurity using the Household Food Insecurity Access Scale (HFIAS) (Coates, Swindale, & Bilinsky, 2007) or the US Food Security Scale (Cook & Frank, 2008), but we only used one item and asked participants whether their food is enough for the whole family. Further studies are warranted to use more comprehensive measure to capture food insecurity and confirm the findings of this study. However, the results in this study could still point pinning the food insecurity issues and provide some insights for the future intervention programs.

In this study, food insecurity affected CD4 counts indirectly through depressive symptoms. Based on an established conceptual framework (Weiser et al., 2011), the impacts of food insecurity on clinical outcomes may be explained by behavioral pathways (delayed



**Figure 1.** The mediation effects of mental health problems on the relationship between food insecurity and immunologic outcome (i.e. CD4 count). Notes: Covariates (age, gender, employment status, marital status, household registration and whether on ART) with *p*-values less than 0.05 in bivariate analysis were adjusted in the final model. \**p* < 0.05, \*\**p* < 0.01, \*\*\**p* < 0.001. All path coefficients were standardized coefficients.

entry into care, poor clinic attendance, interruptions in care), nutritional pathways (macronutrient and micronutrient deficiencies, worse absorption of drugs in the absence of food), and mental health pathways. The results of this study contributed more evidence pertinent to the mental health pathways in that conceptual framework. Food insecurity is a source of significant distress and shame (Mickelson & Williams, 2008). Studies have documented associations between food insecurity and depressive symptoms in HIV population (Palar et al., 2015; Vogenthaler et al., 2011). Previous studies even have found the dose–response relationship between food insecurity and depressive symptoms among women living with HIV (Palar et al., 2018). Depression is common among PLWH and is associated with declining CD4 counts in previous studies (Olisah, Adekeye, & Sheikh, 2014). The negative effects of depression on immune status may consequently result in HIV disease progression. The WHO recommends that attention to psychosocial needs of PLWH should be an integral part of HIV care (World Health Organization, 2005). PLWH need extra emotional care and psychological support to prevent and cope with mental health problems in order to promote immunological recovery. Further investigation of this pathway is needed in larger studies including detailed measures of macronutrient and micronutrient deficiencies. Future interventions should pay more attention on PLWH at a risk of food insecurity with depression or anxiety problems to improve their immune system.

There are some limitations to be aware of in the current study. First, the one-item measure of food insecurity may not accurately capture the practical situation of food insufficiency among PLWH. Second, this is a cross-sectional study, the casual relationships cannot be drawn from this study. Further studies should estimate causality between food insecurity and CD4 counts employing longitudinal study designs. Third, self-report bias, including recall bias and social desirability bias may exist during the data collection. Thus, the rate of food insecurity in this study might be underestimated. Fourth, the effect size of the associations in this study was small. There might be some other covariates or confounders that mediate or moderate the relationships we did not consider. We called for future research to investigate the mediators or moderators in the relationships between psychosocial variables and immunologic outcome. Finally, our results are based on a sample of HIV patients in Guangxi, China. It is necessary to be cautious when extending these findings to other settings.

Despite these limitations, the findings in this study revealed the food insecurity issues among PLWH in China and suggested that food insecurity might compromise the immunologic outcome of HIV patients through depressive symptom. This link sheds light on the importance of addressing food insecurity as part of comprehensive care to improve health outcomes among PLWH. Improving mental health among PLWH with food insecurity may be beneficial on treatment outcome. Future research is needed to better understand the underlying mechanisms and explore other potential pathways.

### Author contributions

Conceptualization, X.Y.Y. and X.M.L.; methodology, X.Y.Y. and C.B.Z.; software, C.B.Z.; validation, X.M.L.; formal analysis, C.C. and C.B.Z.; writing—original draft preparation,

C.C.; writing-review & editing, X.Y.Y.; project administration, S.Q. All authors have read and agreed to the published version of the manuscript.

## Acknowledgements

The study was supported by the National Institute of Child Health and Human Development (grant number#R01HD074221).

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

This work was supported by National Institute of Child Health and Human Development [grant number R01HD074221].

## ORCID

Cheng Chen  <http://orcid.org/0000-0003-1053-3217>  
Xueying Yang  <http://orcid.org/0000-0001-6788-0688>  
Chengbo Zeng  <http://orcid.org/0000-0002-3512-1115>  
Xiaoming Li  <http://orcid.org/0000-0002-5555-9034>  
Shan Qiao  <http://orcid.org/0000-0003-1834-1834>

## References

- Aibibula, W., Cox, J., Hamelin, A. M., Mamiya, H., Klein, M. B., & Brassard, P. (2016). Food insecurity and low CD4 count among HIV-infected people: A systematic review and meta-analysis. *AIDS Care, 28*(12), 1577–1585.
- Anderson, S. (1990). Core indicators of nutritional state for difficult-to-sample populations. *The Journal of Nutrition, 120*(Suppl 11), 1559–1600.
- Andresen, E. M., Malmgren, J. A., Carter, W. B., & Patrick, D. L. (1994). Screening for depression in well older adults: Evaluation of a short form of the CES-D (Center for Epidemiologic Studies Depression Scale). *American Journal of Preventive Medicine, 10*(2), 77–84.
- Antiretroviral Therapy Cohort Collaboration. (2017). Survival of HIV-positive patients starting antiretroviral therapy between 1996 and 2013: A collaborative analysis of cohort studies. *The Lancet HIV, 4*(8), e349–ee56.
- Atkinson, J. H., Higgins, J. A., Vigil, O., Dubrow, R., Remien, R. H., Steward, W. T., et al. (2009). Psychiatric context of acute/early HIV infection. The NIMH Multisite Acute HIV Infection Study: IV. *AIDS and Behavior, 13*(6), 1061–1067.
- Bentwich, Z. (2005). CD4 measurements in patients with HIV: Are they feasible for poor settings? *PLoS Medicine, 2*(7), e214.; quiz e54.
- Benzekri, N. A., Sambou, J. F., Diaw, B., Sall, E. H. I., Sall, F., Niang, A., et al. (2017). The dimensions of food insecurity and malnutrition among people living with HIV in Senegal, West Africa. *AIDS Care, 29*(12), 1510–1516.
- Bing, E. G., Burnam, M. A., Longshore, D., Fleishman, J. A., Sherbourne, C. D., London, A. S., et al. (2001). Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Archives of General Psychiatry, 58*(8), 721–728.

- Burack, J. H., Barrett, D. C., Stall, R. D., Chesney, M. A., Ekstrand, M. L., & Coates, T. J. (1993). Depressive symptoms and CD4 lymphocyte decline among HIV-infected men. *Jama*, *270*(21), 2568–2573.
- China National Health Economics. (2005). *China National Health Accounts Report*. China National Health Economics Institute Beijing.
- Chinese Centers for Disease Prevention and Control. (2018). <http://www.chinacdc.cn/en/>
- Coates, J., Swindale, A., & Bilinsky, P. (2007). *Household food insecurity access scale (HFIAS) for measurement of food access: Indicator guide*. Washington, DC: Food and nutrition technical assistance project, academy for educational Development; 34.
- Cook, J. T., & Frank, D. A. (2008). Food security, poverty, and human development in the United States. *Annals of the New York Academy of Sciences*, *1136*, 193–209.
- Deng, W. (2012). *College of Agriculture*. Nanjing: Guangxi University.
- Frega, R., Duffy, F., Rawat, R., & Grede, N. (2010). Food insecurity in the context of HIV/AIDS: A framework for a new era of programming. *Food and Nutrition Bulletin*, *31*(4), S292–S312.
- Gong, S. (2011). Evaluation of level and quality about regional financial income as well as analysis towards its influencing factors. *Journal of Regional Financial Research*, *461*(4), 78–83.
- Gribble, J. N., Miller, H. G., Cooley, P. C., Catania, J. A., Pollack, L., & Turner, C. F. (2000). The impact of T-ACASI interviewing on reported drug use among men who have sex with men. *Substance Use & Misuse*, *35*(6-8), 869–890.
- Gross, R., Bilker, W. B., Friedman, H. M., & Strom, B. L. (2001). Effect of adherence to newly initiated antiretroviral therapy on plasma viral load. *AIDS (London, England)*, *15*(16), 2109–2117.
- Herbst, A. J., Cooke, G. S., Bärnighausen, T., KanyKany, A., Tanser, F., & Newell, M. (2009). Adult mortality and antiretroviral treatment roll-out in rural KwaZulu-Natal, South Africa. *Bulletin of the World Health Organization*, *87*(10), 754–762.
- Hogg, R. S., O'Shaughnessy, M. V., Gataric, N., Yip, B., Craib, K., Schechter, M. T., et al. (1997). Decline in deaths from AIDS due to new antiretrovirals. *Lancet (London, England)*, *349*(9061), 1294.
- Hu, W., Jiang, H., Chen, W., He, S. H., Deng, B., Wang, W. Y., et al. (2011). Malnutrition in hospitalized people living with HIV/AIDS: Evidence from a cross-sectional study from Chengdu, China. *Asia Pacific Journal of Clinical Nutrition*, *20*(4), 544–550.
- Huang, P., Tan, J., Ma, W., Zheng, H., Lu, Y., Wang, N., et al. (2015). Outcomes of antiretroviral treatment in HIV-infected adults: A dynamic and observational cohort study in Shenzhen, China, 2003-2014. *BMJ Open*, *5*(5), e007508.
- Ickovics, J. R., Hamburger, M. E., Vlahov, D., Schoenbaum, E. E., Schuman, P., Boland, R. J., et al. (2001). Mortality, CD4 cell count decline, and depressive symptoms among HIV-seropositive women: Longitudinal analysis from the HIV epidemiology research study. *Jama*, *285*(11), 1466–1474.
- Kalichman, S. C., Cherry, C., Amaral, C., White, D., Kalichman, M. O., Pope, H., et al. (2010). Health and treatment implications of food insufficiency among people living with HIV/AIDS, Atlanta, Georgia. *Journal of Urban Health: Bulletin of the New York Academy of Medicine*, *87*(4), 631–641.
- Kalichman, S. C., Washington, C., Grebler, T., Hoyt, G., Welles, B., Merely, C., et al. (2015). Treatment outcomes among people living with HIV who are food insecure and prescribed antiretrovirals taken with food. *Journal of Primary Care & Community Health*, *6*(1), 35–40.
- Kaufmann, G. R., Elzi, L., Weber, R., Furrer, H., Giulieri, S., Vernazza, P., et al. (2011). Interruptions of cART limits CD4 T-cell recovery and increases the risk for opportunistic complications and death. *AIDS (London, England)*, *25*(4), 441–451.
- Kaufmann, G. R., Perrin, L., Pantaleo, G., Opravil, M., Furrer, H., Telenti, A., et al. (2003). CD4 T-lymphocyte recovery in individuals with advanced HIV-1 infection receiving potent antiretroviral therapy for 4 years: The Swiss HIV Cohort Study. *Archives of Internal Medicine*, *163*(18), 2187–2195.
- Kemeny, M. E., Weiner, H., Taylor, S. E., Schneider, S., Visscher, B., & Fahey, J. L. (1994). Repeated bereavement, depressed mood, and immune parameters in HIV seropositive and seronegative

- gay men. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association*, 13(1), 14–24.
- Laraia, B. A., Siega-Riz, A. M., Gundersen, C., & Dole, N. (2006). Psychosocial factors and socioeconomic indicators are associated with household food insecurity among pregnant women. *The Journal of Nutrition*, 136(1), 177–182.
- Leserman, J. (2008). Role of depression, stress, and trauma in HIV disease progression. *Psychosomatic Medicine*, 70(5), 539–545.
- Leserman, J., Petitto, J. M., Perkins, D. O., Folds, J. D., Golden, R. N., & Evans, D. L. (1997). Severe stress, depressive symptoms, and changes in lymphocyte subsets in human immunodeficiency virus-infected men. A 2-year follow-up study. *Archives of General Psychiatry*, 54(3), 279–285.
- Levi, J., Raymond, A., Pozniak, A., Vernazza, P., Kohler, P., & Hill, A. (2016). Can the UNAIDS 90-90-90 target be achieved? A systematic analysis of national HIV treatment cascades. *BMJ Global Health*, 1(2), e000010.
- Lewden, C., Chene, G., Morlat, P., Raffi, F., Dupon, M., Dellamonica, P., et al. (2007). HIV-infected adults with a CD4 cell count greater than 500 cells/mm<sup>3</sup> on long-term combination antiretroviral therapy reach same mortality rates as the general population. *Journal of Acquired Immune Deficiency Syndromes* (1999), 46(1), 72–77.
- Mamlin, J., Kimaiyo, S., Lewis, S., Tadayo, H., Jerop, F. K., Gichunge, C., et al. (2009). Integrating nutrition support for food-insecure patients and their dependents into an HIV care and treatment program in Western Kenya. *American Journal of Public Health*, 99(2), 215–221.
- McIntyre, L., Williams, J. V., Lavorato, D. H., & Patten, S. (2013). Depression and suicide ideation in late adolescence and early adulthood are an outcome of child hunger. *Journal of Affective Disorders*, 150(1), 123–129.
- McMahon, J. H., Wanke, C. A., Elliott, J. H., Skinner, S., & Tang, A. M. (2011). Repeated assessments of food security predict CD4 change in the setting of antiretroviral therapy. *Journal of Acquired Immune Deficiency Syndromes* (1999), 58(1), 60–63.
- Mickelson, K. D., & Williams, S. L. (2008). Perceived stigma of poverty and depression: Examination of interpersonal and intrapersonal mediators. *Journal of Social and Clinical Psychology*, 27(9), 903–930.
- Olisah, V. O., Adekeye, O., & Sheikh, T. L. (2014). Depression and CD4 cell count among patients with HIV in a Nigerian University Teaching Hospital. *The International Journal of Psychiatry in Medicine*, 48(4), 253–261.
- Palar, K., Frongillo, E. A., Escobar, J., Sheira, L. A., Wilson, T. E., Adedimeji, A., et al. (2018). Food insecurity, internalized stigma, and depressive symptoms among women living with HIV in the United States. *AIDS and Behavior*, 22(12), 3869–3878.
- Palar, K., Kushel, M., Frongillo, E. A., Riley, E. D., Grede, N., Bangsberg, D., et al. (2015). Food insecurity is longitudinally associated with depressive symptoms among homeless and marginally-housed individuals living with HIV. *AIDS and Behavior*, 19(8), 1527–1534.
- Palella, F. J., Delaney, K. M., Moorman, A. C., Loveless, M. O., Fuhrer, J., Satten, G. A., et al. (1998). Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV outpatient study investigators. *The New England Journal of Medicine*, 338(13), 853–860.
- Patterson, T. L., Shaw, W. S., Semple, S. J., Cherner, M., McCutchan, J. A., Atkinson, J. H., et al. (1996). Relationship of psychosocial factors to HIV disease progression. *Annals of Behavioral Medicine*, 18(1), 30–39.
- Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1(3), 385–401.
- Ruan, Y., Xing, H., Wang, X., Tang, H., Wang, Z., Liu, H., et al. (2010). Virologic outcomes of first-line HAART and associated factors among Chinese patients with HIV in three sentinel antiretroviral treatment sites. *Tropical Medicine & International Health*, 15(11), 1357–1363.
- Singer, A. W., Weiser, S. D., & McCoy, S. I. (2015). Does food insecurity undermine adherence to antiretroviral therapy? A systematic review. *AIDS and Behavior*, 19(8), 1510–1526.

- Vogenthaler, N. S., Hadley, C., Rodriguez, A. E., Valverde, E. E., Del Rio, C., & Metsch, L. R. (2011). Depressive symptoms and food insufficiency among HIV-infected crack users in Atlanta and Miami. *AIDS and Behavior, 15*(7), 1520–1526.
- Vozoris, N. T., & Tarasuk, V. S. (2003). Household food insufficiency is associated with poorer health. *The Journal of Nutrition, 133*(1), 120–126.
- Wang, E. A., McGinnis, K. A., Fiellin, D. A., Goulet, J. L., Bryant, K., Gibert, C. L., et al. (2011). Food insecurity is associated with poor virologic response among HIV-infected patients receiving antiretroviral medications. *Journal of General Internal Medicine, 26*(9), 1012–1018.
- Weiser, S. D., Fernandes, K. A., Brandson, E. K., Lima, V. D., Anema, A., Bangsberg, D. R., et al. (2009a). The association between food insecurity and mortality among HIV-infected individuals on HAART. *Journal of Acquired Immune Deficiency Syndromes (1999), 52*(3), 342–349.
- Weiser, S. D., Frongillo, E. A., Ragland, K., Hogg, R. S., Riley, E. D., & Bangsberg, D. R. (2009b). Food insecurity is associated with incomplete HIV RNA suppression among homeless and marginally housed HIV-infected individuals in San Francisco. *Journal of General Internal Medicine, 24*(1), 14–20.
- Weiser, S. D., Palar, K., Frongillo, E. A., Tsai, A. C., Kumbakumba, E., Depee, S., et al. (2014). Longitudinal assessment of associations between food insecurity, antiretroviral adherence and HIV treatment outcomes in rural Uganda. *AIDS (London, England), 28*(1), 115–120.
- Weiser, S. D., Young, S. L., Cohen, C. R., Kushel, M. B., Tsai, A. C., Tien, P. C., et al. (2011). Conceptual framework for understanding the bidirectional links between food insecurity and HIV/AIDS. *The American Journal of Clinical Nutrition, 94*(6), 1729s–1739s.
- Whitaker, R. C., Phillips, S. M., & Orzol, S. M. (2006). Food insecurity and the risks of depression and anxiety in mothers and behavior problems in their preschool-aged children. *Pediatrics, 118*(3), e859–e868.
- World Health Organization. (2005). World Health Organization Psychosocial Support.
- Yang, X., Li, X., Qiao, S., Zhang, Q., Shen, Z., & Zhou, Y. (2019). Immunological and virologic outcomes of people living with HIV in Guangxi, China: 2012-2017. *PLoS One, 14*(3), e0213205.
- Young, S., Wheeler, A. C., McCoy, S. I., & Weiser, S. D. (2014). A review of the role of food insecurity in adherence to care and treatment among adult and pediatric populations living with HIV and AIDS. *AIDS and Behavior, 18*(Suppl 5), S505–S515.
- Zhang, F., Dou, Z., Ma, Y., Zhao, Y., Liu, Z., Bulterys, M., et al. (2009). Five-year outcomes of the China National Free Antiretroviral Treatment Program. *Annals of Internal Medicine, 151*(4), 241–251. W-52.
- Zung, W. W. (1971). A rating instrument for anxiety disorders. *Psychosomatics, 12*(6), 371–379.