

Oral presentation

Open Access

The Molecular and Cellular Basis of Tumor Rejection After Vaccination With Mammary Adenocarcinoma Cells Transduced With the MHC Class II Transactivator CIITA

Lorenzo Mortara¹, Paola Castellani², Raffaella Meazza³, Giovanna Tosi¹, Andrea De Lerma Barbaro¹, Francesco A Procopio¹, Luciano Zardi⁴, Silvano Ferrini³ and Roberto S Accolla*^{‡1}

Address: ¹Department of Clinical and Biological Sciences, School of Medicine, University of Insubria, Varese, ²Laboratory of Cellular Biology, Istituto Nazionale per la Ricerca sul Cancro, Genova, ³Laboratory of Immunopharmacology, Istituto Nazionale per la Ricerca sul Cancro, Genova and ⁴Philogen, Advanced Biotechnology Center, Genova

Email: Roberto S Accolla* - roberto.accolla@uninsubria.it

* Corresponding author ‡Presenting author

from 2005 International Meeting of The Institute of Human Virology
Baltimore, USA, 29 August – 2 September 2005

Published: 8 December 2005

Retrovirology 2005, **2**(Suppl 1):S55 doi:10.1186/1742-4690-2-S1-S55

CD8⁺ T cell responses are major players of tumor eradication in various vaccination protocols. However, an optimal stimulation of CD4⁺ T helper cells is required for both priming and maintenance of the effector CTL response against the tumor. In this study we show that the murine mammary adenocarcinoma cell line TS/A, a highly malignant MHC-II-negative tumor, is rejected *in vivo* if genetically engineered to express MHC-II molecules by transfer of the MHC-II transactivator CIITA. TS/A-CIITA cells are fully rejected by 93% of the syngeneic recipients and have a significantly lower growth rate in the remaining 7% of animals. Rejection requires CD4⁺ and CD8⁺ cells. CD4⁺ T cells are fundamental in the priming phase, whereas CTLs are the major anti-tumor effectors. All tumor rejecting animals are protected against rechallenge with the parental TS/A tumor. Immunohistochemical data at day 5 post-inoculation showed an higher infiltrate of CD4⁺ T cells in mice bearing TS/A-CIITA, than in mice bearing the TS/A tumor. Subsequently, from day 7 through day 10, TS/A-CIITA tumors showed higher number of both CD4⁺ and CD8⁺ cells, dendritic cells, together with massive necrosis. The frequency of IFN- α -secreting splenocytes early after inoculations was also assessed by an *ex vivo* ELISPOT assay. Only the rejecting TS/A-CIITA animals showed an high frequency of IFN- α -secreting cells (between 80 and 120/10⁶ splenocytes). Importantly, CD4 and CD8 depletion experiments revealed that at the time of tumor resolution the major

cell population recognizing the TS/A-CIITA cells was of CD4 origin. This is the first example of successful tumor vaccination by genetic transfer of CIITA. These results open the way to a possible use of CIITA for increasing both the inducing and the effector phase of the anti-tumor response.