

Post-intensive care syndrome: its pathophysiology, prevention, and future directions



| | |
|------------------------------|---|
| 著者 | INOUE Shigeaki, HATAKEYAMA Junji, KONDO Yutaka, HIFUMI Toru, SAKURAMOTO Hideaki, KAWASAKI Tatsuya, TAITO Shunsuke, NAKAMURA Kensuke, UNOKI Takeshi, KAWAI Yusuke, KENMOTSU Yuji, SAITO Masafumi, YAMAKAWA Kazuma, NISHIDA Osamu |
| 著者別名 | 井上 茂亮, 畠山 淳司, 近藤 豊, 一二三 亨, 櫻本 秀明, 川崎 達也, 對東 俊介, 中村 謙介, 卯野木 健, 河合 佑亮, 劔持 雄二, 齋藤 雅史, 山川 一馬, 西田 修 |
| journal or publication title | Acute Medicine & Surgery |
| volume | 6 |
| number | 3 |
| page range | 233-246 |
| year | 2019-07 |
| URL | http://id.nii.ac.jp/1127/00000830/ |

doi: <https://doi.org/10.1002/ams2.415>



Review Article

Post-intensive care syndrome: its pathophysiology, prevention, and future directions

Shigeaki Inoue,¹  Junji Hatakeyama,² Yutaka Kondo,³ Toru Hifumi,⁴ Hideaki Sakuramoto,⁵ Tatsuya Kawasaki,⁶ Shunsuke Taito,⁷ Kensuke Nakamura,⁸ Takeshi Unoki,⁹  Yusuke Kawai,¹⁰ Yuji Kenmotsu,¹¹ Masafumi Saito,¹ Kazuma Yamakawa,¹² and Osamu Nishida¹³

¹Department of Disaster and Emergency and Critical Care Medicine, Kobe University Graduate School of Medicine, Kobe, Hyogo; ²Department of Emergency and Critical Care Medicine, Yokohama City Minato Red Cross Hospital, Yokohama, Kanagawa; ³Department of Emergency and Critical Care Medicine, Juntendo University Urayasu Hospital, Urayasu, Chiba; ⁴Emergency and Critical Care Medicine, St. Luke's International Hospital, Tokyo; ⁵Department of Adult Health Nursing, College of Nursing, Ibaraki Christian University, Hitachi; ⁶Department of Pediatric Critical Care, Shizuoka Children's Hospital, Shizuoka, ⁷Department of Clinical Practice and Support, Division of Rehabilitation, Hiroshima University Hospital, Hiroshima; ⁸Department of Emergency and Critical Care Medicine, Hitachi General Hospital, Hitachi, Ibaraki; ⁹Department of Adult Health Nursing, School of Nursing, Sapporo City University, Sapporo; ¹⁰Department of Nursing, Fujita Health University Hospital, Toyoake, Aichi; ¹¹Department of Nursing, Tokai University Hachioji Hospital, Hachioji, Tokyo; ¹²Division of Trauma and Surgical Critical Care, Osaka General Medical Center, Osaka City, Osaka; and ¹³Department of Anesthesiology and Critical Care Medicine, Fujita Health University School of Medicine, Toyoake, Aichi, Japan

Expanding elderly populations are a major social challenge in advanced countries worldwide and have led to a rapid increase in the number of elderly patients in intensive care units (ICUs). Innovative advances in medical technology have enabled lifesaving of patients in ICUs, but there remain various problems to improve their long-term prognoses. Post-intensive care syndrome (PICS) refers to physical, cognition, and mental impairments that occur during ICU stay, after ICU discharge or hospital discharge, as well as the long-term prognosis of ICU patients. Its concept also applies to pediatric patients (PICS-p) and the mental status of their family (PICS-F). Intensive care unit-acquired weakness, a syndrome characterized by acute symmetrical limb muscle weakness after ICU admission, belongs to physical impairments in three domains of PICS. Prevention of PICS requires performance of the ABCDEFGH bundle, which incorporates the prevention of delirium, early rehabilitation, family intervention, and follow-up from the time of ICU admission to the time of discharge. Diary, nutrition, nursing care, and environmental management for healing are also important in the prevention of PICS. This review outlines the pathophysiology, prevention, and future directions of PICS.

Key words: Cognitive impairment, end-of-life, ICU-acquired weakness, intensive care unit, mental impairment, physical impairment, PICS-p, post-intensive care syndrome

BACKGROUND

EMERGENCY AND INTENSIVE care medicine have evolved dramatically in the past quarter-century due to technical innovation and guidelines for improving and standardizing auxiliary circulation and respiratory equipment in

the intensive care unit (ICU) and standardization and enhancement of educational programs. For these reasons, the short-term outcomes of ICU patients, including mortality and 28-day survival rates, have dramatically improved; however, the long-term prognosis and quality of life of sepsis patients have not yet improved.¹

Furthermore, the increase in the elderly population is a major social challenge for developed countries. In addition to Japan, the average age of citizens is rising even in advanced countries of Europe as well as the USA, China, Korea, and many other Asian countries. Furthermore, by 2050, the percentage of elderly people over the age of 65 years will be over 20% in most of the world except for countries in Africa and the Middle East, making them super-

Corresponding: Shigeaki Inoue, MD, PhD, MSCI, Department of Disaster and Emergency Medicine, Graduate School of Medicine, Kobe University, Kusunoki-cho 7-5-2, Chuo-ward, Kobe, Japan. E-mail: inoues@med.kobe-u.ac.jp.

Received 17 Feb, 2019; accepted 11 Mar, 2019; online publication 25 Apr, 2019

Funding information

No funding information provided.

aged societies.² The increase in the elderly population is a major social problem in Japan and worldwide and the number of elderly and elderly patients is increasing rapidly. Along with the aging of the population and progress in medical technology, the number of elderly people requiring intensive care increases annually and management of the elderly is essential in the emergency department and ICU.

Age is a poor prognostic factor for mortality in ICU patients, especially those with sepsis.³ Elderly people aged 65 years and older comprise approximately 60% of sepsis patients and account for approximately 80% of deaths.⁴ With aging worldwide, the number of patients with sepsis has increased, greatly influencing the long-term prognosis of ICU patients. Yende and colleagues analyzed two multinational randomized controlled trials (RCTs) including approximately 2,000 sepsis patients, reporting that one-third of patients who left the ICU died within 6 months and the 8 remaining one-third had 6 months. These findings indicate the presence of persisting functional impairment and obstructed activities of daily living.¹ Thus, not only short-term prognoses, such as 28-day and ICU survival rates, but also long-term outcomes in ICU patients should be assessed.

In view of the current and evolving ICU situation, the Society of Critical Care Medicine held a stakeholder conference to address subacute/chronic physical and psychological problems after ICU discharge, in which post-intensive care syndrome (PICS) was proposed.⁵ Post-intensive care syndrome is a physical, cognitive, and mental disorder that occurs during ICU stay or after ICU or hospital discharge

and includes the long-term prognosis of ICU patients and effects on the patient's family (Fig. 1). Furthermore, Nakamura *et al.*⁶ recently proposed the concept of post-acute care syndrome, based on evidence that acute care for elderly people involves difficulties in dysphagia and is related to prolonged ICU stay.

Pathophysiology of PICS

Physical impairments in PICS

With reduced mortality among critically ill patients due to advances in critical care medicine,^{7,8} long-term physical impairment in ICU survivors is a growing concern.^{1,9} Intensive care unit-acquired weakness (ICU-AW) is one factor related to muscle weakness.^{10,11} It is defined as the acute muscle weakness of the extremities in a symmetric pattern, which is caused by critical illness. Intensive care unit-acquired weakness is classified as critical illness polyneuropathy (CIP), critical illness myopathy (CIM), critical illness neuromyopathy (CINM), and muscle deconditioning (Fig. 2).¹²

The diagnosis of ICU-AW is made according to the Medical Research Council scale for grading the strength of various muscle groups in the upper and lower extremities, and a combined score of <48 in all testable muscle groups noted on more than two occasions separated by 24 h is diagnostic of ICU-AW.¹³ The incidence of ICU-AW is 40% in critically ill adult patients.¹⁴ Critical illness polyneuropathy is the most common category, followed by CINM, whereas




Fig. 1. Conceptual framework of post-intensive care syndrome (PICS). ICU, intensive care unit; PICS-F, PICS – family.




Fig. 2. Subclassification of the mechanisms of intensive care unit-acquired weakness (ICU-AW) into two main groups. The first group is ICU-AW with electrophysiologic and histopathologic findings (critical illness polyneuropathy [CIP] and critical illness myopathy [CIM]); the other is ICU-AW with normal diagnostic studies. CIM, abnormal reduction in the amplitude of compound muscle action potentials (CMAPs) and an increase in their duration, normal sensory nerve action potentials (SNAPs), reduced muscle excitability on direct stimulation, and myopathic motor unit potentials on needle electromyography; CINM, critical illness neuromyopathy, coexistence of CIP and CIM; CIP, reduction in the amplitude of CMAPs and SNAPs with normal or mildly reduced nerve conduction velocity; Muscle deconditioning, normal nerve conduction velocity and compound motor action potential, absence of spontaneous activity.

CIP is individually rare.^{15–17} The pathophysiological mechanisms of ICU-AW are considered multifactorial.¹⁰ Microvascular ischemia, catabolism, and immobility can lead to skeletal muscle wasting, while microvascular injury with resulting nerve ischemia, dysfunction of sodium channels, and injury to nerve mitochondria could contribute to critical illness-related neuropathy, myopathy, or both.¹⁰

Intensive care unit-acquired weakness contributes to prolonged mechanical ventilation, increased ICU and hospital lengths of stay, and mortality.^{18–20} The quadriplegia after-effect of disease usually resolves after several weeks to several months; rarely, impairment of motor function in survivors can persist from several months to several years.^{21,22} The risk of ICU-AW is associated with female sex, sepsis, catabolic state, multiorgan failure, systemic inflammatory response syndrome, long duration of mechanical ventilation, immobility, hyperglycemia, glucocorticoids, and neuromuscular blocking agents;¹⁰ however, a systematic review did not link glucocorticoids to this risk.¹⁴ No consensus has been established regarding effective interventions to improve outcomes of patients who develop ICU-AW;

however, preventive measures including early physical rehabilitation,²³ neuromuscular electrical stimulation,²⁴ and glucose control²⁵ have been applied.

Cognitive impairments in PICS

Critically ill patients experience high levels of physical and psychological stress in the ICU; these experiences result in cognitive impairments in patients with PICS. New or worsening impairments in cognitive function persist months to years after hospital discharge and are associated with poor daily functioning and reduced quality of life.^{26,27} Cognitive impairments include impaired memory, executive function, language, attention, and visual-spatial abilities.

Hypoglycemia, hyperglycemia, fluctuations in serum glucose, delirium, and in-hospital acute stress symptoms have been identified as possible risk factors for persistent cognitive impairment after critical illness.^{28–30} There is strong evidence that patients with delirium in the ICU are at a greater risk of long-term outcomes of cognitive dysfunction.^{29,31,32}

Dementia is a relevant disease of cognitive dysfunction (Fig. 3) and a number of studies have reported the association between dementia and ICU treatment.^{33–36} Among 10,348 intensive care patients who survived to hospital discharge, dementia was newly diagnosed in 1,648 (15.0%) over 3 years of follow-up compared to 12.2% in the general population.³³ Furthermore, pre-existing cognitive impairment in ICU populations is widespread. A cross-sectional comparative study reported that 37% of critically ill patients over 65 years of age in the ICU had pre-existing cognitive impairment.³⁴ Pre-existing cognitive impairment also affects cognitive function in PICS.




Fig. 3. Three common cognitive impairments among older adults: delirium, dementia, and depression.

The pathophysiology of cognitive impairment after ICU treatment remains unknown and might be a manifestation of brain dysfunction. However, further research is needed.

Mental impairment in PICS

Depression, anxiety, and post-traumatic stress disorder (PTSD) are the major mental illnesses that comprise PICS. The mental status impairments that can arise among critical illness survivors include depression in approximately 30% of survivors, anxiety in 70%, and PTSD, which is characterized by intrusive memories that arise from a combination of true events after ICU discharge, in 10–50%;³⁷ therefore, every patient with suspected PICS should undergo formal mental assessment if possible. A systematic review by Davydow *et al.*³⁸ showed that two of seven studies indicated female sex to be a significant predictor of PTSD after ICU care. Pre-existing depression, anxiety, PTSD, lower education level, and alcohol abuse also increase the risk of ICU-acquired mental illness.

Regarding prevention and treatment, a systematic review and meta-analysis on the effectiveness of early rehabilitation to prevent PICS in patients with critical illness undertaken by Fuke *et al.*²³ showed that early rehabilitation did not significantly improve patient mental status-related outcomes (hospital anxiety and depression). Among patients who had ICU diaries started on the fourth day of ICU admission, the PTSD symptom scores after 12 months were significantly reduced in surviving patients compared to those in the non-survivors (21 versus 34).³⁹ The diary was a daily record of the patients' ICU stay and was written in plain language by the healthcare staff and/or family, with accompanying photographs. A systematic review on the impact of ICU diaries reported that four of five randomized trials showed a significantly reduced rate of new-onset PTSD after 3 months with the use of ICU diaries (5% versus 13%, $P = 0.02$).⁴⁰

With regard to long-term outcomes in mental illness in PICS, Patel *et al.*⁴¹ prospectively observed 255 patients with shock and acute respiratory distress syndrome (ARDS), reporting an incidence of PTSD associated with ICU admission of 12% within 1 year of discharge; therefore, appropriate recognition of ICU-acquired mental illness followed by early treatment should be considered in ICU care.

Post-intensive care syndrome – family

Critical illness can not only have a significant physical and psychological impact on patients who survive but can also have a psychological impact on their families (Fig. 1). The factors associated with a high risk of adverse psychological

conditions in the families of ICU survivors include anxiety, depression, acute stress disorder, PTSD, and complicated grief. The cluster of such adverse psychological reactions is called post-intensive care syndrome – family (PICS-F).^{5,42}

Prevalence and risk factors

Prevalence of PICS-F conditions in the relatives of adult patients is shown in Table 1.^{9,37,43–54} The wide range of rates is due to differences in each study's patient population, measurement tools, and time frames varying from 1 week to 1 year. The risk factors for PICS-F include female sex, younger relative and patient age, lower educational level, having a critically ill spouse, having more comorbidities, and being an unmarried parent of a critically ill child.^{9,37,48,55,56} Other baseline risk factors include a history of anxiety, depression, or severe mental disease.³⁷

Prevention

Several prevention interventions have focused on adverse psychological reactions, including improving communication, providing family support, family presence in the ICU, and using specific consultations.^{39,57–62} Improving communication during end-of-life care, such as respect for the patient's values, preferences, and expressed needs and shared decision-making, alleviated grief symptoms in relatives of patients who had died in the ICU.^{57,58,62} Emotional and social support involving psychologists,⁵⁹ caseworkers, and social workers^{9,37} in family support can mitigate the impact of the crisis of critical illness and help prepare them for the patient's discharge. Intensive care unit diaries, when

Table 1. Prevalence of the elements of post-intensive care syndrome –family (PICS-F)^{9,37,43–54}

| Elements of PICS-F | Follow-up | Prevalence |
|----------------------|--------------|--|
| Depression | 1 week | 14.6–66.7% |
| | 1–3 months | 8–48.5% |
| | 1–6 months | 17.9% |
| | 1–12 months | 6–43.4% |
| Anxiety | 1 week | 42–66% |
| | 1–3 months | 21–49.3% |
| | 1–6 months | 15–24% |
| PTSD | 3–6 months | 33.1–49.0% |
| Burden/overload | ICU–2 months | 36% |
| Activity restriction | 1–2 months | Activity restriction scale score 22.1–23 |
| Complicated grief | 3–12 months | 5–46% |

ICU, intensive care unit; PTSD, post-traumatic stress disorder.

read by the family members after patient discharge, can fill in memory gaps and help them to understand what happened.^{39,60,61}

Post-intensive care syndrome in pediatrics

Several large critical-care databases have revealed mortality rates of critically ill children of approximately 2–4% in ICUs in developed countries, indicating that most of these children survive. However, some of these pediatric survivors experience long-term morbidity associated with their critical care.^{63–67}

Ong *et al.*⁶⁸ recently undertook a scoping review for critical appraisal of existing published works on functional outcome and physical impairment among pediatric critical care survivors. They found that the rate of acquired functional impairment ranged from 10% to 36% at ICU discharge and from 10% to 13% after 2 years. They also extracted risk factors for acquired functional impairment, including illness severity, the presence of organ dysfunction, length of ICU stay, and younger age.⁶⁸ Pinto *et al.*⁶⁵ reported that some PICU survivors subsequently died after their hospital discharge and developed new morbidity even 3 years later. With a focus on specific diseases and conditions, the SPROUT study, a recent international point prevalence study of pediatric sepsis, showed that 17% of sepsis survivors suffered moderate to severe disability.⁶⁹ Among children who survived ARDS, Ward *et al.*⁷⁰ suggested that up to one-third showed pulmonary dysfunction and that their quality-of-life scores 12 months after discharge were lower than those in asthmatic children.

When the concept of PICS was established in 2012, the necessity for special considerations for children and their

family was also mentioned. Since then, multiple surveys have been undertaken to describe long-term morbidity among children who survived critical illness as well as their families. In 2018, the framework of pediatric PICS (PICS-p) was conceptualized (Fig. 4).⁷¹ The fundamental framework was similar to that of adult PICS, with several unique features.

First, the most important viewpoint is that children's critical illness occurs during the dynamic process of their growth and maturation and that both their outcomes and their family's response (i.e., their parents and siblings) can interdependently influence their subsequent development and quality of life. Second, PICS-p includes a "social health" domain for children and their families in addition to the three conventional domains of physical, cognitive, and emotional health. Critical illness affects the social functioning of both children and their families; that is, reintegration with their friends at school, their social capital, and their parents' unemployment while caring for a sick child. These social health impairments, intertwined with morbidity in other health domains, can negatively impact their development and survival quality. Finally, a variety of recovery trajectories of surviving children and their families are indicated in the PICS-p framework, including improvement, deterioration, vacillation, or plateau over days or decades. Available evidence, although still limited, suggests that their outcomes are more heterogeneous than those for adult ICU survivors and their spouses.^{68,71,72}

Considerable obstacles involving PICS-p still exist, especially in the definition and evaluation of pediatric health status in each domain. Ong *et al.*⁶⁸ identified vast differences in measurement tools and follow-up timings of pediatric functional outcomes among 25 articles included in their




Fig. 4. Proposed framework for post-intensive care syndrome in pediatrics (PICS-p). Compared to the concept of PICS for adult intensive care unit survivors, the unique features of PICS-p include the importance of baseline status, system maturation and psychosocial development, stronger interdependence within the family, and recovery trajectories that can potentially impact a child's life for decades.

scoping review. Children have diverse functional status related to their age and developmental stage. The next step for PICS-p is to establish comprehensive sets of evaluation methods appropriate for each age group in each domain of the framework, which will enable elucidation of the natural history of PICS-p. Thereafter, interventional studies will be warranted to improve long-term outcomes among pediatric critical care survivors.

Prevention of PICS

ABCDEFGH bundle

The ABCDE bundle is widely known as the bundle that addresses the risks of sedation, delirium, and immobility. ABCDE is composed of: A, airway management, assess, prevent, and manage pain; B, breathing trials, including daily interruptions of mechanical ventilation, spontaneous awakening trials, and spontaneous breathing trials; C, choice of analgesia and sedation, coordination of care, and communication; D, delirium assessment, prevention, and management; and E, early mobility and exercise.^{9,73,74} They are also risk factors for PICS. Furthermore, FGH can be added to the list for the prevention of PICS (Fig. 5). FGH includes: F, family involvement, follow-up referrals, and functional reconciliation; G, good handoff communication; and H, handout materials on PICS and PICS-F. The present review focused on early mobility (physical rehabilitation), follow-up referrals (ICU follow-up clinics) with new domains, including nutrition, nursing care, diary, and environment management.

Physical rehabilitation

The main purpose of rehabilitation in the ICU is to improve the quality of life by maintaining, improving, and




Fig. 5. ABCDEFGH bundle for prevention of post-intensive care syndrome.

reacquiring activities of daily living.⁷⁵ Both ICU-AW and delirium, as parts of PICS, are related to a decreased quality of life.^{76,77} The Japanese Clinical Practice Guidelines for Management of Sepsis and Septic Shock 2016 (J-SSCG 2016) suggested implementing early-stage rehabilitation as a PICS preventative measure for sepsis or ICU patients.⁷⁸ Physical rehabilitation in the ICU could improve mobility status and muscle strength.⁷⁹ A recent systematic review clarified that physical rehabilitation decreases ICU-AW but not delirium-free days and did not improve mental health.²³ Additional large randomized controlled trials are needed to clarify the effect of physical rehabilitation on PICS.

The definition of “early” in early rehabilitation practice usually refers to intensive physical rehabilitation that is implemented in addition to regular care at any time during an ICU stay. The term “early” has yet to be defined as, among various studies, the onset of interventions could vary by as much as 1 week.⁸⁰ Many critically ill patients have PICS symptoms following ICU discharge. A previous systematic review reported no clear effect of intensive physical rehabilitation following ICU discharge on clinically relevant outcomes, such as quality of life.⁸¹ Our updated review also clarified no improvement in quality of life or mortality.⁸² Preventing PICS symptoms from ICU admission is more important than intensive treatment of PICS following ICU discharge.

Physical rehabilitation for mobility includes activities such as sitting, standing, and ambulation, as well as passive exercises including range-of-motion exercises and ergometers.⁸⁰ An ICU survey in Japan revealed that sitting on the edge of the bed was routinely provided in ICUs, whereas neuromuscular electrical stimulation and a cycle ergometer were rarely provided.^{83,84} The J-SSCG 2016 recommended against neuromuscular electrical stimulation as an ICU-AW preventative measure.⁷⁸ The dose–response of physical rehabilitation for clinical outcomes is unknown.⁸⁵ High-dose rehabilitation might lead to a higher quality of life than that for low-dose rehabilitation;⁷⁹ however, further studies are needed to clarify this point.

Nutrition

Nutritional therapy is vital for the prevention of PICS, especially ICU-AW. Adequate energy delivery and protein intake are the most important factors for muscle synthesis;^{86,87} moreover, energy debt is covered by catabolism mainly of the muscle, which is associated with lean body mass loss related to risk mortality.⁸⁸ Previous studies on nutrition therapy targeted mortality and infectious complications as outcomes. With the recent opinion that nutrition therapy should target muscle volume and strength,⁸⁹ there is a strong connection between nutritional therapy and PICS

(Fig. 6). Although studies have shown that the securement of minimum energy delivery with supplemental parenteral nutrition from the acute phase was associated with decreased PICS,⁹⁰ overfeeding could induce autophagy impairment and worsen ICU-AW.⁹¹ Therefore, we should target appropriate energy delivery and avoid overfeeding.^{92,93} Here is a paradigm that adequate proteins especially important for critically ill patients. Adequate protein delivery with total energy could reduce PICS;⁹⁴ however, a number of studies have reported that protein delivery alone does not reduce PICS.^{95,96} As muscle protein synthesis is maximized with appropriate exercise in healthy individuals,⁹⁷ not only nutrition therapy alone but also appropriate exercise and rehabilitation together with adequate nutrition are also necessary in critically ill patients (Fig. 6).⁹⁸ As for the particular kind of nutrition, leucine is the amino acid reported to induce muscle protein synthesis.⁹⁹ Unfortunately, administration of specific amino acids including leucine has not shown efficacy in critical care.¹⁰⁰ Approaches to enhance anabolic

power, such as β -hydroxy- β -methylbutyrate¹⁰¹ or oxandrolone,¹⁰² remain to be examined for the prevention of PICS and ICU-AW.¹⁰³

Environmental management for healing

Patients admitted to the ICU experience environmental stimuli, particularly noise and light. Excessive noise in the ICU has been reported in numerous studies.¹⁰⁴

A recent observational study in six ICUs suggested that background noise had a negative impact on sleep quality.¹⁰⁵ Five RCTs^{106–110} examined the effects of noise reduction devices such as earplugs and noise-canceling headphones combined with or without eye masks on sleep quality among patients in the ICU.^{106–110} All RCTs reported better perceived sleep quality in patients with the devices;^{106–110} however, three of five studies were carried out mainly in post-surgical patients and one was for non-ventilated, mostly cardiac patients. Thus, these findings did not have external




Fig. 6. Association between critical illness and intensive care unit-acquired weakness (ICU-AW)/post-intensive care syndrome (PICS) and the importance of nutrition therapy and rehabilitation. Malnutrition and inactivity accelerate ICU-AW/PICS, especially with skeletal muscle volume and strength/functional loss. Nutrition therapy and rehabilitation are essential factors and the basis for the prevention of PICS/ICU-AW.

validity to generalize to all ICU patient populations. Improved sleep quality by using noise reduction devices could reduce the development of delirium in the ICU. Litton *et al.*¹¹¹ undertook a meta-analysis of six RCTs and three non-randomized trials to assess the effects of earplugs on the development of delirium in ICU patients, reporting that earplugs significantly reduced the risk of delirium (relative risk, 0.59; 95% confidence interval, 0.44–0.78).

To date, there is no direct evidence of a relationship between environmental factors and long-term cognitive impairment; however, as mentioned above, some studies have reported a significant relationship between noise reduction strategies and delirium.¹¹¹ Delirium in the ICU was associated with long-term cognitive impairment; therefore, the possibility exists that environmental factors could affect long-term cognitive impairment through delirium during ICU stay due to impaired sleep quality.

Little is known about the contribution of environmental factors to the mental health of patients after intensive care. One RCT reported that music therapy or noise-canceling headphones reduced anxiety during ICU stay compared to usual care in patients with respiratory failure requiring mechanical ventilation.¹¹² These interventions could affect the symptoms of mental health after intensive care, despite the lack of verification.

Nursing care for PICS

One of the most important roles of nurses is the continuous implementation of measures to prevent PICS, including the ABCDEFGH bundle. Nurses spend most of their time on direct patient care.¹¹³ In addition to optimal analgesia, nurses can support safe light sedation by staying near patients.¹¹⁴ Through light sedation, patients can prepare to satisfy the higher levels of human needs.¹¹⁵ To understand and address the patient's needs, nurses need to know the patient's living conditions prior to hospitalization.⁷³ Nurses should assess gaps between the patient's prehospitalization and current functional abilities and should support functional reconciliation. Non-pharmacological interventions can also be important to restore the patient's ordinary daily function in the hospital environment.^{116,117} Family involvement also plays a key role.¹¹⁸ The provision of information, including PICS, to family members and using an ICU diary can strengthen the connection between the patient and family members and medical staff.^{119,120} Moreover, it can also promote family participation in patient care. Early rehabilitation and mobilization interventions can improve physical function in patients with critical illness.²³ Furthermore, short-term and high-frequency rehabilitation and mobilization interventions can improve the functional ability of patients.¹²¹ Nurses

facilitate patient mobility at all hours of the day and night^{113,122} and, therefore, could contribute to improving patient functional ability. The recovery process from PICS is a continuum.¹²³ Functional reconciliation requires continuous and consistent care even after ICU discharge. Thus, good handoff communication including information about PICS is necessary to achieve this consistent care.^{42,73}

Intensive care unit diaries

Intensive care unit diaries are completed by doctors and families of patients to record the patient's status while in the ICU and are kept to describe the patient's experiences. The *icu-diary.org* site has been introduced as one option for keeping an ICU diary.¹²⁴ The ICU diary is written for the patient by a family member or a medical person, such as a nurse, but could also be recorded by the patient. The ICU diaries can help to indicate the orientation of the patient, and could prevent PICS by alleviating anxiety, depression, and PTSD symptoms.^{39,125} Keeping a diary has been shown to reduce PTSD symptoms not only in patients but also in their families.

In these facilities, the nursing team in charge, mainly the main bedside nurse, determine whether an ICU diary is appropriate. If the diary is judged to be useful, then the concept is explained to the patient and their family, and the diary is started after obtaining their consent. The diary is used to periodically record general notes on events and daily occurrences, the patient's life, rehabilitation situation, etc. at the discretion of the nurse in charge. If desired, it can also include photographs. The doctor in charge, physical therapist, and clinical engineering technicians involved in care might also add to the diary. The diary is presented to the patient at ICU discharge.

Intensive care unit follow-up clinics and PICS

Intensive care unit follow-up clinics are specialized clinics for patients who have survived and been discharged from the ICU. They have attracted attention as a place for the diagnosis and treatment of PICS. These follow-up clinics have been spreading mainly in Europe in the last 20 years but have also been gradually spreading in recent years in North America. The ICU follow-up clinics have no fixed form or patient evaluation method and treatment interventions vary between facilities. In addition to supporting rehabilitation, cognitive function, and mental symptoms, clinic pharmacists can also adjust medications.

The RaCTICaL study is a representative RCT that examined the usefulness of ICU follow-up clinics.¹²⁶ This open-label RCT undertaken at three UK institutions targeted all

patients who survived hospitalization after receiving intensive care. In the intervention group, rehabilitation was carried out from hospitalization until discharge to the third month and clinic follow-up observation was carried out at 3 and 9 months after discharge. Neither rehabilitation nor follow-up was carried out in the control group. The primary end-points were the health-related quality of life score and the SF-36 after 12 months, and the secondary end-points were the health-related quality of life score after 6 months and the quality-adjusted life years, PTSD presence and severity of the onset, hospital anxiety and depression scale after 12 months, and cost efficiency after 12 months. However, the nurse-led intensive care follow-up program showed no evidence of the effect or cost-effectiveness in improving patient quality of life in the year after discharge from intensive care. Further work should focus on the roles of early physical rehabilitation, delirium, cognitive dysfunction, and relatives in the recovery from critical illness. Intensive care units should review their follow-up programs based on these findings. In 2015, Jensen and colleagues undertook a lineage review and meta-analysis to examine the effects of ICU follow-up on patient-centric outcomes, that is, quality of life, anxiety, depression, PTSD, physical and cognitive function, and reinstatement.¹²⁷ The results indicated that follow-up consultations do not improve quality of life, anxiety, depression, physical or cognitive function, or return to work. However, consultations appeared to reduce the symptoms of PTSD after ICU admission, perhaps due to individualized interventions aimed at reframing the ICU experience. At present, it is uncertain if the intervention is effective. Post-ICU follow-up needs to be developed in collaboration with patients and their families and the effect should be investigated in larger studies and within comparable settings.

Intensive care unit follow-up clinics are expected to be a place for follow-up of PICS developed during hospitalization as well as for the discovery and treatment of newly developed PICS after discharge. The clinics have been developed mainly in Europe; however, their format and methods for patient evaluations have not been adequately studied and vary between facilities. There is also insufficient evidence regarding the usefulness of ICU follow-up clinics; therefore, further verification is necessary for future development.

End-of-life care in acute medicine

Due to the aging of the general population and declining birth rates, the demographic structure of Japan has become the most super-aged society in the world. Consequently, the disease structure and medical care demand have also changed. In these social backgrounds, ethical concerns regarding life-sustaining therapy often arise in acute care settings, where end-of-life care

is frequently provided. The values and preferences for choice of treatment options vary widely, especially in end-of-life care. The cultural or social backgrounds of a country influence the practice of end-of-life care.^{128–130} Physicians in Asian ICUs tend to avoid withdrawing life-sustaining therapy at the end of life compared to physicians in non-Asian countries.¹³¹ Despite similarities in cultures, there are differences between Japan and other East Asian countries, such as China and Korea, in physician perceptions and practices regarding end-of-life care in intensive care.¹²⁹

Thus, guidance on end-of-life care in acute care settings based on our own cultures and social circumstances is urgently required. In November 2014, Guidelines of the End-of-life Care for Acute Disease and Intensive Care were proposed by three related Japanese associations, namely, the Japanese Association of Acute Medicine, the Japanese Society of Intensive Care Medicine, and the Japanese Circulation Society. These guidelines describe that the withholding, withdrawing, or termination of resuscitation or life-sustaining therapy are chosen according to the directives of the patient and family and the conscience of the medical staff.

Importantly, these guidelines were not based on legislation, which might be of concern for physicians in clinical settings. That is, we Japanese physicians at the front line of end-of-life care feel exposed to personal legal risks when limiting life-sustaining therapy. It is necessary for prefectures, municipalities, and local medical control authorities to take the initiative to establish an ordinance on clinical guidance for end-of-life care.

CONCLUSION

POST-intensive care syndrome includes physical, cognition, and mental impairments that occur during ICU stay or after ICU discharge, as well as the long-term prognosis of ICU patients. For prevention of PICS, it is important to carry out the ABCDEFGH bundle and new therapeutic strategies, including diary, nutrition, nursing care, and environmental management for healing. Additionally, PICS will be a new task for intensive care medicine in the 21st century that has reached the end of mature acute care, including several problems regarding end-of-life care.

DISCLOSURE

Approval of the research protocol: N/A.

Informed consent: N/A.

Registry and the registration no. of the study/trial: N/A.

Animal studies: N/A.

Conflict of interest: None declared.

REFERENCES

- 1 Yende S, Austin S, Rhodes A *et al.* Long-term quality of life among survivors of severe sepsis: analyses of two international trials. *Crit. Care Med.* 2016; 44: 1461–7.
- 2 Petsko GA. A seat at the table. *Genome Biol.* 2008; 9: 113.
- 3 Martin GS, Mannino DM, Moss M. The effect of age on the development and outcome of adult sepsis. *Crit. Care Med.* 2006; 34: 15–21.
- 4 Javadi P, Buchman TG, Stromberg PE *et al.* Iron dysregulation combined with aging prevents sepsis-induced apoptosis. *J. Surg. Res.* 2005; 128: 37–44.
- 5 Needham DM, Davidson J, Cohen H *et al.* Improving long-term outcomes after discharge from intensive care unit: report from a stakeholders' conference. *Crit. Care Med.* 2012; 40: 502–9.
- 6 Nakamura K, Azuhata T, Yokota H. The swallowing problem after acute care in the elderly patients. *J. Jpn Assoc. Acute Med. Health* 2018; 30: 103–109.
- 7 Kaukonen KM, Bailey M, Suzuki S, Pilcher D, Bellomo R. Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000–2012. *JAMA* 2014; 311: 1308–16.
- 8 Zambon M, Vincent JL. Mortality rates for patients with acute lung injury/ARDS have decreased over time. *Chest* 2008; 133: 1120–7.
- 9 Harvey MA, Davidson JE. Postintensive care syndrome: right care, right now... and later. *Crit. Care Med.* 2016; 44: 381–5.
- 10 Kress JP, Hall JB. ICU-acquired weakness and recovery from critical illness. *N. Engl. J. Med.* 2014; 370: 1626–35.
- 11 Latronico N, Bolton CF. Critical illness polyneuropathy and myopathy: a major cause of muscle weakness and paralysis. *Lancet Neurol.* 2011; 10: 931–41.
- 12 Farhan H, Moreno-Duarte I, Latronico N, Zafonte R, Eikermann M. Acquired muscle weakness in the surgical intensive care unit: nosology, epidemiology, diagnosis, and prevention. *Anesthesiology* 2016; 124: 207–34.
- 13 Stevens RD, Marshall SA, Cornblath DR *et al.* A framework for diagnosing and classifying intensive care unit-acquired weakness. *Crit. Care Med.* 2009; 37: S299–308.
- 14 Appleton RT, Kinsella J, Quasim T. The incidence of intensive care unit-acquired weakness syndromes: a systematic review. *J. Intensive Care Soc.* 2015; 16: 126–36.
- 15 Bednarik J, Lukas Z, Vondracek P. Critical illness polyneuropathy: the electrophysiological components of a complex entity. *Intensive Care Med.* 2003; 29: 1505–14.
- 16 Koch S, Spuler S, Deja M *et al.* Critical illness myopathy is frequent: accompanying neuropathy protracts ICU discharge. *J. Neurol. Neurosurg. Psychiatry* 2011; 82: 287–93.
- 17 Lefaucheur JP, Nordine T, Rodriguez P, Brochard L. Origin of ICU acquired paresis determined by direct muscle stimulation. *J. Neurol. Neurosurg. Psychiatry* 2006; 77: 500–6.
- 18 De Jonghe B, Bastuji-Garin S, Sharshar T, Outin H, Brochard L. Does ICU-acquired paresis lengthen weaning from mechanical ventilation? *Intensive Care Med.* 2004; 30: 1117–21.
- 19 Dinglas VD, Aronson Friedman L, Colantuoni E *et al.* Muscle weakness and 5-year survival in acute respiratory distress syndrome survivors. *Crit. Care Med.* 2017; 45: 446–53.
- 20 Stevens RD, Dowdy DW, Michaels RK, Mendez-Tellez PA, Pronovost PJ, Needham DM. Neuromuscular dysfunction acquired in critical illness: a systematic review. *Intensive Care Med.* 2007; 33: 1876–91.
- 21 Guarneri B, Bertolini G, Latronico N. Long-term outcome in patients with critical illness myopathy or neuropathy: the Italian multicentre CRIMYNE study. *J. Neurol. Neurosurg. Psychiatry* 2008; 79: 838–41.
- 22 Koch S, Wollersheim T, Bierbrauer J *et al.* Long-term recovery in critical illness myopathy is complete, contrary to polyneuropathy. *Muscle Nerve* 2014; 50: 431–6.
- 23 Fuke R, Hifumi T, Kondo Y *et al.* Early rehabilitation to prevent postintensive care syndrome in patients with critical illness: a systematic review and meta-analysis. *BMJ Open* 2018; 8: e019998.
- 24 Routsis C, Gerovasili V, Vasileiadis I *et al.* Electrical muscle stimulation prevents critical illness polyneuromyopathy: a randomized parallel intervention trial. *Crit. Care* 2010; 14: R74.
- 25 Hermans G, De Jonghe B, Bruyninckx F, Van den Berghe G. Interventions for preventing critical illness polyneuropathy and critical illness myopathy. *Cochrane Database Syst. Rev.* 2014; 1: CD006832.
- 26 Davidson JE, Harvey MA, Bemis-Dougherty A, Smith JM, Hopkins RO. Implementation of the pain, agitation, and delirium clinical practice guidelines and promoting patient mobility to prevent post-intensive care syndrome. *Crit. Care Med.* 2013; 41: S136–45.
- 27 Wolters AE, Slooter AJ, van der Kooi AW, van Dijk D. Cognitive impairment after intensive care unit admission: a systematic review. *Intensive Care Med.* 2013; 39: 376–86.
- 28 Hopkins RO, Suchyta MR, Snow GL, Jephson A, Weaver LK, Orme JF. Blood glucose dysregulation and cognitive outcome in ARDS survivors. *Brain Inj.* 2010; 24: 1478–84.
- 29 Pandharipande PP, Girard TD, Jackson JC *et al.* Long-term cognitive impairment after critical illness. *N. Engl. J. Med.* 2013; 369: 1306–16.
- 30 Davydow DS, Zatzick D, Hough CL, Katon WJ. In-hospital acute stress symptoms are associated with impairment in cognition 1 year after intensive care unit admission. *Ann. Am. Thorac. Soc.* 2013; 10: 450–7.
- 31 Jackson JC, Gordon SM, Ely EW, Burger C, Hopkins RO. Research issues in the evaluation of cognitive impairment in intensive care unit survivors. *Intensive Care Med.* 2004; 30: 2009–16.
- 32 Katz IR, Curyto KJ, TenHave T, Mossey J, Sands L, Kallan MJ. Validating the diagnosis of delirium and evaluating its

- association with deterioration over a one-year period. *Am. J. Geriatr. Psychiatry* 2001; 9: 148–59.
- 33 Guerra C, Hua M, Wunsch H. Risk of a diagnosis of dementia for elderly medicare beneficiaries after intensive care. *Anesthesiology* 2015; 123: 1105–12.
 - 34 Pisani MA, Redlich C, McNicoll L, Ely EW, Inouye SK. Underrecognition of preexisting cognitive impairment by physicians in older ICU patients. *Chest* 2003; 124: 2267–74.
 - 35 Pisani MA, Redlich CA, McNicoll L, Ely EW, Friedkin RJ, Inouye SK. Short-term outcomes in older intensive care unit patients with dementia. *Crit. Care Med.* 2005; 33: 1371–6.
 - 36 Mandebvu F, Kalman M. The 3 Ds, and newly acquired cognitive impairment: issues for the ICU nurse. *Crit. Care Nurs. Q.* 2015; 38: 317–26.
 - 37 Davidson JE, Jones C, Bienvenu OJ. Family response to critical illness: postintensive care syndrome-family. *Crit. Care Med.* 2012; 40: 618–24.
 - 38 Davydow DS, Gifford JM, Desai SV, Needham DM, Bienvenu OJ. Posttraumatic stress disorder in general intensive care unit survivors: a systematic review. *Gen. Hosp. Psychiatry* 2008; 30: 421–34.
 - 39 Garrouste-Orgeas M, Coquet I, Perier A *et al.* Impact of an intensive care unit diary on psychological distress in patients and relatives*. *Crit. Care Med.* 2012; 40: 2033–40.
 - 40 Mehlhorn J, Freytag A, Schmidt K *et al.* Rehabilitation interventions for postintensive care syndrome: a systematic review. *Crit. Care Med.* 2014; 42: 1263–71.
 - 41 Patel MB, Jackson JC, Morandi A *et al.* Incidence and risk factors for intensive care unit-related post-traumatic stress disorder in veterans and civilians. *Am. J. Respir. Crit. Care Med.* 2016; 193: 1373–81.
 - 42 Elliott D, Davidson JE, Harvey MA *et al.* Exploring the scope of post-intensive care syndrome therapy and care: engagement of non-critical care providers and survivors in a second stakeholders meeting. *Crit. Care Med.* 2014; 42: 2518–26.
 - 43 Haines KJ, Denehy L, Skinner EH, Warrillow S, Berney S. Psychosocial outcomes in informal caregivers of the critically ill: a systematic review*. *Crit. Care Med.* 2015; 43: 1112–20.
 - 44 Netzer G, Sullivan DR. Recognizing, naming, and measuring a family intensive care unit syndrome. *Ann. Am. Thorac. Soc.* 2014; 11: 435–41.
 - 45 Jones C, Skirrow P, Griffiths RD *et al.* Post-traumatic stress disorder-related symptoms in relatives of patients following intensive care. *Intensive Care Med.* 2004; 30: 456–60.
 - 46 Azoulay E, Pochard F, Kentish-Barnes N *et al.* Risk of post-traumatic stress symptoms in family members of intensive care unit patients. *Am. J. Respir. Crit. Care Med.* 2005; 171: 987–94.
 - 47 Anderson WG, Arnold RM, Angus DC, Bryce CL. Posttraumatic stress and complicated grief in family members of patients in the intensive care unit. *J. Gen. Intern. Med.* 2008; 23: 1871–6.
 - 48 Siegel MD, Hayes E, Vanderwerker LC, Loseth DB, Prigerson HG. Psychiatric illness in the next of kin of patients who die in the intensive care unit*. *Crit. Care Med.* 2008; 36: 1722–8.
 - 49 Kentish-Barnes N, Chaize M, Seegers V *et al.* Complicated grief after death of a relative in the intensive care unit. *Eur. Respir. J.* 2015; 45: 1341–52.
 - 50 Torres J, Carvalho D, Molinos E *et al.* The impact of the patient post-intensive care syndrome components upon caregiver burden. *Med. Intensiva* 2017; 41: 454–60.
 - 51 Anderson WG, Arnold RM, Angus DC, Bryce CL. Passive decision-making preference is associated with anxiety and depression in relatives of patients in the intensive care unit. *J. Crit. Care* 2009; 24: 249–54.
 - 52 Matt B, Schwarzkopf D, Reinhart K, König C, Hartog CS. Relatives' perception of stressors and psychological outcomes – results from a survey study. *J. Crit. Care* 2017; 39: 172–7.
 - 53 Petrinc AB, Martin BRJP. Post-intensive care syndrome symptoms and health-related quality of life in family decision-makers of critically ill patients. *Palliat. Support Care* 2017; 16: 719–24.
 - 54 Cameron JI, Chu LM, Matte A *et al.* One-year outcomes in caregivers of critically ill patients. *N. Engl. J. Med.* 2016; 374: 1831–41.
 - 55 Alfheim HB, Rosseland LA, Hofso K, Smastuen MC, Rustoen T. Multiple symptoms in family caregivers of intensive care unit patients. *J. Pain Symptom Manage.* 2018; 55: 387–94.
 - 56 Gries CJ, Engelberg RA, Kross EK *et al.* Predictors of symptoms of posttraumatic stress and depression in family members after patient death in the ICU. *CHEST J.* 2010; 137: 280–7.
 - 57 Lautrette A, Darmon M, Megarbane B *et al.* A communication strategy and brochure for relatives of patients dying in the ICU. *N. Engl. J. Med.* 2007; 356: 469–78.
 - 58 Curtis JR, Treece PD, Nielsen EL *et al.* Randomized trial of communication facilitators to reduce family distress and intensity of end-of-life care. *Am. J. Respir. Crit. Care Med.* 2016; 193: 154–62.
 - 59 Peris A, Bonizzoli M, Iozzelli D *et al.* Early intra-intensive care unit psychological intervention promotes recovery from post traumatic stress disorders, anxiety and depression symptoms in critically ill patients. *Crit. Care* 2011; 15: R41.
 - 60 Jones C, Backman C, Capuzzo M *et al.* Intensive care diaries reduce new onset post traumatic stress disorder following critical illness: a randomised, controlled trial. *Crit. Care* 2010; 14: R168.
 - 61 Jones C, Bäckman C, Griffiths RD. Intensive care diaries and relatives' symptoms of posttraumatic stress disorder after critical illness: a pilot study. *Am. J. Crit. Care* 2012; 21: 172–6.
 - 62 Nelson JE, Puntillo KA, Pronovost PJ *et al.* In their own words: patients and families define high-quality palliative

- care in the intensive care unit. *Crit. Care Med.* 2010; 38: 808.
- 63 Fiser DH, Tilford JM, Roberson PK. Relationship of illness severity and length of stay to functional outcomes in the pediatric intensive care unit: a multi-institutional study. *Crit. Care Med.* 2000; 28: 1173–9.
- 64 Jones S, Rantell K, Stevens K *et al.* Outcome at 6 months after admission for pediatric intensive care: a report of a national study of pediatric intensive care units in the United Kingdom. *Pediatrics* 2006; 118: 2101–8.
- 65 Pinto NP, Rhinesmith EW, Kim TY, Ladner PH, Pollack MM. Long-term function after pediatric critical illness: results from the survivor outcomes study. *Pediatr. Crit. Care Med.* 2017; 18: e122–30.
- 66 Pollack MM, Holubkov R, Funai T *et al.* Pediatric intensive care outcomes: development of new morbidities during pediatric critical care. *Pediatr. Crit. Care Med.* 2014; 15: 821–7.
- 67 Typpo KV, Petersen NJ, Hallman DM, Markovitz BP, Marsiscalco MM. Day 1 multiple organ dysfunction syndrome is associated with poor functional outcome and mortality in the pediatric intensive care unit. *Pediatr. Crit. Care Med.* 2009; 10: 562–70.
- 68 Ong C, Lee JH, Leow MK, Puthuchery ZA. Functional outcomes and physical impairments in pediatric critical care survivors: a scoping review. *Pediatr. Crit. Care Med.* 2016; 17: e247–59.
- 69 Weiss SL, Fitzgerald JC, Pappachan J *et al.* Global epidemiology of pediatric severe sepsis: the sepsis prevalence, outcomes, and therapies study. *Am. J. Respir. Crit. Care Med.* 2015; 191: 1147–57.
- 70 Ward SL, Turpin A, Spicer AC, Treadwell MJ, Church GD, Flori HR. Long-term pulmonary function and quality of life in children after acute respiratory distress syndrome: a feasibility investigation. *Pediatr. Crit. Care Med.* 2017; 18: e48–55.
- 71 Manning JC, Pinto NP, Rennick JE, Colville G, Curley MAQ. Conceptualizing post intensive care syndrome in children—the PICS-p framework. *Pediatr. Crit. Care Med.* 2018; 19: 298–300.
- 72 Nelson LP, Gold JI. Posttraumatic stress disorder in children and their parents following admission to the pediatric intensive care unit: a review. *Pediatr. Crit. Care Med.* 2012; 13: 338–47.
- 73 Davidson JE, Harvey MA, Schuller J, Black G. Post-intensive care syndrome: what is it and how to help prevent it. *Am. Nurse Today* 2013; 8: 32–8.
- 74 Ely EW. The ABCDEF bundle: science and philosophy of how ICU liberation serves patients and families. *Crit. Care Med.* 2017; 45: 321–30.
- 75 Hodgson CL, Udy AA, Bailey M *et al.* The impact of disability in survivors of critical illness. *Intensive Care Med.* 2017; 43: 992–1001.
- 76 Fan E, Dowdy DW, Colantuoni E *et al.* Physical complications in acute lung injury survivors: a two-year longitudinal prospective study. *Crit. Care Med.* 2014; 42: 849–59.
- 77 Naidech AM, Beaumont JL, Rosenberg NF *et al.* Intracerebral hemorrhage and delirium symptoms. Length of stay, function, and quality of life in a 114-patient cohort. *Am. J. Respir. Crit. Care Med.* 2013; 188: 1331–7.
- 78 Nishida O, Ogura H, Egi M *et al.* The Japanese clinical practice guidelines for management of sepsis and septic shock 2016 (J-SSCG 2016). *Acute Med. Surg.* 2018; 5: 3–89.
- 79 Tipping CJ, Harrold M, Holland A, Romero L, Nisbet T, Hodgson CL. The effects of active mobilisation and rehabilitation in ICU on mortality and function: a systematic review. *Intensive Care Med.* 2017; 43: 171–83.
- 80 Taito S, Shime N, Ota K, Yasuda H. Early mobilization of mechanically ventilated patients in the intensive care unit. *J. Intensive Care* 2016; 4: 50.
- 81 Connolly B, Salisbury L, O'Neill B *et al.* Exercise rehabilitation following intensive care unit discharge for recovery from critical illness. *Cochrane Database Syst. Rev.* 2015; CD008632.
- 82 Taito S, Yamauchi K, Tsujimoto Y, Banno M, Tsujimoto H, Kataoka Y. A systematic review and meta-analysis of physical rehabilitation following intensive care unit discharge. 2018; [cited 10 April 2019]. Available from: https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=80532.
- 83 Taito S, Sanui M, Yasuda H, Shime N, Lefor AK. Current rehabilitation practices in intensive care units: a preliminary survey by the Japanese Society of Education for Physicians and Trainees in Intensive Care (JSEPTIC) Clinical Trial Group. *J. Intensive Care* 2016; 4: 66.
- 84 Taito S, Shime N, Yasuda H *et al.* Out-of-bed mobilization of patients undergoing mechanical ventilation with orotracheal tubes: a survey study. *J. Crit. Care* 2018; 47: 173–7.
- 85 Hodgson CL, Capell E, Tipping CJ. Early mobilization of patients in intensive care: organization, communication and safety factors that influence translation into clinical practice. *Crit. Care* 2018; 22: 77.
- 86 Phillips SM. A brief review of critical processes in exercise-induced muscular hypertrophy. *Sports Med.* 2014; 44(Suppl 1): S71–7.
- 87 Kim IY, Schutzler S, Schrader A *et al.* The anabolic response to a meal containing different amounts of protein is not limited by the maximal stimulation of protein synthesis in healthy young adults. *Am. J. Physiol. Endocrinol. Metab.* 2016; 310: E73–80.
- 88 Demling RH. Nutrition, anabolism, and the wound healing process: an overview. *Eplasty* 2009; 9: e9.
- 89 Landi F, Camprubi-Robles M, Bear DE *et al.* Muscle loss: the new malnutrition challenge in clinical practice. *Clin. Nutr.* 2018; S0261–5614(18)32554–8.
- 90 Wischmeyer PE, Hasselmann M, Kummerlen C *et al.* A randomized trial of supplemental parenteral nutrition in

- underweight and overweight critically ill patients: the TOP-UP pilot trial. *Crit. Care* 2017; 21: 142.
- 91 Casaer MP, Wilmer A, Hermans G, Wouters PJ, Mesotten D, Van den Berghe G. Role of disease and macronutrient dose in the randomized controlled EPaNIC trial: a post hoc analysis. *Am. J. Respir. Crit. Care Med.* 2013; 187: 247–55.
- 92 McClave SA, Taylor BE, Martindale RG *et al.* Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: society of critical care medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J. Parenter. Enteral Nutr.* 2016; 40: 159–211.
- 93 Singer P, Blaser AR, Berger MM *et al.* ESPEN guideline on clinical nutrition in the intensive care unit. *Clin. Nutr.* 2018; 38: 48–79.
- 94 Berger MM, Pantet O, Jacquelin-Ravel N *et al.* Supplemental parenteral nutrition improves immunity with unchanged carbohydrate and protein metabolism in critically ill patients: the SPN2 randomized tracer study. *Clin. Nutr.* 2018; S0261–5614(18)32518–4.
- 95 Allingstrup MJ, Kondrup J, Wiis J *et al.* Early goal-directed nutrition versus standard of care in adult intensive care patients: the single-centre, randomised, outcome assessor-blinded EAT-ICU trial. *Intensive Care Med.* 2017; 43: 1637–47.
- 96 Lambell KJ, King SJ, Forsyth AK, Tierney AC. Association of energy and protein delivery on skeletal muscle mass changes in critically ill adults: a systematic review. *JPEN J. Parenter. Enteral Nutr.* 2018; 42: 1112–22.
- 97 Morton RW, Murphy KT, McKellar SR *et al.* A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br. J. Sports Med.* 2018; 52: 376–84.
- 98 Jones C, Eddleston J, McCairn A *et al.* Improving rehabilitation after critical illness through outpatient physiotherapy classes and essential amino acid supplement: a randomized controlled trial. *J. Crit. Care* 2015; 30: 901–7.
- 99 Wolfe RR. The 2017 Sir David P Cuthbertson lecture. Amino acids and muscle protein metabolism in critical care. *Clin. Nutr.* 2018; 37: 1093–100.
- 100 Ginguay A, De Bandt JP, Cynober L. Indications and contraindications for infusing specific amino acids (leucine, glutamine, arginine, citrulline, and taurine) in critical illness. *Curr. Opin. Clin. Nutr. Metab. Care* 2016; 19: 161–9.
- 101 Rahman A, Wilund K, Fitschen PJ *et al.* Elderly persons with ICU-acquired weakness: the potential role for β -hydroxy- β -methylbutyrate (HMB) supplementation? *JPEN J. Parenter. Enteral Nutr.* 2014; 38: 567–75.
- 102 Demling RH, DeSanti L. Oxandrolone induced lean mass gain during recovery from severe burns is maintained after discontinuation of the anabolic steroid. *Burns* 2003; 29: 793–7.
- 103 Stanojevic M, Finnerty CC, Jeschke MG. Anabolic and anti-catabolic agents in critical care. *Curr. Opin. Crit. Care* 2016; 22: 325–31.
- 104 Konkani A, Oakley B. Noise in hospital intensive care units—a critical review of a critical topic. *J. Crit. Care* 2012; 27: 522e1–9.
- 105 Simons KS, Verweij E, Lemmens PMC *et al.* Noise in the intensive care unit and its influence on sleep quality: a multi-center observational study in Dutch intensive care units. *Crit. Care* 2018; 22: 250.
- 106 Hu RF, Jiang XY, Hegadoren KM, Zhang YH. Effects of earplugs and eye masks combined with relaxing music on sleep, melatonin and cortisol levels in ICU patients: a randomized controlled trial. *Crit. Care* 2015; 19: 115.
- 107 Le Guen M, Nicolas-Robin A, Lebard C, Arnulf I, Langeron O. Earplugs and eye masks vs routine care prevent sleep impairment in post-anaesthesia care unit: a randomized study. *Br. J. Anaesth.* 2014; 112: 89–95.
- 108 Litton E, Elliott R, Ferrier J, Webb SAR. Quality sleep using earplugs in the intensive care unit: the QUIET pilot randomised controlled trial. *Crit. Care Resusc.* 2017; 19: 128–33.
- 109 Menger J, Urbanek B, Skhirtladze-Dworschak K *et al.* Earplugs during the first night after cardiothoracic surgery may improve a fast-track protocol. *Minerva Anesthesiol.* 2018; 84: 49–57.
- 110 Scotto CJ, McClusky C, Spillan S, Kimmel J. Earplugs improve patients' subjective experience of sleep in critical care. *Nurs. Crit. Care* 2009; 14: 180–4.
- 111 Litton E, Carnegie V, Elliott R, Webb SA. The efficacy of earplugs as a sleep hygiene strategy for reducing delirium in the ICU: a systematic review and meta-analysis. *Crit. Care Med.* 2016; 44: 992–9.
- 112 Chlan LL, Weinert CR, Heiderscheid A *et al.* Effects of patient-directed music intervention on anxiety and sedative exposure in critically ill patients receiving mechanical ventilatory support: a randomized clinical trial. *JAMA* 2013; 309: 2335–44.
- 113 Young DL, Seltzer J, Glover M *et al.* Identifying barriers to nurse-facilitated patient mobility in the intensive care unit. *Am. J. Crit. Care* 2018; 27: 186–93.
- 114 Strom T, Martinussen T, Toft P. A protocol of no sedation for critically ill patients receiving mechanical ventilation: a randomised trial. *Lancet* 2010; 375: 475–80.
- 115 Jackson JC, Santoro MJ, Ely TM *et al.* Improving patient care through the prism of psychology: application of Maslow's hierarchy to sedation, delirium, and early mobility in the intensive care unit. *J. Crit. Care* 2014; 29: 438–44.
- 116 Bannon L, McGaughey J, Verghis R, Clarke M, McAuley DF, Blackwood B. The effectiveness of non-pharmacological interventions in reducing the incidence and duration of delirium in critically ill patients: a systematic review and meta-analysis. *Intensive Care Med.* 2019; 45: 1–12.

- 117 Kang J, Lee M, Ko H *et al.* Effect of nonpharmacological interventions for the prevention of delirium in the intensive care unit: a systematic review and meta-analysis. *J. Crit. Care* 2018; 48: 372–84.
- 118 Davidson JE, Powers K, Hedayat KM *et al.* Clinical practice guidelines for support of the family in the patient-centered intensive care unit: American College of Critical Care Medicine Task Force 2004–2005. *Crit. Care Med.* 2007; 35: 605–22.
- 119 Azoulay E, Pochard F, Chevret S *et al.* Impact of a family information leaflet on effectiveness of information provided to family members of intensive care unit patients: a multi-center, prospective, randomized, controlled trial. *Am. J. Respir. Crit. Care Med.* 2002; 165: 438–42.
- 120 Garrouste-Orgeas M, Perier A, Mouricou P *et al.* Writing in and reading ICU diaries: qualitative study of families' experience in the ICU. *PLoS ONE* 2014; 9: e110146.
- 121 Bernhardt J, Churilov L, Ellery F *et al.* Prespecified dose-response analysis for A Very Early Rehabilitation Trial (AVERT). *Neurology* 2016; 86: 2138–45.
- 122 Cortes OL, DiCenso A, McKelvie R. Mobilization patterns of patients after an acute myocardial infarction: a pilot study. *Clin. Nurs. Res.* 2015; 24: 139–55.
- 123 Kang J, Jeong YJ. Embracing the new vulnerable self: a grounded theory approach on critical care survivors' post-intensive care syndrome. *Intensive Crit. Care Nurs.* 2018; 49: 44–50.
- 124 [cited 3 Feb 2019]. Available from: <http://www.icu-diary.org/diary/start.html>.
- 125 Petrinc AB, Mazanec PM, Burant CJ, Hoffer A, Daly BJ. Coping strategies and posttraumatic stress symptoms in post-ICU family decision makers. *Crit. Care Med.* 2015; 43: 1205–12.
- 126 Cuthbertson BH, Rattray J, Campbell MK *et al.* The PRaCTICaL study of nurse led, intensive care follow-up programmes for improving long term outcomes from critical illness: a pragmatic randomised controlled trial. *BMJ* 2009; 339: b3723.
- 127 Jensen JF, Thomsen T, Overgaard D, Bestle MH, Christensen D, Egerod I. Impact of follow-up consultations for ICU survivors on post-ICU syndrome: a systematic review and meta-analysis. *Intensive Care Med.* 2015; 41: 763–75.
- 128 Curtis JR, Vincent JL. Ethics and end-of-life care for adults in the intensive care unit. *Lancet* 2010; 376: 1347–53.
- 129 Park SY, Phua J, Nishimura M *et al.* End-of-life care in ICUs in East Asia: a comparison among China, Korea, and Japan. *Crit. Care Med.* 2018; 46: 1114–24.
- 130 Vincent JL. Cultural differences in end-of-life care. *Crit. Care Med.* 2001; 29: N52–5.
- 131 Phua J, Joynt GM, Nishimura M *et al.* Withholding and withdrawal of life-sustaining treatments in intensive care units in Asia. *JAMA Intern. Med.* 2015; 175: 363–71.