

Abstracts for a conference on trace elements in diet, nutrition, and health: essentiality and toxicity

A Joint Conference Constituting the:

VIIIth Conference of the International Society for Trace Element Research in Humans (ISTERH)
IXth Conference of the Nordic Trace Element Society (NTES)
VIth Conference of the Hellenic Trace Element Society (HTES)
Hersonissos (Crete-Greece)
October 21–26, 2007
Creta Maris Conference Center

Members of planning committee:

Curtiss Hunt, Ph.D., Chair, Planning Committee; President, ISTERH
Sophie Ermidou-Pollet, Ph.D., President, HTES
Ole Andersen, Ph.D., Representative, NTES
Dorothy Klimis-Zacas, Ph.D., Chair, Local Organizing Committee; Member, ISTERH
Jeanne Freeland-Graves, Ph.D., Chair, Fundraising Committee; Council Member, ISTERH
Monica Nordberg, Ph.D., Chair, Abstract Committee; Vice President, ISTERH
Hiroko Kodama, Ph.D., Chair, Publicity Committee; Secretary, ISTERH
Serge Pollet, Ph.D., Member at Large; Vice President, HTES
George Brewer, M.D., Member at Large; Council Member, ISTERH
Harold Sandstead, M.D., Member at Large; Immediate Past President, ISTERH

Members of abstract committee:

Monica Nordberg (Sweden), Chair; Ole Andersen, (Denmark), Greg Anderson (Australia), Richard Anderson (USA), Mario Barbagallo (Italy), Ramon Barnes (USA), John Beattie (UK), Muriel Bost (France), Sophie

Ermidou-Pollet (Greece), Susan Fairweather-Tait (UK), Bruce Fowler (USA), Göran Friman (Sweden), Rosalind Gibson (New Zealand), W. Thomas Johnson (USA), Joe Landolph (USA), Bo Lonnerdal (USA), Wolfgang Maret (USA), Jesper Nielsen (Denmark), Serge Pollet (Greece), Prem Ponka (Canada), Manju Reddy (USA), Per Roos (Norway), Anne Roussel (France), Swapan Kumar Roy (Bangladesh), Manuel Ruz (Chile), Hiroshi Satoh (Japan), Ugur Sayli (Turkey), Sunil Sazawal (USA), Songsak Sriamujata (Thailand), John Joseph Strain (UK), Tore Syversen (Norway), Anna Viegas-Crespo (Portugal), Neil Ward (UK), Richard Wood (USA), Katsuhiko Yokoi (Japan)

Acknowledgements

Funding for this conference was made possible in part by 1 R13 DK080637-01 from the National Institutes of Health. The views expressed in written conference materials or publications and by speakers and moderators do not necessarily reflect the official policies of the Department of Health and Human Services, nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government. The project was supported by the National Research Initiative of the USDA Cooperative State Research, Education, and Extension Service Grant number 2007-35200-18235

Category 1: Trace element intakes, dietary patterns, bioavailability, and tissue distributions

1.P01

The effect of enrichment with sunflower seed, sesame seed and alpha tocopherol acetate to linoleic acid quantity in cookies

Sumalika Piammongkol Patcharin Pakdeechanuan; Prince of Songkla University (Pattani, Thailand)

findings from modern basic scientific approaches can greatly improve risk assessments for toxic trace elements and help to delineate populations at special risk within the general population.

14.B.3

Neurological and renal effects of low dose exposure to elemental mercury exposure from amalgam in children

Lars Barregard

Sahlgrenska University Hospital and Academy (Göteborg, Sweden)

Dental amalgam constitutes the most common source of exposure to elemental mercury in the general population, and the possible health risks have been debated for a long time. Many studies have been performed in adults exposed to low levels of elemental mercury, but until recently, a few data were available on this topic in children, who might be more vulnerable. In 2006, two large randomized controlled trials were published, comparing neuropsychological (full scale IQ, memory, attention, visuomotor tests) and renal outcomes (e.g., albumin excretion) between children treated with dental amalgam or composites over a 5-year trial period. The studies included more than 1,000 children from northeast USA or Portugal. These studies failed to demonstrate any statistically significant differences in adverse effects between treatment groups. The increase in urinary mercury in the amalgam-treated groups compared to controls treated with composites was, however, limited, on average about 0.5–1 µg/g creatinine. The results of these two trials will be reviewed, together with subsequent studies using this rich data set. The strengths and limitations of the two trials will be discussed as well as children's exposure to inorganic mercury in general and the results of some recent observational studies of possible adverse effects.

14.B.4

Manganese exposure as a determinant of parkinsonian damage

Roberto Lucchini, Elisa Albini, Laura Benedetti, Stefano Borghesi, Annalisa Caruso, Eleonora Nan, Giovanni Parrinello, and Lorenzo Alessio
University of Brescia (Brescia, Italy)

Manganese is an essential element for humans and animals and plays an important role in bone mineralization, protein and energy metabolism, metabolic regulation, cellular protection from damaging free radical species, and the formation of glycosaminoglycans. Homeostatic mechanisms regulate the absorption and excretion rates to keep manganese concentration within a strict range. Nevertheless, short-term exposure to high doses or prolonged exposure to low doses can determine a manganese overload in the central nervous system, given the slow elimination rate from this organ. In cases of overload, manganese accumulates in the globus pallidus of the basal ganglia, where it can cause cellular damage on the GABAergic and dopaminergic pathways. As a consequence, motor function and coordination of fine movements are affected, and mood regulation as well with marked aggressivity. Human exposure to air concentration higher than 1 mg/m³ can determine the clinical picture of manganism, an atypical Parkinsonism that shows clinical differences from the typical features of Parkinson's Disease. After prolonged exposure at much lower levels, manganese may also act as an environmental trigger and favor the onset of typical Parkinsonian disturbances. This can be determined by a damage of the dopaminergic neurons of the substantia nigra-pars compacta, which is located very closely to the globus pallidus and shares various interconnections within the basal ganglia. This hypothesis is presented based on experimental animal studies and human epidemiological studies on exposed workers and the general population resident in the proximities of polluting sources.

14.B.5

Kidney and bone effects of low-dose cadmium exposure in Sweden

Agneta Åkesson

Karolinska Institutet (Stockholm, Sweden)

Cadmium is a toxic environmental pollutant that all are exposed to via foods considered healthy, such as whole grains, vegetables, and potatoes. High-cadmium exposure is well-known to cause kidney and bone damage, but these associations at low-level cadmium exposure need further attention, and the critical exposure level in the general population needs to be defined. We investigated health effects of cadmium exposure