



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: January 9, 2016

REVISED: March 5, 2016

ACCEPTED: March 8, 2016

PUBLISHED: March 21, 2016

De Sitter from T-branes

Michele Cicoli,^{a,b,c} Fernando Quevedo^{c,d} and Roberto Valandro^{e,f,c}

^a*Dipartimento di Fisica e Astronomia, Università di Bologna,
via Irnerio 46, Bologna, 40126 Italy*

^b*INFN, Sezione di Bologna,
via Irnerio 46, Bologna, 40126 Italy*

^c*ICTP,
Strada Costiera 11, Trieste, 34151 Italy*

^d*DAMTP, Centre for Mathematical Sciences,
Wilberforce Road, Cambridge, CB3 0WA U.K.*

^e*Dipartimento di Fisica, Università di Trieste,
Strada Costiera 11, Trieste, 34151 Italy*

^f*INFN, Sezione di Trieste,
Via Valerio 2, Trieste, 34127 Italy*

E-mail: mcicoli@ictp.it, f.quevedo@damtp.cam.ac.uk,
roberto.valandro@ts.infn.it

ABSTRACT: Hidden sector D7-branes with non-zero gauge flux are a generic feature of type IIB compactifications. A non-vanishing Fayet-Iliopoulos term induced by non-zero gauge flux leads to a T-brane configuration. Expanding the D7-brane action around this T-brane background in the presence of three-form supersymmetry breaking fluxes, we obtain a positive definite contribution to the moduli scalar potential which can be used as an uplifting source for de Sitter vacua. In this way we provide a higher-dimensional understanding of known 4D mechanisms of de Sitter uplifting based on hidden sector F-terms which are non-zero because of D-term stabilisation.

KEYWORDS: Superstring Vacua, D-branes, F-Theory, Flux compactifications

ARXIV EPRINT: [1512.04558](https://arxiv.org/abs/1512.04558)

Contents

1	Introduction	1
2	4D point of view: dS from hidden matter F-terms	3
2.1	Hidden D7-branes and gauge fluxes	3
2.2	Uplifting term from hidden matter F-terms	5
2.3	Gauge fluxes and non-perturbative effects	6
2.4	dS LVS vacua	7
2.5	Open questions in the 4D approach	10
3	8D point of view: dS from T-branes	11
3.1	T-brane background due to gauge fluxes	12
3.2	4D uplifting term	13
4	Conclusions	16
A	AdS KKLT vacua	18
B	D-term potential from D7-branes	19

1 Introduction

Chiral 4D string compactifications have several generic features, like extra dimensions, D-branes, background and gauge fluxes. In recent years global string models compatible with full closed string moduli stabilisation and a semi-realistic chiral visible sector have been constructed [1–4]. These models have to satisfy some global consistency constraints, like RR tadpole cancellation, Freed-Witten anomaly cancellation and discrete K-theory charge cancellation [5, 6]. Moreover the presence of chiral visible sector D-branes can reduce the effects that can be used to fix the moduli [7].

One of the brane configurations should host a visible sector that includes the Standard Model while other brane configurations give rise to hidden sectors. These hidden sectors are in general needed in order to guarantee the global consistency of the underlying compactification. Even though the coupling of these hidden sectors to the Standard Model is only gravitational, they may still play an important rôle to determine the physics of the visible sector by participating in the process of moduli stabilisation and by providing candidates for inflation, dark matter, dark radiation and dark energy.

The strongest global constraints come from the cancellation of D7- and D3-brane charges. The typical presence of O7-planes induces a D7-brane charge that has to be cancelled by some D7-branes which typically belong to the hidden sector, especially in

models where the visible sector is realised on D3-branes at singularities [2–4, 8, 9]. In order to cancel also the D3-brane charge in the presence of a large number of positively contributing three-form fluxes [10], the D7-branes should in general wrap a high degree divisor in the compact manifold. In Large Volume Scenario (LVS) models [11, 12], these cycles have very large volume. Moreover, in the models studied so far with small $h^{1,1}(\text{CY})$ [1–4], this divisor has another peculiar feature: the pull-back of the harmonic $(1, 1)$ -forms from the Calabi-Yau to the D7-brane divisor has a 1D image, implying that any two-form of this kind is proportional to the pull-back of the Kähler form. A gauge flux on such a D7-brane has a remarkable effect: it induces a never-vanishing (inside the Kähler cone) Fayet-Iliopoulos (FI) term in the 4D effective action obtained by compactifying the D7-brane worldvolume theory. An image of the pull-back with dimension one is a sufficient condition to have a non-vanishing FI-term. For more generic cases, the FI term is a linear combination of several Kähler moduli and might vanish inside the Kähler cone; other competing effects in the moduli stabilisation process could however fix the Kähler moduli such that the FI term is non-zero.

In order to have a supersymmetric configuration at leading order, the D-term potential must vanish inducing a non-zero VEV for some charged matter fields if the corresponding FI-term is also non-vanishing. This 4D result, that has been studied in concrete examples in [2–4], can also be derived from an 8D perspective: by solving the 8D equations of motion, a non-zero flux induces a non-zero VEV for the adjoint scalar Φ living on the D7-brane stack, so that $[\Phi, \Phi^\dagger] \neq 0$. This condition is solved by a T-brane background [13–15].

T-branes are particular brane configurations for which the non-Abelian Higgs field Φ describing D-brane deformations is not diagonalisable but takes an upper triangular form. This configuration generates a positive definite contribution to the scalar potential if non-supersymmetric three-form fluxes are switched on. The volume dependence of this new contribution can be computed explicitly, and so its influence on moduli stabilisation can be explicitly analysed. The presence of this uplifting terms allows for dS vacua by tuning background three-form fluxes.

The paper is organised as follows. In section 2 we present the 4D mechanism to generate a dS minimum from a hidden sector D7-brane. The flux on such a brane induces a moduli-dependent FI term in the effective field theory. Vanishing D-terms in turn induce a non-zero VEV for a hidden sector matter field that when substituted in the F-term scalar potential gives rise to a positive definite moduli-dependent term that plays the rôle of an uplifting term. The positivity of the F-term is understood as a soft mass term for the corresponding hidden sector matter field. We also study in detail the tuning of microscopic parameters needed to achieve a viable dS vacuum. In section 3 we present instead the main result of this article: we show that from a higher dimensional point of view the 4D mechanism can be understood in terms of the presence of a T-brane. We consider supersymmetry breaking imaginary self-dual (ISD) three-form and gauge field fluxes in (hidden sector) D7-branes. The gauge fluxes lead to a T-brane configuration. We expand the D7 action and find a positive definite contribution to the scalar potential that precisely reproduces the uplifting term found from the 4D effective field theory point of view. This provides a higher dimensional implementation of this mechanism. A concrete example is based on the

LVS scenario of moduli stabilisation in which three-form fluxes break supersymmetry. We present the equivalent analysis for the KKLT case [16] in appendix A where the vacua, as expected, are only AdS. Finally in appendix B we briefly review the derivation of the 4D D-term potential from the D7 brane action summarising some useful conventions used in the literature.

2 4D point of view: dS from hidden matter F-terms

In this section we review a mechanism for obtaining dS vacua based on F-terms of hidden sector matter fields which are non-zero due to D-term stabilisation.

2.1 Hidden D7-branes and gauge fluxes

Hidden sector D7-branes are a generic feature of globally consistent compact Calabi-Yau (CY) models because of D7-tadpole cancellation [1–4]. In these backgrounds, the orientifold involution typically generates O7-plane fixed points. These objects have a RR D7-brane charge that is measured by $-8[O7]$, where $[O7]$ is the homology class of the four-cycle wrapped by the O7-plane. The simplest way to cancel this charge is to place four D7-branes (plus their images) on top of the O7-plane locus. The resulting D7-worldvolume gauge group is $SO(8)$. Here, without affecting our final results, we consider a simpler situation: we recombine the four D7-branes into a D7-brane wrapping the invariant divisor D_h in the homology class $4[O7]$ (where h stays for ‘hidden sector’). Its image will wrap the same divisor, creating an $SU(2) \cong USp(2)$ stack (the diagonal $U(1)$ is projected out by the orientifold action on the Chan-Paton factors). These branes can support a flux \mathcal{F} along the Cartan generator of $SU(2)$:¹

$$\mathcal{F} = (2\pi\alpha')F - \iota^*B, \tag{2.1}$$

where ι^*B is the pullback of the NS-NS B -field on D_h . In order to cancel the Freed-Witten anomaly [17, 18], the gauge flux $(2\pi\alpha')F$ has to satisfy the following quantisation condition:

$$(2\pi\alpha')F + \frac{c_1(D_h)}{2} \in H^2(D_h, \mathbb{Z}). \tag{2.2}$$

In the chosen configuration D_h is an even cycle, and so $c_1(D_h)$ is an even integral two-form (in a CY $c_1(D) = -[\hat{D}]$). This means that the gauge flux is integrally quantised and in principle can be set to zero (for a different situation with non-spin D7-brane divisor, i.e. $c_1(D_h)$ odd, the gauge flux cannot vanish because of (2.2)). On the other hand, \mathcal{F} may be forced to be non-zero by the presence of a half-integrally quantised B -field. This is typically the case for compactifications with non-perturbative moduli stabilisation.² Therefore \mathcal{F}

¹As the two divisors are in the same homology class, the flux on one brane is equal in absolute value to the flux on the other brane but with opposite sign.

²A non-perturbative superpotential W_{np} can be generated by D-branes wrapping four-cycles D_{np} that are typically non-spin. In order to have a non-zero W_{np} , the flux \mathcal{F}_{np} on these branes should vanish. If F_{np} is half-integral, the B -field must be half-integral as well to have $\mathcal{F}_{np} = 0$. Hence the B -field is in general chosen in such a way to cancel the flux \mathcal{F}_{np} on $O(1)$ E3-instantons wrapped around different divisors since they contribute to W_{np} only if they are invariant under the orientifold involution. Another option is to consider rank-2 E3-instantons and set $B = 0$ [19]. This often produces a too large volume of the compact manifold in the LVS scenario (as discussed in [4]).

is generically non-zero, causing the SU(2) gauge group to break down to U(1), where the U(1) group is anomalous. This anomaly gets cancelled by the Green-Schwarz mechanism and the U(1) gauge boson becomes massive by eating up a combination of an open and a closed string axion.

The closed string modulus T_j whose real part parameterises the Einstein-frame volume of the divisor D_j in units of $\ell_s = 2\pi\sqrt{\alpha'}$ gets a flux-dependent U(1)-charge of the form:

$$q_{hj} = \frac{1}{\ell_s^4} \int_{D_h} \hat{D}_j \wedge \mathcal{F}, \tag{2.3}$$

where \hat{D}_h is the two-form Poincaré dual to the divisor D_h (here we take the extra-dimensional coordinates dimensionful). Moreover the gauge flux \mathcal{F} yields a moduli-dependent Fayet-Iliopoulos (FI) term which looks like (see appendix B for more details):

$$\frac{\xi_h}{M_{\text{P}}^2} = \frac{e^{-\phi/2}}{4\pi\mathcal{V}} \frac{1}{\ell_s^4} \int_{D_h} J \wedge \mathcal{F} = \frac{1}{2\pi} \sum_j \frac{q_{hj}}{2} \frac{t_j}{\mathcal{V}} = -\frac{1}{2\pi} \sum_j q_{hj} \frac{\partial K}{\partial T_j}, \tag{2.4}$$

where ϕ is the dilaton ($e^{\langle\phi\rangle} = g_s$), \mathcal{V} is the Einstein-frame CY volume in units of $\ell_s = M_s^{-1}$ where the string scale is related to the Planck scale as $M_s = g_s^{1/4} M_{\text{P}}/\sqrt{4\pi\mathcal{V}}$, t_j are two-cycle moduli, $J = t_j \hat{D}_j$ is the Kähler form expanded in a basis of (1,1)-forms \hat{D}_j and $K/M_{\text{P}}^2 = -2\ln\mathcal{V}$ is the tree-level Kähler potential for the T -moduli.

A non-zero \mathcal{F}_h induces also D7 matter fields $\phi_j = |\phi_j| e^{i\theta_j}$ charged under the anomalous U(1). These are in the symmetric representation of the U(1) group, i.e. their charges are $q_{\phi_j} = \pm 2$. States with charges of both signs are typically generated for D7-branes wrapping large degree divisors, like in the cases under study.³ In this brane setup (brane and image-brane on top of a divisor D_h in the same class as $4D_{O7}$, with D_{O7} the divisor wrapped by the O7-plane), the number of chiral fields with charge +2 is [21]:⁴

$$I_{U(1)} = \frac{1}{2\ell_s^2} \left(\int_{D_h \cap D_h} (\mathcal{F} - \mathcal{F}') - 2 \int_{D_{O7} \cap D_h} \mathcal{F} \right) = \frac{12}{\ell_s^2} \int_{D_{O7} \cap D_{O7}} \mathcal{F}. \tag{2.5}$$

There are also neutral chiral fields which are counted by $h^{0,2}(D_h)$ and parameterise deformations of D_h . The resulting D-term potential takes the form [22–24]:

$$V_D = \frac{1}{2\text{Re}(f_h)} \left(\sum_j q_{\phi_j} \phi_j \frac{\partial K}{\partial \phi_j} + \frac{M_{\text{P}}^2}{2\pi} \sum_j q_{hj} \frac{\partial K}{\partial T_j} \right)^2 = \frac{\pi}{\text{Re}(T_h)} \left(\sum_j q_{\phi_j} \frac{|\phi_j|^2}{s} - \xi_h \right)^2, \tag{2.6}$$

since $f_h = T_h/(2\pi)$ and the Kähler metric \tilde{K} for charged D7 matter fields depends just on the real part of the axio-dilaton $\text{Re}(S) = s$ [25, 26]:

$$K \supset \tilde{K} \sum_j |\phi_j|^2 = \sum_j \frac{|\phi_j|^2}{s}. \tag{2.7}$$

³See [20] for concrete computations in analogous situations.

⁴Notice that the form (2.5) makes the chiral index manifestly invariant under the orientifold involution ($\mathcal{F} \mapsto -\mathcal{F}$ and the charge changes sign). It can alternatively be written as $I_{U(1)} \propto \int_{D_{O7} \cap D_{O7}} (F - F')$. Remember that the B -field takes part in defining what F' is.

2.2 Uplifting term from hidden matter F-terms

Writing the Kähler moduli as $T_i = \tau_i + i\psi_i$, the vanishing D-term condition reads:

$$\sum_j \frac{q_{\phi_j}}{s} |\phi_j|^2 = \xi(\tau_i). \tag{2.8}$$

When $\xi(\tau_i)$ is non-zero,⁵ the equation (2.8) forces at least one chiral field ϕ with positive charge to get a non-zero vev (assuming that $\xi > 0$; when ξ has opposite sign, the roles of positive and negative charge states are exchanged). If there were only modes with negative charge (pure chiral spectrum) this equation would not allow a supersymmetric configuration and in particular it would set $\langle \phi_j \rangle = 0 \forall j$ [35]. On the other hand, when the D7-brane is wrapping a large cycle (like in the models we are considering), there will be modes with charge of both signs.

The relation (2.8) fixes the combination of $|\phi_j|$ and τ_i corresponding to the combination of θ_j and ψ_i eaten up by the anomalous U(1), which acquires a mass of the form [24]:

$$M_{U(1)}^2 \simeq \frac{M_{\text{P}}^2}{\text{Re}(f_{D7})} (f_{\theta}^2 + f_{\psi}^2), \tag{2.9}$$

where the open and the closed string modes are charged under the anomalous U(1). Here for simplicity we are considering the case when only one open string modulus ϕ gets a non-zero vev⁶

The two terms in (2.9) are proportional respectively to the open and closed string axion decay constants f_{θ} and f_{ψ} which are given by:

$$f_{\theta}^2 = |\phi|^2 \simeq \xi \simeq \frac{\partial K}{\partial \tau} M_{\text{P}}^2 \quad \text{and} \quad f_{\psi}^2 \simeq \frac{\partial^2 K}{\partial \tau^2} M_{\text{P}}^2 \simeq \xi^2. \tag{2.10}$$

Since $\xi/M_{\text{P}}^2 \simeq t/\mathcal{V} \simeq 1/\tau \ll 1$ for cycles in the geometric regime, we realise that $f_{\theta} \gg f_{\psi}$, and so the combination of axions that is eaten up is mainly given by θ , which is then taking a large mass. Since at this level of approximation we are stabilising the moduli supersymmetrically, this tells us which combination of moduli (partner of the axions) is taking a large mass: the relation (2.8) fixes a combination that is mainly given by $|\phi|$ leaving a flat direction that is mostly given by τ . Consequently the mass of the anomalous U(1) is of order the Kaluza-Klein scale

$$M_{U(1)} \simeq \frac{f_{\theta}}{\tau^{1/2}} M_{\text{P}} \simeq \frac{M_{\text{P}}}{\mathcal{V}^{2/3}} \simeq \frac{M_s}{\mathcal{V}^{1/6}} \simeq \frac{1}{\text{Vol}^{1/6}}, \tag{2.11}$$

since via dimensional reduction the Planck scale M_{P} is related to the string scale M_s as $M_{\text{P}} \simeq M_s \mathcal{V}^{1/2}$. By substituting the VEV

$$\frac{|\phi|^2}{s} = \frac{\xi}{q_{\phi}} = \frac{q_T}{q_{\phi}} \frac{t}{4\pi\mathcal{V}} M_{\text{P}}^2, \tag{2.12}$$

⁵This happens in models we want to consider, i.e. a D7-brane wrapping a large cycle with one-dimensional second homology group: when the volume of the CY is non-zero, the FI-term on such brane does not vanish as soon as the flux is different from zero.

⁶At this level of approximation all the other combinations (leaving $\sum_j q_{\phi_j} |\phi_j|^2$ invariant) are flat directions. However these are fixed by positive definite soft mass terms at subleading order in the $1/\mathcal{V}$ expansion (this happens in the considered setup, see (2.14)).

into the F-term scalar potential for the matter field ϕ , one obtains a moduli-dependent positive definite contribution to the total scalar potential which can be used as an uplifting term. In fact, the main contribution to the F-term potential for ϕ comes from supersymmetry breaking effects which generate scalar soft masses of the form

$$m_0^2 = m_{3/2}^2 - F^I F^{\bar{J}} \partial_I \partial_{\bar{J}} \ln \tilde{K}, \tag{2.13}$$

where $m_{3/2} = e^{K/2} |W|$ is the gravitino mass expressed in terms of the 4D Kähler potential K and superpotential W . Given that the Kähler metric for D7 matter fields \tilde{K} depends just on the axio-dilaton S which is fixed supersymmetrically by turning on three-form background fluxes H_3 and F_3 , i.e. $F^S = 0$, the soft masses for D7-matter fields from (2.13) are simply given by:

$$m_0^2 = m_{3/2}^2 > 0. \tag{2.14}$$

This important relation ensures that the F-term scalar potential for the canonically normalised hidden matter field $\varphi = s^{-1/2} \phi$ is positive definite and can indeed play the rôle of the uplifting term:

$$V_{\text{up}} = m_0^2 |\phi|^2 = m_{3/2}^2 |\phi|^2 = \frac{c_{\text{up}}}{\mathcal{V}^{8/3}} M_{\text{P}}^4, \tag{2.15}$$

where (writing $6\mathcal{V} = kt^3$)

$$c_{\text{up}} = e^{K_{\text{cs}}} \left(\frac{6}{k}\right)^{1/3} \frac{q_T}{q_\phi} \frac{|W|^2}{8\pi s} > 0. \tag{2.16}$$

Note that the three-form fluxes H_3 and F_3 , which fix the dilaton S as well as the complex structure moduli, are also responsible for breaking supersymmetry by inducing non-zero F-terms for the Kähler moduli T .

2.3 Gauge fluxes and non-perturbative effects

Let us show how the uplifting term (2.15) can be successfully combined with the effects which fix the Kähler moduli T to obtain in the end a viable dS vacuum. A crucial ingredient to freeze the T -moduli is the presence of a non-perturbative superpotential W_{np} generated by either gaugino condensation on D7-branes or E3-instantons.

Note that a non-vanishing gauge flux \mathcal{F} on D_{h} might induce chiral states between the hidden sector D7-stack and any E3-instanton (the situation is very similar for the gaugino condensation case) which transform in the fundamental representation of the hidden SU(2) gauge group. If the E3-instanton wraps the divisor D_{E3} and carries no gauge flux, the number of these E3 zero-modes is given by [27–30]:⁷

$$I_{D7-E3} = \frac{1}{\ell_s^2} \int_{D_{\text{h}} \cap D_{E3}} \mathcal{F}. \tag{2.17}$$

If present, these charged E3 zero-modes can prevent the contribution of the E3-instanton to the superpotential if their VEV is zero, since they appear as prefactors in W_{np} [5, 7].⁸ Let

⁷The index is actually proportional to $\int_{D_{\text{h}} \cap D_{E3}} F - F_{E3}$, but since $\mathcal{F}_{E3} = 0$ then $2\pi\alpha' F_{E3} = B$ and $2\pi\alpha'(F - F_{E3}) = \mathcal{F}$.

⁸If the E3-instanton wraps the same four-cycle wrapped by the magnetised D7-brane, i.e. $D_{E3} = D_{\text{h}}$, the presence of charged E3 zero-modes is almost unavoidable.

us consider the simplest case with just one charged matter field ϕ and a Kähler modulus T with the following U(1) transformations:

$$\delta\phi = iq_\phi\bar{\phi} \quad \text{and} \quad \delta T = i\frac{q_T}{2\pi}. \quad (2.18)$$

The corresponding non-perturbative superpotential takes the form (setting from now on $M_{\text{P}} = 1$)

$$W_{\text{np}} = A\phi^n e^{-\frac{2\pi}{N}T}, \quad (2.19)$$

and its U(1) transformation looks like

$$\delta W_{\text{np}} = W_{\text{np}} \left(n\frac{\delta\phi}{\phi} - \frac{2\pi}{N}\delta T \right) = iW_{\text{np}} \left(nq_\phi - \frac{q_T}{N} \right). \quad (2.20)$$

Hence W_{np} is U(1)-invariant only if

$$nq_\phi = \frac{q_T}{N}. \quad (2.21)$$

This condition can be seen as a constraint that removes the tuning freedom to obtain a dS vacuum in models where T is fixed by non-perturbative effects as in the KKLT case [16]. On the other hand, if T is fixed via perturbative corrections like the volume modulus in LVS models, the condition (2.21) does not impose any restriction on the possibility to obtain dS vacua. The problem to uplift KKLT-like vacua using F-terms of hidden matter fields which are non-zero due to D-term stabilisation, can be seen also by noting that (2.21) can be rewritten as

$$q_\phi \frac{\partial W}{\partial \phi} \phi + \frac{q_T}{2\pi} \frac{\partial W}{\partial T} = 0. \quad (2.22)$$

Recalling that $D_I W = W_I + W K_I$ and the expression (2.6) for the D-term potential $V_D = D^2 / (2\text{Re}(f_i))$, (2.22) becomes

$$q_\phi \phi \frac{D_\phi W}{W} + \frac{q_T}{2\pi} \frac{D_T W}{W} = \left(q_\phi \phi \frac{\partial K}{\partial \phi} + \frac{q_T}{2\pi} \frac{\partial K}{\partial T} \right) = D, \quad (2.23)$$

showing how the D-term is proportional to a combination of F-terms. Hence vanishing F-terms necessarily imply a vanishing D-term. KKLT vacua are characterised by the fact that $D_T W = 0$, and so (2.23) with $D = 0$ implies $D_\phi W = 0$, showing the impossibility to obtain a dS vacuum following this mechanism. On the other hand, if $D_T W \neq 0$ as in LVS models [11, 12], $D_\phi W$ can also be non-vanishing giving a viable uplifting term even if $D = 0$ at leading order. We shall discuss the LVS case in some more detail in the next subsection and describe the KKLT case in appendix A.

2.4 dS LVS vacua

Let us now consider LVS vacua [11, 12] with magnetised hidden sector D7-branes and E3-instantons where K and W take the form:

$$K = -2 \ln \left(\mathcal{V} + \frac{\zeta s^{3/2}}{2} \right) + c \frac{\phi \bar{\phi}}{s} \quad \text{and} \quad W = W_0 + A_s e^{-a_s T_s}, \quad (2.24)$$

where $\zeta = -\frac{\chi(CY)\zeta(3)}{2(2\pi)^3}$ is a constant controlling the leading order α' correction and the CY volume depends on the big cycle τ_b and the small cycle τ_s as:

$$\mathcal{V} = \lambda_b \tau_b^{3/2} - \lambda_s \tau_s^{3/2}, \tag{2.25}$$

where λ_b and λ_s depend on the triple intersection numbers. The crucial difference with the KKLT case is that only T_s is fixed by non-perturbative effects while T_b is stabilised thanks to perturbative α' corrections to K leading to a supersymmetry breaking AdS vacuum for $c = 0$. Hence we can perturb this vacuum with the inclusion of magnetised branes and hidden matter fields which can lead to a dS solution.

In order to generate a non-vanishing T_s -dependent contribution to W , the B -field is generically chosen to cancel the FW anomaly on the small cycle T_s . This leads to a non-vanishing gauge flux on the stack of hidden sector D7-branes wrapping the big cycle T_b [2]. Thus T_b acquires a non-zero U(1)-charge q_b generating a moduli-dependent FI-term. The corresponding D-term potential reads:

$$V_D = \frac{\pi}{\tau_b} \left(\frac{q_\phi}{s} |\phi|^2 - \xi_b \right)^2, \tag{2.26}$$

where the FI-term takes the form

$$\xi_b = -\frac{q_b}{2\pi} \frac{\partial K}{\partial T_b} = \frac{3q_b}{4\pi\tau_b}. \tag{2.27}$$

Therefore the total scalar potential is

$$V_{\text{tot}} = V_D + V_F = \frac{\pi}{\tau_b} \left(\frac{q_\phi}{s} |\phi|^2 - \frac{3q_b}{4\pi\tau_b} \right)^2 + \frac{1}{s} m_{3/2}^2 |\phi|^2 + \frac{V_F(T)}{2s}, \tag{2.28}$$

where $m_{3/2}^2 = e^K |W| \simeq W_0^2 / (2s\mathcal{V}^2)$ is the gravitino mass and $V_F(T)$ is given by [11, 12]:⁹

$$V_F(T) = \frac{8}{3\lambda_s} (a_s A_s)^2 \sqrt{\tau_s} \frac{e^{-2a_s \tau_s}}{\mathcal{V}} - 4a_s A_s W_0 \tau_s \frac{e^{-a_s \tau_s}}{\mathcal{V}^2} + \frac{3\zeta s^{3/2} W_0^2}{4\mathcal{V}^3}. \tag{2.29}$$

Note that the prefactor of the non-perturbative effect does not depend on the matter field ϕ since T_s does not get charged under the anomalous U(1). Minimising with respect to ϕ we find:

$$\frac{q_\phi}{s} |\phi|^2 = \xi_b - \frac{m_{3/2}^2 \tau_b}{2\pi q_\phi}. \tag{2.30}$$

Substituting this VEV back in (2.28) we obtain (writing $\tau_b = (\mathcal{V}/\lambda_b)^{2/3}$ and setting $s = g_s^{-1}$):

$$V_{\text{tot}} = \frac{g_s}{2} \left[c_{\text{up}} \frac{W_0^2}{\mathcal{V}^{8/3}} \left(1 - \frac{c_{\text{sub}}}{\mathcal{V}^{2/3}} \right) + V_F(T) \right], \tag{2.31}$$

⁹Notice that in LVS, for $W_0 \neq 0$, the AdS minimum is already non-supersymmetric. The flux induced SUSY breaking is captured in the effective field theory by a non-vanishing F-term for the large modulus field T_b which is proportional to W_0 . Other fields such as the dilaton and ϕ itself will also contribute to the breaking of supersymmetry but their F-terms appear at higher orders in the $1/\mathcal{V}$ expansion. The leading contribution to the goldstino comes from the fermionic component of T_b .

with

$$c_{\text{up}} = \frac{3q_b \lambda_b^{2/3}}{4\pi q_\phi} \quad \text{and} \quad c_{\text{sub}} = \frac{W_0^2 g_s}{6q_\phi q_b \lambda_b^{4/3}}. \quad (2.32)$$

The first term in (2.31) is the uplifting term (neglecting corrections proportional to c_{sub} which are subleading for $\mathcal{V} \gg 1$):

$$V_{\text{up}} = \frac{g_s c_{\text{up}}}{2} \frac{W_0^2}{\mathcal{V}^{8/3}}. \quad (2.33)$$

This term modifies the scalar potential such that it now admits a dS minimum. In fact, minimising (2.31) with respect to τ_s we obtain:

$$\mathcal{V} = \frac{3\lambda_s W_0 \sqrt{\tau_s}}{4a_s A_s} e^{a_s \tau_s} \frac{(1-\epsilon)}{(1-\epsilon/4)} \quad \text{with} \quad \epsilon \equiv \frac{1}{a_s \tau_s} \sim \mathcal{O}\left(\frac{1}{\ln \mathcal{V}}\right) \ll 1. \quad (2.34)$$

Substituting this result back in (2.31) we find an effective potential for the volume \mathcal{V} :

$$V_{\text{tot}}(\mathcal{V}) = \frac{g_s W_0^2}{2\mathcal{V}^3} \left[c_{\text{up}} \mathcal{V}^{1/3} - \frac{3\lambda_s}{2a_s^{3/2}} \epsilon^{-3/2} \left(1 - \left(\frac{3\epsilon}{4}\right)^2 \right) + \frac{3\zeta}{4g_s^{3/2}} \right], \quad (2.35)$$

where

$$\epsilon^{-1} = \ln\left(c_s \frac{\mathcal{V}}{W_0}\right) - \ln \sqrt{\tau_s} + \ln \frac{(1-\epsilon)}{(1-\epsilon/4)} = \ln\left(c_s \frac{\mathcal{V}}{W_0}\right) \left[1 - \frac{\epsilon}{2} \ln \tau_s + \mathcal{O}(\epsilon^2) \right], \quad (2.36)$$

with $c_s = 4a_s A_s / (3\lambda_s)$. Given that $\ln \tau_s \sim \mathcal{O}(1)$, (2.36) scales as $\epsilon^{-1} = \ln\left(c_s \frac{\mathcal{V}}{W_0}\right) (1 + \mathcal{O}(\epsilon))$, and so it can be rewritten iteratively as:

$$\epsilon^{-1} = \ln\left(c_s \frac{\mathcal{V}}{W_0}\right) - \frac{1}{2} \ln\left[\frac{1}{a_s} \ln\left(c_s \frac{\mathcal{V}}{W_0}\right)\right] + \mathcal{O}(\epsilon). \quad (2.37)$$

The uplifting term (2.35) does not have a large tuning freedom since, as can be seen from (2.32), c_{up} is an $\mathcal{O}(1)$ number which depends just on triple intersection numbers and gauge flux quanta. On the other hand, the second and the third term in (2.35) have a strong dependence on g_s and $a_s = 2\pi/N$ (where N is the rank of the condensing gauge group) while a milder logarithmic dependence on W_0 . Hence a vanishing vacuum energy can be obtained by tuning the depth of the original AdS minimum by a proper choice of background fluxes, i.e. of g_s and W_0 . In general, for $N = 1$ and large values of W_0 of order 100 the scalar potential has a runaway behaviour unless A_s is tuned to values of order 100 to compensate the large value of W_0 . In this case the potential has a dS minimum around $\mathcal{V} \sim 10^6$ if c_{up} is of order 0.01 and g_s is appropriately tuned so that the third term in (2.35) is of the same order of the first two. Larger values of N and smaller values of W_0 give rise to dS minima for larger values of \mathcal{V} .

Let us see the tuning of the vacuum energy more in detail by minimising the potential (2.35) with respect to \mathcal{V} that leads to:

$$\frac{3\zeta}{4g_s^{3/2}} = \frac{3\lambda_s}{2} \tau_s^{3/2} \left[1 - \frac{\epsilon}{2} + \mathcal{O}(\epsilon^2) \right] - \frac{8}{9} c_{\text{up}} \mathcal{V}^{1/3}, \quad (2.38)$$

which substituted in (2.35) gives a vacuum energy of the form

$$\langle V_{\text{tot}} \rangle = \frac{g_s W_0^2}{18 \mathcal{V}^3} \left[c_{\text{up}} \mathcal{V}^{1/3} - \frac{27\lambda_s}{4a_s^{3/2} \sqrt{\epsilon}} \left(1 - \frac{9}{8}\epsilon + \mathcal{O}(\epsilon^2) \right) \right], \quad (2.39)$$

where \mathcal{V} (and hence ϵ given in (2.37)) should be meant as a function of the underlying parameters (like W_0 and g_s) following the minimizing equation (2.38). Setting this expression equal to zero and plugging the result for c_{up} back in (2.38), we find:

$$\tau_s = \left(\frac{\zeta}{2\lambda_s} \right)^{2/3} \frac{1}{g_s} [1 + 3\epsilon + \mathcal{O}(\epsilon^2)]. \quad (2.40)$$

Comparing this result with the location of the AdS vacuum for $c_{\text{up}} = 0$ we find that the shift of the VEV of τ_s is proportional to ϵ :

$$\frac{\Delta\tau_s}{\tau_s|_{c_{\text{up}}=0}} = \frac{8\epsilon}{3} (1 + \mathcal{O}(\epsilon)). \quad (2.41)$$

Plugging this result in (2.34) the shift of the volume VEV is instead of order

$$\frac{\Delta\mathcal{V}}{\mathcal{V}|_{c_{\text{up}}=0}} = e^{8/3} (1 + \mathcal{O}(\epsilon)) - 1 \simeq 14.4 (1 + \mathcal{O}(\epsilon)) - 1, \quad (2.42)$$

showing that for $\epsilon \ll 1$, i.e. in the regime where higher order instanton contributions to W can be safely neglected, the new dS vacuum is at values of the volume which are about one order of magnitude larger than those of the old AdS vacuum.

We conclude this section by showing in figure 1 how the position of a leading order Minkowski minimum changes as a function of g_s for three values of N , for fixed $W_0 = 1$ and natural values of the underlying parameters, while in figure 2 we present the same behaviour for fixed $N = 6$ and different values of W_0 . To do this, we plot what is inside the square bracket in eq. (2.39) and we call it V_0 (i.e. $V_0 \equiv \frac{18\mathcal{V}^3}{g_s W_0^2} \langle V_{\text{tot}} \rangle$).

2.5 Open questions in the 4D approach

The uplifting mechanism described above from the 4D effective point of view is based on two crucial ingredients, gauge and background fluxes, which induce non-zero hidden sector F-terms via D-term stabilisation. This mechanism for obtaining dS vacua seems to be rather generic and elegant but it raises some open questions which we list below:

- Is it consistent to include an anomalous U(1) in the 4D effective field theory if it acquires a mass of order the Kaluza-Klein scale?
- What is the correct higher-dimensional understanding of this uplifting mechanism? Are we actually expanding the effective theory around a solution which at leading order is supersymmetric?
- Does a non-zero VEV for ϕ correspond to a brane/image-brane recombination (or better a bound-state) that makes the whole procedure inconsistent since the flux $-\mathcal{F}$ on the image brane would cancel off with the flux \mathcal{F} on the brane? This would mean that we have expanded around a supersymmetry breaking solution which might not be under control.




Figure 1. Vacuum energy (V_0) as a function of g_s for $\lambda_s = \lambda_b = W_0 = A_s = 1$, $q_\phi = 2q_b$, $\zeta = 2$ and $a_s = 2\pi/N$. The intersection of the three curves with the abscissa shows the value of g_s that gives a minimum where at leading order $\langle V \rangle = 0$ for $N = 2$ with $\mathcal{V} \simeq 7.5 \cdot 10^7$ (green), $N = 6$ with $\mathcal{V} \simeq 6.5 \cdot 10^6$ (blue) and $N = 10$ with $\mathcal{V} \simeq 3 \cdot 10^4$ (red). Notice that we have plot $V_0 \equiv \frac{18\mathcal{V}^3}{g_s W_0^2} \langle V_{\text{tot}} \rangle$ instead of $\langle V_{\text{tot}} \rangle$, as we are interested in the zeros.




Figure 2. Vacuum energy (V_0) as a function of g_s for $\lambda_s = \lambda_b = A_s = 1$, $q_\phi = 2q_b$, $\zeta = 2$ and $a_s = 2\pi/6$. The intersection of the three curves with the abscissa shows the value of g_s that gives a minimum where at leading order $\langle V \rangle = 0$ for $W_0 = 1$ with $\mathcal{V} \simeq 6.5 \cdot 10^6$ (blue), $W_0 = 10^{-5}$ with $\mathcal{V} \simeq 1.8 \cdot 10^7$ (green) and $W_0 = 10^{-10}$ with $\mathcal{V} \simeq 3.5 \cdot 10^7$ (red).

These questions can be clearly answered only by developing a higher-dimensional understanding which we present in the following section.

3 8D point of view: dS from T-branes

In this section we present an 8D description of the 4D dS vacua presented in section 2. The key-ingredient of this higher-dimensional understanding is the presence of T-branes induced by gauge fluxes.

3.1 T-brane background due to gauge fluxes

We consider a stack of N D7-branes wrapping a compact four-cycle D_h in the CY three-fold X_3 . Since D_h must be holomorphically embedded in X_3 , it has an inherited Kähler form and it is a Kähler manifold. When the closed string fields have vanishing VEVs, the 8D theory living on their worldvolume is a twisted $\mathcal{N} = 1$ SYM theory [31]. The field content is made up of an 8D vector and a complex scalar Φ in the adjoint representation of the gauge group (in this case $U(N)$).

The compactification on D_h breaks the 8D Poincaré invariance and breaks the $\mathcal{N} = 1$ 8D supersymmetry. Thanks to the topological twist, four supercharges are still preserved. Hence the fields organise themselves in 4D $\mathcal{N} = 1$ supermultiplets. We have a 4D vector multiplet with gauge field A_μ , and two chiral multiplets with complex scalars A and Φ in the adjoint representation. Due to the twist, the scalar Φ transforms in the canonical bundle of D_h like a $(2, 0)$ -form, while A is a $(0, 1)$ -form on D_h . The 8D supersymmetric equations of motion also split according to the breaking of the 8D Poincaré group. Along the compact direction they are given by:

$$\mathcal{F}^{0,2} = \mathcal{F}^{2,0} = 0, \quad \bar{\partial}_A \Phi = \partial_A \Phi^\dagger = 0, \quad (3.1)$$

where \mathcal{F} is the curvature of the gauge bundle along D_h and ∂_A is the holomorphic covariant differential relative to its connection and

$$J_{(h)} \wedge \mathcal{F} - [\Phi, \Phi^\dagger] = 0, \quad (3.2)$$

where J is the Kähler form on D_h (in this case it is the pullback of the Kähler form of X_3 , i.e. $J_{(h)} = \iota^* J$). Note that this relation is valid for a canonically normalised 8D scalar field. In 4D these equations boil down to F- and D-term conditions respectively. The second one will receive perturbative corrections. However in the cases we shall consider, the volume of D_h is very large and these corrections are negligible.

The structure group of the gauge bundle on D_h breaks the 4D gauge group to its commutant. As an illustrative example, we consider a stack of two D7-branes with gauge group $U(2)$ and we choose \mathcal{F} to be an Abelian flux along the Cartan of $SU(2)$:

$$\mathcal{F} = f \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (3.3)$$

where f is a $(1, 1)$ -form on D_h . We will consider the case when f is the (non-trivial) pull-back of a two-form of X_3 . This solves automatically the F-term eq. (3.1).

Regarding the D-term eq. (3.2), the flux \mathcal{F} and the Kähler form $J_{(h)}$ determine if Φ can be zero or not. In particular, for simplicity (but without affecting our final results) we will restrict to the case when the pull-back map from two-forms on X_3 to two-forms on D_h has a 1D image (typical of the CY Swiss-cheese explicit models of [2–4]) generated by the $(1, 1)$ -form ω on D_h . This implies that for any combination $a_j \hat{D}_j$ its pull-back is $\iota^* a_j \hat{D}_j = a \omega$. Then, $J_{(h)} = t \omega$ (where t is a linear combination of the coefficient of the CY Kähler form $J = t_i \hat{D}_i$), $f = n \omega$ and

$$J_{(h)} \wedge f = (tn) \omega \wedge \omega. \quad (3.4)$$

This is different from zero for non-zero volume of D_h (in our cases this always happens if we want to keep the volume of X_3 different from zero as well, since $\text{vol}D_h \propto t \sim \tau_b$).¹⁰ Hence (3.2) implies that the commutator $[\Phi, \Phi^\dagger]$ is non-zero. The simplest solution is given by the nilpotent matrix:

$$\Phi = \begin{pmatrix} 0 & \varphi \\ 0 & 0 \end{pmatrix}, \tag{3.5}$$

that solves (3.2) if $\varphi \wedge \bar{\varphi} = tn \omega \wedge \omega$. Recall that φ is a $(2,0)$ -form. On a local patch of D_h we have $\varphi = \hat{\varphi} dz_1 \wedge dz_2$, where $\hat{\varphi}$ is a scalar field and z_1, z_2 are complex local coordinates. The solution (3.5) is called a *T-brane* (because of the triangular form of the matrix) [13–15]. This VEV for Φ does not change the D7-brane locus but anyway breaks the gauge group to the diagonal $U(1)$ of the original $U(2)$. This is interpreted as a bound state of the two branes, even though the locus is made up of two copies of D_h . In particular the broken (Cartan) $U(1)$ disappears from the low energy spectrum, where no FI term will appear.

3.2 4D uplifting term

As we will claim later, the solution described in the previous section is the leading piece of the solution in the presence of supersymmetry breaking ISD three-form fluxes for large volumes of D_h and X_3 . We can then plug this into the 8D action of the D7-branes and expand around this solution. When the supersymmetry breaking three-form fluxes are equal to zero, no dS uplift term is generated (by consistency). In fact the scalar potential at tree level is positive definite or identically zero (depending if we switch on supersymmetric fluxes).

To see how the situation is modified by the presence of three-form fluxes, we need to compute the coupling of the relevant closed string p -form potential to the D7-worldvolume. Eventually we switch on a T-brane background and see how this interplays with three-form fluxes.

We start by reviewing what happens to a single D7-brane. Its 8D action is given by expanding the DBI and CS actions

$$S = -\mu_7 \text{STr} \left\{ \int d^8\xi e^{-\phi} \sqrt{-\det(\iota^* e^{\phi/2} G_{AB} + \mathcal{F}_{AB})} - \int (\iota^* C_6 \wedge \mathcal{F} + \iota^* C_8) \right\}, \tag{3.6}$$

where $\mu_7^{-1} = (2\pi)^3 (2\pi\alpha')^4$, ι^* is the pull-back map from the bulk to the D7-brane world-volume and $\mathcal{F} = (2\pi\alpha')F - \iota^*B$. We want to understand what is the coupling of the bulk fields to the open string scalar Φ . For a single D7-brane, this field controls the deformations of the D7-brane worldvolume in the orthogonal directions in the bulk. For a D7-brane we have one (complex) dimensional normal bundle. In local coordinates for an open patch of the CY three-fold X_3 , z^1 and z^2 are complex coordinates tangent to the D7-brane and z^3 is orthogonal to it. We then make the identification $z^3 = (2\pi\alpha')\Phi$ and expand the

¹⁰The expression $J_{(h)} \wedge f$ is always different from zero under the sufficient assumption that the pull-back has a 1D image on $H^2(D_h)$ and that D_h is a large cycle. However it is a quite generic feature of a background with Kähler moduli stabilised by several effects whenever they imply $J_{(h)}$ not orthogonal to the flux f .

closed string fields around $z^3 = 0$, keeping only the leading terms and making the given identification. We will follow [32, 33] and refer to them for more details.

The basic ingredients are three-form fluxes encoded into the complex three-form $G_3 = F_3 - \tau H_3$, where F_3 and H_3 are the field strengths of the (respectively) RR and NSNS two-form potentials and $\tau = e^{-\phi} + iC_0$ is the axio-dilaton. The three-form G_3 can be decomposed into imaginary selfdual (ISD) and anti-selfdual (IASD) pieces:

$$G_3^\pm = \frac{1}{2}(G_3 \mp i *_6 G_3), \quad *_6 G_3^\pm = \pm i G_3^\pm. \quad (3.7)$$

These components can be decomposed in irreducible pieces of the SU(3) structure group of the CY manifold. For example the ISD component in $\overline{\mathbf{10}}$ representation of SO(6) splits into $\overline{\mathbf{10}} = \overline{\mathbf{1}} \oplus \overline{\mathbf{3}} \oplus \overline{\mathbf{6}}$ where the singlet is the (0, 3) piece $G_{\overline{k}\overline{j}\overline{l}}$ and the antisymmetric and symmetric representations are

$$A_{\overline{i}\overline{j}} = \frac{1}{2}(\epsilon_{ikl} G_{\overline{k}\overline{l}\overline{j}} - \epsilon_{jkl} G_{\overline{k}\overline{l}\overline{i}}), \quad S_{\overline{i}\overline{j}} = \frac{1}{2}(\epsilon_{\overline{i}\overline{k}\overline{l}} G_{\overline{j}\overline{k}\overline{l}} + \epsilon_{\overline{j}\overline{k}\overline{l}} G_{\overline{i}\overline{k}\overline{l}}). \quad (3.8)$$

For the IASD $\mathbf{10}$ component the definitions are analogous (after taking the complex conjugate). Following [32, 33] we neglect the non-primitive components $A_{\overline{i}\overline{j}}$ and A_{ij} as they are incompatible with the CY topology (even though there can be a local component) and set to zero S_{3i} and $S_{\overline{3}\overline{i}}$ since these components would generate a FW anomaly on the worldvolume of the D7-brane (that happens when $\iota^* G_3$ is a non-trivial three-form on the D7-brane worldvolume).

We can now expand the axio-dilaton and the B -field in powers of Φ :

$$\tau = \frac{i}{g_s} \left[1 + \frac{\tau_{33}}{2} (2\pi\alpha')^2 \Phi^2 + \frac{\tau_{\overline{3}\overline{3}}}{2} (2\pi\alpha')^2 \overline{\Phi}^2 + (2\pi\alpha')^2 \tau_{3\overline{3}} |\Phi|^2 \right], \quad (3.9)$$

$$B_{12} = -i g_s \pi \alpha' \left((G_{\overline{1}\overline{2}\overline{3}})^* \Phi - \frac{1}{2} S_{\overline{3}\overline{3}} \overline{\Phi} - G_{123} \Phi + \frac{1}{2} (S_{33})^* \overline{\Phi} \right), \quad (3.10)$$

$$B_{\overline{1}\overline{2}} = -i \frac{g_s \pi \alpha'}{2} \left(-S_{\overline{2}\overline{2}} \Phi + (S_{\overline{1}\overline{1}})^* \overline{\Phi} - S_{11} \overline{\Phi} + (S_{22})^* \Phi \right), \quad (3.11)$$

where the B -field has been derived from $dB_2 = -\frac{\text{Im}G_3}{\text{Im}\tau}$ and $\tau_{3\overline{3}}$ can be computed from the 10D supergravity equation of motion $\tau_{3\overline{3}} = \frac{1}{2i}|G|^2$ (assuming that localised sources give a negligible contribution). Here and in the following we define (setting $A = 0$)

$$|G|^2 \equiv \frac{1}{12} g^{mm'} g^{nn'} g^{kk'} G_{3|mnk} G_{3|m'n'k'} = |G_{123}|^2 + \frac{1}{4} \sum_{k=1}^3 |S_{kk}|^2. \quad (3.12)$$

Finally, from

$$dC_6 = H_3 \wedge C_4 - * \text{Re}G_3 \quad \text{and} \quad dC_8 = H_3 \wedge C_6 - * \text{Red}\tau, \quad (3.13)$$

we obtain

$$C_{\mu\nu\rho\sigma 12} = -i\pi\alpha' \left((G_{\bar{1}\bar{2}\bar{3}})^* \Phi - G_{123} \Phi + \frac{1}{2} S_{\bar{3}\bar{3}} \bar{\Phi} + \frac{1}{2} (S_{33})^* \bar{\Phi} \right), \quad (3.14)$$

$$C_{\mu\nu\rho\sigma 1\bar{2}} = -i\frac{\pi\alpha'}{2} (S_{\bar{2}\bar{2}} \Phi + (S_{22})^* \Phi - (S_{\bar{1}\bar{1}})^* \bar{\Phi} - S_{11} \bar{\Phi}), \quad (3.15)$$

$$\begin{aligned} C_{\mu\nu\rho\sigma 1\bar{1}\bar{2}\bar{2}} &= -\frac{g_s \pi^2 \alpha'^2}{4} \{ [-2G_{123} S_{33} + (S_{11} S_{22})^* - S_{\bar{1}\bar{1}} S_{\bar{2}\bar{2}} + 2(G_{\bar{1}\bar{2}\bar{3}} S_{\bar{3}\bar{3}})^*] \Phi^2 + \text{h.c.} \\ &\quad + [|S_{33}|^2 - |S_{\bar{3}\bar{3}}|^2 - 4|G_{\bar{1}\bar{2}\bar{3}}|^2 + |S_{\bar{1}\bar{1}}|^2 + |S_{\bar{2}\bar{2}}|^2 + 4|G_{123}|^2 - |S_{22}|^2 - |S_{11}|^2] |\Phi|^2 \\ &\quad + ig_s \pi \alpha' [\tau_{33} + (\tau_{\bar{3}\bar{3}})^*] \Phi^2 + \text{h.c.} \}. \end{aligned} \quad (3.16)$$

We now plug these expressions in (3.6) and expand up to quadratic order, obtaining the following contribution to the 8D Lagrangian (we do not write the kinetic terms):

$$\begin{aligned} \mathcal{L} \supset & -2g_s |G|^2 |\Phi|^2 + \frac{g_s}{4} \left\{ (S_{\bar{3}\bar{3}})^* [(G_{\bar{1}\bar{2}\bar{3}})^* - G_{123}] + \frac{1}{2} (S_{11})^* [(S_{22})^* - S_{\bar{2}\bar{2}}] \right. \\ & \left. - (G_{\bar{1}\bar{2}\bar{3}})^* S_{33} - \frac{1}{2} (S_{22})^* S_{\bar{1}\bar{1}} - 2i\tau_{33} \right\} \Phi^2 + \text{h.c.} + [\Phi \partial A], \end{aligned} \quad (3.17)$$

where the terms in the last line are linear in Φ and in the derivatives of the connections.

This is the result for a single D7-brane. When we have several D7-branes on top of each other, each one will contribute with a term like (3.17). Moreover, the gauge group is enhanced from $U(1)^N$ to $U(N)$, due to the massless strings stretching between different branes. The gauge field A and the scalar field Φ live in the adjoint representation of $U(N)$. The terms in the Lagrangian that vanish for zero three-form fluxes are

$$\begin{aligned} \mathcal{L} \supset & -2g_s |G|^2 \text{Tr} |\Phi|^2 + \frac{g_s}{4} \left\{ (S_{\bar{3}\bar{3}})^* [(G_{\bar{1}\bar{2}\bar{3}})^* - G_{123}] + \frac{1}{2} (S_{11})^* [(S_{22})^* - S_{\bar{2}\bar{2}}] \right. \\ & \left. - (G_{\bar{1}\bar{2}\bar{3}})^* S_{33} - \frac{1}{2} (S_{22})^* S_{\bar{1}\bar{1}} - 2i\tau_{33} \right\} \text{Tr} \Phi^2 + \text{h.c.} + \text{Tr}(\Phi \partial A). \end{aligned} \quad (3.18)$$

We now want to insert in this Lagrangian the T-brane background (for simplicity we take $N = 2$ D7-branes in the stack). Since the matrix Φ is nilpotent and A is diagonal (it is along the Cartan), we have:

$$\text{Tr} \Phi^2 = 0 \quad \text{and} \quad \text{Tr}(\Phi \partial A + \Phi [A, A]) = 0. \quad (3.19)$$

The only contribution to the vacuum energy is then given by the first term in (3.18). Hence the term contributing to the dS uplift is

$$\mathcal{L}_{\text{up}} = 2g_s |G|^2 \varphi \wedge \bar{\varphi}. \quad (3.20)$$

Let us see what is the volume dependence of this term. First of all, consider the three-form fluxes. Since they are quantised according to $\frac{1}{(2\pi)^2 \alpha'} \int_{\Sigma_3} G_3 = n_F - \tau n_H$ (with $n_F, n_H \in \mathbb{Z}$), the field strength has to go like (using $M_s = \ell_s^{-1}$)

$$G_{mnp} \sim (n_F - \tau n_H) M_s \sim (n_F - \tau n_H) \frac{M_{\text{P}}}{\mathcal{V}^{1/2}}. \quad (3.21)$$

Hence,

$$|G|^2 \sim (g^{-1})^3 M_s^2 \sim \frac{1}{\mathcal{V}} \frac{M_{\text{P}}^2}{\mathcal{V}} = \frac{M_{\text{P}}^2}{\mathcal{V}^2}. \quad (3.22)$$

Finally, from the T-brane equation of motion (3.2) and assuming that the D7-brane wraps a large cycle (whose volume is $\frac{1}{2} \int_{D_{\text{h}}} J_{(\text{h})}^2 \simeq t^2 \simeq \tau_b \simeq \mathcal{V}^{2/3}$), we have

$$|\varphi|^2 \sim \mathcal{V}^{1/3} M_s^2 \sim \frac{M_{\text{P}}^2}{\mathcal{V}^{2/3}}. \quad (3.23)$$

Hence the uplift term in the 4D potential takes the form:¹¹

$$V_{\text{up}} = \frac{C_{\text{up}}}{\mathcal{V}^{8/3}} M_{\text{P}}^4, \quad (3.24)$$

where $C_{\text{up}} \sim 2g_s n \int_{D_{\text{h}}} |\hat{G}|^2 \omega \wedge \omega$ is a constant that depends just on three-form and gauge fluxes (the dilaton is also a flux dependent function after stabilisation) since we set $\hat{G} = \frac{G}{M_{\text{P}}} \mathcal{V}$. This term is *positive definite* and is therefore suitable to uplift an AdS LVS vacuum to a dS solution. Note that (3.24) has the same volume scaling as the uplifting term (2.33) which we found using the 4D effective action.

Ref. [33] also studied corrections to the soft terms due to a non-zero gauge flux on the D7-brane worldvolume. In principle we should consider this effect as well, as the T-brane solution contains a non-zero flux. However, the extra contribution given by the presence of non-zero gauge fluxes is suppressed with respect to (3.24). In fact, following [33], we find a correction of the form $|G|^2 |\Phi|^2 |\mathcal{F}|^2 M_s^{-4}$. The factor $|\mathcal{F}|^2 M_s^{-4}$ goes like $\mathcal{V}^{-2/3}$ (due to two powers of the inverse metric) and provides a suppression factor of order $\mathcal{V}^{-2/3}$ for the flux correction to the uplift potential.¹²

One could also question the procedure of expanding around the T-brane solution when three-form fluxes are switched on since they modify the equations of motion for Φ . However a careful analysis reveals that the new solution is only a small perturbation around the used T-brane background and that this perturbation does not affect the uplift term.¹³

4 Conclusions

In this paper we have shown how a dS minimum can arise in type IIB CY compactifications if a T-brane is present among the hidden sector D7-branes.

We have first reviewed the 4D effective field theory picture that has been studied in [2–4]: the dS uplift term is generated by a fluxed D7-brane in the hidden sector. The flux on the D7-brane induces an FI-term. When this is different from zero, the D-term potential forces

¹¹No factor $\mathcal{V}^{2/3}$ appears after integrating over D_{h} . This is because we are taking a volume one surface and treating the volume modulus as a parameter. The canonical normalised 8D scalar that appears in (3.2) has already absorbed the factor coming from this modulus.

¹²As we have a non-abelian stack, one should have in principle included terms from the Myers action, proportional to $\text{Tr} i_{\Phi} C$ and $\text{Tr} i_{\Phi} i_{\Phi} C$. For the solution (3.5) these are zero, as $\text{Tr} \Phi = 0$ and $\text{Tr} [\text{Re} \Phi, \text{Im} \Phi] = 0$.

¹³Here we have considered the expansion of the 8D action in Φ up to quadratic order. This reproduces the controlled expansion in ϕ in the 4D approach, where it is more manifest that the higher order terms are suppressed by powers of the volume \mathcal{V} .

some charged hidden sector modes ϕ to get a non-zero VEV. When three-form fluxes are switched on, supersymmetry is broken and a positive soft supersymmetry-breaking mass-squared is produced for ϕ . Since the VEV of ϕ is different from zero, due to the D-term condition, this mass term produces a constant positive definite term that can be used to achieve dS vacua. We have explained how this term can lead to a dS minimum by tuning W_0 and g_s .

This description can raise some concerns. First of all, in the effective field theory we have a non-vanishing FI-term. This means that the fluxed D7-brane is not supersymmetric. In fact, from the 4D point of view we see that it is the VEV of ϕ that restores supersymmetry (at the leading order). The field ϕ describes deformations of the D7-brane. The effective theory we have used is defined for the D7-brane on a given locus and corresponds to $\phi = 0$. On the other hand, this effective theory is telling us that the VEV of ϕ is non-zero. This means that the right minimum is not the configuration around which we are expanding. Moreover, note that to cancel the FI term it is enough that only one charged hidden field takes a non-vanishing VEV. In particular one could wonder whether open string modes with opposite charge could get non-zero vev: in our case, where all the soft mass terms are equal to $m_{3/2}^2 > 0$ (see (2.14)) this does not happen, as it would not lead to a minimum of the full scalar potential. In other setups, with a different susy breaking pattern, there could be tachyonic mass terms for some open string modes; these could allow a minimum of the full potential where modes with opposite charge get non-zero vev: however, as one can check, one obtains a de Sitter uplift term also in this case if at least one open string mode with positive charge has positive mass term. The case of pure chiral spectrum (where open string modes have negative charge and $\xi > 0$) does not lead to a supersymmetric configuration and we have not considered it here.

All these considerations led us to conjecture that the right supersymmetric configuration, that one should expand around, is a T-brane solution of the 8D equations of motion for the theory living on the D7-brane worldvolume. This takes place indeed when a non-zero gauge flux prevents the adjoint field Φ to commute with its conjugate. One can check that this is realised if an off-diagonal element of Φ gets a non-zero VEV that corresponds to a chiral state in the 4D effective field theory. To check this conjecture, we considered a simple SU(2) model that has a T-brane solution, i.e. a proper VEV for Φ . Three-form fluxes induce some particular couplings for Φ in the 8D worldvolume theory. After substituting the non-zero VEV of Φ in these couplings, we obtain exactly the uplift term found by using the 4D approach.

We have therefore succeeded to obtain metastable de Sitter string vacua starting from standard manifestly supersymmetric configurations with supersymmetry broken spontaneously. The mechanism is understood from both the 4D effective field theory and the higher dimensional one. The phenomenological implications of this class of de Sitter scenarios have been studied in [34]. It would be interesting to construct a consistent compact setup with a global T-brane solution, as those constructed via tachyon condensation in [35] and where moduli stabilisation is obtained in a dS LVS minimum. In principle this could be done in the global models presented in [2–4] (where the uplift mechanism was studied only from the 4D point of view) by constructing the tachyon matrix realising the T-brane

configuration and studying its stability and its low energy spectrum like in [35]. We leave this for future work.

Acknowledgments

We would like to thank D. Gallego, A. Hebecker, C. Mayrhofer and T. Weigand for useful conversations.

A AdS KKLT vacua

In this appendix we consider for completeness the case of KKLT vacua. Unlike the LVS vacua discussed in the text in this case, as expected, there are only supersymmetric AdS vacua. We focus on KKLT-like vacua with magnetised D7-branes and E3-instantons where K and W take the form:

$$K = -3\ln(T + \bar{T}) + c \frac{\phi\bar{\phi}}{(T + \bar{T})^\alpha} \quad \text{and} \quad W = W_0 + A\phi^n e^{-aT}. \quad (\text{A.1})$$

where $\alpha \geq 0$. The original KKLT model with just a closed string mode T can easily be recovered by setting $c = n = 0$. The two Kähler covariant derivatives read:

$$\begin{aligned} D_T W &= -aA\phi^n e^{-aT} - \frac{3}{T + \bar{T}} \left(1 + \frac{\alpha c}{3} \frac{\phi\bar{\phi}}{(T + \bar{T})^\alpha} \right) W \\ D_\phi W &= nA\phi^{n-1} e^{-aT} + c \frac{\bar{\phi}}{(T + \bar{T})^\alpha} W. \end{aligned} \quad (\text{A.2})$$

Imposing $D_T W = 0$ implies:

$$aA\phi^n e^{-aT} = -\frac{3}{T + \bar{T}} \left(1 + \frac{\alpha c}{3} \frac{\phi\bar{\phi}}{(T + \bar{T})^\alpha} \right) W. \quad (\text{A.3})$$

Substituting this result into $D_\phi W$ we obtain:

$$D_\phi W = \left[-\frac{3n}{T + \bar{T}} + ac \frac{\phi\bar{\phi}}{(T + \bar{T})^\alpha} \left(1 - \frac{n\alpha}{a(T + \bar{T})} \right) \right] \frac{W}{a\phi} \quad (\text{A.4})$$

$D_\phi W = 0$ has a solution at:

$$c \frac{\phi\bar{\phi}}{(T + \bar{T})^\alpha} = \frac{3n}{a(T + \bar{T})} \left(1 - \frac{n\alpha}{a(T + \bar{T})} \right)^{-1}. \quad (\text{A.5})$$

Plugging this solution back in (A.3) we find:

$$\begin{aligned} aA\phi^n e^{-aT} &= -\frac{3}{T + \bar{T}} \left[1 + \frac{n\alpha}{a(T + \bar{T})} \left(1 - \frac{n\alpha}{a(T + \bar{T})} \right)^{-1} \right] W \\ &= -\frac{3}{T + \bar{T}} \left(1 + \sum_{k=0}^{\infty} \epsilon^{k+1} \right) W \quad \text{with} \quad \epsilon \equiv \frac{n\alpha}{a(T + \bar{T})} \ll 1. \end{aligned} \quad (\text{A.6})$$

The parameter ϵ is very small since $a(T + \bar{T}) \gg 1$ in order to neglect higher order instanton contributions to W . It is clear that the solution with $n \neq 0$ is just a small perturbation with respect to the supersymmetric solution of the original KKLT model with $c = n = 0$. Because of the relation (2.23), the D-terms are automatically zero. Thus the new system keeps featuring the same supersymmetric solution of the original KKLT model. Let us study the form of the scalar potential to see if it can admit also dS solutions. Writing $T = \tau + i\psi$ and $\phi = |\phi| e^{i\theta}$ and minimising with respect to the axions, the F-term potential at leading order in an ϵ^{-1} expansion reads:

$$V_F = \frac{a^2 A^2 |\phi|^{2n} e^{-2a\tau}}{6\tau} - \frac{aAW_0 |\phi|^n e^{-a\tau}}{2\tau^2} + c \frac{W_0^2}{(2\tau)^3} \frac{|\phi|^2}{(2\tau)^\alpha}. \quad (\text{A.7})$$

On the other hand, the D-term potential at first order in ϵ^{-1} looks like:

$$V_D = \frac{1}{\tau} \left(q_\phi c \frac{|\phi|^2}{(2\tau)^\alpha} - q_T \frac{3}{2\tau} \right)^2. \quad (\text{A.8})$$

Minimising $V_{\text{tot}} = V_D + V_F$ with respect to $|\phi|$ we find at leading order:

$$\frac{\partial V_{\text{tot}}}{\partial |\phi|} \simeq cq_\phi \frac{4}{\tau} \frac{|\phi|}{(2\tau)^\alpha} \left(q_\phi c \frac{|\phi|^2}{(2\tau)^\alpha} - q_T \frac{3}{2\tau} \right). \quad (\text{A.9})$$

Setting this to zero gives (we discard the solution $|\phi| = 0$ since it gives a run-away for τ when substituted in V_{tot}):

$$q_\phi c \frac{|\phi|^2}{(2\tau)^\alpha} = q_T \frac{3}{2\tau}, \quad (\text{A.10})$$

which has the same leading order form of (A.5) once we impose the U(1)-invariance condition (2.21). Plugging this solution back in V_{tot} we end up with:

$$V_{\text{tot}} \simeq \frac{a^2 A^2 |\phi|^{2n} e^{-2a\tau}}{6\tau} - \frac{aAW_0 |\phi|^n e^{-a\tau}}{2\tau^2} + \frac{q_T}{q_\phi} \frac{3W_0^2}{16\tau^4}. \quad (\text{A.11})$$

Minimising with respect to τ we find at leading order:

$$\frac{\partial V_{\text{tot}}}{\partial \tau} \simeq -\frac{a^3 A^2 |\phi|^{2n} e^{-2a\tau}}{3\tau} + \frac{a^2 AW_0 |\phi|^n e^{-a\tau}}{2\tau^2} = 0. \quad (\text{A.12})$$

This equation admits a solution at:

$$aA |\phi|^n e^{-a\tau} = \frac{3W_0}{2\tau}, \quad (\text{A.13})$$

which reproduces (A.6) at leading order. Hence we conclude that this model can admit only supersymmetric AdS vacua.

B D-term potential from D7-branes

In this appendix we shall work out the exact form of the gauge kinetic function and the FI-terms following [22] and appendix A.3 of [24]. The 4D D-term potential that arises from the dimensional reduction of the D7-brane action takes the form:

$$V_D = \frac{\mu_7}{2} e^{-\phi} \frac{I_1^2}{I_2}, \quad (\text{B.1})$$

where:

$$\mu_7 = (2\pi)^{-3}(2\pi\alpha')^{-4}, \quad I_1 = \int_{D_i} J \wedge \mathcal{F}, \quad I_2 = \frac{1}{2} \int_{D_i} J \wedge J. \quad (\text{B.2})$$

Given that the real part of the gauge kinetic function is [22]:

$$\text{Re}(f_i) = g_i^{-2} = \mu_7(2\pi\alpha')^2 e^{-\phi} I_2, \quad (\text{B.3})$$

the D-term potential can be rewritten as:

$$V_D = \frac{g^2}{2} \left(\mu_7 2\pi\alpha' e^{-\phi} I_1 \right)^2 = \frac{g^2}{2} \xi_i^2. \quad (\text{B.4})$$

Let us work out the expression of the FI-term ξ . If we expand J and \mathcal{F} in a basis of (1,1)-forms as $J = t_j^{(s)} \hat{D}_j$ ($t_j^{(s)}$ are two-cycle volumes in the string frame and in units of ℓ_s) and $\mathcal{F} = f_k \hat{D}_k$, we get:

$$I_1 = t_j^{(s)} f_k \int_{D_i} \hat{D}_j \wedge \hat{D}_k = t_j^{(s)} f_k \frac{1}{\ell_s^2} \int_{X_3} \hat{D}_j \wedge \hat{D}_j \wedge \hat{D}_k = k_{ijk} t_j^{(s)} f_k \ell_s^4, \quad (\text{B.5})$$

where the triple intersection numbers k_{ijk} are defined as:

$$k_{ijk} = \frac{1}{\ell_s^6} \int_{X_3} \hat{D}_j \wedge \hat{D}_j \wedge \hat{D}_k. \quad (\text{B.6})$$

Given that the gauge flux is quantised as (\tilde{D} denotes the two-cycle dual to the divisor D):

$$\int_{\tilde{D}_k} F = 2\pi n_k \Leftrightarrow \frac{1}{2\pi\alpha'} \int_{\tilde{D}_k} \mathcal{F} = 2\pi n_k \quad n_k \in \mathbb{Z}, \quad (\text{B.7})$$

we find that the flux quanta f_k take the form:

$$\frac{f_j}{2\pi\alpha'} \int_{\tilde{D}_k} \hat{D}_j = 2\pi n_k \Rightarrow f_k = \frac{4\pi^2\alpha'}{\ell_s^2} n_k, \quad (\text{B.8})$$

and so the FI-term becomes (using $\ell_s = M_s^{-1}$):

$$\xi_i = \frac{e^{-\phi}}{(2\pi)^2} \left(\frac{\ell_s^2}{2\pi\alpha'} \right)^2 k_{ijk} t_j^{(s)} n_k M_s^2. \quad (\text{B.9})$$

From the dimensional reduction of the 10D action we have:

$$M_s^2 = \frac{(2\pi)^3}{2} e^{\phi/2} \left(\frac{2\pi\alpha'}{\ell_s^2} \right)^4 \frac{M_{\text{P}}^2}{\mathcal{V}}, \quad (\text{B.10})$$

where $\mathcal{V} = e^{-3\phi/2} \mathcal{V}_{(s)}$ is the CY volume in Einstein-frame. Hence (B.9) can be rewritten in Planck units as (where $t_j = e^{-\phi/2} t_j^{(s)}$):

$$\frac{\xi_i}{M_{\text{P}}^2} = \left(\frac{2\pi\alpha'}{\ell_s^2} \right) \frac{e^{-\phi/2}}{\ell_s^4} \frac{1}{2\mathcal{V}} \int_{D_i} J \wedge \mathcal{F} = 2\pi \left(\frac{2\pi\alpha'}{\ell_s^2} \right)^2 k_{ijk} n_k \frac{t_j}{2\mathcal{V}}. \quad (\text{B.11})$$

The quantities $k_{ijk}n_k$ are integers and give the charge of the j -th Kähler modulus under the anomalous $U(1)$ living on the i -th divisor:

$$q_{ij} = k_{ijk}n_k = \frac{1}{2\pi} \left(\frac{\ell_s^2}{2\pi\alpha'} \right) \frac{1}{\ell_s^4} \int_{D_i} \hat{D}_j \wedge \mathcal{F}. \quad (\text{B.12})$$

Moreover, using four-cycle moduli:

$$\text{Re}(T_i) = \frac{1}{2} \int_{D_i} J \wedge J = \frac{1}{2} k_{ijk} t_j t_k, \quad (\text{B.13})$$

and the fact that $K/M_{\text{P}}^2 = -2 \ln \mathcal{V}$ we have that:

$$\frac{\partial K}{\partial T_j} = -\frac{t_j}{2\mathcal{V}}, \quad (\text{B.14})$$

and so (B.11) takes the form:

$$\frac{\xi_i}{M_{\text{P}}^2} = -2\pi \left(\frac{2\pi\alpha'}{\ell_s^2} \right)^2 q_{ij} \frac{\partial K}{\partial T_j}. \quad (\text{B.15})$$

On the other hand, the gauge kinetic function (B.3) can be rewritten in a more compact form using (B.13) as:

$$\text{Re}(f_i) = g_i^{-2} = \frac{e^{-\phi}}{(2\pi)^3 (2\pi\alpha')^2} \frac{1}{2} \int_{D_i} J \wedge J = \frac{1}{(2\pi)^3} \left(\frac{\ell_s^2}{2\pi\alpha'} \right)^2 \text{Re}(T_i). \quad (\text{B.16})$$

Therefore the exact expressions of ξ_i and f_i depend on the choice of the fundamental unit of lengths ℓ_s . Let us give two examples:

1. For $\ell_s = \sqrt{\alpha'}$ we have [22]:

$$f_k = (2\pi)^2 n_k, \quad n_k \in \mathbb{Z} \quad \text{and} \quad M_s^2 = \frac{(2\pi)^7}{2} g_s^{1/2} \frac{M_{\text{P}}^2}{\mathcal{V}}, \quad (\text{B.17})$$

together with:

$$f_i = \frac{T_i}{(2\pi)^5} \quad \text{and} \quad \frac{\xi_i}{M_{\text{P}}^2} = -(2\pi)^3 q_{ij} \frac{\partial K}{\partial T_j}. \quad (\text{B.18})$$

This is the convention we followed in this paper. If the Kähler moduli are instead redefined as $\frac{T_i}{(2\pi)^5} \rightarrow T_i$ [22], the new expression of f_i , ξ_i and M_s become:

$$f_i = T_i, \quad \frac{\xi_i}{M_{\text{P}}^2} = -\frac{q_{ij}}{(2\pi)^2} \frac{\partial K}{\partial T_j} \quad \text{and} \quad M_s^2 = \frac{1}{2} \sqrt{\frac{g_s}{2\pi}} \frac{M_{\text{P}}^2}{\mathcal{V}}. \quad (\text{B.19})$$

2. For $\ell_s = 2\pi\sqrt{\alpha'}$ we have [24]:

$$f_k = n_k, \quad n_k \in \mathbb{Z} \quad \text{and} \quad M_s^2 = \frac{g_s^{1/2}}{4\pi} \frac{M_{\text{P}}^2}{\mathcal{V}}, \quad (\text{B.20})$$

together with:

$$f_i = \frac{T_i}{2\pi} \quad \text{and} \quad \frac{\xi_i}{M_{\text{P}}^2} = -\frac{q_{ij}}{2\pi} \frac{\partial K}{\partial T_j}. \quad (\text{B.21})$$

If the Kähler moduli are redefined as $\frac{T_i}{2\pi} \rightarrow T_i$ [23], the new expression of f_i , ξ_i and M_s become:

$$f_i = T_i, \quad \frac{\xi_i}{M_{\text{P}}^2} = -\frac{q_{ij}}{(2\pi)^2} \frac{\partial K}{\partial T_j} \quad \text{and} \quad M_s^2 = \frac{1}{2(2\pi)^2} \sqrt{\frac{g_s}{2\pi}} \frac{M_{\text{P}}^2}{\mathcal{V}}. \quad (\text{B.22})$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] M. Cicoli, C. Mayrhofer and R. Valandro, *Moduli Stabilisation for Chiral Global Models*, *JHEP* **02** (2012) 062 [[arXiv:1110.3333](https://arxiv.org/abs/1110.3333)] [[INSPIRE](#)].
- [2] M. Cicoli, S. Krippendorff, C. Mayrhofer, F. Quevedo and R. Valandro, *D-Branes at del Pezzo Singularities: Global Embedding and Moduli Stabilisation*, *JHEP* **09** (2012) 019 [[arXiv:1206.5237](https://arxiv.org/abs/1206.5237)] [[INSPIRE](#)].
- [3] M. Cicoli, S. Krippendorff, C. Mayrhofer, F. Quevedo and R. Valandro, *D3/D7 Branes at Singularities: Constraints from Global Embedding and Moduli Stabilisation*, *JHEP* **07** (2013) 150 [[arXiv:1304.0022](https://arxiv.org/abs/1304.0022)] [[INSPIRE](#)].
- [4] M. Cicoli, D. Klevers, S. Krippendorff, C. Mayrhofer, F. Quevedo and R. Valandro, *Explicit de Sitter Flux Vacua for Global String Models with Chiral Matter*, *JHEP* **05** (2014) 001 [[arXiv:1312.0014](https://arxiv.org/abs/1312.0014)] [[INSPIRE](#)].
- [5] R. Blumenhagen, V. Braun, T.W. Grimm and T. Weigand, *GUTs in Type IIB Orientifold Compactifications*, *Nucl. Phys. B* **815** (2009) 1 [[arXiv:0811.2936](https://arxiv.org/abs/0811.2936)] [[INSPIRE](#)].
- [6] A. Collinucci, M. Kreuzer, C. Mayrhofer and N.O. Walliser, *Four-modulus ‘Swiss Cheese’ chiral models*, *JHEP* **07** (2009) 074 [[arXiv:0811.4599](https://arxiv.org/abs/0811.4599)] [[INSPIRE](#)].
- [7] R. Blumenhagen, S. Moster and E. Plauschinn, *Moduli Stabilisation versus Chirality for MSSM like Type IIB Orientifolds*, *JHEP* **01** (2008) 058 [[arXiv:0711.3389](https://arxiv.org/abs/0711.3389)] [[INSPIRE](#)].
- [8] G. Aldazabal, L.E. Ibáñez, F. Quevedo and A.M. Uranga, *D-branes at singularities: A Bottom up approach to the string embedding of the standard model*, *JHEP* **08** (2000) 002 [[hep-th/0005067](https://arxiv.org/abs/hep-th/0005067)] [[INSPIRE](#)].
- [9] J.P. Conlon, A. Maharana and F. Quevedo, *Towards Realistic String Vacua*, *JHEP* **05** (2009) 109 [[arXiv:0810.5660](https://arxiv.org/abs/0810.5660)] [[INSPIRE](#)].
- [10] S.B. Giddings, S. Kachru and J. Polchinski, *Hierarchies from fluxes in string compactifications*, *Phys. Rev. D* **66** (2002) 106006 [[hep-th/0105097](https://arxiv.org/abs/hep-th/0105097)] [[INSPIRE](#)].
- [11] V. Balasubramanian, P. Berglund, J.P. Conlon and F. Quevedo, *Systematics of moduli stabilisation in Calabi-Yau flux compactifications*, *JHEP* **03** (2005) 007 [[hep-th/0502058](https://arxiv.org/abs/hep-th/0502058)] [[INSPIRE](#)].
- [12] J.P. Conlon, F. Quevedo and K. Suruliz, *Large-volume flux compactifications: Moduli spectrum and D3/D7 soft supersymmetry breaking*, *JHEP* **08** (2005) 007 [[hep-th/0505076](https://arxiv.org/abs/hep-th/0505076)] [[INSPIRE](#)].
- [13] R. Donagi, S. Katz and E. Sharpe, *Spectra of D-branes with Higgs vevs*, *Adv. Theor. Math. Phys.* **8** (2004) 813 [[hep-th/0309270](https://arxiv.org/abs/hep-th/0309270)] [[INSPIRE](#)].
- [14] S. Cecotti, C. Cordova, J.J. Heckman and C. Vafa, *T-Branes and Monodromy*, *JHEP* **07** (2011) 030 [[arXiv:1010.5780](https://arxiv.org/abs/1010.5780)] [[INSPIRE](#)].
- [15] R. Donagi and M. Wijnholt, *Gluing Branes, I*, *JHEP* **05** (2013) 068 [[arXiv:1104.2610](https://arxiv.org/abs/1104.2610)] [[INSPIRE](#)].
- [16] S. Kachru, R. Kallosh, A.D. Linde and S.P. Trivedi, *de Sitter vacua in string theory*, *Phys. Rev. D* **68** (2003) 046005 [[hep-th/0301240](https://arxiv.org/abs/hep-th/0301240)] [[INSPIRE](#)].

- [17] R. Minasian and G.W. Moore, *K theory and Ramond-Ramond charge*, *JHEP* **11** (1997) 002 [[hep-th/9710230](#)] [[INSPIRE](#)].
- [18] D.S. Freed and E. Witten, *Anomalies in string theory with D-branes*, *Asian J. Math.* **3** (1999) 819 [[hep-th/9907189](#)] [[INSPIRE](#)].
- [19] P. Berglund and I. Garcia-Etxebarria, *D-brane instantons on non-Spin cycles*, *JHEP* **01** (2013) 056 [[arXiv:1210.1221](#)] [[INSPIRE](#)].
- [20] A.P. Braun, M. Rummel, Y. Sumitomo and R. Valandro, *de Sitter vacua from a D-term generated racetrack potential in hypersurface Calabi-Yau compactifications*, *JHEP* **12** (2015) 033 [[arXiv:1509.06918](#)] [[INSPIRE](#)].
- [21] R. Blumenhagen, B. Körs, D. Lüst and S. Stieberger, *Four-dimensional String Compactifications with D-branes, Orientifolds and Fluxes*, *Phys. Rept.* **445** (2007) 1 [[hep-th/0610327](#)] [[INSPIRE](#)].
- [22] M. Haack, D. Krefl, D. Lüst, A. Van Proeyen and M. Zagermann, *Gaugino Condensates and D-terms from D7-branes*, *JHEP* **01** (2007) 078 [[hep-th/0609211](#)] [[INSPIRE](#)].
- [23] D. Cremades, M.-P. Garcia del Moral, F. Quevedo and K. Suruliz, *Moduli stabilisation and de Sitter string vacua from magnetised D7 branes*, *JHEP* **05** (2007) 100 [[hep-th/0701154](#)] [[INSPIRE](#)].
- [24] M. Cicoli, M. Goodsell, J. Jaeckel and A. Ringwald, *Testing String Vacua in the Lab: From a Hidden CMB to Dark Forces in Flux Compactifications*, *JHEP* **07** (2011) 114 [[arXiv:1103.3705](#)] [[INSPIRE](#)].
- [25] L. Aparicio, D.G. Cerdeno and L.E. Ibáñez, *Modulus-dominated SUSY-breaking soft terms in F-theory and their test at LHC*, *JHEP* **07** (2008) 099 [[arXiv:0805.2943](#)] [[INSPIRE](#)].
- [26] J.P. Conlon, D. Cremades and F. Quevedo, *Kähler potentials of chiral matter fields for Calabi-Yau string compactifications*, *JHEP* **01** (2007) 022 [[hep-th/0609180](#)] [[INSPIRE](#)].
- [27] R. Blumenhagen, M. Cvetič and T. Weigand, *Spacetime instanton corrections in 4D string vacua: The Seesaw mechanism for D-brane models*, *Nucl. Phys. B* **771** (2007) 113 [[hep-th/0609191](#)] [[INSPIRE](#)].
- [28] L.E. Ibáñez and A.M. Uranga, *Neutrino Majorana Masses from String Theory Instanton Effects*, *JHEP* **03** (2007) 052 [[hep-th/0609213](#)] [[INSPIRE](#)].
- [29] B. Florea, S. Kachru, J. McGreevy and N. Saulina, *Stringy Instantons and Quiver Gauge Theories*, *JHEP* **05** (2007) 024 [[hep-th/0610003](#)] [[INSPIRE](#)].
- [30] R. Blumenhagen, M. Cvetič, S. Kachru and T. Weigand, *D-Brane Instantons in Type II Orientifolds*, *Ann. Rev. Nucl. Part. Sci.* **59** (2009) 269 [[arXiv:0902.3251](#)] [[INSPIRE](#)].
- [31] C. Beasley, J.J. Heckman and C. Vafa, *GUTs and Exceptional Branes in F-theory — I*, *JHEP* **01** (2009) 058 [[arXiv:0802.3391](#)] [[INSPIRE](#)].
- [32] P.G. Cámara, L.E. Ibáñez and A.M. Uranga, *Flux-induced SUSY-breaking soft terms on D7-D3 brane systems*, *Nucl. Phys. B* **708** (2005) 268 [[hep-th/0408036](#)] [[INSPIRE](#)].
- [33] P.G. Cámara, L.E. Ibáñez and I. Valenzuela, *Flux-induced Soft Terms on Type IIB/F-theory Matter Curves and Hypercharge Dependent Scalar Masses*, *JHEP* **06** (2014) 119 [[arXiv:1404.0817](#)] [[INSPIRE](#)].
- [34] L. Aparicio, M. Cicoli, S. Krippendorf, A. Maharana, F. Muia and F. Quevedo, *Sequestered de Sitter String Scenarios: Soft-terms*, *JHEP* **11** (2014) 071 [[arXiv:1409.1931](#)] [[INSPIRE](#)].
- [35] A. Collinucci and R. Savelli, *T-branes as branes within branes*, *JHEP* **09** (2015) 161 [[arXiv:1410.4178](#)] [[INSPIRE](#)].