

Dynamical effects in the vibrationally resolved C 2s⁻¹ photoionization cross section ratios of Methane

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 112048

(<http://iopscience.iop.org/1742-6596/635/11/112048>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 87.30.54.86

This content was downloaded on 31/01/2017 at 14:04

Please note that [terms and conditions apply](#).

You may also be interested in:

[Optically Forbidden Excitations of 2s Electron of Neon Studied by Fast Electron Impact](#)
Ge Min, Zhu Lin-Fan, Liu Cun-Ding et al.

[Universality classes of critical antiferromagnets](#)
H Johannesson

[Spin-flop transition in random antiferromagnets](#)
R A Tahir-Kheli, S M Rezende, F G B Moreira et al.

[A vibrationally resolved experimental study of the sulfur L-shell photoelectron spectrum of CS₂](#)
H Wang, M Bässler, I Hjelte et al.

[Lower-hybrid-resonance heating of a plasma in a parallel-plate waveguide](#)
S. Puri and M. Tutter

[The medical applications of ultrasonics](#)
P N T Wells

Dynamical effects in the vibrationally resolved C 2s⁻¹ photoionization cross section ratios of Methane

Saikat Nandi*¹, Minna Patanen*, Catalin Miron*[§], John D. Bozek*, Etienne Plésiat[†], Fernando Martín[‡], Daniele Toffoli[‡] and Piero Decleva^{‡ 2}

*Synchrotron SOLEIL, L'Orme des Merisiers, Saint-Aubin, BP 48, 91192 Gif-sur-Yvette Cedex, France

[†]Departamento de Química, Módulo 13, Universidad Autónoma de Madrid, Cantoblanco 28049, Madrid, Spain

[‡]Dipartimento di Scienze Chimiche e Farmaceutiche, Università di Trieste, Via L. Giorgieri 1, 34127 Trieste, Italy

[§]Extreme Light Infrastructure - Nuclear Physics (ELI-NP), 'Horia Hulubei' National Institute for Physics and Nuclear Engineering, 30 Reactorului Street, RO-077125 Măgurele, Jud. Ilfov, Romania

Synopsis The vibrationally resolved C 2s photoionization cross-section of methane was investigated both theoretically and experimentally. When compared to that of C 1s photoionization, a rather different pattern has been observed, suggesting a strong interplay between the electron diffraction and interference effects.

Interference and diffraction effects in molecular photoionization have been recently revealed and subjected to a number of studies [1-5]. The experimental observation of such effects is very difficult when measuring absolute cross sections, but they can be evidenced in a much easier way by monitoring the photon energy dependence of the appropriate ratios of partial cross sections. One such possibility is the study of ν -ratios that is cross section ratios of different vibrational levels in the same electronic channel. In absence of any diffraction and interference effects, the ν -ratios should be essentially energy-independent, a constant equal to the Franck-Condon ratios. However, the experimentally studied ν -ratios of valence [1] and core ionized [4,5] show apparent energy-dependent oscillations, which can be qualitatively predicted from the Cohen-Fano model [1,3] and simple diffraction effects [2,4,5]. The method to study ν -ratios has proven to be a very sensitive probe, as it can reveal even the tiny diffraction associated with the light hydrogen atoms, as has been shown in the core ionization of CH₄ [2].

A key feature of the oscillations is their relationship to the molecular geometry, and in particular the bond length between emitting and scattering centers [5]. While the observed patterns are quite accurately reproduced by high quality ab-initio theoretical calculations, a full understanding of all factors involved is still incomplete. To this end we have compared the behavior of ν -ratios in the photoionization of the inner-valence 2a₁ orbital of methane which is a linear combination of C 2s and H 1s orbitals, with the completely localized C 1s core orbital. The vibrationally resolved photoionization

cross section ratios of the symmetric stretching mode in the C 1s and C 2s photoionization of CH₄ were measured at PLÉIADES beam-line [6] using a VG-Scienta R4000 spectrometer. Unexpectedly, both experimental and theoretical ν -ratios of 2a₁ photoionization exhibit a quite different pattern compared to the C1s photoionization, indicating that in case of the 2a₁ state the relative cross section oscillations are also modulated by other effects in addition to the diffraction of an electron issued from a localized emitter. A possible explanation lays in the more delocalized and spatially extended nature of the initial 2a₁ (C 2s) orbital, although additional complications (e.g. apparent predissociation effects in the experimental spectra) may also come into play. A better understanding of the behavior of the ν -ratios of an inner-valence orbital may give us not only structural information (e.g., the C-H distance in this case) but also reveals information of other contributing factors such as the molecular orbital composition (i.e. electron density).

References

- [1] S. E. Canton, E. Plésiat, J. D. Bozek, B. S. Rude, P. Decleva, and F. Martín 2011 *Proc Natl Acad Sci USA* **108** 7302
- [2] E. Plésiat, L. Argenti, C. Miron, E. Kukuk, K. Ueda, P. Decleva and F. Martín 2012 *Phys. Rev. A* **85** 023409
- [3] R. K. Kushawaha M. Patanen, R. Guillemin, L. Journal, C. Miron, M. Simon, M. N. Piancastelli, P. Decleva 2013 *Proc Natl Acad Sci USA* **110** 15201
- [4] K. Ueda *et al* 2013 *J. Chem Phys.* **139** 124306
- [5] E. Kukuk *et al* 2013 *Phys. Rev. A* **88** 033412
- [6] <http://www.synchrotron-soleil.fr/Recherche/LignesLumiere/PLEIADES>

¹ E-mail: saikat.nandi@synchrotron-soleil.fr

² E-mail: decleva@units.it

