

Relationship between the gut microbiota and bile acid composition in the ileal mucosa of Crohn's disease

著者	BAMBA Shigeki, INATOMI Osamu, NISHIDA Atsushi, OHNO Masashi, IMAI Takayuki, TAKAHASHI Kenichiro, NAITO Yuji, IWAMOTO Junichi, HONDA Akira, INOHARA Naohiro, ANDOH Akira
journal or publication title	Intestinal Research
volume	20
number	3
page range	370-380
year	2022-07
URL	http://hdl.handle.net/10422/00013392

doi: 10.5217/ir.2021.00054(<https://doi.org/10.5217/ir.2021.00054>)

See “Relationship between the gut microbiota and bile acid composition in the ileal mucosa of Crohn’s disease” on pages 370-380.

Supplementary Table 1. Background Characteristics of Crohn's Disease

Characteristics	Inactive (n = 18)	Active (n = 9)
Sex (male/female)	14/4	7/2
Age (yr), median (IQR)	35.1 (31.9–47.0)	47.4 (35.3–60.7)
Body mass index (kg/m ²), median (IQR)	21.2 (18.6–24.5)	21.7 (20.2–23.5)
Smoking status (never/previous/current)	12/3/3	7/2/0
History of intestinal resection (yes/no)	3/15	0/9
Disease duration (yr), median (IQR)	9.5 (6.6–14.5)	14.6 (1.6–33.0)
Disease location (L1/L2/L3)	9/1/8	4/0/5
Disease behavior (B1/B2/B3)	9/7/2	1/5/3
CDAI, median (IQR)	62.0 (46.5–83.7)	66.0 (48.0–96.0)
Disease		
Crohn's disease	18	9
Gastrointestinal bleeding	-	-
Intestinal neoplasia	-	-
Other	-	-
Medication, No. (%)		
Proton pump inhibitors	3 (16.7)	2 (22.2)
Probiotics	6 (33.3)	4 (44.4)
5-ASA/SASP	11 (61.1)	7 (77.8)
Prednisolone	1 (5.6)	0
Immunomodulators	10 (55.6)	3 (33.3)
Biologics	6 (33.3)	3 (33.3)

IQR, interquartile range; L1, ileal; L2, colonic; L3, ileocolonic; B1, non-stricturing, non-penetrating; B2, stricturing; B3, penetrating; CDAI, Crohn's disease activity index; 5-ASA, 5-aminosalicylates; SASP, sulfasalazine.