

Grigoras, C.; Grigoriev, V.; Grigoryan, A.; Grigoryan, S.; Groettvik, O. S.; Grossa, F.; Grosse-Oetringhaus, J. F.; Grosso, R.; Guardiano, G. G.; Guernane, R.; Guilbaud, M.; Guittiere, M.; Gulbrandsen, K.; Gunji, T.; Gupta, A.; Gupta, R.; Guzman, I. B.; Gyulai, L.; Habib, M. K.; Hadjidakis, C.; Hamagaki, H.; Hamar, G.; Hamid, M.; Hannigan, R.; Haque, M. R.; Harlenderova, A.; Harris, J. W.; Harton, A.; Hasenbichler, J. A.; Hassan, H.; Hatzifotiadou, D.; Hauer, P.; Havener, L. B.; Hayashi, S.; Heckel, S. T.; Hellbär, E.; Helstrup, H.; Herman, T.; Hernandez, E. G.; Herrera Corral, G.; Herrmann, F.; Hetland, K. F.; Hillemanns, H.; Hills, C.; Hippolyte, B.; Hohlweber, B.; Honermann, J.; Hong, G. H.; Horak, D.; Hornung, S.; Hosokawa, R.; Hristov, P.; Huang, C.; Hughes, C.; Huhn, P.; Humanic, T. J.; Hushnud, H.; Husova, L. A.; Hussain, N.; Hutter, D.; Iddon, J. P.; Ilkaev, R.; Ilyas, H.; Inaba, M.; Innocenti, G. M.; Ippolitov, M.; Isakov, A.; Islam, M. S.; Ivanov, M.; Ivanov, V.; Izucheev, V.; Jacak, B.; Jacazio, N.; Jacobs, P. M.; Jadlovska, S.; Jadlovsky, J.; Jaelani, S.; Jahnke, C.; Jakubowska, M. J.; Janik, M. A.; Janson, T.; Jercic, M.; Jevons, O.; Jonas, F.; Jones, P. G.; Jowett, J. M.; Jung, J.; Jung, M.; Junique, A.; Jusko, A.; Kalinak, P.; Kalweit, A.; Kaplin, V.; Kar, S.; Karasu Uysal, A.; Karatovic, D.; Karavichev, O.; Karavicheva, T.; Karczmarczyk, P.; Karpechev, E.; Kazantsev, A.; Kebschull, U.; Keidel, R.; Keil, M.; Ketzer, B.; Khabanova, Z.; Khan, A. M.; Khan, S.; Khanzadeev, A.; Kharlov, Y.; Khatun, A.; Khuntia, A.; Kileng, B.; Kim, B.; Kim, D.; Kim, D. J.; Kim, E. J.; Kim, J.; Kim, J. S.; Kim, J.; Kim, J.; Kim, M.; Kim, S.; Kim, T.; Kirsch, S.; Kisiel, I.; Kiselev, S.; Kisiel, A.; Klay, J. L.; Klein, J.; Klein, S.; Klein-Bösing, C.; Kleiner, M.; Klemenz, T.; Kluge, A.; Knospe, A. G.; Kobdaj, C.; Köhler, M. K.; Kollegger, T.; Kondratyev, A.; Kondratyeva, N.; Kondratyuk, E.; Konig, J.; Konigstorfer, S. A.; Konopka, P. J.; Kornakov, G.; Koryciak, S. D.; Koska, L.; Kovalenko, O.; Kovalenko, V.; Kowalski, M.; Králik, I.; Kraváková, A.; Kreis, L.; Krivda, M.; Krizek, F.; Krizkova Gajdosova, K.; Kroesen, M.; Krüger, M.; Kryshen, E.; Krzewicki, M.; Kuera, V.; Kuhn, C.; Kuijer, P. G.; Kumaoka, T.; Kumar, D.; Kumar, L.; Kundu, S.; Kurashvili, P.; Kurepin, A.; Kurepin, A. B.; Kuryakin, A.; Kushpil, S.; Kvapil, J.; Kweon, M. J.; Kwon, J. Y.; Kwon, Y.; La Pointe, S. L.; La Rocca, P.; Lai, Y. S.; Lakrathok, A.; Lamanna, M.; Langoy, R.; Lapidus, K.; Larionov, P.; Laudi, E.; Lautner, L.; Lavicka, R.; Lazareva, T.; Lea, R.; Lee, J.; Lehrbach, J.; Lemmon, R. C.; León Monzón, I.; Lesser, E. D.; Lettrich, M.; Lévai, P.; Li, X.; Li, X. L.; Lien, J.; Lietava, R.; Lim, B.; Lim, S. H.; Lindenstruth, V.; Lindner, A.; Lippmann, C.; Liu, A.; Liu, J.; Lofnes, I. M.; Loginov, V.; Loizides, C.; Loncar, P.; Lopez, J. A.; Lopez, X.; López Torres, E.; Luhder, J. R.; Lunardon, M.; Luparello, G.; Ma, Y. G.; Maevskaya, A.; Mager, M.; Mahmoud, T.; Maire, A.; Malaev, M.; Malik, Q. W.; Malinina, L.; Mal'Kevich, D.; Mallick, N.; Malzacher, P.; Mandaglio, G.; Manko, V.; Manso, F.; Manzari, V.; Mao, Y.; Mareš, J.; Margagliotti, G. V.; Margotti, A.; Marín, A.; Markert, C.; Marquard, M.; Martin, N. A.; Martinengo, P.; Martinez, J. L.; Martínez, M. I.; Martínez García, G.; Masciocchi, S.; Masera, M.; Masoni, A.; Massacrier, L.; Mastroserio, A.; Mathis, A. M.; Matonoha, O.; Matuoka, P. F. T.; Matyja, A.; Mayer, C.; Mazuecos, A. L.; Mazzaschi, F.; Mazzilli, M.; Mazzoni, M. A.; Mechler, A. F.; Meddi, F.; Melikyan, Y.; Menchaca-Rocha, A.; Meninno, E.; Menon, A. S.; Meres, M.; Mhlanga, S.; Miake, Y.; Micheletti, L.; Migliorin, L. C.; Mihaylov, D. L.; Mikhaylov, K.; Mishra, A. N.; Mikowiec, D.; Modak, A.; Mohanty, A. P.; Mohanty, B.; Mohisin Khan, M.; Moravcova, Z.; Mordasini, C.; Moreira De Godoy, D. A.; Moreno, L. A. P.; Morozov, I.; Morsch, A.; Mrnjavac, T.; Muccifora, V.; Mudnic, E.; Mühlheim, D.; Muhuri, S.; Mulligan, J. D.; Mulliri, A.; Munhoz, M. G.; Munzer, R. H.; Murakami, H.; Murray, S.; Musa, L.; Musinsky, J.; Myers, C. J.; Myrcha, J. W.; Naik, B.; Nair, R.; Nandi, B. K.; Nania, R.; Nappi, E.; Naru, M. U.; Nassirpour, A. F.; Nattrass, C.; Neagu, A.; Nellen, L.; Nesbo, S. V.; Neskovic, G.; Nesterov, D.; Nielsen, B. S.; Nikolaev, S.; Nikulin, S.; Nikulin, V.; Noferini, F.; Noh, S.; Nomokonov, P.; Norman, J.; Novitzky, N.; Nowakowski, P.; Nyandin, A.; Nystrand, J.; Ogino, M.; Ohlson, A.; Oleniacz, J.; Oliveira Da Silva, A. C.; Oliver, M. H.; Onnerstad, A.; Oppedisano, C.; Ortiz Velasquez, A.; Osako, T.; Oskarsson, A.; Otwinowski, J.; Oyama, K.; Pachmayer, Y.; Pacik, V.; Padhan, S.; Pagano, D.; Pai, G.; Palasciano, A.; Pan, J.; Panebianco, S.; Pareek, P.; Park, J.; Parkkila, J. E.; Pathak, S. P.; Patra, R. N.; Paul, B.; Pazzini, J.; Pei, H.; Peitzmann, T.; Peng, X.; Pereira, L. G.; Pereira Da Costa, H.; Peresunko, D.; Perez, G. M.; Perrin, S.; Pestov, Y.; Petráek, V.; Petrovici, M.; Pezzi, R. P.; Piano, S.; Pikna, M.; Pillot, P.; Pinazza, O.; Pinsky, L.; Pinto, C.; Pisano, S.; Posko, M.; Planinic, M.; Pliquet, F.; Poghosyan, M. G.; Polichtchouk, B.; Politano', S.; Poljak, N.; Pop, A.; Porteboeuf-Houssais, S.; Porter, J.; Pozdniakov, V.; Prasad, S. K.; Preghenella, R.; Prino, F.; Pruneau, C. A.; Pshenichnov, I.; Puccio, M.; Qiu, S.; Quaglia, L.; Quishpe, R. E.; Ragoni, S.; Rakotozafindrabe, A.; Ramello, L.; Rami, F.; Ramirez, S. A. R.; Ramos, A. G. T.; Raniwala, R.; Raniwala, S.; Räsänen, S. S.; Rath, R.; Ravasenga, I.; Read, K. F.; Redelbach, A. R.; Redlich, K.; Rehman, A.; Reichelt, P.; Reidt, F.; Reme-ness, H. A.; Renfordt, R.; Rescakova, Z.; Reygers, K.; Riabov, A.; Riabov, V.; Richert, T.; Richter, M.; Riegler, W.; Riggi, F.; Ristea, C.; Rode, S. P.; Rodríguez Cahuantzi, M.; Røed, K.; Rogalev, R.; Rogochaya, E.; Rogoschinski, T. S.; Rohr, D.; Röhrlrich, D.; Rojas, P. F.; Rokita, P. S.; Ronchetti, F.; Rosano, A.; Rosas, E. D.; Rossi, A.; Rotondi, A.; Roy, A.; Roy, P.; Roy, S.; Rubini, N.; Rueda, O. V.; Rui, R.; Rumyantsev, B.; Rustamov, A.; Ryabinkin, E.; Ryabov, Y.; Rybicki, A.; Rytkonen, H.; Rzesz, W.; Saarimaki, O. A. M.; Sadek, R.; Sadovsky, S.; Saetre, J.; Šafaík, K.; Saha, S. K.; Saha, S.; Sahoo, B.; Sahoo, P.; Sahoo, R.; Sahoo, S.; Sahu, D.; Sahu, P. K.; Saini, J.; Sakai, S.; Sambyal, S.; Samsonov, V.; Sarkar, D.; Sarkar, N.; Sarma, P.; Sarti, V. M.; Sas, M. H. P.; Schambach, J.; Scheid, H. S.; Schiaua, C.; Schicker, R.; Schmah, A.; Schmidt, C.; Schmidt, H. R.; Schmidt, M. O.; Schmidt, M.; Schmidt, N. V.; Schmier, A. R.; Schotter, R.; Schukraft, J.; Schutz, Y.; Schwarz, K.; Schweda, K.; Sciolli, G.; Scomparin, E.; Seger, J. E.; Sekiguchi, Y.; Sekihata, D.; Selyuzhenkov, I.; Senyukov, S.; Seo, J. J.; Serebryakov, D.; Šerkšnyt, L.; Sevcenco, A.; Shaba, T. J.; Shabanov, A.; Shabetai, A.; Shahoyan, R.; Shaikh, W.; Shangaraev, A.; Sharma, A.; Sharma, H.; Sharma, M.; Sharma, N.; Sharma, S.; Sheibani, O.; Shigaki, K.; Shimomura, M.; Shirinkin, S.; Shou, Q.; Sibiriak, Y.; Siddhanta, S.; Siemiarzuk, T.; Silva, T. F.; Silvermyr, D.; Simonetti, G.; Singh, B.; Singh, R.; Singh, R.; Singh, V. K.; Singhal, V.; Sinha, T.; Sitar, B.; Sitta, M.; Skaali, T. B.; Skorodumovs, G.; Slupecki, M.; Smirnov, N.; Snellings, R. J. M.; Soncco, C.; Song, J.; Songmoolnak, A.; Soramel, F.; Sorensen, S.; Sputowska, I.; Stachel, J.; Stan, I.; Steffanic, P. J.; Stiefelmaier, S. F.; Stocco, D.; Storetvedt, M. M.;

Stylianidis, C. P.; Suaide, A. A. P.; Sugitate, T.; Suire, C.; Suljic, M.; Sultanov, R.; Šumbera, M.; Sumberia, V.; Sumowidagdo, S.; Swain, S.; Szabo, A.; Szarka, I.; Tabassam, U.; Taghavi, S. F.; Taillepied, G.; Takahashi, J.; Tambave, G. J.; Tang, S.; Tang, Z.; Tarhini, M.; Tarzila, M. G.; Tauro, A.; Tejeda Muñoz, G.; Telesca, A.; Terlizzi, L.; Terrevoli, C.; Tersimonov, G.; Thakur, S.; Thomas, D.; Tieulent, R.; Tikhonov, A.; Timmins, A. R.; Tkacik, M.; Toia, A.; Topilskaya, N.; Toppi, M.; Torales-Acosta, F.; Torres, S. R.; Trifiró, A.; Tripathy, S.; Tripathy, T.; Trogolo, S.; Trombetta, G.; Trubnikov, V.; Trzaska, W. H.; Trzcinski, T. P.; Trzeciak, B. A.; Tumkin, A.; Turrisi, R.; Tveter, T. S.; Ullaland, K.; Uras, A.; Urioni, M.; Usai, G. L.; Vala, M.; Valle, N.; Vallero, S.; van der Kolk, N.; van Doremalen, L. V. R.; van Leeuwen, M.; Vande Vyvre, P.; Varga, D.; Varga, Z.; Varga-Kofarago, M.; Vargas, A.; Vasileiou, M.; Vasiliev, A.; Vázquez Doce, O.; Vechernin, V.; Vercellin, E.; Vergara Limón, S.; Vermunt, L.; Vértesi, R.; Verweij, M.; Vickovic, L.; Vilakazi, Z.; Villalobos Baillie, O.; Vino, G.; Vinogradov, A.; Virgili, T.; Vislavicius, V.; Vodopyanov, A.; Volkel, B.; Völkli, M. A.; Voloshin, K.; Voloshin, S. A.; Volpe, G.; von Haller, B.; Vorobyev, I.; Voscek, D.; Vrláková, J.; Wagner, B.; Wang, C.; Wang, D.; Weber, M.; Wegrynek, A.; Wenzel, S. C.; Wessels, J. P.; Wiechula, J.; Wikne, J.; Wilk, G.; Wilkinson, J.; Willems, G. A.; Willsher, E.; Windelband, B.; Winn, M.; Witt, W. E.; Wright, J. R.; Wu, W.; Wu, Y.; Xu, R.; Yalcin, S.; Yamaguchi, Y.; Yamakawa, K.; Yang, S.; Yano, S.; Yin, Z.; Yokoyama, H.; Yoo, I. -K.; Yoon, J. H.; Yuan, S.; Yuncu, A.; Zaccolo, V.; Zaman, A.; Zampolli, C.; Zanolli, H. J. C.; Zardoshti, N.; Zarochentsev, A.; Závada, P.; Zaviyalov, N.; Zbroszczyk, H.; Zhalov, M.; Zhang, S.; Zhang, X.; Zhang, Y.; Zherebchevskii, V.; Zhi, Y.; Zhou, D.; Zhou, Y.; Zhu, J.; Zhu, Y.; Zichichi, A.; Zinovjev, G.; Zurlo, N.. - In: PHYSICS LETTERS. SECTION B. - ISSN 0370-2693. - ELETTRONICO. - 827:(2022), pp. 1-15. [10.1016/j.physletb.2022.136984]



Production of Λ and K_S^0 in jets in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and pp collisions at $\sqrt{s} = 7$ TeV



ALICE Collaboration*

ARTICLE INFO

Article history:

Received 8 September 2021

Received in revised form 11 February 2022

Accepted 18 February 2022

Available online 25 February 2022

Editor: M. Doser

ABSTRACT

The production of Λ baryons and K_S^0 mesons (V^0 particles) was measured in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and pp collisions at $\sqrt{s} = 7$ TeV with ALICE at the LHC. The production of these strange particles is studied separately for particles associated with hard scatterings and the underlying event to shed light on the baryon-to-meson ratio enhancement observed at intermediate transverse momentum (p_T) in high multiplicity pp and p-Pb collisions. Hard scatterings are selected on an event-by-event basis with jets reconstructed with the anti- k_T algorithm using charged particles. The production of strange particles associated with jets $p_{T,jet}^{ch} > 10$ and $p_{T,jet}^{ch} > 20$ GeV/c in p-Pb collisions, and with jet $p_{T,jet}^{ch} > 10$ GeV/c in pp collisions is reported as a function of p_T . Its dependence on angular distance from the jet axis, $R(V^0, jet)$, for jets with $p_{T,jet}^{ch} > 10$ GeV/c in p-Pb collisions is reported as well. The p_T -differential production spectra of strange particles associated with jets are found to be harder compared to that in the underlying event and both differ from the inclusive measurements. In events containing a jet, the density of the V^0 particles in the underlying event is found to be larger than the density in the minimum bias events. The Λ/K_S^0 ratio associated with jets in p-Pb collisions is consistent with the ratio in pp collisions and follows the expectation of jets fragmenting in vacuum. On the other hand, this ratio within jets is consistently lower than the one obtained in the underlying event and it does not show the characteristic enhancement of baryons at intermediate p_T often referred to as "baryon anomaly" in the inclusive measurements.

© 2022 European Organization for Nuclear Research, ALICE. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

High-energy heavy-ion collisions provide a unique opportunity to study properties of the hot and dense medium composed of deconfined partons, known as the quark-gluon plasma (QGP) [1–6]. A cross-over transition from hadronic matter to the QGP at zero baryochemical potential is expected to take place once the temperature reaches values of about $T_c = 156$ MeV based on quantum chromodynamics (QCD) calculations performed on a lattice [7–9]. The measurements indicate that collisions of lead ions at the Large Hadron Collider (LHC) at a centre-of-mass energy per nucleon-nucleon collision of $\sqrt{s_{NN}} = 2.76$ TeV create conditions well above T_c at approximately zero baryochemical potential [10].

The interpretation of nucleus-nucleus (AA) collision results requires the understanding of results from smaller collision systems such as proton-proton (pp) or proton-nucleus (pA). To separate initial state effects, linked to the use of nuclear beams or targets, from final-state effects, associated with the presence of hot and dense matter, particle production is compared in pp,

pA, and AA reactions. However, the measurements at the LHC in high-multiplicity pp and p-Pb collisions have revealed unexpectedly strong long-range correlations of produced particles typical of Pb-Pb collisions [11–21]. Measurements of identified light-flavour hadrons [22–25], strange particles [26–29], and heavy-flavour particles [30,31] in small systems have also shown qualitatively similar features as in AA collisions [22,32–36]. In particular, the baryon-to-meson yield ratio as a function of transverse momentum (p_T) shows a pronounced maximum at intermediate p_T (2–5 GeV/c) [23,26,28]. The p_T dependence of the ratio was discussed in terms of particle production within a common velocity field (collective flow) [37], soft-hard parton recombination [38] and high-energy parton shower (jet) hadronization at high p_T . On the other hand, the jet suppression ascribed to the parton energy loss in the QGP observed in central AA collisions is not observed in p-Pb collisions [39–50]. The measurements show that the impact of the initial-state nuclear effects such as shadowing and potential gluon saturation effects, e.g., Color Glass Condensate (CGC) [51,52], or multiple scatterings and hadronic re-interactions in the initial and final states [53,54] is small on the jet production in p-Pb collisions. To understand particle production mechanisms in small systems, the separation of particles produced in hard processes

* E-mail address: alice-publications@cern.ch.

(jets) from those produced in the underlying event is important. It allows one to investigate similarities and expose differences in particle production mechanisms in high-multiplicity pp, p-Pb events, and heavy-ion collisions.

In this letter, measurements of K_S^0 and Λ ($\bar{\Lambda}$), the V^0 particles, in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and pp collisions at $\sqrt{s} = 7$ TeV are reported. The production of V^0 particles is studied separately within the region associated with a hard parton scattering and the underlying event. Hard scatterings are tagged by selecting a reconstructed jet with transverse momentum $p_{T,jet}^{ch} > 10$ or 20 GeV/c using charged particles with the anti- k_T algorithm [55] and the resolution parameter $R = 0.4$. The baryon-to-meson ratio of V^0 particles associated with jets is reported as a function of particle transverse momentum and distance to the jet axis. To contrast the strangeness production associated with a hard scattering and subsequent jet fragmentation with the production in the underlying event we report the ratio for the case of particles not associated with jets. The p_T -differential ratio is also compared with a Pythia 8 (version 8.2.43; Tune 4C) [56] simulation.

2. Data analysis

2.1. The ALICE detector and data sample

The ALICE apparatus consists of central barrel detectors covering the pseudorapidity interval $|\eta_{lab}| < 0.9$, a forward muon spectrometer covering $-4.0 < \eta_{lab} < -2.5$, and a set of detectors at forward and backward rapidities used for triggering and event characterization. Further information can be found in Ref. [57]. Tracking and particle identification in the context of this analysis are performed using the information provided by the Inner Tracking System (ITS) [58] and the Time Projection Chamber (TPC) [59], which have full azimuthal coverage in the pseudorapidity interval $|\eta_{lab}| < 0.9$. These central barrel detectors are located inside a large solenoidal magnet, which provides a magnetic field of 0.5 T along the beam direction (z-axis in the ALICE reference frame). The ITS is composed of six cylindrical layers of silicon detectors, with radial distances from the beam axis ranging from 3.9 cm to 43.0 cm. The two innermost layers are equipped with Silicon Pixel Detectors (SPD) covering the pseudorapidity ranges of $|\eta_{lab}| < 2.0$ and $|\eta_{lab}| < 1.4$, respectively. The two intermediate layers are made of Silicon Drift Detectors (SDD), while Silicon Strip Detectors (SSD) equip the two outermost layers. The high spatial resolution of the silicon sensors, together with the low material budget (on average 7.7% of a radiation length for tracks crossing the ITS perpendicularly to the detector surfaces, i.e., $\eta_{lab} = 0$) and the small distance of the innermost layer from the beam pipe, allow for the measurement of the track impact parameter d_{DCA} in the transverse plane. The d_{DCA} is defined by the distance of closest approach (DCA) of the track to the primary vertex in the plane transverse to the beam direction, and is measured with a resolution better than 75 μm for transverse momenta $p_T > 1$ GeV/c, including the contribution from the primary vertex position resolution [58]. At larger radii ($85 < r < 247$ cm), the 500 cm long cylindrical TPC provides track reconstruction with up to 159 three-dimensional space points per track, as well as particle identification via the measurement of the specific energy deposit dE/dx in the gas. The overall p_T resolution given by combining ITS and TPC information is typically 1% for momenta of 1 GeV/c and 7% for momenta of 10 GeV/c [60].

The data sample used in this analysis was recorded by the ALICE detector [57] during the LHC p-Pb run at $\sqrt{s_{NN}} = 5.02$ TeV and pp run at $\sqrt{s} = 7$ TeV in 2013 and 2010, respectively. Because of the 2-in-1 magnet design of the LHC [61], the energies of the two beams are not independent and their ratio is fixed to be equal to the ratio of the charge-to-mass ratios of each beam. Consequently, for p-Pb collisions, the nucleon-nucleon centre-of-mass

system is shifted in rapidity by $\Delta y_{NN} = 0.465$ in the direction of the proton beam. In the analyzed data sample the Pb beam circulated in the “counter-clockwise” direction travelling from negative to positive rapidity in the laboratory reference frame. The setup of the detector, trigger, and the analysis strategy is identical in both collision systems unless explicitly stated otherwise.

The data samples presented in this letter were recorded using the minimum bias trigger implemented by the VZERO detector [62]. The VZERO system consists of two arrays of 32 scintillator tiles each, placed around the beam vacuum pipe on either side of the interaction region covering the pseudorapidity intervals $2.8 < \eta_{lab} < 5.1$ (VZERO-A) and $-3.7 < \eta_{lab} < -1.7$ (VZERO-C). In addition, in p-Pb collisions, two neutron Zero Degree Calorimeters (ZDCs), located at +112.5 m (ZNA) and -112.5 m (ZNC) from the interaction point, are used in the offline event selection for rejecting of beam-background events, exploiting the correlation between the arrival times measured in ZNA and ZNC. In pp collisions, a logical OR between the requirement of at least one hit in the SPD and a hit in one of the two VZERO scintillator arrays is used for event selection. In p-Pb collisions, a coincidence of signals in both VZERO-A and VZERO-C is required to remove contamination from single-diffractive and electromagnetic events [63]. The events are further selected to require a reconstructed vertex within 10 cm ($|v_z| < 10$ cm) of the nominal centre of the detector along the beam axis and vertices built from the SPD tracklets, which are the short track segments measured with SPD, and from the tracks measured with combined information from ITS and TPC are compatible. The fraction of events with the vertex selection criteria is about 98.2% of all triggered events. In total, about 96×10^6 (177×10^6) events, corresponding to an integrated luminosity of $\mathcal{L} \approx 46 \mu\text{b}^{-1}$ (2.9 nb^{-1}), are used in the analysis of the p-Pb (pp) data sample.

2.2. Charged-particle and jet reconstruction

The charged-particle reconstruction and jet reconstruction in this letter follow the approach described in detail in Refs. [44,64]. Here only a brief review of the most relevant points is given. Charged-particle tracks, reconstructed in the ITS and the TPC with $p_T > 0.15$ GeV/c and within the TPC acceptance $|\eta_{lab}| < 0.9$ that satisfy a DCA requirement $d_{DCA} < 2.4$ cm, are used as input to the jet reconstruction. The azimuthal distribution of these tracks is not completely uniform due to inefficient regions in the SPD. This is compensated by considering in addition tracks with less than three reconstructed track points in the ITS or no points in the SPD. To improve the momentum resolution for those tracks, the primary vertex is used as an additional constraint in the track fitting. This approach yields a uniform tracking efficiency within the acceptance. These complementary tracks constitute approximately 4.3% and 5% of the overall used track sample in p-Pb and pp collisions, respectively. The efficiency for charged-particle detection, including the effect of tracking efficiency as well as the geometrical acceptance, is 70% (60%) at $p_T = 0.15$ GeV/c and increases to 85% (87%) at $p_T = 1$ GeV/c and above for p-Pb (pp) collisions.

The jets are reconstructed using the anti- k_T algorithm [55] from the FastJet package [65,66] with resolution parameter $R = 0.4$. Only those jets for which the jet-axis is found within the acceptance window $|\eta_{lab}| < 0.35$ are used in this analysis. This condition ensures the jet cone is fully overlapping with the acceptances of both charged-particle tracks ($|\eta_{lab}| < 0.9$) and the V^0 particles ($|\eta_{lab}| < 0.75$, as explained in detail in section 2.3). The jet transverse momentum is calculated with FastJet using the p_T recombination scheme.

In general, the transverse-momentum density of the background (ρ^{ch}), originating from the underlying event and/or pile-up, contributes to the jet energy reconstructed by the jet finder. The

correction of the jet-energy scale accounting for the background contribution can be estimated on an event-by-event basis using the median of the transverse momentum density of all the clusters reconstructed with the k_T algorithm [67]. In pp and p-Pb collisions, an estimate adequate for the more sparse environment than Pb-Pb collisions is employed by scaling ρ^{ch} with an additional factor to account for event regions without particles [44]. The resulting mean of the background p_T density in p-Pb collisions is $\langle \rho^{\text{ch}} \rangle = 1.02 \text{ GeV}/c \text{ rad}^{-1}$ (with negligible statistical uncertainty) for unbiased events and $\langle \rho^{\text{ch}} \rangle = 2.2 \pm 0.01 \text{ GeV}/c \text{ rad}^{-1}$ for events containing a jet with uncorrected transverse momentum $p_{T,\text{jet}}^{\text{ch, raw}} > 20 \text{ GeV}/c$ [44]. In pp collisions, the background density is around $1 \text{ GeV}/c \text{ rad}^{-1}$ and not subtracted on a jet-by-jet basis but the related uncertainty on the jet p_T scale is absorbed into the systematic uncertainty.

The jet finding efficiency, which encodes the effects of single-particle momentum resolution and reconstruction efficiency on the jet reconstruction, is estimated using a PYTHIA 6 [68] + GEANT 3 [69] simulation by comparing the generated jets to reconstructed ones and found to be larger than 96% in the considered momentum range ($p_{T,\text{jet}}^{\text{ch}} > 10 \text{ GeV}/c$).

2.3. Reconstruction of V^0 particles

The V^0 particles, K_S^0 and $\Lambda(\bar{\Lambda})$, are identified by taking advantage of the characteristics of their weak decay topologies in the channels $K_S^0 \rightarrow \pi^+ \pi^-$ and $\Lambda(\bar{\Lambda}) \rightarrow p \pi^-(p \pi^+)$, which have branching ratios of 69.2% and 63.9%, respectively [70]. The reconstruction and the selection criteria of the V^0 particles follow the analysis in Ref. [23] with the exception of the rapidity selection of the particles and their decay products. The decay products of the V^0 particles, π^\pm and $p(\bar{p})$, are identified in the central barrel with the TPC using the specific energy loss dE/dx in the gas by measuring up to 159 samples per track with a resolution of about 6% [71]. Since the V^0 daughter tracks are displaced from the primary vertex and tracks in the jet are selected by criteria optimized for particles produced at the primary vertex, only about 0.1% of the V^0 daughter tracks contribute to the charged-particle jet reconstruction. The V^0 decay daughter tracks are selected in the acceptance window $|\eta_{\text{lab}}| < 0.8$ following the criteria used in the inclusive analysis [23,72]. To avoid the fiducial effect, only the V^0 candidates found in $|\eta_{\text{lab}}| < 0.75$ are retained. This ensured that the reconstruction efficiency is approximately constant throughout the selected pseudorapidity range. The topological selection of V^0 candidates within the kinematic range of this analysis yields almost background-free invariant mass spectra with the lowest signal-to-background ratio among all of the V^0 particles still exceeding 10. The p_T -differential yields of the V^0 particles are extracted using the invariant-mass method, described in Ref. [23], where the combinatorial background is interpolated from the side bands defined in terms of the mass peak width σ in intervals $[-12\sigma, -6\sigma]$ and $[6\sigma, 12\sigma]$ with respect to the mean of the peak.

2.4. Matching of V^0 particles to jets and underlying event

To obtain the yield of V^0 particles within a jet cone, the V^0 particles are selected based on their distance from the jet centroid in the pseudorapidity (η_{lab}) and azimuthal angle (φ) plane

$$R(V^0, \text{jet}) = \sqrt{(\eta_{\text{lab}}^{\text{jet}} - \eta_{\text{lab}}^{V^0})^2 + (\varphi^{\text{jet}} - \varphi^{V^0})^2}. \quad (1)$$

A V^0 particle with a radial distance from a given jet $R(V^0, \text{jet}) < R_{\text{match}}$ is considered matched to the jet and referred to as the “ V^0 inside the jet cone” (JC V^0). In p-Pb collisions the probability for a

particle with $p_T > 0.5 \text{ GeV}/c$ to lie in the overlapping region of two different jets with $p_{T,\text{jet}}^{\text{ch}} > 10 \text{ GeV}/c$ is less than 1% and in these cases the higher-energy jet is preferred. Moreover, removal of the events with the same particle matching to two or more jets did not alter the result of the analysis. The procedure for extracting the yield of V^0 particles, associated with a jet within a cone defined by R_{match} , can be summarized as follows. For each p_T interval the JC V^0 yield is extracted using the invariant mass technique, where the combinatorial background is interpolated from the side bands. Then the raw JC V^0 yield is corrected for the contribution of particles from the underlying event (the UE V^0).

Conceptually, the UE V^0 particles represent the particles that are not associated with the hard scatterings tagged by the charged jets considered in this analysis. To extract the UE V^0 yield several estimators were investigated: i) an *outside cone* (OC) selection, composed of the V^0 particles that satisfy the condition of $R(V^0, \text{jet}) > R_{\text{cut}}$ (e.g. $R_{\text{cut}} = 0.6$) within events containing a jet; ii) the *perpendicular cone* (PC) selection, composed of the V^0 particles found in a range with radius $R = 0.4$ in η and φ space perpendicular to the jet axis at the same η ; and iii) the *non-jet event* (NJ) selection, composed of the V^0 particles found in events that do not contain a jet with $p_{T,\text{jet}}^{\text{ch}} > 5 \text{ GeV}/c$.

In practice, a useful quantity for performing the subtraction of the non-jet contribution of the V^0 particles is their density per unit area

$$\rho^{V^0}(p_T) = N^{V^0}(p_T)/A^{V^0}, \quad (2)$$

where N^{V^0} is the number of V^0 particles and A^{V^0} is the acceptance in pseudorapidity and azimuthal angle. Consequently, the number of the UE V^0 particles within a jet cone can be calculated as $N = \rho^{V^0} A_{\text{jet}}$ for each estimator separately. The jet area $A_{\text{jet}} = \pi R_{\text{match}}^2$ is considered in this analysis. In general the density of V^0 particles within jets can be defined as

$$\rho_{\text{JC}}^{V^0} - \rho_{\text{UE}}^{V^0}, \quad (3)$$

where UE can be any of the OC, PC, or NJ background estimators. In this analysis, PC is chosen as the default background estimator, while OC and NJ are used to quantify the systematic uncertainty.

2.5. Corrections for finite V^0 reconstruction efficiency and feed-down

The reconstruction efficiencies of V^0 particles are estimated using the DPMJET [73] and PYTHIA 6 [68] Monte Carlo generators in p-Pb and pp collisions, respectively, with the same selection criteria as in the data except the daughter track particle identification with dE/dx in the TPC (see more details in [23]). These simulations are based on the GEANT 3 transport code [69] for the detector description and response.

Due to differences in the experimental acceptance for V^0 particles associated with jets and those extracted through the various estimators of the underlying event, the efficiencies of V^0 particles are estimated separately for every case. Fig. 1 shows the reconstruction efficiencies for inclusive V^0 s and those for JC V^0 s with $R(V^0, \text{jet}) < 0.4$ and UE V^0 s. The UE V^0 s are estimated with the OC estimator with $R(V^0, \text{jet}) > 0.6$ in p-Pb collisions and with the PC estimator in pp collisions. In particular, for $R(V^0, \text{jet}) < 0.4$ the efficiency at $p_T < 2 \text{ GeV}/c$ is about 20% larger than in the inclusive case while it approaches the inclusive case at higher p_T . This is due to the fact that the η -differential reconstruction efficiency of V^0 particles decreases with $|\eta_{\text{lab}}|$ and the pseudorapidity distribution of V^0 particles matched with jets is narrower than that of inclusive ones. This results in a higher η -integrated efficiency of JC V^0 s than inclusive V^0 s. This effect is more pronounced at low p_T .




Fig. 1. Reconstruction efficiency of V^0 particles in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV (left panel) and in pp collisions at $\sqrt{s} = 13$ TeV (right panel) for three selection criteria: inclusive, within $R(V^0, \text{jet}) < 0.4$ and V^0 's in UE (upper panels) and the ratio relative to inclusive selection (lower panels). UE V^0 's are estimated with the OC estimator ($R(V^0, \text{jet}) > 0.6$) in p-Pb collisions and with the PC estimator in pp collisions.

Table 1

Relative systematic uncertainties in percent for K_S^0 , Λ , and $\bar{\Lambda}$ in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. The right three columns in the last two rows represent the uncertainties of $\Lambda + \bar{\Lambda}$. For each particle, the reported values correspond to the uncertainties at $p_T = 0.6$, 2, and 10 GeV/c. See text for details.

	K_S^0	Λ		$\bar{\Lambda}$	
Particle identification		< 1	negl.	3.2	negl.
Track selection	negl.	negl.	1.5	negl.	1
Topological selection	0.3	1.1	0.7	0.7	0.6
Proper lifetime		negl.	2.9	3.7	negl.
Competing V^0 selection		< 1	negl.	3.4	negl.
Signal extraction	1.8	4.2	2.4	2.1	1.8
Jet p_T scale	1.4	5.4	37.1	3.2	4.2
UE subtraction	21	9.6	1.2	40	24
					0.6

The p_T -differential yields of Λ and $\bar{\Lambda}$ reconstructed for JC and UE selections are also corrected for the feed-down from the decays of Ξ^0 and Ξ^- particles and their respective anti-particles. The Ξ production in jets is estimated based on measurements of the multi-strange baryons and their decays at high p_T performed in pp collisions [74] and extrapolated to low p_T using the PYTHIA 8 event generator. The applied correction amounts to 15% and is independent of the Λ and $\bar{\Lambda}$ momenta. Conversely, the Λ yields are not corrected for the feed-down from Ω^- baryons as this contribution is negligible compared to the systematic uncertainties of the present measurement. Since Λ from non-weak decays of the Σ^0 and $\Sigma^*(1385)$ family cannot be distinguished from the direct ones, the identified Λ yield includes these contributions [72].

2.6. Systematic uncertainties

The main sources of systematic uncertainty in the V^0 particle reconstruction are uncertainties on the material budget (4%), the track selection (up to 4%), feed-down correction for the Λ (5% for $p_T < 3.7$ GeV/c and 7% for $p_T > 3.7$ GeV/c), proper lifetime selection criteria (up to $\sim 4\%$), and topological selections depending on transverse momentum and particle species (up to 1.6%). The systematic uncertainties on the extracted yields for K_S^0 mesons and Λ and $\bar{\Lambda}$ baryons in p-Pb and pp collisions are reported as point-to-point uncertainties in Table 1 and Table 2. The “negl.” in the table denotes an uncertainty of less than 0.1%. The total uncertainty on the yields is calculated by adding the individual uncertainties on track selection, material budget, feed-down corrections and the listed V^0 selections in quadrature.

Particle identification (PID). The uncertainty due to the particle identification is estimated by varying the selection criteria of the dE/dx in the TPC from a default 5σ to 4, 6 and 7 standard deviations from the nominal dE/dx for pions and protons normalized to the detector resolution.

Track selection. The uncertainty originating from the track selection is estimated by repeating the analysis with an increased number of required TPC space points per track by about 7% and 15% from the nominal requirement of 70 points.

Topological selection. The uncertainty associated with the topological selection of the V^0 candidates (the two-dimensional decay radius, daughter track DCA to primary vertex, DCA of V^0 daughters, and cosine of the pointing angle) is obtained by varying the parameters of the selections for each of the V^0 species separately as described in detail in Ref. [23].

Proper lifetime selection. The uncertainty due to the selection on the proper lifetime of V^0 candidates, defined as the product of the mass m_0 , decay length L , and the inverse of the particle momentum p ($m_0 L c / p < 20$ cm for K_S^0 and $m_0 L c / p < 30$ cm for Λ and $\bar{\Lambda}$), is obtained by redoing the analysis with different selection criteria (12 and 40 cm for K_S^0 and 20 and 40 cm for Λ and $\bar{\Lambda}$).

Competing V^0 selection. The invariant mass of each candidate can be calculated either under the K_S^0 or the Λ ($\bar{\Lambda}$) mass hypothesis. A K_S^0 candidate is rejected if its invariant mass under the hypothesis of a Λ or $\bar{\Lambda}$ lies in the window of ± 10 MeV/ c^2 around the mass of the Λ or $\bar{\Lambda}$, and a Λ ($\bar{\Lambda}$) candidate is rejected if its invariant mass under the K_S^0 hypothesis lies in the window of ± 5 MeV/ c^2 around the K_S^0 mass. To assess the uncertainty related to this selection the analysis is repeated varying the invariant mass

Table 2

Relative systematic uncertainties in percent for K_S^0 , Λ , and $\bar{\Lambda}$ in pp collisions at $\sqrt{s} = 7$ TeV. The right three columns in the last two rows represent the uncertainties of $\Lambda + \bar{\Lambda}$. For each particle, the reported values correspond to the uncertainties at $p_T = 0.6, 2$ and 10 GeV/c. See text for details.

	K_S^0			Λ			$\bar{\Lambda}$		
Particle identification	negl.	negl.	2.3	negl.	negl.	5	negl.	negl.	2.5
Track selection	1.8	3.6	2.8	1.7	4	3.6	1.5	4.3	2.5
Topological selection	2.4	negl.	negl.	3.5	0.7	5.1	3.7	1.5	2.5
Proper lifetime		negl.		2.6	3	negl.	2.7	2.5	negl.
Competing V^0 selection		< 1		negl.	negl.	9.5	negl.	negl.	10.6
Signal extraction	negl.	negl.	1.4		< 1		negl.	negl.	1.3
Jet p_T scale	0.5	1	9.9			5.3	2	10	
UE subtraction	4.8	3.6	negl.			8.9	11.2	negl.	

Table 3

Relative systematic uncertainties in percent for the $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio of the spectrum of K_S^0 and Λ ($\bar{\Lambda}$) for $p_{T,jet}^{ch} > 10$ and 20 GeV/c in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, and for $p_{T,jet}^{ch} > 10$ GeV/c in pp collisions at $\sqrt{s} = 7$ TeV. For each case, the reported values correspond to the uncertainties at $p_T = 0.6, 2$, and 10 GeV/c.

	p-Pb			pp		
	$p_T > 10$ GeV/c	$p_T > 20$ GeV/c	$p_T > 10$ GeV/c	$p_T > 20$ GeV/c	$p_T > 10$ GeV/c	$p_T > 20$ GeV/c
V^0 reconstruction	8.3	8.9	9.2	8.8	8.1	9.2
Jet p_T scale	1.5	4.2	3.3	9.2	2.4	7.4
UE subtraction	26.3	10.3	negl.	21.1	7.1	negl.
					7.5	8.2
						negl.

window of 3 and 6 MeV/c² for K_S^0 and with no rejection for Λ ($\bar{\Lambda}$) baryons.

Underlying event subtraction. Two main sources of uncertainties originating from the mis-association of V^0 particles with the UE are considered: i) the V^0 particle is found outside the selected jet and is classified as an UE particle; however, it may have originated from a physical jet outside the fiducial acceptance of jets considered in the analysis and/or from a *true* low- p_T jet, below the considered thresholds; and ii) the V^0 particle originates from a true high- p_T jet; however, due to the finite detector efficiency the jet has not been reconstructed above the considered p_T threshold.

The uncertainty on the UE V^0 density is estimated using the OC and NJ selections as alternatives for the density calculation, since the former is sensitive to particles outside the jet cone but originating from a physical jet and the latter is sensitive to those signals contributing to the UE due to the finite detector efficiency. The standard deviation of the difference of the reconstructed V^0 yields in OC and NJ is included as an additional systematic uncertainty on the density of particles within the jets. In p-Pb collisions the uncertainty is largest for low-momentum particles ($p_T < 2$ GeV/c) reaching up to 20% (40%) for K_S^0 (Λ) but drops rapidly with p_T to negligible values for $p_T > 6$ GeV/c. For pp collisions the trend of the uncertainty is similar to the trend seen in p-Pb, however the magnitude is smaller, reaching values up to 5% (9%) for K_S^0 (Λ).

Jets p_T scale. The systematic uncertainty originating from the selection of the jet p_T is estimated by varying the jet p_T around the chosen thresholds of 10 and 20 GeV/c by 2 GeV/c. This variation accounts for jet resolution effects due to detector effects and the fluctuations of the event background density as reported in Ref. [44]. For jets with $p_{T,jet}^{ch} > 10$ GeV/c at low momenta ($p_{T,V^0} < 2$ GeV/c) it reaches up to 10%, while it is about 20% for jets of $p_{T,jet}^{ch} > 20$ GeV/c. It remains almost constant at about 3% for $p_{T,V^0} > 2$ GeV/c for jets $p_{T,jet}^{ch} > 10$ GeV/c and about 5% for jets $p_{T,jet}^{ch} > 20$ GeV/c.

Uncertainty of the $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio. The uncertainties on V^0 yields, material budget and feed-down correction are propagated to the ratio quadratically. The uncertainties related to the jet p_T and UE estimation are obtained by calculating the deviation of ratios between the default analysis and various selection criteria. Table 3 shows the point-to-point relative systematic uncertain-

ties on the $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio reconstructed within $R = 0.4$ jets with $p_{T,jet}^{ch} > 10$ GeV/c (left column) and $p_{T,jet}^{ch} > 20$ GeV/c (middle column) in p-Pb collisions. For $p_{T,jet}^{ch} > 20$ GeV/c, the total uncertainty is about 16% and is largely independent of particle p_T with the largest contribution of 8–9% originating from the uncertainty on the V^0 reconstruction. The relative systematic uncertainties on the $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio for pp collisions are shown in the right column in Table 3.

3. Results

In the following, results for V^0 particles with four different selections are discussed. Their labels in the figures are defined as follows: i) V^0 s obtained from the unbiased events without any jet veto are labelled as “Inclusive” particles; ii) V^0 s matched to jets in a cone with a radius of 0.4 are labelled as particles within “ $R(V^0, jet) < 0.4$ ”, the remaining underlying event background is not subtracted from this sample; iii) the label “ V^0 s in jets” refers to V^0 s produced in jets obtained by subtracting the underlying event background from the previous sample; iv) V^0 s from the underlying event estimated in cones perpendicular to the jet axis are labelled as “Perp. cone” particles.

The fully corrected densities of K_S^0 and the sum of Λ and $\bar{\Lambda}$ particles associated with a hard scattering, tagged by a jet, are shown in Figs. 2 and 3 for p-Pb and pp collisions, respectively. The per-jet density of V^0 particles within jets is compared with that of inclusive particles (irrespective of their association with a hard scattering) and with underlying event V^0 s obtained using the PC selection. In the case of inclusive particles the distribution is normalized to the product of the total number of events and the acceptance of the V^0 particles in a single event (full azimuth and $|\eta_{lab}| < 0.75$). As expected, the p_T dependence of the density of both K_S^0 and Λ particles within jets, as defined by Eq. (3), is considerably less steep than in the case of inclusive particles. The density distribution of inclusive V^0 s is lower than that of the PC selection since the latter are obtained from events contain jets with $p_{T,jet}^{ch} > 10$ GeV/c. But the density distribution of the PC selection shows a strong, steeply falling p_T dependence with respect to the inclusive one. Both the inclusive and the PC distributions show a rapid decrease with p_T , reaching values more than an order




Fig. 2. The p_T -differential density of particles $d\rho/V^0/dp_T$ (see Eq. (2)) in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV for K_S^0 (upper panel) and the sum of Λ and $\bar{\Lambda}$ (lower panel). The density is shown for three selection criteria: inclusive particles from minimum bias events (black full circle), particles associated with the underlying event production estimated with PC selection (blue open circle, labelled as “Perp. cone” in the figure), JC V^0 s with $R(V^0, \text{jet}) < 0.4$ (green full square). The density distribution of V^0 s in jets with UE background subtracted (defined by Eq. (3)) is shown as the red full triangle. Statistical uncertainties and systematic uncertainties are shown as vertical bars and open boxes, respectively.

of magnitude lower than the JC density for particle p_T exceeding $4 \text{ GeV}/c$. This is consistent with the expectation that the high- p_T particles originate from jet fragmentation.

Ratios of Λ and K_S^0 yields can be obtained by dividing the normalized density distributions. Here, the sum of the Λ and $\bar{\Lambda}$ densities is divided by twice the density of K_S^0 . Fig. 4 shows the ratio for the JC selection (without the UE background subtraction) as a function of the distance from the jet axis $R(V^0, \text{jet})$ in p-Pb collisions. The ratio is shown for three p_T intervals: low p_T ($0.6 < p_T < 1.8 \text{ GeV}/c$), intermediate p_T ($2.2 < p_T < 3.7 \text{ GeV}/c$), and high p_T ($4.2 < p_T < 12 \text{ GeV}/c$). The sources of the systematic uncertainties (open boxes) are summarized in Table 3. The uncertainty on V^0 yields extraction is uncorrelated with V^0 p_T but correlated with $R(V^0, \text{jet})$; the uncertainties on jet p_T scale and on UE subtraction are uncorrelated on both V^0 p_T and $R(V^0, \text{jet})$. The ratio as a function of $R(V^0, \text{jet})$ at low p_T , dominated by the UE contribution, is approximately constant at about 0.2. It is independent of the distance to the jet axis even at large distances of $R(V^0, \text{jet}) > 1.2$. This value is consistent with the inclusive measurements in p-Pb collisions, but also in pp and peripheral Pb-Pb collisions where effects related to the collective expansion of the system are either not present or small [24].

Conversely, the intermediate- p_T selection shows an increase of the ratio from about 0.3 when evaluated close to the jet axis to values of about 0.6 at $R(V^0, \text{jet})$ distances of about 0.5. For distances $R(V^0, \text{jet}) > 0.5$ the ratio remains constant. The ratio of 0.6 is consistent with the inclusive measurement in p-Pb collisions




Fig. 3. The p_T -differential density of particles $d\rho/V^0/dp_T$ (see Eq. (2)) in pp collisions at $\sqrt{s} = 7$ TeV for K_S^0 (upper), and the sum of Λ and $\bar{\Lambda}$ (lower). The density is shown for four selection criteria with the same definitions as Fig. 2.




Fig. 4. The $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV as a function of $R(V^0, \text{jet})$ for three different V^0 -particle p_T intervals associated with charged jets with $p_{T,\text{jet}}^{\text{ch}} > 10 \text{ GeV}/c$. The data points of the ratios in $0.6 < p_{T,V^0} < 1.8 \text{ GeV}/c$ and in $4.2 < p_{T,V^0} < 12 \text{ GeV}/c$ are shifted to the left and right sides from the centre, along the $R(V^0, \text{jet})$ -axis for better visibility. Statistical uncertainties (vertical bars) and systematic uncertainties (open boxes) are shown. The sources of the systematic uncertainty are summarized in Table 3. The uncertainty on V^0 yields extraction is uncorrelated with V^0 p_T but correlated with $R(V^0, \text{jet})$, the uncertainties on jet p_T scale and on UE subtraction are uncorrelated on both V^0 p_T and $R(V^0, \text{jet})$.

[23] and this p_T region is where the enhanced $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio in the inclusive measurements is found to be the largest. It is worthwhile to stress that for the results shown in Fig. 4, the UE backgrounds are not subtracted. Therefore, the evolution of the ratio as a function of the distance from the jet axis demonstrates how the two sources, UE and jet, compete. The lack of enhance-




Fig. 5. The $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV (upper panel) and pp collisions at $\sqrt{s} = 7$ TeV (lower panel) as a function of V^0 -particle p_T , associated with charged jets with $p_{T,jet}^{ch} > 10$ GeV/c (for both pp and p-Pb collisions) and 20 GeV/c (for p-Pb collisions only) together with that in inclusive and PC selection, and JC selection in case of pp collisions. The systematic uncertainties (open boxes) are fully uncorrelated with p_T . In both upper and lower panels, the black dashed curves are the results for inclusive V^0 s from PYTHIA 8 simulations. The jet selection within PYTHIA 8 is made using the generator level information with $p_{T,jet}^{ch} > 10$ GeV/c shown as the red curves.

ment close to the jet axis indicates that the enhanced $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio is not associated with jets.

In each p_T interval the ratio is dominated by the lower side of the selection window due to the steeply falling particle p_T spectrum. This is especially the case for $4.2 < p_T < 12$ GeV/c where the dominating component originates from p_T of about 4.5 GeV/c and the $R(V^0, jet)$ dependence at high p_T is similar to that for $2.2 < p_T < 3.7$ GeV/c. The ratio at high p_T associated with jets is discussed below.

Fig. 5 shows the ratio of Λ to K_S^0 as a function of particle p_T in both pp and p-Pb collisions for the different selection criteria. The systematic uncertainties (open boxes) are fully uncorrelated with p_T . In the case of p-Pb collisions, the ratio of the inclusive particles, the particles from the PC selection, and for those within jet with resolution parameter $R = 0.4$ and $p_{T,jet}^{ch} > 10$ and > 20 GeV/c are shown. Prior to forming the ratio, the UE density contribution obtained with the PC selection is subtracted for each particle species separately. Additionally, the p-Pb results are shown for the case where every V^0 particle is required to be close to the jet axis with its distance $R(V^0, jet) < 0.4$. The inclusive and the PC distributions show the enhancement at a p_T of about 3 GeV/c. The measurement of the inclusive case differs from that in Ref. [23] as the region $|\eta_{lab}| < 0.75$ is used here instead of the rapidity region in centre-of-mass frame $0 < y_{CMS} < 0.5$. The two measurements are otherwise consistent with each other. The PC distribution above 2 GeV/c reaches systematically higher values than the inclusive.

The ratio within jets is consistently lower than the inclusive one and approximately independent of p_T beyond 2 GeV/c. In particular, for particles associated with the jet it does not show a maximum at intermediate p_T . Clearly the enhancement of the ratio seen in the inclusive measurement is not present within jets. This conclusion holds not only for jets with $p_T > 10$ GeV/c but also for higher p_T (> 20 GeV/c) jets.

The results for pp collisions shown in Fig. 5 are obtained with jets reconstructed with $R = 0.4$ and for the same value of the matching radius $R(V^0, jet) < 0.4$. Apart from the inclusive particle selection and UE selection, the figure shows the ratio for particles within jets for the UE subtracted in the JC and UE unsubtracted case, demonstrating the small magnitude of background effects. Qualitatively similar features of the ratio are seen in both collision systems.

Selecting hard scatterings according to the jet energy carried exclusively by the primary charged particles induces biases and inefficiencies in the selection of the parton showers. The bias is related to the probabilistic process of fragmentation and hadronization. The analysis presented here tags only parton showers fragmenting into a configuration of hadrons that produce a charged particle jet with $p_{T,jet}^{ch} > 10$ GeV/c with a given R with a finite efficiency. Therefore, there can be cases of V^0 particles that originated from a parton shower but are rejected in the analysis based on the energy carried only by the primary charged particles. The same analysis performed using the PYTHIA 8 event generator shows that the most probable p_T of the full jet with $R = 0.4$ is larger by about 40% as compared to the $p_{T,jet}^{ch}$. Moreover, since the daughters of the V^0 particles are not included in the jet energy calculation there are cases of jets containing V^0 particles but not included in the JC selection. On the other hand, Fig. 5 shows that the inclusive $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio at high p_T is fully consistent with the ratio from particles associated with jets in this analysis. This suggests that the conclusion on the absence of the baryon-to-meson enhancement in jets made with the charged jets alone holds for all energetic parton showers and hadron configurations within jets.

Fig. 5 shows also the results compared with those obtained with the PYTHIA 8 [56] event generator with tune 4C (the dashed curves) run for pp collisions at $\sqrt{s} = 5.02$ TeV (top panel) and $\sqrt{s} = 7$ TeV (bottom panel). The comparison shows that the characteristic maximum at intermediate p_T in the inclusive ratio is not reproduced by the generator. However, for both collision systems the ratio within jets after the subtraction of the underlying event is consistent with the data points within uncertainties for $p_T > 6$ GeV/c. Note that PYTHIA 8 was chosen here merely as an example and the aim is not for a thorough review of the strangeness production in the Monte Carlo generators. The comparison with experimental data is found to be sufficient to demonstrate the clear similarities of the baryon-to-meson ratio within jets.

Fig. 6 shows the comparison of the ratio obtained in jets in pp and p-Pb collisions for the same selection of the matching radius $R(V^0, jet) < 0.4$ in both systems. The ratio obtained in p-Pb collisions is systematically higher for $2 < p_T < 8$ GeV/c with respect to that in pp collisions. However, the difference between the two collision systems is less than 2σ . The deviation between pp and p-Pb collisions has to be studied with higher precision in the future.

4. Summary

The production of V^0 particles (Λ baryons and K_S^0 mesons) is measured separately for particles associated with hard scatterings, tagged by reconstructed charged-particle jets, and the underlying event in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV and pp collisions at $\sqrt{s} = 7$ TeV for the first time at the LHC. The p_T -differential density distributions of V^0 particles associated within jets are com-




Fig. 6. The $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio in pp collisions at $\sqrt{s} = 7$ TeV and in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV as a function of V^0 -particle p_T associated with charged particle jets with $p_{T,jet}^{ch} > 10$ GeV/c reconstructed using the anti- k_T jet finder with resolution parameter $R = 0.4$. The ratio is shown for the same selection of the matching radius $R(V^0, jet) < 0.4$ in both systems. The systematic uncertainties (open boxes) are uncorrelated between the systems.

pared with those obtained from inclusive analysis and the underlying event. In both collision systems, the distribution of particles associated within jets is harder than that obtained in the underlying event since the high- p_T particles originate from jet fragmentation. The density of particles in the UE is larger than in the inclusive case as the former is obtained from events requiring a presence of a jet with $p_{T,jet}^{ch} > 10$ GeV/c. The $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio (without the UE subtracted) is studied as a function of $R(V^0, jet)$, defined as the distance between the jet axis and the V^0 particle, in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. At intermediate p_T , the ratio increases with $R(V^0, jet)$ from a value about 0.3 to 0.6 up to $R(V^0, jet) = 0.5$ reaching a constant value of about 0.6 for $R(V^0, jet) > 0.5$. This demonstrates that the enhanced $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio at intermediate- p_T observed in the inclusive analysis is not associated with jets since the underlying event contribution is more significant at larger $R(V^0, jet)$. The $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio associated with jets (with the UE subtracted) is consistent with the inclusive case within uncertainties for $p_T > 6$ GeV/c. The results in p-Pb collisions for $R(V^0, jet) < 0.4$ are consistent with the ratio measured in pp collisions. Finally, the enhancement in the $(\Lambda + \bar{\Lambda})/2K_S^0$ ratio at intermediate p_T found in the inclusive measurements in p-Pb and Pb-Pb collisions is not present for particles associated with hard scatterings tagged by jets reconstructed from charged particles for $p_{T,jet}^{ch} > 10$ GeV/c in p-Pb and pp collisions. As the baryon-to-meson enhancement (“baryon anomaly”) found in the inclusive measurements has been linked to the interplay of radial flow and parton recombination at intermediate p_T , its absence within the jet cone demonstrates that these effects are indeed limited to the soft particle production processes.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid

(WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Universidade Federal do Rio Grande do Sul (UFRGS), Brazil; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the Villum Fonden and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l'Énergie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy, Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; Indonesian Institute of Science, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Institute for Innovative Science and Technology, Nagasaki Institute of Applied Science (IIIST), Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Education and Science, National Science Centre and WUT ID-UB, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics and Ministry of Research and Innovation and Institute of Atomic Physics, Romania; Joint Institute for Nuclear Research (JINR), Ministry of Education and Science of the Russian Federation, National Research Centre Kurchatov Institute, Russian Science Foundation and Russian Foundation for Basic Research, Russia; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSDTA) and Office of the Higher Education Commission under NRU project of Thailand, Thailand; Turkish Atomic Energy Agency (TAEK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department

of Energy, Office of Nuclear Physics (DOE NP), United States of America.

References

- [1] E.V. Shuryak, Theory and phenomenology of the QCD vacuum. 7. Macroscopic excitations, Phys. Rep. 115 (1984) 151.
- [2] J. Cleymans, R. Gavai, E. Suhonen, Quarks and gluons at high temperatures and densities, Phys. Rep. 130 (1986) 217.
- [3] S. Bass, M. Gyulassy, H. Stoecker, W. Greiner, Signatures of quark-gluon plasma formation in high energy heavy-ion collisions: a critical review, J. Phys. G 25 (1999) R1–R57, arXiv:hep-ph/9810281 [hep-ph].
- [4] H. Satz, Color deconfinement in nuclear collisions, Rep. Prog. Phys. 63 (2000) 1511, arXiv:hep-ph/0007069 [hep-ph].
- [5] B.V. Jacak, B. Müller, The exploration of hot nuclear matter, Science 337 (6092) (2012) 310–314.
- [6] B. Müller, J. Schukraft, B. Wyslouch, First results from Pb+Pb collisions at the LHC, Annu. Rev. Nucl. Part. Sci. 62 (2012) 361–386, arXiv:1202.3233 [hep-ex].
- [7] S. Borsanyi, G. Endrodi, Z. Fodor, A. Jakovac, S.D. Katz, et al., The QCD equation of state with dynamical quarks, J. High Energy Phys. 1011 (2010) 077, arXiv: 1007.2580 [hep-lat].
- [8] T. Bhattacharya, M.I. Buchoff, N.H. Christ, H.-T. Ding, R. Gupta, et al., QCD phase transition with chiral quarks and physical quark masses, Phys. Rev. Lett. 113 (8) (2014) 082001, arXiv:1402.5175 [hep-lat].
- [9] P. Braun-Munzinger, V. Koch, T. Schäfer, J. Stachel, Properties of hot and dense matter from relativistic heavy ion collisions, Phys. Rep. 621 (Mar 2016) 76–126, <https://doi.org/10.1016/j.physrep.2015.12.003>.
- [10] ALICE Collaboration, J. Adam, et al., Direct photon production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 754 (2016) 235–248, arXiv:1509.07324 [nucl-ex].
- [11] CMS Collaboration, V. Khachatryan, et al., Observation of long-range near-side angular correlations in proton-proton collisions at the LHC, J. High Energy Phys. 1009 (2010) 091, arXiv:1009.4122 [hep-ex].
- [12] CMS Collaboration, S. Chatrchyan, et al., Observation of long-range near-side angular correlations in proton-lead collisions at the LHC, Phys. Lett. B 718 (2013) 795–814, arXiv:1210.5482 [nucl-ex].
- [13] ALICE Collaboration, B. Abelev, et al., Long-range angular correlations on the near and away side in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 719 (2013) 29–41, arXiv:1212.2001 [nucl-ex].
- [14] ATLAS Collaboration, G. Aad, et al., Observation of associated near-side and away-side long-range correlations in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector, Phys. Rev. Lett. 110 (18) (2013) 182302, arXiv:1212.5198 [hep-ex].
- [15] ATLAS Collaboration, G. Aad, et al., Measurement with the ATLAS detector of multi-particle azimuthal correlations in p + Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 725 (2013) 60–78, arXiv:1303.2084 [hep-ex].
- [16] CMS Collaboration, S. Chatrchyan, et al., Multiplicity and transverse momentum dependence of two- and four-particle correlations in pPb and PbPb collisions, Phys. Lett. B 724 (2013) 213–240, arXiv:1305.0609 [nucl-ex].
- [17] ALICE Collaboration, S. Acharya, et al., Azimuthal correlations of prompt D mesons with charged particles in pp and p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C 80 (10) (2020) 979, arXiv:1910.14403 [nucl-ex].
- [18] ALICE Collaboration, J. Adam, et al., Forward-central two-particle correlations in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 753 (2016) 126–139, arXiv: 1506.08032 [nucl-ex].
- [19] ATLAS Collaboration, G. Aad, et al., Observation of long-range elliptic azimuthal anisotropies in $\sqrt{s} = 13$ and 2.76 TeV pp collisions with the ATLAS detector, Phys. Rev. Lett. 116 (17) (2016) 172301, arXiv:1509.04776 [hep-ex].
- [20] CMS Collaboration, V. Khachatryan, et al., Measurement of long-range near-side two-particle angular correlations in pp collisions at $\sqrt{s} = 13$ TeV, Phys. Rev. Lett. 116 (17) (2016) 172302, arXiv:1510.03068 [nucl-ex].
- [21] LHCb Collaboration, R. Aaij, et al., Measurements of long-range near-side angular correlations in $\sqrt{s_{NN}} = 5$ TeV proton-lead collisions in the forward region, Phys. Lett. B 762 (2016) 473–483, arXiv:1512.00439 [nucl-ex].
- [22] ALICE Collaboration, B. Abelev, et al., Long-range angular correlations of π , K and p in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 726 (2013) 164–177, arXiv:1307.3237 [nucl-ex].
- [23] ALICE Collaboration, B. Abelev, et al., Multiplicity dependence of pion, kaon, proton and lambda production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 728 (2014) 25–38, arXiv:1307.6796 [nucl-ex].
- [24] ALICE Collaboration, S. Acharya, et al., Multiplicity dependence of light-flavor hadron production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. C 99 (2) (2019) 024906, arXiv:1807.11321 [nucl-ex].
- [25] CMS Collaboration, V. Khachatryan, et al., Long-range two-particle correlations of strange hadrons with charged particles in pPb and PbPb collisions at LHC energies, Phys. Lett. B 742 (2015) 200–224, arXiv:1409.3392 [nucl-ex].
- [26] ZEUS Collaboration, S. Chekanov, et al., Measurement of K_S^0 , Λ , $\bar{\Lambda}$ production at HERA, Eur. Phys. J. C 51 (2007) 1–23, arXiv:hep-ex/0612023.
- [27] ALICE Collaboration, J. Adam, et al., Enhanced production of multi-strange hadrons in high-multiplicity proton-proton collisions, Nat. Phys. 13 (2017) 535–539, arXiv:1606.07424 [nucl-ex].
- [28] CMS Collaboration, V. Khachatryan, et al., Multiplicity and rapidity dependence of strange hadron production in pp, pPb, and PbPb collisions at the LHC, Phys. Lett. B 768 (2017) 103–129, arXiv:1605.06699 [nucl-ex].
- [29] CMS Collaboration, A.M. Sirunyan, et al., Elliptic flow of charm and strange hadrons in high-multiplicity pPb collisions at $\sqrt{s_{NN}} = 8.16$ TeV, Phys. Rev. Lett. 121 (8) (2018) 082301, arXiv:1804.09767 [hep-ex].
- [30] LHCb Collaboration, R. Aaij, et al., Measurement of B^+ , B^0 and Λ_b^0 production in pPb collisions at $\sqrt{s_{NN}} = 8.16$ TeV, Phys. Rev. D 99 (5) (2019) 052011, arXiv: 1902.05599 [hep-ex].
- [31] LHCb Collaboration, R. Aaij, et al., Measurement of b hadron fractions in 13 TeV pp collisions, Phys. Rev. D 100 (3) (2019) 031102, arXiv:1902.06794 [hep-ex].
- [32] PHENIX Collaboration, S. Adler, et al., Scaling properties of proton and anti-proton production in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions, Phys. Rev. Lett. 91 (2003) 172301, arXiv:nucl-ex/0305036.
- [33] STAR Collaboration, H. Long, Nuclear modification of identified strange particles at moderate p_T in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV at RHIC, J. Phys. G 30 (2004) S193–S197.
- [34] STAR, STAR RICH Collaboration, J. Adams, et al., Measurements of identified particles at intermediate transverse momentum in the STAR experiment from Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, arXiv:nucl-ex/0601042.
- [35] R.J. Fries, V. Greco, P. Sorensen, Coalescence models for hadron formation from quark-gluon plasma, Annu. Rev. Nucl. Part. Sci. 58 (2008) 177–205, arXiv:0807.4939 [nucl-th].
- [36] ALICE Collaboration, B. Abelev, et al., K_S^0 and Λ production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. Lett. 111 (2013) 222301, arXiv:1307.5530 [nucl-ex].
- [37] E. Schnedermann, J. Sollfrank, U.W. Heinz, Thermal phenomenology of hadrons from 200A GeV S+S collisions, Phys. Rev. C 48 (1993) 2462–2475, arXiv:nucl-th/9307020 [nucl-th].
- [38] R. Fries, B. Müller, C. Nonaka, S. Bass, Hadronization in heavy ion collisions: recombination and fragmentation of partons, Phys. Rev. Lett. 90 (2003) 202303, arXiv:nucl-th/0301087 [nucl-th].
- [39] ATLAS Collaboration, G. Aad, et al., Observation of a centrality-dependent dijet asymmetry in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC, Phys. Rev. Lett. 105 (2010) 252303, arXiv:1011.6182 [hep-ex].
- [40] CMS Collaboration, S. Chatrchyan, et al., Jet momentum dependence of jet quenching in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Lett. B 712 (2012) 176–197, arXiv:1202.5022 [nucl-ex].
- [41] ATLAS Collaboration, G. Aad, et al., Measurement of the jet radius and transverse momentum dependence of inclusive jet suppression in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector, Phys. Lett. B 719 (2013) 220–241, arXiv:1208.1967 [hep-ex].
- [42] ALICE Collaboration, B. Abelev, et al., Measurement of charged jet suppression in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, J. High Energy Phys. 03 (2014) 013, arXiv:1311.0633 [nucl-ex].
- [43] ATLAS Collaboration, G. Aad, et al., Measurements of the nuclear modification factor for jets in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector, Phys. Rev. Lett. 114 (7) (2015) 072302, arXiv:1411.2357 [hep-ex].
- [44] ALICE Collaboration, J. Adam, et al., Measurement of charged jet production cross sections and nuclear modification in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 749 (2015) 68–81, arXiv:1503.00681 [nucl-ex].
- [45] ALICE Collaboration, J. Adam, et al., Measurement of dijet k_T in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 746 (2015) 385–395, arXiv:1503.03050 [nucl-ex].
- [46] CMS Collaboration, V. Khachatryan, et al., Measurement of inclusive jet production and nuclear modifications in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C 76 (7) (2016) 372, arXiv:1601.02001 [nucl-ex].
- [47] ALICE Collaboration, B. Abelev, et al., Transverse momentum dependence of inclusive primary charged-particle production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C 74 (9) (2014) 3054, arXiv:1405.2737 [nucl-ex].
- [48] ATLAS Collaboration, G. Aad, et al., Transverse momentum, rapidity, and centrality dependence of inclusive charged-particle production in $\sqrt{s_{NN}} = 5.02$ TeV p + Pb collisions measured by the ATLAS experiment, Phys. Lett. B 763 (2016) 313–336, arXiv:1605.06436 [hep-ex].
- [49] CMS Collaboration, V. Khachatryan, et al., Nuclear effects on the transverse momentum spectra of charged particles in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C 75 (5) (2015) 237, arXiv:1502.05387 [nucl-ex].
- [50] ALICE Collaboration, J. Adam, et al., Centrality dependence of charged jet production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Eur. Phys. J. C 76 (5) (2016) 271, arXiv:1603.03402 [nucl-ex].
- [51] L.D. McLerran, The color glass condensate and small-x physics, Lect. Notes Phys. 583 (2002) 291–334, arXiv:hep-ph/0104285 [hep-ph].
- [52] C. Salgado, J. Alvarez-Muniz, F. Arleo, N. Armesto, M. Botje, et al., Proton-nucleus collisions at the LHC: scientific opportunities and requirements, J. Phys. G 39 (2012) 015010, arXiv:1105.3919 [hep-ph].
- [53] A. Krzywicki, J. Engels, B. Petersson, U. Sukhatme, Does a nucleus act like a gluon filter?, Phys. Lett. B 85 (1979) 407.

- [54] A. Accardi, Final state interactions and hadron quenching in cold nuclear matter, Phys. Rev. C 76 (2007) 034902, arXiv:0706.3227 [nucl-th].
- [55] M. Cacciari, G.P. Salam, G. Soyez, The anti- k_t jet clustering algorithm, J. High Energy Phys. 0804 (2008) 063, arXiv:0802.1189 [hep-ph].
- [56] T. Sjöstrand, S. Ask, J.R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C.O. Rasmussen, P.Z. Skands, An introduction to PYTHIA 8.2, Comput. Phys. Commun. 191 (Jun 2015) 159–177, <https://doi.org/10.1016/j.cpc.2015.01.024>.
- [57] ALICE Collaboration, K. Aamodt, et al., The ALICE experiment at the CERN LHC, J. Instrum. 3 (2008), S08002.
- [58] ALICE Collaboration, K. Aamodt, et al., Alignment of the ALICE inner tracking system with cosmic-ray tracks, J. Instrum. 5 (2010) P03003, arXiv:1001.0502 [physics.ins-det].
- [59] J. Alme, Y. Andres, H. Appelshauser, S. Bablok, N. Bialas, et al., The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events, Nucl. Instrum. Methods A 622 (2010) 316–367, arXiv:1001.1950 [physics.ins-det].
- [60] ALICE Collaboration, K. Aamodt, et al., Transverse momentum spectra of charged particles in proton-proton collisions at $\sqrt{s} = 900$ GeV with ALICE at the LHC, Phys. Lett. B 693 (2010) 53–68, arXiv:1007.0719 [hep-ex].
- [61] L. Evans, P. Bryant, LHC machine, J. Instrum. 3 (2008), S08001.
- [62] ALICE Collaboration, E. Abbas, et al., Performance of the ALICE VZERO system, J. Instrum. 8 (2013) P10016, arXiv:1306.3130 [nucl-ex].
- [63] ALICE Collaboration, B. Abelev, et al., Pseudorapidity density of charged particles in p + Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 110 (3) (2013) 032301, arXiv:1210.3615 [nucl-ex].
- [64] ALICE Collaboration, B. Abelev, et al., Charged jet cross sections and properties in proton–proton collisions at $\sqrt{s} = 7$ TeV, Phys. Rev. D 91 (11) (2015) 112012, arXiv:1411.4969 [nucl-ex].
- [65] M. Cacciari, G.P. Salam, G. Soyez, FastJet user manual, Eur. Phys. J. C 72 (2012) 1896, arXiv:1111.6097 [hep-ph].
- [66] M. Cacciari, G.P. Salam, Dispelling the N^3 myth for the k_t jet-finder, Phys. Lett. B 641 (2006) 57–61, arXiv:hep-ph/0512210 [hep-ph].
- [67] M. Cacciari, G.P. Salam, G. Soyez, The catchment area of jets, J. High Energy Phys. 0804 (2008) 005, arXiv:0802.1188 [hep-ph].
- [68] T. Sjöstrand, S. Mrenna, P.Z. Skands, PYTHIA 6.4 physics and manual, J. High Energy Phys. 05 (2006) 026, arXiv:hep-ph/0603175.
- [69] R. Brun, F. Bruyant, F. Carminati, S. Giani, M. Maire, A. McPherson, G. Patrick, L. Urban, GEANT Detector Description and Simulation Tool, CERN-W5013, CERN-W-5013, W5013, 1994.
- [70] Particle Data Group Collaboration, M. Tanabashi, et al., Review of particle physics, Phys. Rev. D 98 (3) (2018) 030001.
- [71] ALICE Collaboration, B. Abelev, et al., Performance of the ALICE experiment at the CERN LHC, Int. J. Mod. Phys. A 29 (2014) 1430044, arXiv:1402.4476 [nucl-ex].
- [72] ALICE Collaboration, K. Aamodt, et al., Strange particle production in proton–proton collisions at $\sqrt{s} = 0.9$ TeV with ALICE at the LHC, Eur. Phys. J. C 71 (2011) 1594, arXiv:1012.3257 [hep-ex].
- [73] S. Roesler, R. Engel, J. Ranft, The Monte Carlo event generator DPMJET-III, arXiv:hep-ph/0012252 [hep-ph].
- [74] ALICE Collaboration, B. Abelev, et al., Multi-strange baryon production in pp collisions at $\sqrt{s} = 7$ TeV with ALICE, Phys. Lett. B 712 (2012) 309–318, arXiv:1204.0282 [nucl-ex].

ALICE Collaboration

S. Acharya ¹⁴², D. Adamová ⁹⁷, A. Adler ⁷⁵, J. Adolfsson ⁸², G. Aglieri Rinella ³⁵, M. Agnello ³¹, N. Agrawal ⁵⁵, Z. Ahammed ¹⁴², S. Ahmad ¹⁶, S.U. Ahn ⁷⁷, Z. Akbar ⁵², A. Akindinov ⁹⁴, M. Al-Turany ¹⁰⁹, D. Aleksandrov ⁹⁰, B. Alessandro ⁶⁰, H.M. Alfanda ⁷, R. Alfaro Molina ⁷², B. Ali ¹⁶, Y. Ali ¹⁴, A. Alici ²⁶, N. Alizadehvandchali ¹²⁶, A. Alkin ³⁵, J. Alme ²¹, T. Alt ⁶⁹, L. Altenkamper ²¹, I. Altsybeev ¹¹⁴, M.N. Anaam ⁷, C. Andrei ⁴⁹, D. Andreou ⁹², A. Andronic ¹⁴⁵, M. Angeletti ³⁵, V. Anguelov ¹⁰⁶, F. Antinori ⁵⁸, P. Antonioli ⁵⁵, C. Anuj ¹⁶, N. Apadula ⁸¹, L. Aphecetche ¹¹⁶, H. Appelshäuser ⁶⁹, S. Arcelli ²⁶, R. Arnaldi ⁶⁰, I.C. Arsene ²⁰, M. Arslanbekci ^{147,106}, A. Augustinus ³⁵, R. Averbeck ¹⁰⁹, S. Aziz ⁷⁹, M.D. Azmi ¹⁶, A. Badalà ⁵⁷, Y.W. Baek ⁴², X. Bai ¹⁰⁹, R. Bailhache ⁶⁹, Y. Bailung ⁵¹, R. Bala ¹⁰³, A. Balbino ³¹, A. Baldisseri ¹³⁹, M. Ball ⁴⁴, D. Banerjee ⁴, R. Barbera ²⁷, L. Barioglio ^{107,25}, M. Barlou ⁸⁶, G.G. Barnaföldi ¹⁴⁶, L.S. Barnby ⁹⁶, V. Barret ¹³⁶, C. Bartels ¹²⁹, K. Barth ³⁵, E. Bartsch ⁶⁹, F. Baruffaldi ²⁸, N. Bastid ¹³⁶, S. Basu ^{82,144}, G. Batigne ¹¹⁶, B. Batyunya ⁷⁶, D. Bauri ⁵⁰, J.L. Bazo Alba ¹¹³, I.G. Bearden ⁹¹, C. Beattie ¹⁴⁷, I. Belikov ¹³⁸, A.D.C. Bell Hechavarria ¹⁴⁵, F. Bellini ³⁵, R. Bellwied ¹²⁶, S. Belokurova ¹¹⁴, V. Belyaev ⁹⁵, G. Bencedi ^{70,146}, S. Beole ²⁵, A. Bercuci ⁴⁹, Y. Berdnikov ¹⁰⁰, A. Berdnikova ¹⁰⁶, D. Berenyi ¹⁴⁶, L. Bergmann ¹⁰⁶, M.G. Besoiu ⁶⁸, L. Betev ³⁵, P.P. Bhaduri ¹⁴², A. Bhasin ¹⁰³, I.R. Bhat ¹⁰³, M.A. Bhat ⁴, B. Bhattacharjee ⁴³, P. Bhattacharya ²³, L. Bianchi ²⁵, N. Bianchi ⁵³, J. Bielčík ³⁸, J. Bielčíková ⁹⁷, J. Biernat ¹¹⁹, A. Bilandžić ¹⁰⁷, G. Biro ¹⁴⁶, S. Biswas ⁴, J.T. Blair ¹²⁰, D. Blau ⁹⁰, M.B. Blidaru ¹⁰⁹, C. Blume ⁶⁹, G. Boca ²⁹, F. Bock ⁹⁸, A. Bogdanov ⁹⁵, S. Boi ²³, J. Bok ⁶², L. Boldizsár ¹⁴⁶, A. Bolozdynya ⁹⁵, M. Bombara ³⁹, P.M. Bond ³⁵, G. Bonomi ¹⁴¹, H. Borel ¹³⁹, A. Borissov ⁸³, H. Bossi ¹⁴⁷, E. Botta ²⁵, L. Bratrud ⁶⁹, P. Braun-Munzinger ¹⁰⁹, M. Bregant ¹²², M. Broz ³⁸, G.E. Bruno ^{108,34}, M.D. Buckland ¹²⁹, D. Budnikov ¹¹⁰, H. Buesching ⁶⁹, S. Bufalino ³¹, O. Bugnon ¹¹⁶, P. Buhler ¹¹⁵, O. Busch ^{135,I}, Z. Buthelezi ^{73,133}, J.B. Butt ¹⁴, S.A. Bysiak ¹¹⁹, D. Caffarri ⁹², M. Cai ^{28,7}, A. Caliva ¹⁰⁹, E. Calvo Villar ¹¹³, J.M.M. Camacho ¹²¹, R.S. Camacho ⁴⁶, P. Camerini ²⁴, F.D.M. Canedo ¹²², A.A. Capon ¹¹⁵, F. Carnesecchi ²⁶, R. Caron ¹³⁹, J. Castillo Castellanos ¹³⁹, E.A.R. Casula ²³, F. Catalano ³¹, C. Ceballos Sanchez ⁷⁶, P. Chakraborty ⁵⁰, S. Chandra ¹⁴², W. Chang ⁷, S. Chapeland ³⁵, M. Chartier ¹²⁹, S. Chattopadhyay ¹⁴², S. Chattopadhyay ¹¹¹, A. Chauvin ²³, T.G. Chavez ⁴⁶, C. Cheshkov ¹³⁷, B. Cheynis ¹³⁷, V. Chibante Barroso ³⁵, D.D. Chinellato ¹²³, S. Cho ⁶², P. Chochula ³⁵, P. Christakoglou ⁹², C.H. Christensen ⁹¹, P. Christiansen ⁸², T. Chujo ¹³⁵, C. Cicalo ⁵⁶, L. Cifarelli ²⁶, F. Cindolo ⁵⁵, M.R. Ciupak ¹⁰⁹, G. Clai ^{55,II}, J. Cleymans ^{125,I}, F. Colamaria ⁵⁴, J.S. Colburn ¹¹², D. Colella ^{108,54,34,146}, A. Collu ⁸¹, M. Colocci ^{35,26}, M. Concas ^{60,III}, G. Conesa Balbastre ⁸⁰, Z. Conesa del Valle ⁷⁹, G. Contin ²⁴, J.G. Contreras ³⁸, T.M. Cormier ⁹⁸, P. Cortese ³², M.R. Cosentino ¹²⁴, F. Costa ³⁵, S. Costanza ²⁹, P. Crochet ¹³⁶, E. Cuautle ⁷⁰, P. Cui ⁷, L. Cunqueiro ⁹⁸, A. Dainese ⁵⁸, F.P.A. Damas ^{116,139}, M.C. Danisch ¹⁰⁶, A. Danu ⁶⁸, I. Das ¹¹¹, P. Das ⁸⁸, P. Das ⁴, S. Das ⁴, S. Dash ⁵⁰, S. De ⁸⁸, A. De Caro ³⁰, G. de Cataldo ⁵⁴, L. De Cilladi ²⁵, J. de Cuveland ⁴⁰, A. De Falco ²³, D. De Gruttola ³⁰,

- N. De Marco ⁶⁰, C. De Martin ²⁴, S. De Pasquale ³⁰, S. Deb ⁵¹, H.F. Degenhardt ¹²², K.R. Deja ¹⁴³,
 L. Dello Stritto ³⁰, S. Delsanto ²⁵, W. Deng ⁷, P. Dhankher ¹⁹, D. Di Bari ³⁴, A. Di Mauro ³⁵, R.A. Diaz ⁸,
 T. Dietel ¹²⁵, Y. Ding ⁷, R. Divià ³⁵, D.U. Dixit ¹⁹, Ø. Djupsland ²¹, U. Dmitrieva ⁶⁴, J. Do ⁶², A. Dobrin ⁶⁸,
 B. Dönigus ⁶⁹, O. Dordic ²⁰, A.K. Dubey ¹⁴², A. Dubla ^{109,92}, S. Dudi ¹⁰², M. Dukhishyam ⁸⁸, P. Dupieux ¹³⁶,
 T.M. Eder ¹⁴⁵, R.J. Ehlers ⁹⁸, V.N. Eikeland ²¹, D. Elia ⁵⁴, B. Erazmus ¹¹⁶, F. Ercolelli ²⁶, F. Erhardt ¹⁰¹,
 A. Erokhin ¹¹⁴, M.R. Ersdal ²¹, B. Espagnon ⁷⁹, G. Eulisse ³⁵, D. Evans ¹¹², S. Evdokimov ⁹³, L. Fabbietti ¹⁰⁷,
 M. Faggion ²⁸, J. Faivre ⁸⁰, F. Fan ⁷, A. Fantoni ⁵³, M. Fasel ⁹⁸, P. Fecchio ³¹, A. Feliciello ⁶⁰, G. Feofilov ¹¹⁴,
 A. Fernández Téllez ⁴⁶, A. Ferrero ¹³⁹, A. Ferretti ²⁵, V.J.G. Feuillard ¹⁰⁶, J. Figiel ¹¹⁹, S. Filchagin ¹¹⁰,
 D. Finogeev ⁶⁴, F.M. Fionda ²¹, G. Fiorenza ¹⁰⁸, F. Flor ¹²⁶, A.N. Flores ¹²⁰, S. Foertsch ⁷³, P. Foka ¹⁰⁹,
 S. Fokin ⁹⁰, E. Fragiocomo ⁶¹, E. Frajna ¹⁴⁶, U. Fuchs ³⁵, N. Funicello ³⁰, C. Furget ⁸⁰, A. Furs ⁶⁴,
 J.J. Gaardhøje ⁹¹, M. Gagliardi ²⁵, A.M. Gago ¹¹³, A. Gal ¹³⁸, C.D. Galvan ¹²¹, P. Ganoti ⁸⁶, C. Garabatos ¹⁰⁹,
 J.R.A. Garcia ⁴⁶, E. Garcia-Solis ¹⁰, K. Garg ¹¹⁶, C. Gargiulo ³⁵, A. Garibbi ⁸⁹, K. Garner ¹⁴⁵, P. Gasik ¹⁰⁹,
 E.F. Gauger ¹²⁰, A. Gautam ¹²⁸, M.B. Gay Ducati ⁷¹, M. Germain ¹¹⁶, J. Ghosh ¹¹¹, P. Ghosh ¹⁴², S.K. Ghosh ⁴,
 M. Giacalone ²⁶, P. Gianotti ⁵³, P. Giubellino ^{109,60}, P. Giubilato ²⁸, A.M.C. Glaenzer ¹³⁹, P. Glässel ¹⁰⁶,
 V. Gonzalez ¹⁴⁴, L.H. González-Trueba ⁷², S. Gorbunov ⁴⁰, L. Görlich ¹¹⁹, S. Gotovac ³⁶, V. Grabski ⁷²,
 L.K. Graczykowski ¹⁴³, L. Greiner ⁸¹, A. Grelli ⁶³, C. Grigoras ³⁵, V. Grigoriev ⁹⁵, A. Grigoryan ^{1,1},
 S. Grigoryan ^{76,1}, O.S. Groettvik ²¹, F. Grossa ⁶⁰, J.F. Grosse-Oetringhaus ³⁵, R. Grossi ¹⁰⁹,
 G.G. Guardiano ¹²³, R. Guernane ⁸⁰, M. Guilbaud ¹¹⁶, M. Guittiere ¹¹⁶, K. Gulbrandsen ⁹¹, T. Gunji ¹³⁴,
 A. Gupta ¹⁰³, R. Gupta ¹⁰³, I.B. Guzman ⁴⁶, L. Gyulai ¹⁴⁶, M.K. Habib ¹⁰⁹, C. Hadjidakis ⁷⁹, H. Hamagaki ⁸⁴,
 G. Hamar ¹⁴⁶, M. Hamid ⁷, R. Hannigan ¹²⁰, M.R. Haque ^{143,88}, A. Harlenderova ¹⁰⁹, J.W. Harris ¹⁴⁷,
 A. Harton ¹⁰, J.A. Hasenbichler ³⁵, H. Hassan ⁹⁸, D. Hatzifotiadou ⁵⁵, P. Hauer ⁴⁴, L.B. Havener ¹⁴⁷,
 S. Hayashi ¹³⁴, S.T. Heckel ¹⁰⁷, E. Hellbär ⁶⁹, H. Helstrup ³⁷, T. Herman ³⁸, E.G. Hernandez ⁴⁶,
 G. Herrera Corral ⁹, F. Herrmann ¹⁴⁵, K.F. Hetland ³⁷, H. Hillemanns ³⁵, C. Hills ¹²⁹, B. Hippolyte ¹³⁸,
 B. Hohlweber ^{92,107}, J. Honermann ¹⁴⁵, G.H. Hong ¹⁴⁸, D. Horak ³⁸, S. Hornung ¹⁰⁹, R. Hosokawa ¹⁵,
 P. Hristov ³⁵, C. Huang ⁷⁹, C. Hughes ¹³², P. Huhn ⁶⁹, T.J. Humanic ⁹⁹, H. Hushnud ¹¹¹, L.A. Husova ¹⁴⁵,
 N. Hussain ⁴³, D. Hutter ⁴⁰, J.P. Iddon ^{35,129}, R. Ilkaev ¹¹⁰, H. Ilyas ¹⁴, M. Inaba ¹³⁵, G.M. Innocenti ³⁵,
 M. Ippolitov ⁹⁰, A. Isakov ^{38,97}, M.S. Islam ¹¹¹, M. Ivanov ¹⁰⁹, V. Ivanov ¹⁰⁰, V. Izucheev ⁹³, B. Jacak ⁸¹,
 N. Jacazio ³⁵, P.M. Jacobs ⁸¹, S. Jadlovska ¹¹⁸, J. Jadlovsky ¹¹⁸, S. Jaelani ⁶³, C. Jahnke ^{123,122},
 M.J. Jakubowska ¹⁴³, M.A. Janik ¹⁴³, T. Janson ⁷⁵, M. Jercic ¹⁰¹, O. Jevons ¹¹², F. Jonas ^{98,145}, P.G. Jones ¹¹²,
 J.M. Jowett ^{35,109}, J. Jung ⁶⁹, M. Jung ⁶⁹, A. Junique ³⁵, A. Jusko ¹¹², P. Kalinak ⁶⁵, A. Kalweit ³⁵, V. Kaplin ⁹⁵,
 S. Kar ⁷, A. Karasu Uysal ⁷⁸, D. Karatovic ¹⁰¹, O. Karavichev ⁶⁴, T. Karavicheva ⁶⁴, P. Karczmarczyk ¹⁴³,
 E. Karpechev ⁶⁴, A. Kazantsev ⁹⁰, U. Kebschull ⁷⁵, R. Keidel ⁴⁸, M. Keil ³⁵, B. Ketzer ⁴⁴, Z. Khabanova ⁹²,
 A.M. Khan ⁷, S. Khan ¹⁶, A. Khanzadeev ¹⁰⁰, Y. Kharlov ⁹³, A. Khatun ¹⁶, A. Khuntia ¹¹⁹, B. Kileng ³⁷,
 B. Kim ^{17,62}, D. Kim ¹⁴⁸, D.J. Kim ¹²⁷, E.J. Kim ⁷⁴, J. Kim ¹⁴⁸, J.S. Kim ⁴², J. Kim ¹⁰⁶, J. Kim ¹⁴⁸, J. Kim ⁷⁴,
 M. Kim ¹⁰⁶, S. Kim ¹⁸, T. Kim ¹⁴⁸, S. Kirsch ⁶⁹, I. Kisiel ⁴⁰, S. Kiselev ⁹⁴, A. Kisiel ¹⁴³, J.L. Klay ⁶, J. Klein ³⁵,
 S. Klein ⁸¹, C. Klein-Bösing ¹⁴⁵, M. Kleiner ⁶⁹, T. Klemenz ¹⁰⁷, A. Kluge ³⁵, A.G. Knospe ¹²⁶, C. Kobdaj ¹¹⁷,
 M.K. Köhler ¹⁰⁶, T. Kollegger ¹⁰⁹, A. Kondratyev ⁷⁶, N. Kondratyeva ⁹⁵, E. Kondratyuk ⁹³, J. Konig ⁶⁹,
 S.A. Königstorfer ¹⁰⁷, P.J. Konopka ^{35,2}, G. Kornakov ¹⁴³, S.D. Koryciak ², L. Koska ¹¹⁸, O. Kovalenko ⁸⁷,
 V. Kovalenko ¹¹⁴, M. Kowalski ¹¹⁹, I. Králik ⁶⁵, A. Kravčáková ³⁹, L. Kreis ¹⁰⁹, M. Krivda ^{112,65}, F. Krizek ⁹⁷,
 K. Krizkova Gajdosova ³⁸, M. Kroesen ¹⁰⁶, M. Krüger ⁶⁹, E. Kryshen ¹⁰⁰, M. Krzewicki ⁴⁰, V. Kučera ³⁵,
 C. Kuhn ¹³⁸, P.G. Kuijer ⁹², T. Kumaoka ¹³⁵, D. Kumar ¹⁴², L. Kumar ¹⁰², S. Kundu ^{35,88}, P. Kurashvili ⁸⁷,
 A. Kurepin ⁶⁴, A.B. Kurepin ⁶⁴, A. Kuryakin ¹¹⁰, S. Kushpil ⁹⁷, J. Kvapil ¹¹², M.J. Kweon ⁶², J.Y. Kwon ⁶²,
 Y. Kwon ¹⁴⁸, S.L. La Pointe ⁴⁰, P. La Rocca ²⁷, Y.S. Lai ⁸¹, A. Lakrathok ¹¹⁷, M. Lamanna ³⁵, R. Langoy ¹³¹,
 K. Lapidus ³⁵, P. Larionov ⁵³, E. Laudi ³⁵, L. Lautner ^{35,107}, R. Lavicka ³⁸, T. Lazareva ¹¹⁴, R. Lea ^{141,24},
 J. Lee ¹³⁵, J. Lehrbach ⁴⁰, R.C. Lemmon ⁹⁶, I. León Monzón ¹²¹, E.D. Lesser ¹⁹, M. Lettrich ^{35,107}, P. Lévali ¹⁴⁶,
 X. Li ¹¹, X.L. Li ⁷, J. Lien ¹³¹, R. Lietava ¹¹², B. Lim ¹⁷, S.H. Lim ¹⁷, V. Lindenstruth ⁴⁰, A. Lindner ⁴⁹,
 C. Lippmann ¹⁰⁹, A. Liu ¹⁹, J. Liu ¹²⁹, I.M. Lofnes ²¹, V. Loginov ⁹⁵, C. Loizides ⁹⁸, P. Loncar ³⁶, J.A. Lopez ¹⁰⁶,
 X. Lopez ¹³⁶, E. López Torres ⁸, J.R. Luhder ¹⁴⁵, M. Lunardon ²⁸, G. Luparello ⁶¹, Y.G. Ma ⁴¹, A. Maeviskaya ⁶⁴,
 M. Mager ³⁵, T. Mahmoud ⁴⁴, A. Maire ¹³⁸, M. Malaev ¹⁰⁰, Q.W. Malik ²⁰, L. Malinina ^{76,IV}, D. Mal'Kevich ⁹⁴,
 N. Mallick ⁵¹, P. Malzacher ¹⁰⁹, G. Mandaglio ^{33,57}, V. Manko ⁹⁰, F. Manso ¹³⁶, V. Manzari ⁵⁴, Y. Mao ⁷,
 J. Mareš ⁶⁷, G.V. Margagliotti ²⁴, A. Margotti ⁵⁵, A. Marín ¹⁰⁹, C. Markert ¹²⁰, M. Marquard ⁶⁹,
 N.A. Martin ¹⁰⁶, P. Martinengo ³⁵, J.L. Martinez ¹²⁶, M.I. Martínez ⁴⁶, G. Martínez García ¹¹⁶,
 S. Masciocchi ¹⁰⁹, M. Masera ²⁵, A. Masoni ⁵⁶, L. Massacrier ⁷⁹, A. Mastroserio ^{140,54}, A.M. Mathis ¹⁰⁷,

- O. Matonoha ⁸², P.F.T. Matuoka ¹²², A. Matyja ¹¹⁹, C. Mayer ¹¹⁹, A.L. Mazuecos ³⁵, F. Mazzaschi ²⁵, M. Mazzilli ^{35,54}, M.A. Mazzoni ⁵⁹, A.F. Mechler ⁶⁹, F. Meddi ²², Y. Melikyan ⁶⁴, A. Menchaca-Rocha ⁷², E. Meninno ^{115,30}, A.S. Menon ¹²⁶, M. Meres ¹³, S. Mhlanga ^{125,73}, Y. Miake ¹³⁵, L. Micheletti ²⁵, L.C. Migliorin ¹³⁷, D.L. Mihaylov ¹⁰⁷, K. Mikhaylov ^{76,94}, A.N. Mishra ^{146,70}, D. Miśkowiec ¹⁰⁹, A. Modak ⁴, A.P. Mohanty ⁶³, B. Mohanty ⁸⁸, M. Mohisin Khan ¹⁶, Z. Moravcova ⁹¹, C. Mordasini ¹⁰⁷, D.A. Moreira De Godoy ¹⁴⁵, L.A.P. Moreno ⁴⁶, I. Morozov ⁶⁴, A. Morsch ³⁵, T. Mrnjavac ³⁵, V. Muccifora ⁵³, E. Mudnic ³⁶, D. Mühlheim ¹⁴⁵, S. Muhuri ¹⁴², J.D. Mulligan ⁸¹, A. Mulliri ²³, M.G. Munhoz ¹²², R.H. Munzer ⁶⁹, H. Murakami ¹³⁴, S. Murray ¹²⁵, L. Musa ³⁵, J. Musinsky ⁶⁵, C.J. Myers ¹²⁶, J.W. Myrcha ¹⁴³, B. Naik ⁵⁰, R. Nair ⁸⁷, B.K. Nandi ⁵⁰, R. Nania ⁵⁵, E. Nappi ⁵⁴, M.U. Naru ¹⁴, A.F. Nassirpour ⁸², C. Nattrass ¹³², A. Neagu ²⁰, L. Nellen ⁷⁰, S.V. Nesbo ³⁷, G. Neskovic ⁴⁰, D. Nesterov ¹¹⁴, B.S. Nielsen ⁹¹, S. Nikolaev ⁹⁰, S. Nikulin ⁹⁰, V. Nikulin ¹⁰⁰, F. Noferini ⁵⁵, S. Noh ¹², P. Nomokonov ⁷⁶, J. Norman ¹²⁹, N. Novitzky ¹³⁵, P. Nowakowski ¹⁴³, A. Nyanin ⁹⁰, J. Nystrand ²¹, M. Ogino ⁸⁴, A. Ohlson ⁸², J. Oleniacz ¹⁴³, A.C. Oliveira Da Silva ¹³², M.H. Oliver ¹⁴⁷, A. Onnerstad ¹²⁷, C. Oppedisano ⁶⁰, A. Ortiz Velasquez ⁷⁰, T. Osako ⁴⁷, A. Oskarsson ⁸², J. Otwinowski ¹¹⁹, K. Oyama ⁸⁴, Y. Pachmayer ¹⁰⁶, V. Pacik ⁹¹, S. Padhan ⁵⁰, D. Pagano ¹⁴¹, G. Paić ⁷⁰, A. Palasciano ⁵⁴, J. Pan ¹⁴⁴, S. Panebianco ¹³⁹, P. Pareek ¹⁴², J. Park ⁶², J.E. Parkkila ¹²⁷, S.P. Pathak ¹²⁶, R.N. Patra ¹⁰³, B. Paul ²³, J. Pazzini ¹⁴¹, H. Pei ⁷, T. Peitzmann ⁶³, X. Peng ⁷, L.G. Pereira ⁷¹, H. Pereira Da Costa ¹³⁹, D. Peresunko ⁹⁰, G.M. Perez ⁸, S. Perrin ¹³⁹, Y. Pestov ⁵, V. Petráček ³⁸, M. Petrovici ⁴⁹, R.P. Pezzi ⁷¹, S. Piano ⁶¹, M. Pikna ¹³, P. Pillot ¹¹⁶, O. Pinazza ^{55,35}, L. Pinsky ¹²⁶, C. Pinto ²⁷, S. Pisano ⁵³, M. Płoskoń ⁸¹, M. Planinic ¹⁰¹, F. Pliquet ⁶⁹, M.G. Poghosyan ⁹⁸, B. Polichtchouk ⁹³, S. Politano ³¹, N. Poljak ¹⁰¹, A. Pop ⁴⁹, S. Porteboeuf-Houssais ¹³⁶, J. Porter ⁸¹, V. Pozdniakov ⁷⁶, S.K. Prasad ⁴, R. Preghenella ⁵⁵, F. Prino ⁶⁰, C.A. Pruneau ¹⁴⁴, I. Pshenichnov ⁶⁴, M. Puccio ³⁵, S. Qiu ⁹², L. Quaglia ²⁵, R.E. Quishpe ¹²⁶, S. Ragoni ¹¹², A. Rakotozafindrabe ¹³⁹, L. Ramello ³², F. Rami ¹³⁸, S.A.R. Ramirez ⁴⁶, A.G.T. Ramos ³⁴, R. Raniwala ¹⁰⁴, S. Raniwala ¹⁰⁴, S.S. Räsänen ⁴⁵, R. Rath ⁵¹, I. Ravasenga ⁹², K.F. Read ^{98,132}, A.R. Redelbach ⁴⁰, K. Redlich ^{87,V}, A. Rehman ²¹, P. Reichelt ⁶⁹, F. Reidt ³⁵, H.A. Reme-ness ³⁷, R. Renfordt ⁶⁹, Z. Rescakova ³⁹, K. Reygers ¹⁰⁶, A. Riabov ¹⁰⁰, V. Riabov ¹⁰⁰, T. Richert ^{82,91}, M. Richter ²⁰, W. Riegler ³⁵, F. Riggi ²⁷, C. Ristea ⁶⁸, S.P. Rode ⁵¹, M. Rodríguez Cahuantzi ⁴⁶, K. Røed ²⁰, R. Rogalev ⁹³, E. Rogochaya ⁷⁶, T.S. Rogoschinski ⁶⁹, D. Rohr ³⁵, D. Röhrich ²¹, P.F. Rojas ⁴⁶, P.S. Rokita ¹⁴³, F. Ronchetti ⁵³, A. Rosano ^{33,57}, E.D. Rosas ⁷⁰, A. Rossi ⁵⁸, A. Rotondi ²⁹, A. Roy ⁵¹, P. Roy ¹¹¹, S. Roy ⁵⁰, N. Rubini ²⁶, O.V. Rueda ⁸², R. Rui ²⁴, B. Rumyantsev ⁷⁶, A. Rustamov ⁸⁹, E. Ryabinkin ⁹⁰, Y. Ryabov ¹⁰⁰, A. Rybicki ¹¹⁹, H. Rytkonen ¹²⁷, W. Rzesz ¹⁴³, O.A.M. Saarimaki ⁴⁵, R. Sadek ¹¹⁶, S. Sadovsky ⁹³, J. Saetre ²¹, K. Šafařík ³⁸, S.K. Saha ¹⁴², S. Saha ⁸⁸, B. Sahoo ⁵⁰, P. Sahoo ⁵⁰, R. Sahoo ⁵¹, S. Sahoo ⁶⁶, D. Sahu ⁵¹, P.K. Sahu ⁶⁶, J. Saini ¹⁴², S. Sakai ¹³⁵, S. Sambyal ¹⁰³, V. Samsonov ^{100,95,I}, D. Sarkar ¹⁴⁴, N. Sarkar ¹⁴², P. Sarma ⁴³, V.M. Sarti ¹⁰⁷, M.H.P. Sas ¹⁴⁷, J. Schambach ^{98,120}, H.S. Scheid ⁶⁹, C. Schiaua ⁴⁹, R. Schicker ¹⁰⁶, A. Schmah ¹⁰⁶, C. Schmidt ¹⁰⁹, H.R. Schmidt ¹⁰⁵, M.O. Schmidt ¹⁰⁶, M. Schmidt ¹⁰⁵, N.V. Schmidt ^{98,69}, A.R. Schmier ¹³², R. Schotter ¹³⁸, J. Schukraft ³⁵, Y. Schutz ¹³⁸, K. Schwarz ¹⁰⁹, K. Schweda ¹⁰⁹, G. Scioli ²⁶, E. Scomparin ⁶⁰, J.E. Seger ¹⁵, Y. Sekiguchi ¹³⁴, D. Sekihata ¹³⁴, I. Selyuzhenkov ^{109,95}, S. Senyukov ¹³⁸, J.J. Seo ⁶², D. Serebryakov ⁶⁴, L. Šerkšnytė ¹⁰⁷, A. Sevcenco ⁶⁸, T.J. Shaba ⁷³, A. Shabanov ⁶⁴, A. Shabetai ¹¹⁶, R. Shahoyan ³⁵, W. Shaikh ¹¹¹, A. Shangaraev ⁹³, A. Sharma ¹⁰², H. Sharma ¹¹⁹, M. Sharma ¹⁰³, N. Sharma ¹⁰², S. Sharma ¹⁰³, O. Sheibani ¹²⁶, K. Shigaki ⁴⁷, M. Shimomura ⁸⁵, S. Shirinkin ⁹⁴, Q. Shou ⁴¹, Y. Sibiriak ⁹⁰, S. Siddhanta ⁵⁶, T. Siemianczuk ⁸⁷, T.F. Silva ¹²², D. Silvermyr ⁸², G. Simonetti ³⁵, B. Singh ¹⁰⁷, R. Singh ⁸⁸, R. Singh ¹⁰³, R. Singh ⁵¹, V.K. Singh ¹⁴², V. Singhal ¹⁴², T. Sinha ¹¹¹, B. Sitar ¹³, M. Sitta ³², T.B. Skaali ²⁰, G. Skorodumovs ¹⁰⁶, M. Slupecki ⁴⁵, N. Smirnov ¹⁴⁷, R.J.M. Snellings ⁶³, C. Soncco ¹¹³, J. Song ¹²⁶, A. Songmoolnak ¹¹⁷, F. Soramel ²⁸, S. Sorensen ¹³², I. Sputowska ¹¹⁹, J. Stachel ¹⁰⁶, I. Stan ⁶⁸, P.J. Steffanic ¹³², S.F. Stiefelmaier ¹⁰⁶, D. Stocco ¹¹⁶, M.M. Storetvedt ³⁷, C.P. Stylianidis ⁹², A.A.P. Suaide ¹²², T. Sugitate ⁴⁷, C. Suire ⁷⁹, M. Suljic ³⁵, R. Sultanov ⁹⁴, M. Šumbera ⁹⁷, V. Sumberia ¹⁰³, S. Sumowidagdo ⁵², S. Swain ⁶⁶, A. Szabo ¹³, I. Szarka ¹³, U. Tabassam ¹⁴, S.F. Taghavi ¹⁰⁷, G. Taillepied ¹³⁶, J. Takahashi ¹²³, G.J. Tambave ²¹, S. Tang ^{136,7}, Z. Tang ¹³⁰, M. Tarhini ¹¹⁶, M.G. Tarzila ⁴⁹, A. Tauro ³⁵, G. Tejeda Muñoz ⁴⁶, A. Telesca ³⁵, L. Terlizzi ²⁵, C. Terrevoli ¹²⁶, G. Tersimonov ³, S. Thakur ¹⁴², D. Thomas ¹²⁰, R. Tieulent ¹³⁷, A. Tikhonov ⁶⁴, A.R. Timmins ¹²⁶, M. Tkacik ¹¹⁸, A. Toia ⁶⁹, N. Topilskaya ⁶⁴, M. Toppi ⁵³, F. Torales-Acosta ¹⁹, S.R. Torres ³⁸, A. Trifiró ^{33,57}, S. Tripathy ^{55,70}, T. Tripathy ⁵⁰, S. Trogolo ^{35,28}, G. Trombetta ³⁴, V. Trubnikov ³, W.H. Trzaska ¹²⁷, T.P. Trzcinski ¹⁴³, B.A. Trzeciak ³⁸, A. Tumkin ¹¹⁰, R. Turrisi ⁵⁸, T.S. Tveter ²⁰, K. Ullaland ²¹, A. Uras ¹³⁷, M. Urioni ¹⁴¹, G.L. Usai ²³, M. Vala ³⁹, N. Valle ²⁹, S. Vallero ⁶⁰, N. van der Kolk ⁶³, L.V.R. van Doremalen ⁶³,

M. van Leeuwen ⁹², P. Vande Vyvre ³⁵, D. Varga ¹⁴⁶, Z. Varga ¹⁴⁶, M. Varga-Kofarago ¹⁴⁶, A. Vargas ⁴⁶,
 M. Vasileiou ⁸⁶, A. Vasiliev ⁹⁰, O. Vázquez Doce ¹⁰⁷, V. Vechernin ¹¹⁴, E. Vercellin ²⁵, S. Vergara Limón ⁴⁶,
 L. Vermunt ⁶³, R. Vértesi ¹⁴⁶, M. Verweij ⁶³, L. Vickovic ³⁶, Z. Vilakazi ¹³³, O. Villalobos Baillie ¹¹²,
 G. Vino ⁵⁴, A. Vinogradov ⁹⁰, T. Virgili ³⁰, V. Vislavicius ⁹¹, A. Vodopyanov ⁷⁶, B. Volkel ³⁵,
 M.A. Völkl ^{106, 105}, K. Voloshin ⁹⁴, S.A. Voloshin ¹⁴⁴, G. Volpe ³⁴, B. von Haller ³⁵, I. Vorobyev ¹⁰⁷,
 D. Voscek ¹¹⁸, J. Vrláková ³⁹, B. Wagner ²¹, C. Wang ⁴¹, D. Wang ⁴¹, M. Weber ¹¹⁵, A. Wegrzynek ³⁵,
 S.C. Wenzel ³⁵, J.P. Wessels ¹⁴⁵, J. Wiechula ⁶⁹, J. Wikne ²⁰, G. Wilk ⁸⁷, J. Wilkinson ¹⁰⁹, G.A. Willems ¹⁴⁵,
 E. Willsher ¹¹², B. Windelband ¹⁰⁶, M. Winn ¹³⁹, W.E. Witt ¹³², J.R. Wright ¹²⁰, W. Wu ⁴¹, Y. Wu ¹³⁰,
 R. Xu ⁷, S. Yalcin ⁷⁸, Y. Yamaguchi ⁴⁷, K. Yamakawa ⁴⁷, S. Yang ²¹, S. Yano ^{47, 139}, Z. Yin ⁷, H. Yokoyama ⁶³,
 I.-K. Yoo ¹⁷, J.H. Yoon ⁶², S. Yuan ²¹, A. Yuncu ¹⁰⁶, V. Zaccolo ²⁴, A. Zaman ¹⁴, C. Zampolli ³⁵, H.J.C. Zanolli ⁶³,
 N. Zardoshti ³⁵, A. Zarochentsev ¹¹⁴, P. Závada ⁶⁷, N. Zaviyalov ¹¹⁰, H. Zbroszczyk ¹⁴³, M. Zhalov ¹⁰⁰,
 S. Zhang ⁴¹, X. Zhang ⁷, Y. Zhang ¹³⁰, Y. Zhang ⁷, V. Zherebchevskii ¹¹⁴, Y. Zhi ¹¹, D. Zhou ⁷, Y. Zhou ⁹¹,
 J. Zhu ^{7, 109}, Y. Zhu ⁷, A. Zichichi ²⁶, G. Zinovjev ³, N. Zurlo ¹⁴¹

¹ A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia² AGH University of Science and Technology, Cracow, Poland³ Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine⁴ Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India⁵ Budker Institute for Nuclear Physics, Novosibirsk, Russia⁶ California Polytechnic State University, San Luis Obispo, CA, United States⁷ Central China Normal University, Wuhan, China⁸ Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba⁹ Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico¹⁰ Chicago State University, Chicago, IL, United States¹¹ China Institute of Atomic Energy, Beijing, China¹² Chungbuk National University, Cheongju, Republic of Korea¹³ Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovakia¹⁴ COMSATS University Islamabad, Islamabad, Pakistan¹⁵ Creighton University, Omaha, NB, United States¹⁶ Department of Physics, Aligarh Muslim University, Aligarh, India¹⁷ Department of Physics, Pusan National University, Pusan, Republic of Korea¹⁸ Department of Physics, Sejong University, Seoul, Republic of Korea¹⁹ Department of Physics, University of California, Berkeley, CA, United States²⁰ Department of Physics, University of Oslo, Oslo, Norway²¹ Department of Physics and Technology, University of Bergen, Bergen, Norway²² Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy²³ Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy²⁴ Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy²⁵ Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy²⁶ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy²⁷ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy²⁸ Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy²⁹ Dipartimento di Fisica e Nucleare e Teorica, Università di Pavia and Sezione INFN, Pavia, Italy³⁰ Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy³¹ Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy³² Dipartimento di Scienze e Innovazione Tecnologica dell'Università del Piemonte Orientale and INFN Sezione di Torino, Alessandria, Italy³³ Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy³⁴ Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy³⁵ European Organization for Nuclear Research (CERN), Geneva, Switzerland³⁶ Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia³⁷ Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway³⁸ Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic³⁹ Faculty of Science, P.J. Šafárik University, Košice, Slovakia⁴⁰ Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany⁴¹ Fudan University, Shanghai, China⁴² Gangneung-Wonju National University, Gangneung, Republic of Korea⁴³ Gauhati University, Department of Physics, Guwahati, India⁴⁴ Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany⁴⁵ Helsinki Institute of Physics (HIP), Helsinki, Finland⁴⁶ High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico⁴⁷ Hiroshima University, Hiroshima, Japan⁴⁸ Hochschule Worms, Zentrum für Technologietransfer und Telekommunikation (ZTT), Worms, Germany⁴⁹ Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania⁵⁰ Indian Institute of Technology Bombay (IIT), Mumbai, India⁵¹ Indian Institute of Technology Indore, Indore, India⁵² Indonesian Institute of Sciences, Jakarta, Indonesia⁵³ INFN, Laboratori Nazionali di Frascati, Frascati, Italy⁵⁴ INFN, Sezione di Bari, Bari, Italy⁵⁵ INFN, Sezione di Bologna, Bologna, Italy⁵⁶ INFN, Sezione di Cagliari, Cagliari, Italy⁵⁷ INFN, Sezione di Catania, Catania, Italy⁵⁸ INFN, Sezione di Padova, Padova, Italy⁵⁹ INFN, Sezione di Roma, Rome, Italy⁶⁰ INFN, Sezione di Torino, Turin, Italy

- ⁶¹ INFN, Sezione di Trieste, Trieste, Italy
⁶² Inha University, Incheon, Republic of Korea
⁶³ Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands
⁶⁴ Institute for Nuclear Research, Academy of Sciences, Moscow, Russia
⁶⁵ Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia
⁶⁶ Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India
⁶⁷ Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic
⁶⁸ Institute of Space Science (ISS), Bucharest, Romania
⁶⁹ Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany
⁷⁰ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico
⁷¹ Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
⁷² Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
⁷³ iThemba LABS, National Research Foundation, Somerset West, South Africa
⁷⁴ Jeonbuk National University, Jeonju, Republic of Korea
⁷⁵ Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany
⁷⁶ Joint Institute for Nuclear Research (JINR), Dubna, Russia
⁷⁷ Korea Institute of Science and Technology Information, Daejeon, Republic of Korea
⁷⁸ KTO Karatay University, Konya, Turkey
⁷⁹ Laboratoire de Physique des 2 Infinis, Irène Joliot-Curie, Orsay, France
⁸⁰ Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France
⁸¹ Lawrence Berkeley National Laboratory, Berkeley, CA, United States
⁸² Lund University Department of Physics, Division of Particle Physics, Lund, Sweden
⁸³ Moscow Institute for Physics and Technology, Moscow, Russia
⁸⁴ Nagasaki Institute of Applied Science, Nagasaki, Japan
⁸⁵ Nara Women's University (NWU), Nara, Japan
⁸⁶ National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece
⁸⁷ National Centre for Nuclear Research, Warsaw, Poland
⁸⁸ National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India
⁸⁹ National Nuclear Research Center, Baku, Azerbaijan
⁹⁰ National Research Centre Kurchatov Institute, Moscow, Russia
⁹¹ Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark
⁹² Nikhef, National institute for subatomic physics, Amsterdam, Netherlands
⁹³ NRC Kurchatov Institute IHEP, Protvino, Russia
⁹⁴ NRC «Kurchatov» Institute - ITEP, Moscow, Russia
⁹⁵ NRNU Moscow Engineering Physics Institute, Moscow, Russia
⁹⁶ Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom
⁹⁷ Nuclear Physics Institute of the Czech Academy of Sciences, Řež u Prahy, Czech Republic
⁹⁸ Oak Ridge National Laboratory, Oak Ridge, TN, United States
⁹⁹ Ohio State University, Columbus, OH, United States
¹⁰⁰ Petersburg Nuclear Physics Institute, Gatchina, Russia
¹⁰¹ Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia
¹⁰² Physics Department, Panjab University, Chandigarh, India
¹⁰³ Physics Department, University of Jammu, Jammu, India
¹⁰⁴ Physics Department, University of Rajasthan, Jaipur, India
¹⁰⁵ Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany
¹⁰⁶ Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany
¹⁰⁷ Physik Department, Technische Universität München, Munich, Germany
¹⁰⁸ Politecnico di Bari and Sezione INFN, Bari, Italy
¹⁰⁹ Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
¹¹⁰ Russian Federal Nuclear Center (VNIIEF), Sarov, Russia
¹¹¹ Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India
¹¹² School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom
¹¹³ Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru
¹¹⁴ St. Petersburg State University, St. Petersburg, Russia
¹¹⁵ Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria
¹¹⁶ SUBATECH, IMT Atlantique, Université de Nantes, CNRS-IN2P3, Nantes, France
¹¹⁷ Suranaree University of Technology, Nakhon Ratchasima, Thailand
¹¹⁸ Technical University of Košice, Košice, Slovakia
¹¹⁹ The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland
¹²⁰ The University of Texas at Austin, Austin, TX, United States
¹²¹ Universidad Autónoma de Sinaloa, Culiacán, Mexico
¹²² Universidade de São Paulo (USP), São Paulo, Brazil
¹²³ Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil
¹²⁴ Universidade Federal do ABC, Santo Andre, Brazil
¹²⁵ University of Cape Town, Cape Town, South Africa
¹²⁶ University of Houston, Houston, TX, United States
¹²⁷ University of Jyväskylä, Jyväskylä, Finland
¹²⁸ University of Kansas, Lawrence, KS, United States
¹²⁹ University of Liverpool, Liverpool, United Kingdom
¹³⁰ University of Science and Technology of China, Hefei, China
¹³¹ University of South-Eastern Norway, Tønsberg, Norway
¹³² University of Tennessee, Knoxville, TN, United States
¹³³ University of the Witwatersrand, Johannesburg, South Africa
¹³⁴ University of Tokyo, Tokyo, Japan
¹³⁵ University of Tsukuba, Tsukuba, Japan
¹³⁶ Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand, France
¹³⁷ Université de Lyon, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, Lyon, France
¹³⁸ Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France
¹³⁹ Université Paris-Saclay Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPhN), Saclay, France
¹⁴⁰ Università degli Studi di Foggia, Foggia, Italy

¹⁴¹ Università di Brescia and Sezione INFN, Brescia, Italy¹⁴² Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India¹⁴³ Warsaw University of Technology, Warsaw, Poland¹⁴⁴ Wayne State University, Detroit, MI, United States¹⁴⁵ Westfälische Wilhelms-Universität Münster, Institut für Kernphysik, Münster, Germany¹⁴⁶ Wigner Research Centre for Physics, Budapest, Hungary¹⁴⁷ Yale University, New Haven, CT, United States¹⁴⁸ Yonsei University, Seoul, Republic of Korea^I Deceased.^{II} Also at: Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Bologna, Italy.^{III} Also at: Dipartimento DET del Politecnico di Torino, Turin, Italy.^{IV} Also at: M.V. Lomonosov Moscow State University, D.V. Skobeltsyn Institute of Nuclear, Physics, Moscow, Russia.^V Also at: Institute of Theoretical Physics, University of Wroclaw, Poland.