

## References

- [1] Behzad, M. Graphs and their chromatic number. Doctoral thesis, Michigan State University, East Lansing, MI, 1965.
- [2] Behzad, M., Chartrand, G. and Cooper Jr, J.K. The colour numbers of complete graphs. *J. Lond. Math. Sci.*, 42:226–228, 1967. <https://doi.org/10.1112/jlms/s1-42.1.226>
- [3] Campos, C.N. and de Mello, C.P. The total chromatic number of some bipartite graphs. *Electronic Notes in Discrete Mathematics*, 22:557–561, 2005. <https://doi.org/10.1016/j.endm.2005.06.082>
- [4] Campos, C.N., Dantas, S. and de Mello, C.P. The total chromatic number of some families of snarks. *Discrete Mathematics*, 311:984–988, 2011. <https://doi.org/10.1016/j.disc.2011.02.013>
- [5] Campos, C.N. and de Mello, C.P. A result on the total colouring of powers of cycles. *Discrete Applied Mathematics*, 155:585–597, 2007. <https://doi.org/10.1016/j.dam.2006.08.010>
- [6] Colin McDiarmid, J.H. and Sanchez-Arroyo, A. Total colouring regular bipartite graphs is NP-hard. *Discrete Mathematics*, 124:155–162, 1994. [https://doi.org/10.1016/0012-365X\(92\)00058-Y](https://doi.org/10.1016/0012-365X(92)00058-Y)
- [7] Douglas B. West. Introduction to graph theory. Second Edition, Prentice Hall, Upper Saddle River, NJ, 2001.
- [8] Rosenfeld, M. On the total chromatic number of a graph. *Israel J. Math.*, 9:396–402, 1971. <https://doi.org/10.1007/BF02771690>
- [9] Sanchez-Arroyo, A. Determining the total colouring number is NP-hard. *Discrete Math.* 78:315–319, 1989. [https://doi.org/10.1016/0012-365X\(89\)90187-8](https://doi.org/10.1016/0012-365X(89)90187-8)
- [10] Sasaki, D., Dantas, S., de Figueiredo, C.M.H. and Preissmann, M. The hunting of a snark with total chromatic number 5. *Discrete Applied Mathematics*, 164:470–481, 2014. <https://doi.org/10.1016/j.dam.2013.04.006>
- [11] Vijayaditya, N. On total chromatic number of a graph. *J. London Math. Soc.*, 2:405–408, 1971. <https://doi.org/10.1112/jlms/s2-3.3.405>