



12-2002

Detrimental effects of prostaglandin F₂[alpha] on preimplantation bovine embryos

Fernando Nester Scenna

Follow this and additional works at: https://trace.tennessee.edu/utk_gradthes

Recommended Citation

Scenna, Fernando Nester, "Detrimental effects of prostaglandin F₂[alpha] on preimplantation bovine embryos. " Master's Thesis, University of Tennessee, 2002.
https://trace.tennessee.edu/utk_gradthes/6544

This Thesis is brought to you for free and open access by the Graduate School at TRACE: Tennessee Research and Creative Exchange. It has been accepted for inclusion in Masters Theses by an authorized administrator of TRACE: Tennessee Research and Creative Exchange. For more information, please contact trace@utk.edu.

To the Graduate Council:

I am submitting herewith a thesis written by Fernando Nester Scenna entitled "Detrimental effects of prostaglandin F₂[alpha] on preimplantation bovine embryos." I have examined the final electronic copy of this thesis for form and content and recommend that it be accepted in partial fulfillment of the requirements for the degree of Master of Science, with a major in Animal Science.

F. Neal Schrick, Major Professor

We have read this thesis and recommend its acceptance:

Lannett Edwards, Patty Tithof, Jay Whelan

Accepted for the Council:


Carolyn R. Hodges

Vice Provost and Dean of the Graduate School

(Original signatures are on file with official student records.)



To the Graduate Council:

I am submitting herewith a thesis written by Fernando Néstor Scenna entitled "Detrimental Effects of Prostaglandin $F_{2\alpha}$ on Preimplantation Bovine Embryos". I have examined the final paper copy of this thesis for form and content and recommend that it be accepted in partial fulfillment of the requirements for the degree of Master of Science, with a major in Animal Science.



F. Neal Schrick, Major Professor

We have read this thesis
and recommend its acceptance:



Accepted for the Council:



Vice Provost and Dean of Graduate Studies

AG-VET-MED.

Thesis
2002
.S345

**DETRIMENTAL EFFECTS OF PROSTAGLANDIN F_{2α} ON
PREIMPLANTATION BOVINE EMBRYOS**

A Thesis

Presented for the

Master of Science

Degree

The University of Tennessee, Knoxville

Fernando Néstor Scenna

December, 2002

ACKNOWLEDGEMENTS

First of all, I would like to thank my major professor, Dr. F. Neal Schrick, for providing me the opportunity of pursuing this degree under his guidance. He trusted me without knowing me personally and when he had never had an international student before. More than anything, I thank him for his friendship and patience, for all the things he taught me, and the way he promotes thinking on my own. Additionally, I thank Dr. Lannett Edwards for her guidance, constant support, and for teaching me how to perform *in vitro* culture of bovine oocytes and embryos in her laboratory. I also wish to thank Drs. Patty Tithof and Jay Whelan for serving on my graduate committee, critical review of this thesis, and helpful suggestions.

I would also like to thank all my friends: Tracy and Bob, Mitch, Tim, Heather, Roberto, Paul, Jimmy, Howard, Susan, Kim, Debbie, Patty, Alan, TJ, Janelle, Becca, Melody, Kristen, Nancy and the “lunch room gang” for the happiness and fun they brought into my life.

Finally, I dedicate this thesis to my wife, Magdalena, whose love and support have filled my life throughout these years, and to all my family in Argentina, especially to my parents, Nestor and Graciela, my sister, Silvana, her husband, Fabricio, and my little niece, Maria Lujan. Thank you for your encouragement and love that never wavers.

ABSTRACT

Two studies were performed to determine effects of prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) on continued embryonic development of pre-compacted (*in vitro*-produced) and compacted (*in vivo*-derived) bovine embryos. In Experiment 1, pre-compacted (16-32 cell) *in vitro*-produced embryos were randomly assigned to receive either: 1) control (KSOM-PVA; n= 168); 2) PGF-1 (1 ng/mL $PGF_{2\alpha}$ in KSOM-PVA; n= 143); 3) PGF-10 (10 ng/mL $PGF_{2\alpha}$ in KSOM-PVA; n= 168); 4) PGF-100 (100 ng/mL $PGF_{2\alpha}$ in KSOM-PVA; n= 136), or 5) PGE-5 (5 ng/mL PGE_2 in KSOM-PVA; n= 62). Following four days of culture (5.5% CO_2 , 7% O_2 , and 87.5% N_2 at 38.5°C) in assigned treatments, development of embryos was determined by experienced technicians uninformed of treatments. Continued development of *in vitro*-produced 16- to 32-cell embryos (pre-compacted) to blastocyst was reduced by addition of $PGF_{2\alpha}$ in culture media (PGF-1, 30.4%; PGF-10, 41.4%; PGF-100, 33.3%, and control, 51.8%; $P=0.002$). In addition, culture of embryos in medium containing 1 ng/mL of $PGF_{2\alpha}$ had a more detrimental effect than addition of 10 ng/mL of $PGF_{2\alpha}$ ($P<0.05$). Furthermore, blastocyst development did not differ between control and PGE_2 treatments ($P>0.10$).

In Experiment 2, compacted (*in vivo*-derived) glycerol-frozen embryos were thawed and then allowed to regain normal morphology in holding medium for 30 min. Embryos were then sorted by stage of development and quality according with IETS guidelines for classification of bovine embryos. Compacted morula in the quality score of 1, 2 or 3 (n= 436) were randomly assigned to one of four treatments: 1) control (KSOM-PVA; n= 110); 2) PGF-0.1 (0.1 ng/mL $PGF_{2\alpha}$ in KSOM-PVA; n=108); 3) PGF-1 (1

ng/mL PGF_{2α} in KSOM-PVA; n=109); 4) PGF-10 (10 ng/mL PGF_{2α} in KSOM-PVA; n=109). After 24 h in culture (5.5% CO₂, 7% O₂, and 87.5% N₂ at 38.5°C), embryos were washed and placed in KSOM containing 0.5% BSA without PGF_{2α} for an additional 48 h. Embryo development was evaluated by experienced technicians uninformed of treatments. Continued development of *in vivo*-derived compacted morula to blastocyst was not affected by addition of PGF_{2α} to the culture medium (control, 68.1%; PGF-0.1, 58.9%; PGF-1, 61%, and PGF-10, 60%; $P>0.10$). However, hatching rates of embryos cultured with 0.1, 1 and 10 ng/mL PGF_{2α} were significantly lower than controls (24, 29.1, 24.5, and 44.5, respectively; $P=0.05$). No significant differences in hatching rates between doses of PGF_{2α} were observed.

In conclusion, these studies suggest a direct negative effect of PGF_{2α} on continued embryonic development of pre-compacted and compacted bovine embryos. Responses to PGF_{2α} of pre-compacted (decreased blastocyst development) and compacted (decreased hatching rates) embryos may involve different mechanisms of action of PGF_{2α} depending on stage of development. In compacted embryos, PGF_{2α} may act through inhibiting completion of blastocoel formation and hatching; whereas, PGF_{2α} may have more dramatic effects by altering compaction in pre-compacted embryos.

TABLE OF CONTENTS

1. INTRODUCTION	1
2. LITERATURE REVIEW	3
Early embryonic development	3
Embryonic genome activation	7
Compaction process in embryos	8
Involvement of cadherins in embryonic development.....	10
Role of gap junctions in embryonic development	11
Importance of tight junctions in embryo development	15
Cell death during embryonic development	16
Prostaglandin synthesis and action	18
Mechanism of action of PGF _{2α}	21
Prostaglandin F _{2α} association with early embryonic loss.....	24
Summary of literature	27
3. EXPERIMENTAL PROCEDURES	28
Materials	28
Preliminary experiment.....	28
Preparation of prostaglandin stocks	29
Experimental methods for Experiment 1	30
Oocyte collection	30
Sperm preparation and <i>in vitro</i> fertilization.....	31
<i>In vitro</i> culture.....	31
Experimental methods for Experiment 2	34
Radioimmunoassay for PGF _{2α} concentrations in culture media	35
Guide for classification of bovine embryos.....	37
Statistical analysis.....	37
4. RESULTS AND DISCUSSION	41

LIST OF REFERENCES.....	51
VITA.....	66

LIST OF FIGURES

Figure 2-1. Onset of key regulatory events during development of bovine embryos	5
Figure 2-2. Location and timing of development in bovine embryos.....	6
Figure 2-3. Schematic representation of prostaglandin F _{2α} synthesis	20
Figure 3-1. Schematic illustration of experimental design for Experiment 1	33
Figure 3-2. Schematic illustration of experimental design for Experiment 2.....	36
Figure 3-3. Concentrations of PGF _{2α} in the culture medium at the beginning (day 5) and at the end (day 9) of experimental period for Experiment 1.....	38
Figure 3-4. Concentrations of PGF _{2α} in the culture medium for Experiment 2	39
Figure 4-1. Percentage of pre-compacted embryos that continued development to blastocyst in Experiment 1	42
Figure 4-2. Percentage of compacted morula embryos that continued development to blastocyst in Experiment 2	47
Figure 4-3. Percentage of compacted morula embryos that continued development to blastocyst and hatched from the zona pellucida in Experiment 2.....	48

CHAPTER 1

INTRODUCTION

Inadequate reproductive performance in dairy and beef cattle reduces production efficiency and results in a profound economic impact to the cattle industry. Net calf crop (percent of beef cows weaning a calf) in the United States has been estimated to be between 65 to 85% (Bellows et al., 1979), which indicates a high incidence of embryonic or fetal wastage. Early embryonic death accounts for 75 to 80% of reproductive losses between fertilization and calving (Sreenan and Diskin, 1983). Embryonic losses appear to occur soon after the embryo enters the uterus (5-8 days after mating or insemination), when the morula stage embryo is developing into blastocyst (Ayalon, 1978; Maurer and Chenault, 1983; Wiebold, 1988; Dunne et al., 2000).

Numerous factors such as heat stress (Putney et al., 1989; Malayer et al., 1990), nutrition (Butler, 1998), mastitis (Hockett et al., 1998), manipulation of the reproductive tract during embryo transfer and artificial insemination (Wann and Randel, 1990; Velez et al., 1991), and plant toxins (tall fescue; J. W. Oliver, personal communication) may contribute to early embryonic losses through premature release of uterine luminal concentrations of prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) in cattle. Schrick and coworkers (1993) reported that embryo quality in beef cattle tended to be negatively correlated with elevated uterine luminal concentrations of $PGF_{2\alpha}$. Furthermore, addition of $PGF_{2\alpha}$ to culture medium has been shown to inhibit *in vitro* development of rabbit (Maurer and Beier, 1976) and rat embryos (Breuel et al., 1993a). In addition, several studies have shown that administration of $PGF_{2\alpha}$ on days 5 through 8 following artificial insemination in cows receiving supplemental progestogen resulted in reduced pregnancy rates and

retarded development of embryos (Buford et al., 1996; Hockett et al., 1998; Seals et al., 1998; Lemaster et al., 1999). These results suggest a direct and/or indirect effect of $\text{PGF}_{2\alpha}$ on embryonic survival in the cow.

The mechanisms by which $\text{PGF}_{2\alpha}$ reduces embryonic survival and development are unknown. Therefore, the objective of this thesis was to determine if direct effects of $\text{PGF}_{2\alpha}$ occur on embryonic development during culture of pre-compacted (*in vitro*-produced) or compacted (*in vivo*-derived) bovine embryos.

CHAPTER 2

LITERATURE REVIEW

Early embryonic development

Establishment of a viable pregnancy in cattle depends on many complex processes. Functional gametes must be produced by both sexes, the female must show estrus, estrus must be detected by the male, mating must be completed within the functional lifespan of the gametes, and proper uterine environment and corpus luteum functionality must ensure embryonic and fetal development.

In the cow, a population of primordial follicles containing an oocyte arrested at prophase of the first meiotic division (immature oocyte) is established during fetal life (van Wezel and Rodgers, 1996). During each estrous cycle, two or three follicular waves occur in which a group of follicles are recruited to initiate growth (Savio et al, 1988; Ginther et al., 1989; Fortune, 1993). During each follicular wave, only one of these follicles becomes dominant and continues to grow, while subordinate follicles regress (Knopf et al., 1989). Upon luteolysis (destruction of the corpus luteum by $\text{PGF}_{2\alpha}$ of endometrial or exogenous origin), the dominant follicle increases size and produces estradiol 17β to “trigger” a preovulatory peak of luteinizing hormone (LH) and estrous behavior. Luteinizing hormone causes the oocyte in the dominant follicle to mature (progresses to metaphase of the second meiotic division) and ovulation to take place (Espey, 1994).

After expulsion from the ovulatory follicle, the oocyte is transported to the oviduct. The oviduct is comprised of four distinct regions: infundibulum, ampulla, ampullary-isthmic junction, and isthmus (reviewed by Ellington, 1991). The mature

oocyte is captured by the infundibulum and transported to the ampullary-isthmus junction where capacitated sperm binds to the zona pellucida surrounding the oocyte, undergoes acrosome reaction and penetrates the oocyte's plasma membrane (reviewed by Saling, 1991). After fertilization and organization of the male and female pronuclei, the zygote becomes an embryo that undergoes a series of synchronized cell divisions (also called cleavage divisions; Figure 2-1).

The first cleavage division generates a two-cell embryo, in which cells are called blastomeres. As the early embryo is transported through the oviduct (isthmus region), each blastomere undergoes subsequent divisions yielding 4-, 8-, 16- and 32-cell embryos (Figures 2-1; 2-2). The embryo reaches the uterine horn on day 5-6 after estrus; at which time, blastomeres flatten against each other forming a "solid ball" of cells called morula (Figure 2-2). The morula stage embryo begins to form a fluid filled cavity (the blastocoel) and the embryo becomes a blastocyst (Watson et al., 1992). Throughout all these developmental stages, blastomeres become smaller with no net increase in size of the embryo. As a result of additional fluid accumulation within the blastocoel, the embryo starts to expand (expanded blastocyst) and finally hatches from the zona pellucida (hatched blastocyst) on approximately day 9 after estrus (Figure 2-2; Van Soom et al., 1997).

For attachment to the uterine endometrium, the blastocyst must hatch (escape) from the zona pellucida (Figure 2-1). Thus, embryo hatching is a critical step for a successful pregnancy in cattle. Furthermore, low implantation rates of *in vitro*-fertilized human embryos are largely due to impaired development and hatching of blastocyst (Magli et al., 1998).

5




Figure 2-1. Onset of key regulatory events during development of bovine embryos. EGA= embryonic genome activation.













<u>Location</u>	<u>Day</u>	<u>Development</u>			
Oviduct (Ampulla)	1	Oocyte			
Oviduct (A-I Junction)	2	1 to 2 cell			
Oviduct (Isthmus)	2-3	4 cell			
Oviduct (Isthmus)	3-5	8 cell			
Uterus	5	16 to 32 cell (pre-compacted)			
Uterus	6	Morula, Compacted Morula			
Uterus	7,8,9	Blastocyst Expanded Blastocyst, Hatched Blastocyst			

Figure 2-2. Location and timing of development in bovine embryos.

The process of hatching in bovine embryos has been well documented by Massip and Mulnard (1980). In cattle, accumulation of fluid in the blastocoel of *in vivo*-derived embryos distends the zona pellucida (ZP), which becomes gradually thinner. Then, the ZP breaks rapidly and hatching begins immediately by protrusion of the blastocyst through the opening in the ZP. The ZP does not show signs of lysis but is perforated in one location only. Moreover, zona-pellucida-intact embryos, as well as empty ZP, can be found in uterine flushings up to day 16 after ovulation (Betteridge et al., 1980) indicating that enzymatic lysis of the ZP does not occur during hatching in cattle.

The fully expanded bovine blastocyst must exert hydrostatic pressure to create a hole in the ZP through which the blastocyst egresses. Fluid accumulation in the blastocoel is achieved by the presence of a $\text{Na}^+\text{-K}^+\text{-ATPase}$ pump and tight junctions in the trophoctoderm epithelium. On the other hand, *in vitro* studies in hamsters and mouse indicated that blastocyst expansion is followed by a focal ZP thinning preceding a focal ZP rupture, protrusion of the embryo, and complete ZP dissolution. These findings indicate that enzymatic degradation of the ZP is essential for hatching of hamsters and mouse embryos (Kane and Bavister, 1988; Gonzalez and Bavister, 1995; O'Sullivan, et al., 2001). In fact, hatching of *in vitro*-produced mouse blastocyst has been related to a trypsin-like proteinase associated with cells of the trophoctoderm (Perona and Wassarman, 1986; Sawada et al., 1990).

Embryonic genome activation

Maternal mRNAs and proteins accumulated in the oocyte's cytoplasm during the growth phase within the follicle are able to support early embryonic divisions (Arends

and Willye, 1991). Further development is achieved by activation of the embryonic genome, which differs in time of initiation between species: 1-cell stage in the mouse (Aoki et al., 1997), 4-cell stage in humans (Braude et al., 1988), and 8- to 16-cell stage in cattle (Camous et al., 1984; Figure 2-1), sheep (Crosby et al., 1988), and rabbits (Manes, 1973). The majority of maternal proteins and mRNA are degraded during transition from maternal to embryonic genome activation (Latham et al., 1991). Memili and coworkers (1998) suggested minor genome activation between the 1- and the late 4-cell stages and a major gene activation at the 8-cell stage in bovine embryos. During embryonic genome activation, several important genes for cell proliferation, compaction and blastocyst formation are activated. Some of those genes include expression of growth factors and their receptors (Rappolee et al., 1990), connexins (Kidder, 1987) and Na⁺-K⁺ ATPase enzyme (Watson and Kidder, 1988).

Compaction process in embryos

Compaction represents one of the first key critical events important for continued development. It allows polarization and differentiation of blastomeres into two different cell populations (by induction of cell adhesion molecules between blastomeres that enhance cell communication and adherence) and facilitates fluid accumulation within the embryo that directly contributes to blastocyst formation and subsequent hatching from the zona pellucida (Watson et al., 1992). Compaction appears to occur in all mammalian embryos but timing differences exist between species: mouse at 8-cell stage (Ducibella and Anderson, 1975), humans at 16-cell stage (Edwards et al., 1981), and bovine (Van Soom et al., 1997; Figure 2-1) and rabbit (Koyama et al., 1994) at 32-cell stage.

During compaction, blastomeres adhere to and flatten against each other so boundaries cannot be recognized between cells (Van Soom et al., 1997). During this period, development of Ca^{++} -dependent cell adhesion molecules such as E-cadherin and establishment of gap and tight junctions lead to cellular and morphogenetic differentiation of blastomeres. Outer blastomeres polarize and form the trophoctoderm (TE), while inner blastomeres remain apolar and produce the inner cell mass (ICM; Koyama et al., 1994). The inner cell mass forms the embryo proper and some extra-embryonic membranes (allantoids and amnion); whereas, trophoctoderm cells will combine with the ICM-derived extra-embryonic membranes to form the fetal placenta (Schlafer et al., 2000).

The free or apical membrane surfaces of the TE develop a microvillus cap with several Na^+ -dependent transport systems (Miller and Schultz, 1985; Fleming and Johnson, 1988). These transport systems play a central role in establishment of an ion concentration gradient across the epithelium that facilitates osmotic accumulation of water into the blastocoelic space to form the nascent blastocoel (Biggers et al., 1988). Basolateral surfaces of TE remain free of microvilli, but become distinguished from apical surfaces by localization of tight junctions (Fleming et al., 1989), gap junctions (Kidder, 1987), adherent junctions (Vestweber et al., 1987), and acquisition of an active Na^+ - K^+ ATPase pump (Watson et al., 1999). Even though all mammalian embryos undergo compaction, some differences exist between *in vivo*-derived and *in vitro*-produced bovine embryos (Van Soom et al., 1997). Both embryo types start compaction at 32-cell stage, but compaction in *in vitro*-produced embryos is shorter and less obvious than in *in vivo*-derived embryos.

In summary, development of cell adhesion molecules and gap junction-mediated interblastomeric cell communication induces cell polarization and differentiation of undifferentiated blastomeres into TE epithelium and ICM during compaction. Moreover, the trophectoderm develops the capacity to initiate and regulate blastocoel formation by the action of $\text{Na}^+\text{-K}^+$ ATPase enzyme (in the basolateral surface TE epithelium). Additionally, presence of a junctional apical complex (formed by tight junction at the apical part of the TE epithelium) prevents uncontrolled leakage of this fluid from the blastocoel cavity (Watson et al., 1992).

Involvement of cadherins in embryonic development

Cell adhesion molecules play a critical role during compaction and blastocyst formation in the early embryo, and therefore in subsequent development and viability of the conceptus. Cadherins are a superfamily of Ca^{++} -dependent adhesion molecules that provide strong mechanical attachment between adjacent cells. Cadherin molecules have three major domains or regions: an extracellular domain responsible for specific recognition of the same cadherin present in neighboring cells, a transmembrane domain that spans the cell membrane, and a cytoplasmic domain that extends into the cell and associates with cytoplasmic proteins called catenins, which in turn bind cadherins to actin filaments in the cytoskeleton (Ivanov et al., 2001).

In the mouse, E-cadherin is expressed in the oocyte and during the early period of embryonic genome activation (at 2-cell stage; Vestweber et al., 1987). Also, E-cadherin-mediated adhesion between blastomeres initiates compaction and plays a critical role in differentiation of the trophectoderm and morphogenesis of the mouse blastocyst (Fleming

et al., 2001). E-cadherin is uniformly distributed on the surface of all mouse blastomeres during early cleavage but redistributes to cell-cell contact sites upon activation at the 8-cell stage (when compaction begins) suggesting post-translational regulations. In fact, E-cadherin becomes phosphorylated for the first time at compaction (Sefton et al., 1992; Figure 2-1). Activation of PKC with phorbol ester or treatment with diacylglycerides caused 4-cell mouse embryos to compact ahead of normal timing, while PKC inhibitors prevented compaction (Winkel et al., 1990). Moreover, mouse embryos lacking E-cadherin and alpha catenin failed to generate a normal trophectoderm and died before implantation (Laure et al., 1994; Riethmacher et al., 1995). In cattle, E-cadherin is evenly distributed around cell margins of 1-cell zygotes and cleavage stage embryos and becomes restricted to basolateral membranes of trophectoderm cells during morula stage, while maintaining apolar distributions in the ICM (Barcroft et al., 1998).

Role of gap junctions in embryonic development

Gap junctions are channels present in opposing plasma membranes of neighboring cells (Kidder and Winterghager, 2001). The cell membranes come close together, but are separated by 2-4 nanometers. These junctions are composed of at least 13 different proteins called connexins (Bruzzone and Goodenough, 1996; De Sousa et al., 1997); however, several studies identified connexin 43 as the most important gap junctional protein expressed in mouse (De Sousa et al., 1993; Valdimarsson and Kidder, 1995), bovine (Wrenzycki et al., 1996; 1999), and rat (Reuss et al., 2002) preimplantation embryos.

Intercellular communication through gap junctions is believed to be essential for cellular growth, cellular differentiation and development (Warner, 1992; Zhang and Thorgeirsson, 1994). Recently, a targeted gene inactivation approach has been used to explore the functional significance of connexin diversity. For example, connexin 26 is expressed in the labyrinth of the mouse placenta and its absence causes impaired metabolite transfer from maternal blood to the developing embryo, leading to death in mid-gestation (Gabriel et al., 1998). Female mice lacking connexin 37, which is present in the ovary, develop a block in ovarian follicular development and are sterile (Simon et al., 1997). Although connexin 43 is the most widely expressed connexin, offspring lacking connexin 43 survive until birth and die soon after as a result of morphological heart defects (Reaume et al., 1995), but they do not show any other abnormalities when compared to animals with normal expression of connexin 43. De Sousa et al. (1997) showed that preimplantation mouse embryos deficient in connexin 43 retained the capacity for cell-to-cell transfer of fluorescent dye (dye coupling), but at a severely reduced dye coupling. Researchers suggested that loss of connexin 43 can be compensated by other connexins such as connexin 45, a protein also assembled into membrane plaques at the time of compaction.

At the plasma membrane, six copies of the same connexin form a hexagonal ring-shaped channel called connexon. Connexons in adjacent cell membranes align together to form a hydrophilic channel that allows passage of ions and small molecules such as cyclic AMP, Ca^{++} and inositol tri-phosphate (IP_3) between cells and thereby coordinate metabolic and electrical activities (Lowenstein, 1981). Studies using Lucifer yellow have shown that blastomeres of *in vivo*- and *in vitro*-produced mouse embryos are coupled by

functional gap junctions (De Sousa et al., 1993). Boni et al. (1999) showed that diffusion of dye between ICM cells started at morula stage in *in vivo*-derived and at the blastocyst stage in *in vitro*-produced bovine embryos. In the same study, gap junctions were found in the ICM as well as between ICM and TE, but not between TE cells of *in vitro*-derived embryos. However, gap junctions were present through all cells and in a higher density in *in vivo*-derived embryos.

Although assembly of functional gap junctions in mammalian preimplantation embryos is a temporally regulated event occurring during compaction (Figure 2-1), both mRNA and protein corresponding to connexin 43 are being synthesized as early as the 4-cell stage in mouse embryos (one cell cycle before compaction; De Sousa et al., 1993). Injection of gap junction antisense RNA to 2- and 4-cell mouse embryos reduced the percentage of compacted embryos to less than 20%, while 90% of uninjected embryos compacted (Bevilacqua et al., 1989). Also, when the same antisense was injected to 8-cell stage mouse embryos, only 5% of the embryos developed to blastocyst stage, while blastulation occurred in 90% of control embryos. Similarly, injection of gap junction antibodies inhibited dye transfer and dramatically reduced electrical coupling of 8-cell mouse embryos (Lee et al., 1987).

In mouse embryos, De Sousa et al. (1993) demonstrated that connexin 43 shows different localization between blastomeres of morula and blastocyst embryos: a zonular distribution predominated between outside blastomeres and trophectoderm cells; whereas, plaque-like localizations predominated between inside blastomeres and cells of the inner cell mass. In the same study, treatment of uncompact 8-cell mouse embryos with monensin or brefeldin-A (inhibitors of protein trafficking) prevented the appearance

of gap junction-like structures. In another study, inhibition of DNA synthesis with aphidicolin during the third and fourth cell cycles of mouse embryos had no effect on establishment of gap junctional coupling, but administration of aphidicolin for 10 h during the second cell cycle resulted in failure of gap junctional coupling (Valdimarsson and Kidder, 1995). Thus, timing of gap junctional coupling is linked to DNA replication in the 2-cell mouse embryo in the mouse.

Data on intercellular communication through gap junctions in early bovine embryos are limited. Utilizing polymerase chain reaction (PCR), Wrenzycki and colleagues (1996) reported that transcripts for connexin 43 were present in immature oocytes and all developmental stages of *in vitro*-produced embryos through morula stage; whereas, only *in vivo*-derived morula and blastocyst stages expressed connexin 43. This findings correlate with reports from Prather et al. (1993) reporting that *in vivo*-derived morula and blastocyst embryos showed dye transfer indicating the existence of gap junctions, while *in vitro*-produced morula and blastocyst embryos did not show spreading of dye.

In cultured vascular cells, gap junctional intercellular communication is inhibited by PKC-dependent excessive phosphorylation of connexin 43, which is the primary gap junction in vascular cells. In addition, excessive phosphorylation of the same protein induces impairment of ventricular conduction in the heart (Inoguchi et al., 2001). Furthermore, Criswell (1995) demonstrated that high intracellular levels of Ca^{++} , addition of Ca^{++} ionophore 4-br-A23187, increased PKC activity and inclusion of TPA (an activator of PKC) resulted in inhibition of gap junctional communication in cultured uterine smooth muscle cells.

Importance of Tight junctions in embryonic development

Tight junctions (zonula occludens) seal the trophectoderm epithelium in a narrow band just beneath their apical surfaces and right above zonula adherens junctions. Tight junctions represent a site of close intercellular adhesion including partial membrane fusion that forms a barrier against paracellular diffusion, generates a trans-epithelial resistance, and allows blastocoel formation (Biggers et al., 1988). Tight junctions comprise several proteins (both transmembrane and cytoplasmic) with the most important being zonula occludens 1 (ZO-1) and cingulin (Citi, 1993), which are only present within the trophectoderm. Following compaction, polar cells adhere to and gradually envelope nonpolar cells causing ZO-1 in the nonpolar cells to disappear. Thus, symmetrically cell contact appears to initiate ZO-1 down regulation in the ICM lineage (Fleming and Hay, 1991). In cattle, ZO-1 is undetectable until the morula stage when it first appears as punctuate points between outer cells (Barcroft et al., 1998; Figure 2-1). In the blastocyst, ZO-1 is localized as a continuous ring at the apical points of trophectoderm cell contact, whereas it remains undetectable in the ICM.

Although tight junction formation does not begin until compaction of the 8-cell stage mouse embryo (a process dependent upon E-cadherin activation), immunoblotting of unfertilized eggs and preimplantation mouse embryos showed high levels of cingulin in eggs and early cleavage stages, a decline during later cleavage, and an increase in cingulin levels during the stage of expanding blastocyst (Javed et al., 1993). In cattle, ZO-1 is undetectable until morula stage when first appears as punctuate points between outer cells. In the blastocyst, ZO-1 is localized as a continuous ring at the apical points of

trophectoderm cell contact, whereas remains undetectable in the ICM (Barcroft et al., 1998).

Cell death during embryonic development

Cell death occurs as a normal feature of early embryonic development in both *in vivo* and *in vitro*-derived embryos (Jurisicova et al., 1996; Hardy, 1997; Matwee et al., 2000). Two forms of cell death can be distinguished based on morphological and molecular criteria (Willye et al., 1980; Granville et al., 1998).

Necrosis is characterized by nuclear disintegration, cellular swelling, rupture of internal and external membranes with release of lytic enzymes, and damage to surrounding cells by stimulation of an inflammatory reaction (Majno and Joris, 1995). Necrosis is considered to be an “accidental” cell death initiated by gross environmental insults such as severe hypoxia, heat shock and exposure to metabolic toxins (Willye et al., 1980). In contrast, apoptosis (also called programmed cell death) occurs in single rather than groups of cells and is thought to be a self-directed death or “cell suicide” to assure normal development and differentiation by removal of damaged or defective cells (Edwards, 1998).

Apoptosis is characterized by a series of morphological changes. First, chromatin aggregates into large compact granular masses on the inner nuclear membrane, the cytoplasm condenses and the nuclear and cytoplasmic membranes become indented. Then, DNA fragments and the nucleus separate into discrete masses. During the final phases of the apoptotic processes there is extensive blebbing of the plasma membrane

with the formation of membrane-bound apoptotic bodies, which can be phagocytosed by surrounding cells or extruded into an adjacent lumen (Willye et al., 1980).

In addition to morphological changes, biochemical alterations at the plasma membrane or at the DNA structure are also observed. In the earliest stages of apoptosis, phosphatidyl-serine, a membrane phospholipid generally associated to the inner cytoplasmic leaflet, is exposed to the outer leaflet and promotes phagocytosis by neighboring cells (Martin et al., 1995). This feature is useful in determining apoptosis by utilizing the anticoagulant, protein annexin V, which has a specific and high affinity for phosphatidyl-serine. Healthy cells incubated in fluorescein isothiocyanate (FITC)-conjugated annexin V remain unlabelled, as annexin V can not access to the inner leaflet of the plasma membrane. However, cells undergoing apoptosis show a marked increase in annexin V binding, indicating that phosphatidyl-serine is translocated to the outer leaflet (Martin et al., 1995).

Rounding and separation of apoptotic cells from neighboring cells is associated to a lack of gap junction communication between cells (Hardy et al., 1996). Deoxyribonucleic acid between nucleosomes is cleaved by endonucleases into oligonucleosomal fragments (Arends and Willye, 1991; Rueda et al., 1995). These fragments can be visualized as a typical "DNA ladder" with bands at 180-200 bp intervals by agarose gel electrophoresis.

Due to the small number of cells in preimplantation embryos, it is impossible to use electrophoretic techniques to observe for DNA laddering typical of apoptotic nucleus (Hardy, 1999). Fortunately, development of TdT-mediated dUTP nick-end labelling (TUNEL) enables assessment of nuclear DNA fragmentation *in situ* (Gavrieli et al.,

1992). Matwee et al. (2000) observed TUNEL labelling in bovine oocytes, 8- to 16-cell embryos, morula, and blastocyst stage embryos, but labelling was not observed in zygotes or early cleavage stage embryos. Similar results were observed in human (Jurisicova et al., 1996) and porcine embryos (Long et al., 1998) suggesting that occurrence of apoptosis is dependent on developmental stage of the embryo and early stage embryos (before compaction) seem to be more resistant to apoptosis than later stages.

Levels of transcripts involved in the apoptotic pathway might be below a specific threshold before embryonic genome activation and cellular differentiation into the trophoctoderm and inner cell mass. Undifferentiated tissues during fetal development in humans possess an inhibitor of caspases activity (Adida et al., 1998). Early stage bovine embryos may possess a similar mechanism (maternally produced proteins and mRNA) to avoid apoptosis until morula stage. Apoptosis may also be an efficient mechanism to eliminate blastomeres that did not begin transcription of the embryonic genome after the 8-cell stage in cattle.

Prostaglandin synthesis and action

Prostaglandins are biochemical compounds derived from the polyunsaturated fatty acid, arachidonic acid. Prostaglandins are involved in several physiological processes such as inflammation, production of pain and fever, control of blood vessel tone and arterial blood pressure, luteolysis, and blood clotting (Nett et al., 1976; Seibert et al., 1994; Blatteis and Sehic, 1997). With the exemption of red blood cells, prostaglandins are synthesized and released by all mammalian cells and tissues. However, unlike other

hormones, prostaglandins are not stored in cells but are produced and released immediately after stimulation (Bito, 1975).

Phospholipase A₂ (PLA₂) catalyzes the hydrolysis of the sn-2 position of membrane glycerophospholipids to liberate arachidonic acid (AA), a precursor of eicosanoids including prostaglandins and leukotrienes. So far, at least 19 enzymes that possess PLA₂ activity have been identified in mammals (reviewed by Murakami and Kudo, 2002). Since arachidonic acid contains 20 carbon atoms, prostaglandins are referred to eicosanoids. Arachidonic acid is derived from membrane phospholipids primarily by the action of cytoplasmic phospholipase A₂ (Flint et al., 1986). This cleavage state is the rate-limiting step in prostaglandin synthesis (Figure 2-3). Free arachidonic acid is then converted to prostaglandin H₂ (PGH₂) by the action of cyclooxygenase enzymes at the membranes of the endoplasmic reticulum (Smith and DeWitt, 1996) or the nucleus (Parfenova et al., 2001).

There are two forms of cyclooxygenase: COX-1, a constitutive enzyme found in almost all cells; and COX-2, an inducible form in response to inflammatory reactions, hormonal and environmental stimulus (Smith and DeWitt, 1996). Cyclooxygenase-1 and -2 catalyze the conversion of free arachidonic acid to PGH₂ by two sequential enzymatic steps: oxygenation, which converts free arachidonic acid to the unstable intermediate PGG₂, followed by a peroxidation that converts PGG₂ to PGH₂ (Smith and DeWitt, 1996).

Different specific enzymes, whose presence varies depending on the cell type, mediate formation of the D, E, I, and F series from PGH₂ in different tissues. Prostaglandin production can be blocked by administration of nonsteroidal anti-




Figure 2-3. Schematic representation of prostaglandin $F_{2\alpha}$ synthesis.

inflammatory drugs (NSAIDs) such as aspirin, indomethacin and phenylbutazone and by steroidal anti-inflammatory drugs like hydrocortisone, prednisone and prednisolone. Nonsteroidal anti-inflammatory drugs act through inhibition of COX-1 and COX-2 (Vane and Botting, 1996), while steroidal anti-inflammatory drugs interfere with mobilization of arachidonic acid from the plasma membrane by inhibition of phospholipase A₂ (Rothhut and Russo-Marie, 1984).

Mechanism of action of PGF_{2α}

In nonpregnant cows, PGF_{2α} is released from the uterus in a pulsatile manner to cause regression of the corpus luteum (Kindahl et al., 1976) and allows ovulation of the dominant follicle. Oxytocin stimulates uterine secretion of PGF_{2α} when administered *in vivo* (Silvia and Taylor, 1989) and *in vitro* (Lafrance and Goff, 1990). Oxytocin released from large luteal cells binds to its receptors in the uterus endometrium and stimulates uterine secretion of PGF_{2α} by activation of phospholipase A₂ and conversion of arachidonic acid to PGF_{2α}. Then, uterine PGF_{2α} stimulates further oxytocin release from the corpus luteum (CL; Flint and Sheldrick, 1982). Thus, a positive feedback loop between PGF_{2α} and oxytocin seems responsible for the generation of each pulse of PGF_{2α} during luteolysis.

The PGF_{2α} receptor is a seven transmembrane G-protein coupled receptor (Sakamoto et al., 1994). After binding to its receptor, PGF_{2α} activates a membrane-bound phospholipase C, which causes hydrolysis of membrane phosphatidylinositol bisphosphate (PIP₂) to inositol triphosphate (IP₃) and diacylglycerol (DAG) (Berridge, 1987). Increased concentrations of IP₃ result in a release of free Ca⁺⁺ from intracellular

stores to the cytosol, while DAG enhances the affinity of protein kinase C (PKC) for Ca^{++} , which results in PKC activation.

Although information regarding the presence of $\text{PGF}_{2\alpha}$ receptors in early stage embryos has not been found in the literature, the bovine CL represents an excellent model for understanding the mechanisms of action of $\text{PGF}_{2\alpha}$. Protein kinase C is believed to mediate many of the antisteroidogenic actions of $\text{PGF}_{2\alpha}$ in large luteal cells (McGuire et al., 1994). In addition to this, $\text{PGF}_{2\alpha}$ may interfere with the ability of LH to activate PKA, an enzyme needed for phosphorylation and stimulation of other key enzymes in the luteal steroidogenic pathway, such as cholesterol esterase (hydrolyzes the stored cholesterol) and steroidogenic acute regulatory protein (StAR), which transports cholesterol to the mitochondria (Garverick et al., 1985). In contrast to early signaling events, downstream signals in response to $\text{PGF}_{2\alpha}$ are poorly understood. Chen et al. (1998) reported that $\text{PGF}_{2\alpha}$ activates the Raf/Mek1/mitogen-activated protein kinase (MAPK) signaling cascade in bovine luteal cells. Therefore, signals on the cell surface may be responsible for transcription of luteal genes.

Prostaglandin $\text{F}_{2\alpha}$ promotes apoptosis in luteal cells by activation of endonucleases after influx of Ca^{++} from intracellular stores (Juengel et al., 1993). Also during luteolysis in cattle, mRNA encoding Bax (a pro-apoptotic gene) is elevated resulting in an increased ratio of Bax to Bcl-2 (anti-apoptotic gene), an event consistent with Bax-mediated apoptosis (Rueda et al., 1997). Prostaglandin $\text{F}_{2\alpha}$ also induces apoptosis in luteal cells by generation of oxygen radicals such as superoxide anion, hydroxyl radical and hydrogen peroxide (Riley and Behrman, 1991; Carlson et al., 1993). Exposure of rat luteal cells to these oxidants resulted in oxidation and loss of fluidity in

cellular membranes (Sawada and Carlson, 1991). Decreased membrane fluidity affects cellular function including receptor binding and membrane-bound enzyme activity as demonstrated by Wu et al. (1992), who suggested that superoxide radical production inhibited LH stimulation of cyclic AMP and decreased activity of protein kinase A. In pigs, $\text{PGF}_{2\alpha}$ depleted the CL of vitamin C, a natural antioxidant that prevents oxygen radical formation (Petroff et al., 1999).

Although $\text{PGF}_{2\alpha}$ signals cells through G-protein coupled receptors, other prostaglandins such as the PGJ_2 series are actively incorporated into the nucleus and associate with nuclear proteins by unknown mechanisms (Narumiya et al., 1987). 15-deoxy-12, 14- PGJ_2 is the final metabolite of PGD_2 degradation and is a potent ligand for a nuclear hormone receptor known as peroxisome proliferator-activated receptor γ (PPAR- γ). Peroxisome proliferator-activated receptor- γ is a member of a superfamily of nuclear hormone receptors that function as ligand-dependent transcription factors (Schoonjans et al., 1996). Three receptor subtypes of PPARs (termed α , δ , and γ) have been identified. The transcriptional activity of PPAR subtypes is enhanced by numerous chemical compounds including fatty acids, selected prostaglandins, and peroxisome proliferators (Schoonjans et al., 1996).

After activation, PPARs control expression of genes implicated in extra- and intracellular lipid metabolism involved in peroxisomal β -oxidation pathways (Schoonjans, 1996). Padilla et al. (2000) reported that PPAR- γ agonist and PGJ_2 produced cytotoxicity and antiproliferative effects on normal and malignant B cells. Cytotoxicity was mediated via apoptosis as shown by positive annexin V staining and

confirmed by DNA fragmentation detected using TUNEL assay. Prostaglandin I₂, a key prostaglandin during mouse implantation, also mediates its actions through activation of PPARs (Hyunjung et al., 1999).

Prostaglandin F_{2α} is a potent, bioactive compound. As such, its production and metabolism must be tightly regulated. The lungs appear to play an important role in inactivating prostaglandins. In the ewe, PGF_{2α} in the blood stream has a half-life of less than 1 min, since 99% of it is metabolized to 13,14-dihydro-15-keto-PGF_{2α} (PGFM) in a single passage through the lung (Hansel and Dowd, 1986). In the cow, only 16% of PGF_{2α} remained in the blood after three passages through the lung (Davis et al., 1984).

Prostaglandin F_{2α} association with early embryonic loss

The first indications that PGF_{2α} were involved in embryonic loss occurred in a series of experiments in the “short cycle” cow at West Virginia University. In the postpartum cow, the first ovulation is followed by a short luteal phase (short cycle) due to premature release of PGF_{2α} (Cooper et al., 1991). Cows with short luteal phase failed to maintain pregnancy even when exogenous progestogen was provided (Breuel et al., 1993b). Moreover, when good quality frozen-thawed embryos were transferred to progestogen-supplemented animals on day 7 after estrus, pregnancy was maintained in 28% of short cycle cows compared to 58% of cows with normal luteal phase (Butcher et al., 1992). However, pregnancy rates did not differ when day-6 embryos from short cycle or normal cycle cows were transferred to normally cycling recipients (Schrick et al., 1993), suggesting that an improper uterine environment appears to play a role in lower survival of embryos in cows with short cycle as compared to cows with normal luteal phases.

Schrick et al. (1993) observed that concentrations of PGF_{2α} in flushing media from cows with short luteal phases were more than double those from cows with normal luteal phases (636 ± 82 and 288 ± 90 pg/ml, respectively). Furthermore, embryo quality tended to be negatively correlated with concentrations of PGF_{2α} in the flushing media ($r = -0.42$; $P = 0.07$). Thus, a direct negative effect of PGF_{2α} may be implicated in lowering embryonic survival as demonstrated *in vitro* in the mouse (Harper and Skarnes, 1972), rabbit (Maurer and Beier, 1976), and rat (Breuel et al., 1993a).

Effects of PGF_{2α} on embryo survival have been examined in several studies using progestogen-supplemented cows to replace the regressing corpus luteum. Buford et al. (1996) demonstrated that PGF_{2α} was detrimental to embryonic survival when 15 mg of PGF_{2α} every 8 h were injected to nonlactating cycling beef cows during days 4 through 7 or 5 through 8 after insemination, but pregnancy rates remained high when lutectomy (removal of the corpus luteum) was performed before PGF_{2α} treatment begun. Similarly, PGF_{2α} administration to progestogen-supplemented beef cows on days 5 to 8 postbreeding reduced pregnancy rates (23%) compared to control (72%), whereas lutectomy prior PGF_{2α} administration tended to improve pregnancy rates (55%; Seals et al., 1998). Removal of the corpus luteum eliminates oxytocin release from the corpus luteum and blocks the positive feedback loop between oxytocin and PGF_{2α}. As a result, PGF_{2α} concentration in the uterine lumen would remain low, allowing embryonic development to occur. Furthermore, Seals et al. (1998) reported that detrimental action(s) of PGF_{2α} occurred during early embryonic development (days 5 to 8) since administration of PGF_{2α} during filamentous embryo development (beyond day 10) had no effect on maintenance of pregnancy.

Buford et al. (1996) demonstrated that postpartum cows bred at their first estrus after weaning had higher pregnancy rates when lutectomy was used in conjunction with flunixin meglumine (an inhibitor of COX-1 and COX-2), but lutectomy alone failed to maintain pregnancy rates. The authors suggested that both premature secretion of PGF_{2α} and a factor released by the regressing corpus luteum may contribute to embryonic death in the postpartum cow with a short luteal phase.

Oxytocin, which is released by large cells of the regressing corpus luteum in response to PGF_{2α}, can increase secretion of PGF_{2α} from the uterus (Milvae and Hansel, 1980). Lemaster et al. (1999) treated progestogen-supplemented cows with oxytocin rather than PGF_{2α} in the same type of experiment performed by Seals et al. (1998). Administration of oxytocin on days 5 through 8 every 8 h reduced pregnancy rates to 33% compared to the control group (80%). In the same study, administration of flunixin meglumine blocked the negative effects of oxytocin, while lutectomy did not; thus, the role of oxytocin on embryonic death may involve further release of PGF_{2α} rather than having a direct negative effect on the embryo.

Numerous factors such as heat stress (Putney et al., 1989; Malayer et al., 1990), nutrition (Butler, 1998), mastitis (Hockett et al., 2000), manipulation of the reproductive tract during embryo transfer and artificial insemination (Odensvik et al., 1993), and plant toxins (tall fescue; J. W. Oliver, personal communication) may contribute to early embryonic losses (before day 8) through premature release of uterine luminal concentrations of PGF_{2α} in cattle. Administration of flunixin meglumine or ibuprofen lysinate (inhibitors of COX-1 and COX-2) at time of embryo transfer improved pregnancy rates in cows (Elli et al., 2001; Schrick et al., 2001). Furthermore, low-dose

aspirin treatment improved implantation and pregnancy rates in human patients undergoing *in vitro* fertilization and transfer (Rubinstein et al., 1999). Thus, *in vivo* and *in vitro* studies have shown that $\text{PGF}_{2\alpha}$ decreased embryonic development and survival in several species.

Summary of literature

Early embryonic wastage results in reduced reproductive performance of cattle and in a vast economic loss to beef and dairy producers. Several factors such as heat stress, nutrition, mastitis, plant toxins and manipulation of the reproductive tract may cause early embryonic death by induction of $\text{PGF}_{2\alpha}$ release from the uterine endometrium.

Presence of $\text{PGF}_{2\alpha}$ has been shown to have direct and indirect negative effects on embryo development; however, the exact mechanisms of $\text{PGF}_{2\alpha}$ responsible of lowering embryo survival are still unclear. Possible explanations may involve: 1) stimulation of the apoptotic cascade (by up regulation of pro-apoptotic genes, activation of endonucleases and/or formation of free radicals); and/or 2) unknown downstream signals capable of phosphorylation/dephosphorylation of key regulatory proteins (such as PKC) during compaction, which may lead to failures to either compact or form a blastocoelic cavity.

The objective of this study was to determine any direct effect of $\text{PGF}_{2\alpha}$ on embryonic development of pre-compacted (*in vitro*-produced) and compacted (*in vivo*-derived) bovine embryos.

CHAPTER 3

EXPERIMENTAL PROCEDURES

Materials

Dimethyl sulphoxide (DMSO), polyvinyl alcohol (PVA), bovine serum albumin (BSA), percoll, penicillamine, hypotaurine, epinephrine and the majority of reagents used for *in vitro* production of bovine embryos were purchased from Sigma Chemical, Inc. (St. Louis, MO). Tissue culture medium-199, gentamicin, and penicillin-streptomycin were purchased from Specialty Media, Inc. (Phillipsburg, NJ). Prostaglandin F_{2α} (PGF_{2α}) and prostaglandin E₂ (PGE₂) were purchased from Cayman Chemical Company (Ann Arbor, MI). Fetal bovine serum (FBS) was obtained from BioWhittaker (Walkersville, MD). Follitropin-V was provided by Vetrepharm Canada, Inc. (London, Ontario) and luteinizing hormone (LH) was obtained from the United States Department of Agriculture (Beltsville, MD). Vigro was purchased from AB Technology, Inc. (Pullman, WA). Embryos were generously donated by Trans Ova (Sioux City, IA), Em Tran (Elizabethtown, PA), and Harrogate Genetics Inc. (Harrogate, TN). Ovaries were purchased from Brown's Packing Plant (Gaffney, SC). Media (HEPES-TALP, IVF-TALP, and SPERM-TALP; Parrish et al., 1988) and KSOM (Biggers et al., 2000), with modifications generously provided by Dr. John Hasler, were prepared in the laboratory or purchased from Specialty Media.

Preliminary experiment

A preliminary experiment was performed to determine if differences in embryo development existed between KSOM media supplemented with either PVA (3 mg/mL,

0.3%) or BSA (5 mg/mL, 0.5%). This study was performed due to a concern regarding the possibility that PGF_{2α} might bind to BSA present in KSOM culture medium utilized during embryo development (Tanaka et al., 2000). A total of 187 *in vivo*-derived glycerol frozen bovine embryos were thawed using the three-step glycerol removal procedure (see below). Morula stage embryos were cultured for 3 days in either KSOM supplemented with PVA or BSA in a humidified atmosphere of 5.5% CO₂, 7% O₂, and 87.5% N₂ at 38.5°C. After 3 days of culture (corresponding to day 9 of embryo development; Figure 2-2), blastocyst development was assessed using IETS guidelines for classification of bovine embryos (see below).

Chi-square analysis was used to determine potential differences in development of embryos (SAS 8.02, SAS Institute Inc., Cary, NC). There were no differences ($P > 0.10$) in proportion of morula continuing to blastocyst between KSOM-PVA (45.2%) or KSOM-BSA (54.2%) culture media. As such, PVA was added to KSOM media during PGF_{2α} treatments to avoid any possible binding of PGF_{2α} to BSA.

Preparation of prostaglandin stocks

Prostaglandin F_{2α} (1 mg) and PGE₂ (1 mg) were solubilized in 10 mL of DMSO each (Stock #1 of either PGF_{2α} or PGE₂, 1000 μg/mL). Then, 30 μL aliquots from each prostaglandin solution were stored at -20 C° until needed. For each replicate of Experiments 1 and 2, serial 10-fold dilutions of PGF_{2α} stock #1 in KSOM-PVA were made to yield desired concentrations of PGF_{2α}. Likewise, stock #1 of PGE₂ was diluted to a final concentration of 5 ng/mL in 3 replicates of Experiment 1.

Experimental Methods for Experiment 1

Oocyte collection

The procedures utilized for *in vitro* production of embryos (IVP) were modifications of procedures previously described by Edwards and Hansen (1996). Ovaries obtained from an abattoir were packaged in thermoses and contained within a cooler during air transport to the laboratory. Upon arrival to the laboratory, ovaries were immediately washed with warm water equilibrated to arrival temperature of ovaries (generally between 28 to 30 C°). Extraneous tissue surrounding the ovaries was removed and ovaries were washed with water an additional time.

For oocyte recovery, ovaries were held firmly by clamping the base of the ovary with a hemostat and checkerboard incisions were made across follicles (approximately 3-8 mm in diameter) using a scalpel blade. Ovaries were vigorously washed in oocyte collection medium (OCM) in order to remove cumulus oocyte complexes (COCs) contained within the follicles. Collection medium was filtered and rinsed using an Emcon Filter unit (Vet Concepts, Spring Valley, WI) until medium was clean. Medium containing COCs was poured into a gridded culture dish where searching of oocytes was accomplished. Cumulus oocyte complexes were transferred to an "X" plate with OCM and washed four times to eliminate cellular debris. Cumulus oocyte complexes of good quality were washed in oocyte maturation medium (OMM) and placed in groups of approximately 50 COCs per well in a 4-well plate containing 500 µL OMM. Maturation of oocytes was performed in an incubator at 5.5% CO₂ in air at 38.5° C until time of fertilization (approximately 22 h after placement in OMM). Maturation medium was

equilibrated in the incubator (5.5% CO₂ in air at 38.5°C) for a minimum of 5 h before oocyte collection.

Sperm preparation and in vitro fertilization

At 22-24 h following oocyte collection, maturation medium from each well was carefully removed and 25 µl of penicillamine/hypotaurine/epinephrine (PHE) and 500 µL of fertilization medium (IVF-Talp) were added per well. Two straws of semen from two bulls (140 Se5 and 140 Se55), known to have high motility as well as producing high cleavage and blastocyst percentages were used to fertilize oocytes for every replicate of the study. Briefly, semen straws were removed from liquid nitrogen tank and placed in water at 36.7°C for 45 sec. Each straw was then emptied on top of a discontinuous Percoll density gradient (2 mL 45% Percoll over 2 mL 90% Percoll contained in a 15 mL conical tube) and sperm was centrifuged at 2200 rpm for 15 min in order to remove excess extender, debris, and nonmotile sperm prior to fertilization. Sperm pellet present at the bottom of the 90% fraction was collected and transferred to 10 mL Sperm-Talp and centrifuged at 1100 rpm for 8 min. Supernatant was removed and sperm pellet re-suspended in 500 µL of modified IVF-Talp. Sperm concentration was determined using a hemocytometer and sperm was added to each well at a final concentration of 375,000 sperm per well. Lastly, oocytes and sperm were incubated at 5.5% CO₂ in air at 38.5°C for 18-22 h.

In vitro culture

Approximately 18-22 h post fertilization, putative zygotes (PZ) were denuded of cumulus cells by vortexing. Putative zygotes were transferred to a 15 mL conical tube containing 500 μ L of Hepes-Talp and vortexed for 4 min. Recovered putative zygotes were washed 4 times in Hepes-Talp and once in KSOM-BSA before transferring groups of approximately 50 zygotes to 4-well plates containing 500 μ L of KSOM-BSA per well. Zygotes were placed in a humidified atmosphere of 5.5% CO₂, 7% O₂, and 87.5% N₂ at 38.5°C for 5 days.

On day 5 of embryonic development, 16- to 32-cell stage embryos were rapidly washed in KSOM-PVA four times, evaluated for quality (normal shape, defined blastomeres, extruded blastomeres, cytoplasmic fragmentation, even cytoplasm), evenly sorted in four groups (formed by embryos of good quality), transferred to experimental treatments in a 4-well plate, and placed in the incubator (5.5% CO₂, 7% O₂, and 87.5% N₂ at 38.5°C). Embryos remained in each treatment for 4 additional days (Figure 3-1). Embryo development was assessed using IETS guidelines for classification of bovine embryos.

Utilizing past studies on concentrations of PGF_{2 α} in uterine flush media, dilutions of PGF_{2 α} were prepared to obtain 0, 1, 10 and 100 ng/mL (Figure 3-1). Treatments were placed in each well of a 4-well plate (500 μ L/well) and equilibrated in the incubator (5.5% CO₂, 7% O₂, and 87.5% N₂ at 38.5°C) for at least 5 h prior to embryo placement.

For each replicate, two media samples from each treatment were collected to determine concentrations of PGF_{2 α} by radioimmunoassay. The first sample was collected




Figure 3-1. Schematic illustration of experimental design for Experiment 1. Treatments were control, 1 ng/mL (PGF-1), 10 ng/mL (PGF-10), and 100 ng/mL of prostaglandin $F_{2\alpha}$ (PGF-100), and 5 ng/mL of prostaglandin E_2 (PGE-5).

on the day in which treatments began (day 5 of culture) and the second sample obtained after determination of blastocyst development (day 9 of culture).

A total of 7 replicates were utilized for Experiment 1. In each replicate, embryos at 16- to 32-cell stage receive one of the following culture treatments: 1) control (KSOM-PVA; n= 168); 2) PGF-1 (1 ng/mL PGF_{2α} in KSOM-PVA; n= 143); 3) PGF-10 (10 ng/mL PGF_{2α} in KSOM-PVA; n= 168); 4) PGF-100 (100 ng/mL PGF_{2α} in KSOM-PVA; n= 136), and 5) PGE-5 (5 ng/mL PGE₂ in KSOM-PVA; n= 62; Figure 3-1). Prostaglandin E₂ is thought to be a “beneficial” prostaglandin during embryo development (Biggers et al., 1978). Therefore, embryos cultured in media with addition of 5 ng/mL of PGE₂ were considered to be a positive control group (Gurevich et al., 1993).

Experimental Methods for Experiment 2


In vivo-derived glycerol-frozen bovine embryos were thawed utilizing the three-step glycerol removal procedure. Briefly, after removal from liquid nitrogen, straws containing one or more embryos were held in air for 10 sec and then placed in a water bath at 25°C for 45 sec. Straws were then emptied into an empty well of a 6-well plate. Embryos were subsequently held and washed 5 min in each of the following solutions (present in the same 6-well plate): 1) 1 mL of a solution made of 2 mL of 1M sucrose plus 4 mL 10% glycerol, 2) 1 mL of a solution made of 2 mL of 1M sucrose plus 2 mL 10% glycerol plus 2 mL PBS (with 0.4% BSA), 3) 1 mL of a solution made of 2 mL of 1M sucrose plus 4 mL PBS (with 0.4% BSA), and 4) 1 mL of PBS. After washing in the last solution, embryos were allowed to regain normal morphology for 30 min in Vigro

holding medium and sorted by stage of development and quality according with the IETS guidelines for classification of bovine embryos. For this study, a total of 4 replicates were used.


Schrick and coworkers (1993) reported 0.64 ng/mL of PGF_{2α} in flush media from short cycle cows. Based on these findings, concentrations of 0 (control, KSOM-PVA), 0.1, 1 and 10 ng/mL of PGF_{2α} in culture medium were selected as treatments. Each treatment was placed in a different well of a 4-well plate (500 μL/well) and then maintained in the incubator (5.5% CO₂, 7% O₂, and 87.5% N₂ at 38.5°C) for at least 5 h before placement of embryos to allow equilibration of medium. At the same time, media samples from each treatment were collected and stored at -20° C for determination of PGF_{2α} concentrations by radioimmunoassay.

Morula embryos (corresponding to day 6-7 of embryo development; Figure 2-2) of quality grade 1, 2 or 3 were rapidly washed four times in KSOM-PVA and randomly assigned to one of four treatments: 1) control (KSOM with PVA, n= 110); 2) PGF-0.1 (0.1 ng/mL PGF_{2α} in KSOM-PVA, n=108); 3) PGF-1 (1 ng/mL PGF_{2α} in KSOM-PVA, n=109), 4) PGF-10 (10 ng/mL PGF_{2α} in KSOM-PVA, n=109; Figure 3-2) and placed in the incubator (5.5% CO₂, 7% O₂, and 87.5% N₂ at 38.5°C).

After culturing embryos for 24 h in their respective treatment, embryos were washed and placed in KSOM-BSA without PGF_{2α} for an additional 48 h (Figure 3-2). Evaluation of embryo development using IETS guidelines for classification of bovine embryos was performed by experienced technicians uninformed of treatments.



Compacted Morula



36

Figure 3-2. Schematic illustration of experimental design for Experiment 2. Treatments were control, 0.1 ng/mL (PGF-0.1), 1 ng/mL (PGF-1), and 10 ng/mL of prostaglandin $F_{2\alpha}$ (PGF-10).

Radioimmunoassay for PGF_{2α} concentrations in culture media

Concentrations of prostaglandin F_{2α} in culture media samples were determined by the procedure of Cooper et al. (1991) as modified by Schrick et al. (1993) for flash samples. Inter- and intra-assay coefficients of variation for the PGF_{2α} assay were 14% and 24%, respectively. For Experiment 1, concentrations of PGF_{2α} in culture media at the beginning (day 5) and at the end of culture period (day 9) yielded: 0.3 and 0.4 ng/mL for control, 1.1 and 1.3 ng/mL for PGF-1, 9.6 and 11.0 ng/mL for PGF-10, and 80.9 and 103.5 ng/mL for PGF-100, respectively (Figure 3-3). Prostaglandin E₂ concentrations at the beginning of the treatments (0.27 ng/mL) were analyzed using the same RIA protocol than PGF_{2α} to verify absence of the latter during the culture period. For Experiment 2, concentrations of PGF_{2α} were 0.4, 0.5, 1.3, and 9.9 ng/mL for control, PGF-0.1, PGF-1, and PGF-10, respectively (Figure 3-4).

Guide for classification of bovine embryos

The International Embryo Transfer Society (IETS) has published a guide for classification of bovine embryos according to stage of development and quality (Manual of the International Embryo Transfer Society. 1998. Third Edition). As mentioned, these guidelines for development were used throughout both experiments.




Figure 3-3. Concentrations of PGF_{2α} in the culture medium at the beginning (day 5) and at the end (day 9) of experimental period for Experiment 1. Treatments were control, 5 ng/mL of prostaglandin E₂ (PGE-5), 1 ng/mL of prostaglandin F_{2α} (PGF-1), 10 ng/mL of prostaglandin F_{2α} (PGF-10), and 100 ng/mL of prostaglandin F_{2α} (PGF-100).




Figure 3-4. Concentrations of PGF_{2α} in the culture medium for Experiment 2. Treatments were control, 0.1 ng/mL (PGF-1), 10 ng/mL (PGF-10), and 100 ng/mL of prostaglandin F_{2α} (PGF-100).

No.	Stage
1	Unfertilized
2	2- to 12-cell
3	Early Morula
4	Morula
5	Early Blastocyst
6	Blastocyst
7	Expanded Blastocyst
8	Hatched Blastocyst
9	Expanded Hatched Blastocyst

Statistical analysis

Data for Experiments 1 and 2 were analyzed to test for blastocyst development and hatching rates using an incomplete block design and a randomized block design, respectively. Analysis of variance was performed using mixed models (SAS 8.02, SAS Institute Inc., Cary, NC) and contrasts were tested to identify differences across levels of prostaglandin $F_{2\alpha}$ on embryonic development. Analyses were then validated with Chi-square analysis (SAS 8.02, SAS Institute Inc., Cary, NC).

CHAPTER 4

RESULTS AND DISCUSSION

The aim of this thesis was to investigate effects of prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) on development of *in vitro*-produced and *in vivo*-derived bovine embryos. Utilization of these types of bovine embryos provided a model to ascertain effects of $PGF_{2\alpha}$ in culture on pre-compacted (*in vitro*-produced) and compacted (*in vivo*-derived) embryos.

Results from Experiment 1 indicated that culture of 16- to 32-cell pre-compacted (*in vitro*-produced) embryos with 1, 10 or 100 ng/mL of $PGF_{2\alpha}$ in the culture medium reduced blastocyst development when compared to control (30.4 ± 5.3 , 41.4 ± 5.1 , 33.3 ± 5.3 , and 51.8 ± 5.1 respectively, $P=0.002$, Figure 4-1). In addition, culture of embryos in media containing 1 ng/mL of $PGF_{2\alpha}$ had a more detrimental effect than addition of 10 ng/mL of $PGF_{2\alpha}$. However, hatching rates between treatments did not differ and neither blastocyst development nor hatching rates differed between control and PGE_2 treatments ($P>0.10$). Thus, addition of $PGF_{2\alpha}$ to culture medium had a direct negative effect on development of 16- to 32-cell *in vitro*-produced embryos to blastocyst without altering hatching rates.

These results agree with Breuel et al. (1993a) and Buuck (1997) in which addition of $PGF_{2\alpha}$ during *in vitro* culture decreased development of 8-cell rat embryos to blastocyst. Furthermore, administration of $PGF_{2\alpha}$ to progestogen-supplemented cows on days 5 through 8 after mating decreased pregnancy rates (Seals et al., 1998) and embryo quality and development in cattle (Hockett et al., 1998). Lemaster et al. (1999) reported that administration of oxytocin on days 5 to 8 after mating to progestogen-supplemented




Figure 4-1. Percentage of pre-compacted embryos that continued development to blastocyst in Experiment 1. Treatments were control, 1 ng/mL (PGF-1), 10 ng/mL (PGF-10), and 100 ng/mL of prostaglandin $F_{2\alpha}$ (PGF-100). ^{a,b,c}Values with different superscripts differ; $P=0.002$). Pooled standard errors were ± 5.1 , 5.3, 5.1 and 5.3 for control, PGF-1, PGF-10 and PGF-100, respectively.

cows reduced pregnancy rates by stimulation of uterine release of PGF_{2α}. Therefore, reduction in development to blastocyst stage could be related to detrimental effects of PGF_{2α} on embryonic development prior to and/or during the compaction process.

Compaction, which in bovine embryos occurs at the 16- to 32-cell stage, is an essential step of embryonic development in which formation of cell-to-cell adhesion molecules allows blastomeres to differentiate into two distinctive cell populations (the trophoctoderm and the inner cell mass) and cavitation to occur (Watson et al., 1992). Utilization of 16 to 32-cell embryos in Experiment 1 served as a model to investigate PGF_{2α} action(s) prior to or during compaction. Formation of gap junctions and cadherins during compaction is considered a key event for continued development and blastocyst morphogenesis (Fleming et al., 2001). In fact, injection of gap junction antisense RNA decreased compaction and blastocyst formation in the mouse embryo (Bevilacqua et al., 1989); whereas, mouse embryos lacking E-cadherin failed to generate a normal trophoctoderm and died before implantation (Laure et al., 1994; Riethmacher et al., 1995).

The most understood mechanism by which PGF_{2α} exerts its biological activity is through receptor-mediated events. However, since detrimental effects of PGF_{2α} on embryonic development did not show a “dose response” change in development to blastocyst, actions of PGF_{2α} on bovine embryos may not involve receptor-mediated events. Prostaglandin F_{2α} binds to specific G-protein coupled receptors on targeted cells (Sakamoto et al., 1994). Using the corpus luteum as an example, the ligand-receptor binding activates phospholipase C that produces the hydrolysis of phosphatidylinositol

bisphosphate (PIP₂) into inositol triphosphate (IP₃) and diacylglycerol (DAG). Then, IP₃ induces the release of Ca⁺⁺ from the endoplasmic reticulum increasing intracellular concentrations of calcium, while DAG increases the affinity of PKC for Ca⁺⁺ (Davis et al., 1987). The final outcome is a dramatic increase in intracellular levels of Ca⁺⁺ and activation of PKC. Protein kinase C can then phosphorylate and regulate other proteins. Protein kinase C-dependent excessive phosphorylation of connexin 43 inhibited gap junction communication and impaired ventricular conduction in the heart (Riley and Behrman, 1991; Carlson et al., 1993). Increased intracellular levels of calcium by PGF_{2α} induced apoptosis in luteal cells during corpus luteum regression (Juengel et al., 1993). Calcium stimulates Ca⁺⁺-dependent endonucleases to cut DNA into internucleosomal fragments, characteristic of the apoptotic process (Rueda et al., 1995). Furthermore, Criswell (1995) also demonstrated that high intracellular levels of Ca⁺⁺ and addition of Ca⁺⁺ ionophore 4-br-A23187 decreased gap junction communication. Thus, PGF_{2α} may exert negative effects on pre-compacted embryos by altering the formation and/or function of gap junctions or by inducing apoptosis through activation of endonucleases. Decreased intercellular communication and stimulation of apoptosis in early embryos will result in impaired cellular differentiation and growth, ultimately resulting in embryonic death.

Prostaglandins are also assumed to traverse membranes by simple diffusion. However, prostaglandins are charged anions at physiological pH and may transverse biological membranes inefficiently (Bito, 1975). This limited simple diffusion appears to be augmented by a carrier mediated prostaglandin transport (PGT; (Shuster, 1998). Prostaglandin transporter is a member of the 12-membrane-spanning superfamily of

transporters and has been cloned in rats and humans (Kanai et al., 1995; Lu et al., 1996). Prostaglandin transporters from both species bind $\text{PGF}_{2\alpha}$, PGE_2 , and PGD_2 with high affinity and might mediate transepithelial transport of prostaglandins to cells (Lu et al., 1996) and the efflux of newly synthesized prostaglandins from cells (Chan et al., 1998). Clearance of prostaglandins may also be associated with PGT (Kanai et al., 1995). The gene encoding PGT in humans is regulated by fluid mechanical stimuli in cultured endothelial cells (Topper et al., 1998), suggesting that this is an inducible gene potentially capable of responding to physiological stimuli *in vivo*. In the embryo, stretching of the trophoctoderm during cavitation could induce PGT especially for binding of PGE_2 . Prostaglandin E_2 has been shown to be involved in fluid accumulation during blastocyst formation and hatching of the embryo from the zona pellucida (Biggers et al., 1978). It may be possible that $\text{PGF}_{2\alpha}$ could also bind and be internalized into individual blastomeres through PGT that normally would bind PGE_2 .

Prostaglandin $\text{F}_{2\alpha}$ might also enter the cells by simple diffusion or by binding to PGT and activating peroxisome proliferator-activated receptors (PPAR) in the nucleus. Peroxisome proliferator-activated receptors are a superfamily of nuclear hormone receptors that function as a ligand-dependent transcription factors (Schoonjans et al., 1996). The transcriptional activity of PPAR is enhanced by numerous chemical compounds including fatty acids and prostaglandins (Schoonjans et al., 1996). After activation, PPARs could then regulate transcription of developmentally important genes involved in embryonic development (connexin 43, E-cadherin, $\text{Na}^+ - \text{K}^+ - \text{ATPase}$ pump) or cause cytotoxicity via apoptosis. Padilla et al. (2000) reported that PPAR γ agonist and PGI_2 produced cytotoxicity and antiproliferative effects on normal and malignant B cells.

Another action of $\text{PGF}_{2\alpha}$ to decrease embryonic development could be the generation of free oxygen radicals. Exposure of rat luteal cells to oxygen radicals induced apoptosis (Riley and Behrman, 1991; Carlson et al., 1993) and loss of fluidity in cellular membranes (Sawada and Carlson, 1991).

In Experiment 2, development to blastocyst of compacted (*in vivo*-derived) morula stage embryos (at least two stages beyond morula stage according to IETS guidelines) did not differ between treatments (68.1% for control, 58.9% for PGF -0.1, 61% for PGF -1 and 60% for PGF -10, $P>0.10$, Figure 4-2). However, culture of embryos in 0.1, 1, or 10 ng/mL of $\text{PGF}_{2\alpha}$ for 24 h decreased hatching rates when compared to control (24%, 29.1%, 24.5%, and 44.5% respectively, $P=0.05$, Figure 4-3).

Thus, results from Experiment 2 indicated that continued development of compacted (*in vivo*-derived) morula embryos to the blastocyst stage was not altered by addition of $\text{PGF}_{2\alpha}$ to culture medium. However, $\text{PGF}_{2\alpha}$ significantly reduced hatching rates of *in vivo*-derived morula. This is in agreement with Maurer and Beier (1976) who reported that addition of $\text{PGF}_{2\alpha}$ to culture medium directly decreased the ability of 8-cell rabbit embryos to form expanded and hatched blastocyst, but had no effect on development to early blastocyst. These findings are also consistent with *in vivo* studies in which progestogen-supplemented cows administered $\text{PGF}_{2\alpha}$ (days 5 through 8 after estrous) had decreased pregnancy rates following transfer of morula stage embryos (Hernandez-Fonseca et al., 1997). Whereas, transfer of blastocyst stage embryos to progestogen-supplemented cows receiving $\text{PGF}_{2\alpha}$ had pregnancy rates similar to controls (E. Keith Inskeep, personal communication). Moreover, Schrick et al. (2001) reported that administration of flunixin meglumine (a COX-1 and COX-2 inhibitor) at the time of




Figure 4-2. Percentage of compacted morula embryos that continued development to blastocyst in Experiment 2. Treatments were control, 0.1 ng/mL (PGF-0.1), 1 ng/mL (PGF-1), and 10 ng/mL of prostaglandin $F_{2\alpha}$ (PGF-10). Pooled standard errors were ± 5.4 .




Figure 4-3. Percentage of compacted morula embryos that continued development to blastocyst and hatched from the zona pellucida in Experiment 2. Treatments were control, 0.1 ng/mL (PGF-0.1), 1 ng/mL (PGF-1), and 10 ng/mL of prostaglandin $F_{2\alpha}$ (PGF-10). ^{a,b}Values differ between treatments; $P=0.05$. Pooled standard errors were ± 6.0 .

embryo transfer improved pregnancy rates of morula and blastocyst stage embryos in cattle. Thus, results from past and current studies indicate a negative effect of PGF_{2α} on embryonic development after compaction in cattle.

Although the exact mechanisms by which PGF_{2α} is affecting embryonic development at the morula stage are unknown and evidence of PGF_{2α} receptors in preimplantation bovine embryos have not been located in the literature, some modes of action of PGF_{2α} on embryo development can be hypothesized. The reduction in hatching rates of *in vivo*-derived embryos may have resulted from an interruption of Na⁺ transport across the trophoctoderm epithelium by PGF_{2α}. Kim and Yeoun (1983) observed a reduction of Na⁺-K⁺-ATPase activity after treating luteal membranes with PGF_{2α} (a receptor mediated event). After compaction, accumulation of blastocoelic fluid within the embryo causes expansion and hatching from the zona pellucida (Latham et al., 1999). Localization of an active Na⁺-K⁺-ATPase at the basolateral area of trophoctoderm cell assures movement of Na⁺ against a concentration gradient from the outside to the interior of the embryo (Watson and Kidder, 1988); whereas, tight junction formation between trophoctoderm cells regulates paracellular transport of Cl⁻ and controls leakage of the nascent blastocoelic fluid (Fleming et al., 1989).

Possible explanation of deleterious effects of PGF_{2α} on hatching rates would involve disruption and/or dysfunction of tight junction or adherent junctions (E-cadherin) between trophoctoderm cells. Tsukita et al. (1991) observed an increase in paracellular permeability of tight junctions due to elevated intracellular levels of Ca⁺⁺. Furthermore, Morgado-Diaz and de Souza (2000) reported that tyrosine phosphorylation causes disassembly of adherent junctions in cell culture. Both, high intracellular levels of Ca⁺⁺

and protein phosphorylation are common features associated with the action of PGF_{2α} on target cells.

In conclusion, these studies suggest a direct negative effect of PGF_{2α} on embryonic development of pre-compacted (*in vitro*-produced) and compacted (*in vivo*-derived) bovine embryos. Prostaglandin F_{2α} reduced the ability of 16- to 32-cell pre-compacted embryos to develop to blastocyst and may be related to deleterious actions of PGF_{2α} during or before the time of compaction (during the transition from 32-cell embryo to compacted morula). Furthermore, PGF_{2α} decreased hatching rates of compacted morula embryos possibly through disruption/dysfunction of tight junctions or by alterations in active transport of Na⁺ during blastocoel formation. It was interesting to note that different concentrations of PGF_{2α} did not show a “dose response” effect on development of embryos and hatching rates. Thus, detrimental effects of PGF_{2α} on continued embryonic development does not appear to be a receptor-mediated event in either pre-compacted or compacted bovine embryos. Possible mechanisms of action may involve alterations in gap junction and cell adhesion molecules functionality and/or formation, induction of apoptotic mechanisms, or alterations in gene transcription during embryonic development. Future research should be aimed at determining exact mechanisms by which PGF_{2α} alters embryonic development.

- Adida, C., P. L. Crotty, J. McGrath, D. Berrebi, J. Diebold and D. C. Altieri. 1998. Developmentally regulated expression of the novel cancer anti-apoptosis gene survivin in human and mouse differentiation. *Am. J. Pathol.* 152: 43-49.
- Aoki, F., D. M. Worrall and R. M. Shultz. 1997. Regulation of transcriptional activity during the first and second cell cycles in the preimplantation mouse embryo. *Dev. Biol.* 181: 296-307.
- Arends, M. J. and A. H. Willye. 1991. Apoptosis: mechanisms and roles in pathology. *Int. Rev. Exp. Pathol.* 32: 223-254.
- Ayalon, N. 1978. A review of embryonic mortality in cattle. *J. Reprod. Fertil.* 54: 483-493.
- Barcroft, L. C., A. Hay-Schmidt, A. Caveney, E. Gilfoyle, E. W. Overstrom, P. Hyttel and A. J. Watson. 1998. Trophectoderm differentiation in the bovine embryo: characterization of a polarized epithelium. *J. Reprod. Fertil.* 114: 327-339.
- Barnes, F. L. and N. L. First. 1991. Embryonic transcription in in vitro cultured bovine embryos. *Mol. Reprod. Dev.* 29: 117-123.
- Bellows, R. A., R. E. Short and R. B. Staigmiller. 1976. Research areas in beef cattle reproduction. In: H.W. Hawk, A. Osum (Ed), *Animal reproduction: BARC Symposium 3*. Montclair; 1979: 3-18.
- Berridge, M. J. 1987. Inositol triphosphate and diacylglycerol: two interacting second messengers. *Annu. Rev. Biochem.* 56: 159-193.
- Betteridge, K. J., M. D. Eaglesome, G. C. B. Randall, D. Mitchell. 1980. Collection, description and transfer of embryos from cattle 10-16 days after oestrus. *J. Reprod. Fertil.* 59: 205:216.
- Bevilacqua, A., R. Loch-Caruso and R. P. Erickson. 1989. Abnormal development and dye coupling produced by antisense RNA to gap junction protein in mouse preimplantation embryos. *Proc. Natl. Acad. Sci.* 86: 5444-5448.
- Biggers, J. D., J. E. Bell and D. J. Benos. 1988. Mammalian blastocyst: transport functions in a developing epithelium. *Am. J. Physiol.* 255: C419-C432.
- Biggers, J. D., B. V. Leonov, J. F. Baskar and J. Fried. 1978. Inhibition of hatching of mouse blastocysts in vitro by prostaglandin antagonist. *Biol. Reprod.* 19: 519-533.

- Biggers, J. D., L. K. McGinnis and M. Raffin. 2000. Amino acid and preimplantation development of the mouse in protein-free potassium simplex optimized medium. *Biol. Reprod.* 63: 281-293.
- Bito, L. Z. 1975. Are prostaglandins intracellular, transcellular or extracellular autolysis? *Prostaglandins* 45: 229-235.
- Blatteis, C. M. and E. Sehic. 1997. Fever: How many circulating pyrogens signal the brain? *News Physiol. Sci.* 12: 1-9.
- Boni, R., E. Tosti, S. Roviello and B. Dale. 1999. Intercellular communication in in vivo- and in vitro-produced Bovine Embryos. *Biol. Reprod.* 61: 1050-1055.
- Braude, P., V. Bolton and S. Moore. 1988. Human gene expression first occurs between the four- and eight-cell stages of preimplantation development. *Nature* 332: 459-461.
- Breuel, K. F., A. Fukuda and F. N. Schrick. 1993a. Effect of prostaglandin $F_{2\alpha}$ on development of 8-cell rat embryos in vitro. *Biol. Reprod.* 48: (Supplement 1):173.
- Breuel, K. F., P. E. Lewis, F. N. Schrick, A. W. Lishman, E. K. Inskeep and R. L. Butcher. 1993b. Factors affecting fertility in the postpartum cow: role of the oocyte and follicle in conception rate. *Biol. Reprod.* 48: 655-661.
- Bruzzone, R. and D. A. Goodenough. 1996. The cellular internet: On line with connexins. *Bioessays* 18: 709-718.
- Buford, W. I., N. Ahmad, F. N. Schrick, R. L. Butcher, P. E. Lewis and E. K. Inskeep. 1996. Embryotoxicity of a regressing corpus luteum in beef cows supplemented with progesterone. *Biol. Reprod.* 54: 531-537.
- Butcher, R. L., J. E. Reber, A. W. Lishman, K. F. Breuel, F. N. Schrick, J. C. Spitzer and E. K. Inskeep. 1992. Maintenance of pregnancy in postpartum beef cows that have short-lived corpora lutea. *J. Anim. Sci.* 70: 3831-3837.
- Butler, W. R. 1998. Review: effect of protein nutrition on ovarian and uterine physiology in dairy cattle. *J. Dairy Sci.* 81: 2533-2539.
- Buuck, M. J. 1997. Effects of prostaglandin $F_{2\alpha}$ on in vitro and in vivo development of preimplantation rat embryos. Thesis. University of Tennessee, Knoxville.
- Camous S., V. Kopecky, J. E. Flechon. 1986. Autoradiographic detection of the earliest stage of ^3H -uridine incorporation in the cow embryo. *Biol. Cell* 58: 195-200.

- Carlson, J. C., X. M. Wu and M. Sawada. 1993. Oxygen radicals and the control of ovarian corpus luteum function. *Free Radic. Biol. Med.* 14: 78-84.
- Chan, B. S., J. A. Satriano, M. Pucci and V. L. Schuster. 1998. Mechanism of prostaglandin E₂ transport across the plasma membrane of HeLa cells and *Xenopus* oocytes expressing the prostaglandin transporter "PGT". *J. Biol. Chem.* 273: 6689-6697.
- Chen, D. B., S. D. Westfall, H. W. Fong, M. S. Roberson and J. S. Davis. 1998. Prostaglandin F_{2α} stimulates the Raf/Mek1/mitogen-activated protein kinase (MAPK) signaling cascade in bovine luteal cells. *Endocrinology* 139: 3876-3885.
- Citi, S. 1993. The molecular organization of tight junctions. *J. Cell Biol.* 122: 485-489.
- Cooper, D. A., D. A. Carver, P. Villeneuve, W. J. Silvia and E. K. Inskeep. 1991. Effects of progesterone treatment on concentrations of prostaglandins and oxytocin in plasma from the posterior vena cava of post-partum beef cows. *J. Reprod. Fertil.* 91: 411-421.
- Criswell, K. A., R. Loch-Caruso and E. L. Stuenkel. 1995. Lindane inhibition of gap junctional communication in myometrial myocytes is partially dependent on phosphoinositide-generated second messengers. *Toxicol. Appl. Pharmacol.* 130: 280-293.
- Crosby, I. M., F. Gandolfi and R. M. Moor. 1988. Control of protein synthesis during early cleavage of sheep embryos. *J. Reprod. Fertil.* 82: 769-775.
- Davis, A. J., I. R. Fleet, P. A. Hansford, F. A. Harrison and F. M. M. Walker. 1984. Pulmonary metabolism of prostaglandin F_{2α} in the conscious non-pregnant cow. *Proc. Physiol. Soc., J. Physiol.* 358:107.
- Davis, J. S., L. L. Weakland, R. V. Farese and L. A. West. 1987. Prostaglandin F_{2α} stimulates phosphatidylinositol 4,5-bisphosphate hydrolysis and mobilizes intracellular Ca⁺⁺ in bovine luteal cells. *Proc. Natl. Acad. Sci.* 84: 3728-2732.
- De Sousa, P. A., S. C. Juneja, S. Caveney, F. D. Houghton, T. C. Davies, A. G. Reaume, J. Rossant and G. M. Kidder. 1997. Normal development of preimplantation mouse embryos deficient in gap junctional coupling. *J. Cell Sci.* 110: 1751-1758.
- De Sousa, P. A., G. Valdimarsson, B. J. Nicholson and G. M. Kidder. 1993. Connexin trafficking and the control of gap junctions assembly in mouse preimplantation embryos. *Development* 117: 1355-1367.

- Ducibella, T. and E. Anderson. 1975. Cell shape and membrane changes in the eight-cell mouse cell embryo: prerequisites for morphogenesis of the blastocyst. *Dev. Biol.* 47: 45-58.
- Dunne, L. D., M. G. Diskin and J. M. Sreenan. 2000. Embryo and foetal loss in beef heifers between day 14 of gestation and full term. *Anim. Reprod. Sci.* 58: 39-44.
- Edwards, M. J. 1998. Apoptosis, the heat shock response, hyperthermia, birth defects, disease and cancer. Where are the common links? *Cell Stress Caperones* 3: 213-220.
- Edwards, J. L., and P. J. Hansen. 1996. Elevated temperature increases heat shock protein 70 synthesis in bovine two-cell embryos and compromises function of maturing oocytes. *Biol. Reprod.* 55: 340-346.
- Edwards, R. G., J. M. Purdy, P. C. Steptoe and D. E. Walters. 1981. The growth of human pre-implantation embryos in vitro. *Am. J. Obstet. Gynecol.* 141: 408-416.
- Elli, M., B. Gaffuri, A. Frigerio, M. Zanardelli, D. Covini, M. Candiani and M. Vignali. 2001. Effect of a single dose of ibuprofen lysinate before embryo transfer on pregnancy rates in cows. *Reproduction* 121: 151-154.
- Ellington, J. E. 1991. The bovine oviduct and its role in reproduction: a review of the literature. *Cornell Vet.* 81: 313-328.
- Espey, L. L. 1994. Current status of the hypothesis that the mammalian ovulation is comparable to an inflammatory reaction. *Biol. Reprod.* 50: 233-238.
- Fleming, T. P. and M. J. Hay. 1991. Tissue specific control of expression of the tight junction polypeptide ZO-1 in the mouse early embryo. *Development* 113: 295-304.
- Fleming, T. P. and M. H. Johnson. 1988. From egg to epithelium. *Annu. Rev. Cell Biol.* 4: 459-485.
- Fleming, T. P., J. McConnell, M. H. Johnson and B. R. Stevenson. 1989. Development of tight junctions de novo in the mouse early embryo: control of assembly of the tight junction-specific protein, ZO-1. *J. Cell Biol.* 108: 1407-1418.
- Fleming, T. P., B. Sheth and I. Fesenko. 2001. Cell adhesion in the preimplantation mammalian embryo and its role in trophoctoderm differentiation and blastocyst morphogenesis. *Front. Biosci.* 6: 1000-1007.

- Flint, A. P. F., W. M. F. Leat, E. L. Sheldrick and H. J. Stewart. 1986. Stimulation of phosphoinositide hydrolysis by oxytocin and the mechanism by which oxytocin controls prostaglandin synthesis in the ovine endometrium. *Biochem. J.* 237: 797-805.
- Flint, A. P. F. and E. L. Sheldrick. 1982. Ovarian secretion of oxytocin is stimulated by prostaglandins. *Nature* 297: 587-588.
- Fortune, J. E. 1993. Follicular dynamics during the bovine estrous cycle: a limiting factor in improvement of fertility? *Anim. Reprod. Sci.* 33: 111-125.
- Gabriel, H. D., D. Jung, A. Temme, O. Traub, E. Winterhager and K. Willecke. 1998. Transplacental uptake of glucose is decreased in embryonic lethal connexin 26 deficient mice. *J. Cell Biol.* 140: 1453-1461.
- Garverick, H. A., M. F. Smith, R. G. Elmore, G. L. Morehouse, S. P. Agudo and W. L. Zahler. 1985. Changes and interrelationships among luteal LH receptors, adenylate cyclase activity and phosphodiesterase activity during the bovine estrous cycle. *J. Anim. Sci.* 61: 216-223.
- Gavrieli, Y., Y. Sherman and S. A. Ben-Sasson. 1992. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J. Cell Biol.* 119: 493-501.
- Ginther, O. J., L. Knopf and J. P. Kastelic. 1989. Ovarian follicular dynamics in heifers during early pregnancy. *Biol. Reprod.* 41.
- Gonzalez, D. S., B. D. Bavister. 1995. Zona pellucida escape by hamster blastocysts in vitro is delayed and morphologically different compared with zona escape in vivo. *Biol. Reprod.* 52: 470-480.
- Granville, D. J., C. M. Carthy, D. W. C. Hunt and B. M. McManus. 1998. Apoptosis: molecular aspects of cell death and disease. *Lab. Invest.* 78: 893-913.
- Gurevich, M., E. Harel-Markowitz, S. Marcus, L. S. Shore and M. Shemesh. 1993. Prostaglandin production by the oocyte cumulus complex around the time of fertilization and the effect of prostaglandin E on the development of the early bovine embryo. *Reprod. Fertil. Dev.* 5: 281-283.
- Hansel, W. and J. P. Dowd. 1986. New concepts of the control of corpus luteum function. *J. Reprod. Fertil.* 78: 77-89.
- Hardy, K. 1997. Cell death in the mammalian blastocyst. *Mol. Hum. Reprod.* 3: 919-925.
- Hardy, K. 1999. Apoptosis in the human embryo. *Rev. Reprod.* 4: 125-134.

- Hardy, K., R. M. L. Winston and D. L. Becker. 1996. Expression of intercellular junctions during preimplantation development of the human embryo. *Mol. Hum. Reprod.* 2: 621-632.
- Harper, M. J. C. and R. C. Skarnes. 1972. Inhibition of abortion and fetal death produced by endotoxin or prostaglandin F_{2α}. *Prostaglandins* 2: 295-309.
- Hernandez-Fonseca, H. J., B. L. Sayre, R. L. Butcher and E. K. Inskeep. 1997. Systemic vs local embryotoxic effect of early luteal regression in cattle. *J. Anim. Sci.* 75 (Suppl.1): 221.
- Hockett, M. E., N. R. Rohrbach and F. N. Schrick. 1998. Effect of administration of PGF_{2α} on embryonic development and quality in cows supplemented with exogenous progestogen. *J. Anim. Sci.* 76 (Suppl. 1): 241.
- Hockett, M. E., F. M. Hopkins, M. J. Lewis, A. M. Saxton, H. H. Dowlen, S. P. Oliver, and F. N. Schrick. 2000. Endocrine profiles of dairy cows following experimentally induced clinical mastitis during early lactation. *Anim. Reprod. Sci.* 58: 241-251.
- Hyunjung, L., R. A. Gupta, W. Ma, B. C. Paria, D. E. Moller, J. D. Morrow, R. N. Dubois, J. M. Trzaskos and S. K. Dey. 1999. Cyclo-oxygenase-2 derived prostacyclin mediates embryo implantation in the mouse via PPAR-γ. *Genes Dev.* 13: 1561-1574.
- Inoguchi, T., H. Y. Yu, M. Imamura, M. Kakimoto and T. Kuroki. 2001. Altered gap junction activity in cardiovascular tissues of diabetes. *Medical Electron Microscopy* 34: 86-91.
- Ivanov, D. B., M. P. Philippova and V. A. Tkachuk. 2001. Structure and functions of classical cadherins. *Biochemistry-Moscow* 66: 1174-1186.
- Javed, Q., T. P. Fleming, M. J. Hay and S. Citi. 1993. Tight junction protein cingulin is expressed by maternal and embryonic genomes during early mouse development. *Development* 117: 1145-1151.
- Juengel, J. L., H. A. Garverick, A. L. Johnson, R. S. Youngquist and M. F. Smith. 1993. Apoptosis during luteal regression in cattle. *Endocrinology* 132: 249-254.
- Juriscova, A., S. Varmuza and R. F. Casper. 1996. Programmed cell death and human embryo fragmentation. *Mol. Hum. Reprod.* 2: 93-98.
- Kanai, N., R. Lu, J. A. Satriano, Y. Bao, A. W. Wolkoff and V. L. Schuster. 1995. Identification and characterization of a prostaglandin transporter. *Science* 268: 866-869.

- Kane, M. T., B. D. Bavister. 1988. Vitamin requirements for development of eight-cell hamster embryos to hatching blastocysts in vitro. *Biol. Reprod.* 39: 1137-1143.
- Kidder, G. M. 1987. Intercellular communication during mouse embryogenesis. In: *The mammalian preimplantation embryo: regulation of growth and differentiation in vitro.* B. D. Bavister. New York, Plenum Press: 45-71.
- Kidder, G. M. and E. Winterhager. 2001. Intercellular communication in preimplantation development: the role of gap junctions. *Front. Biosci.* 6: 53-64.
- Kim, I. and D. S. Yeoun. 1983. Effect of prostaglandin $F_{2\alpha}$ on Na-K-ATPase activity in luteal membranes. *Biol. Reprod.* 29: 48-55.
- Kindahl, H., L. E. Edqvist, A. Bane and E. Granstrom. 1976. Blood levels of progesterone and 15-keto-13,14-dihydro-prostaglandin $F_{2\alpha}$ during the normal estrous cycle and early pregnancy in heifers. *Acta Endocrinol. (Copenh).* 82: 134-149.
- Knopf, L., J. P. Kastelic, E. Schallenberger and O. J. Ginther. 1989. Ovarian dynamics in heifers: test of two-wave hypothesis by ultrasonically monitoring individual follicles. *Domest. Anim. Endocrinol.* 6: 111-119.
- Koyama, H., H. Suzuki, X. Yang, S. Jiang and R. H. Foote. 1994. Analysis of polarity of bovine and rabbit embryos by scanning electron microscopy. *Biol. Reprod.* 50: 163-170.
- Lafrance, M. and A. K. Goff. 1990. Control of bovine uterine prostaglandin $F_{2\alpha}$ release in vitro. *Biol. Reprod.* 42: 288-293.
- Latham, K. E., J. I. Garrels, C. Chang and D. Solter. 1991. Quantitative analysis of protein synthesis in mouse embryos. *Development* 112: 921-932.
- Latham, K. E., K. Kutyna and Q. Wang. 1999. Genetic variation in trophectoderm function in parthenogenetic mouse embryos. *Dev. Genet.* 24: 329-335.
- Laure, L., M. Ohsugi, J. Hirchenhain and R. Kemler. 1994. E-cadherin null mutant embryos fail to form a trophectoderm epithelium. *Proc. Natl. Acad. Sci.* 91: 8263-8267.
- Lee, S., N. B. Gilula and A. E. Warner. 1987. Gap junctional communication and compaction during preimplantation stages of mouse development. *Cell* 51: 851-860.

- Lemaster, J. W., R. C. Seals, F. M. Hopkins and F. N. Schrick. 1999. Effects of administration of oxytocin on embryonic survival in progesterone supplemented cattle. *Prostaglandins* 57: 259-268.
- Long, C. R., J. R. Dobrinsky, W. M. Garret and L. A. Johnson. 1998. Dual labeling of the cytoskeleton and DNA strand breaks in porcine embryos produced in vivo and in vitro. *Mol. Reprod. Dev.* 51: 59-65.
- Lowenstein, W. R. 1981. Junctional intercellular communication: the cell-to-cell membrane channel. *Physiol. Rev.* 61: 829-913.
- Lu, R., N. Kanai, Y. Bao and V. L. Schuster. 1996. Cloning, in vitro expression, and tissue distribution of a human prostaglandin transporter cDNA (hPGT). *J. Clin. Invest.* 98: 1142-1149.
- Magli, M. C., L. Gianaroli, A. P. Ferraretti, D. Fortini, G. Aicardi, N. Montanero. 1998. Rescue of implantation potential in embryos with poor prognosis by assisted zona hatching. *Hum. Reprod.* 13: 1331-1335.
- Majno, G. and A. Joris. 1995. Apoptosis, oncosis and necrosis. *Am. J. Pathol.* 146: 3-15.
- Malayer, J. R., P. J. Hansen, T. S. Gross and W. W. Tatcher. 1990. Regulation of heat shock-induced alterations in the release of prostaglandins by the uterine endometrium of cows. *Theriogenology* 34: 219-230.
- Manes, C. 1973. The participation of the embryonic genome during early cleavage in the rabbit. *Dev. Biol.* 32: 453-459.
- Martin, S. J., C. P. M. Reutelingsperger, A. J. McGahon, J. A. Rader, R. C. A. A. van Schie, D. M. LaFace and D. R. Green. 1995. Early redistribution of plasma membrane phosphatidylserine is a general feature of apoptosis regardless of the initiating stimulus: inhibition by overexpression of Bcl-2 and Abl. *J. Exp. Med.* 182: 1545-1556.
- Massip, A., J. Mulnard. 1980. Time-lapse cinematographic analysis of hatching of normal and frozen-thawed cow blastocysts. *J. Reprod. Fertil.* 58: 475-478.
- Matwee, C., D. H. Betts and W. A. King. 2000. Apoptosis in the early bovine embryo. *Zygote* 8: 57-68.
- Maurer, P. R. and H. M. Beier. 1976. Uterine proteins and development in vitro of rabbit preimplantation embryos. *J. Reprod. Fertil.* 48: 33-41.
- Maurer, R. R. and J. R. Chenault. 1983. Fertilization failure and embryonic mortality in parous and nonparous beef cattle. *J. Anim. Sci.* 56: 1186-1189.

- McGuire, W. J., J. L. Juengel and G. D. Niswender. 1994. Protein kinase C second messenger system mediates the antisteroidogenesis effects of prostaglandin F_{2α} in the ovine corpus luteum in vivo. *Biol. Reprod.* 51: 800-806.
- Memili, E., T. Dominko and N. L. First. 1998. Onset of transcription in bovine oocytes, 2-, 4- and 8-cell embryos. *Mol. Reprod. Dev.* 51: 36-41.
- Miller, J. G. O. and G. A. Schultz. 1985. Amino acid transport in mouse blastocyst compartments. *J. Embryol. Exp. Morphol.* 89: 149-158.
- Milvae, R. A. and W. Hansel. 1980. Concurrent uterine venous and ovarian arterial prostaglandin F_{2α} concentrations in heifers treated with oxytocin. *J. Reprod. Fertil.* 60: 7-16.
- Morgado-Diaz, J. A. and W. de Souza. 2000. Evidence that increased tyrosine phosphorylation causes disassembly of adherens junctions but does not perturb paracellular permeability in Caco-2 cells. *Tissue Cell* 33: 500-513.
- Murakami, M., I. Kudo. 2002. Phospholipase A₂. *J. Biochem.* 131: 285-292
- Narumiya, S., K. Ohno, M. Fukushima and M. Fujiwara. 1987. Site and mechanism of growth inhibition by prostaglandins. III. Distribution and binding of prostaglandin A₂ and 12-prostaglandin J₂ in nuclei. *J. Pharmacol. Exp. Ther.* 242: 306.
- Nett, T. M., M. C. McClellan and G. D. Niswender. 1976. Effects of prostaglandins on the ovine corpus luteum: blood flow, secretion of progesterone and morphology. *Biol. Reprod.* 15: 66-78.
- Odensvik, K., M. Duchens and H. Gustafsson. 1993. Does mechanical manipulation of the reproductive organs cause a prostaglandin release in the heifer during embryo transfer? *Acta Vet. Scand.* 34: 219-221.
- O'Sullivan, C. M., S. Y. Liu, D. E. Rancourt. 2001. A novel murine tryptase involved in blastocyst hatching and outgrowth. *Reproduction.* 122: 61-71.
- Padilla, J., K. Kaur, H. J. Cao, T. J. Smith and R. P. Phipps. 2000. Peroxisome proliferator activator receptor-γ agonist and 15-deoxy-12,14-PGJ₂ induce apoptosis in normal and malignant B-lineage cells. *J. Immunol.* 165: 6941-6948.
- Parfenova, H., V. N. Parfenov, B. V. Shlopov, V. Levine, S. Falkos, M. Pourcyrous, C. W. Leffler. 2001. Dynamics of nuclear localization sites for COX-2 in vascular endothelial cells. *Am. J. Physiol-Cell Ph.* 281: C166-C178.
- Parrish, J. J., J. Susko-Parrish, M. A. Winer and N. L. First. 1988. Capacitation of bovine sperm by heparin. *Biol. Reprod.* 38: 1171-1180.

- Perona, R. M., P. M. Wassarman. 1986. Mouse blastocysts hatch in vitro by using a trypsin-like proteinase associated with cells of mural trophoctoderm. *Dev. Biol.* 114: 42-52.
- Petroff, B. K., R. E. Ciereszko, K. Dabrowski, A. C. Ottobre, W. F. Pope and J. S. Ottobre. 1999. Prostaglandin F_{2α} depletes the porcine corpus luteum of vitamin C by inducing secretion of the vitamin into the bloodstream. *Endocrinology* 140: 1018-1021.
- Prather, R. S. and N. L. First. 1993. Cell-to-cell coupling in early-stages bovine embryos: a preliminary report. *Theriogenology* 39: 561-567.
- Putney, D. J., S. Mullins, W. W. Tacher, M. Drost and T. S. Gross. 1989. Embryonic development in superovulated dairy cattle exposed to elevated ambient temperatures between the onset of estrus and insemination. *Anim. Reprod. Sci.* 19: 37-51.
- Rappolee, D. A., K. S. Sturm, G. A. Schultz, R. A. Pederson and Z. Werb. 1990. The expression of growth factor ligand and receptors in preimplantation mouse embryos. *Mol. Cell. Biol.* 117: 11-25.
- Reaume, A. G., P. A. De Sousa, S. Kulkarni, B. L. Langille, D. Zhu, T. C. Davies, S. C. Juneja, G. M. Kidder and J. Rossant. 1995. Cardiac malformation in neonatal mice lacking connexin 43. *Science* 267: 1831-1834.
- Reuss, B., P. Hellmann, O. Traub, A. Butterweck and E. Winterhager. 2002. Expression of connexin 31 and connexin 43 genes in early rat embryos. *Dev. Genet.* 21: 82-90.
- Riethmacher, D., V. Brinkmann and C. Birchmeier. 1995. A targeted mutation in the mouse E-cadherin gene results in defective preimplantation development. *Proc. Natl. Acad. Sci.* 92: 855-859.
- Riley, J. C. M. and H. R. Behrman. 1991. Oxygen radicals and reactive oxygen species in reproduction. *Proc. Soc. Exp. Biol. Med.* 198: 781-791.
- Rothhut, B. and F. Russo-Marie. 1984. Novel concepts in the mode of action of anti-inflammatory steroids. *Agents Actions* 14: 171-180.
- Rubinstein, M., A. Marazzi and E. de Fried. 1999. Low-dose aspirin treatment improves ovarian responsiveness, uterine and ovarian blood flow velocity, implantation and pregnancy rates in patients undergoing in vitro fertilization: a prospective, randomized, double-blind placebo-controlled assay. *Fertil. Steril.* 71: 825-829.

- Rueda, B. R., K. I. Tilly, I. W. Botros, P. D. Jolly, T. R. Hansen, P. B. Hoyer and J. L. Tilly. 1997. Increased bax and interleukin-1beta-converting enzyme messenger ribonucleic acid levels coincide with apoptosis in the bovine corpus luteum during structural regression. *Biol. Reprod.* 56: 186-193.
- Rueda, B. R., K. I. Tilly, T. R. Hansen, P. B. Hoyer and J. L. Tilly. 1995. Expression of superoxide dismutase, catalase and glutathione peroxidase in the bovine corpus luteum: evidence supporting a role for oxidative stress in luteolysis. *Endocrine* 3: 227-232.
- Sakamoto, K., T. Ezaski, K. Miwa, E. Okuda-Ashitaka, T. Hountani, T. Sugimoto, S. Ito and O. Hayaishi. 1994. Molecular cloning and expression of a cDNA of the bovine prostaglandin F_{2α} receptor. *J. Biol. Chem.* 269: 3881-3886.
- Saling, P. M. 1991. How the egg regulates sperm function during gamete interaction: facts and fantasies. *Biol. Reprod.* 44: 246-251.
- Savio, J. D., L. Keenan, M. P. Boland and J. F. Roche. 1988. Pattern of growth of dominant follicles during the oestrus cycle of heifers. *J. Reprod. Fertil.* 83: 663-671.
- Sawada, H., K. Yamazaki, M. Hoshi. 1990. Trypsin-like hatching protease from mouse embryos: evidence for the presence in culture media and its enzymatic properties. *J. Exp. Zool.* 254: 83-87.
- Sawada, M. and J. C. Carlson. 1991. Rapid plasma membrane changes in superoxide radical formation, fluidity, and phospholipase A₂. *Endocrinology* 128: 2992-2998.
- Schlafer, D. H., P. J. Fisher and C. J. Davis. 2000. The bovine placenta before and after birth: placental development and function in health and disease. *Anim. Reprod. Sci.* 60-61: 145-160.
- Schoonjans, K., B. Staels and J. Auwerx. 1996. The peroxisome proliferator activated receptors (PPARS) and their effects on lipid metabolism and adipocyte differentiation. *Biochim. Biophys. Acta*: 1302-1393.
- Schrack, F. N., M. Hockett, T. M. Towns, A. M. Saxton, N. E. Wert, M. Mimms, D. Ezcurra and M. E. Wehrman. 2001. Administration of prostaglandin inhibitor at embryo transfer improves pregnancy rates. *Theriogenology* 55 (Suppl.1): 370.
- Schrack, F. N., E. K. Inskip and R. L. Butcher. 1993. Pregnancy rates for embryos transferred from early postpartum beef cows into recipients with normal estrous cycle. *Biol. Reprod.* 49: 617-621.

- Seals, R. C., J. W. Lemaster, F. M. Hopkins and F. N. Schrick. 1998. Effects of elevated concentrations of prostaglandin $F_{2\alpha}$ on pregnancy rates in progestogen supplemented cattle. *Prostaglandins* 56: 377-389.
- Sefton, M., M. H. Johnson and L. Clayton. 1992. Synthesis and phosphorylation of uvomorulin during mouse early development. *Development* 115: 313-318.
- Seibert, K., Y. Zhang and K. Leahy. 1994. Pharmacological and biochemical demonstration of the role of cyclooxygenase 2 in inflammation and pain. *Proc. Natl. Acad. Sci.* 91: 12013-12017.
- Shuster, V. L. 1998. Molecular mechanisms of prostaglandin transport. *Annu. Rev. Physiol.* 60: 221-242.
- Silvia, W. J. and M. L. Taylor. 1989. Relationship between uterine secretion of prostaglandin $F_{2\alpha}$ induced by oxytocin and endogenous concentrations of estradiol and progesterone at three stages of the bovine estrous cycle. *J. Anim. Sci.* 67: 2347-2353.
- Simon, A. M., D. A. Goodenough and D. L. Paul. 1997. Female infertility in mice lacking connexin 37. *Nature* 385: 525-529.
- Smith, W. L. and D. L. DeWitt. 1996. Prostaglandin endoperoxide H synthase-1 and -2. *Adv. Immunol.* 62: 167-215.
- Sreenan, J. M. and M. G. Diskin. 1983. Early embryonic mortality in the cow: its relationship with progesterone concentration. *Vet. Rec.* 112: 517-521.
- Tanaka, M., T. Miyazaki, S. Tanigaki, K. Kasai, K. Minegishi, K. Miyakoshi, H. Ishimoto and Y. Yoshimura. 2000. Participation of reactive oxygen species in $PGF_{2\alpha}$ -induced apoptosis in rat luteal cells. *J. Reprod. Fertil.* 120: 239-245.
- Telford, N. A., A. J. Watson and G. A. Schultz. 1990. Transition from maternal to embryonic control in early mammalian development: a comparison of several species. *Mol. Reprod. Dev.* 26: 90-100.
- Topper, J. N., J. Cai, G. Stavrakis, K. R. Anderson, E. A. Woolf, B. A. Sampson, F. J. Schoen, D. Falb and M. A. Gimbrone. 1998. Human prostaglandin transporter gene (h PGT) is regulated by fluid mechanical stimuli in cultured endothelial cells and expressed in vascular endothelium in vivo. *Circulation* 98: 2396-2403.
- Tsukita, S., K. Oishi, T. Akiyama, Y. Yamanashi, T. Yamamoto and S. Tsukita. 1991. Specific proto-oncogenic tyrosine kinase of src family are enriched in cell-to-cell adherens junctions where the level of tyrosine phosphorylation is elevated. *J. Cell Biol.* 113: 867-879.

- Valdimarsson, G. and G. M. Kidder. 1995. Temporal control of gap junction assembly in preimplantation mouse embryo. *J. Cell Sci.* 108: 1715-1722.
- Van Soom, A., M. L. Boerjan, P. E. J. Bols, G. Vanroose, A. Lein, M. Coryn and A. de Kruif. 1997. Timing of compaction and inner cell allocation in bovine embryos produced in vivo after superovulation. *Biol. Reprod.* 57: 1041-1049.
- van Wezel, T. and R. J. Rodgers. 1996. Morphological characterization of bovine primordial follicles and their environment in vivo. *Biol. Reprod.* 55: 1003-1111.
- Vane, J. R. and R. M. Botting. 1996. Mechanism of action of anti-inflammatory drugs. *Scand. J. Rheumatol.* 102 (Suppl.): 9-21.
- Velez, J. S., R. D. Randel and D. A. Neundorff. 1991. Effect of uterine manipulation on postpartum fertility and plasma 13,14-dihydro-15-keto-prostaglandin $F_{2\alpha}$ in Brahman cows and first-calf heifers. *Theriogenology* 36: 987-997.
- Vestweber, D., A. Gossler, K. Boller and R. Kemler. 1987. Expression and distribution of cell adhesion molecule uvomorulin in mouse preimplantation embryos. *Dev. Biol.* 124: 451-456.
- Wann, R. A. and R. D. Randel. 1990. Effect of uterine manipulation 35 days after parturition on plasma concentrations of 13, 14-dihydro-15-keto prostaglandin $F_{2\alpha}$ in multiparous and primiparous brahman cows. *J. Anim. Sci.* 68: 1389-1394.
- Warner, A. E. 1992. Gap junction in development, a perspective. *Semin. Cell Biol.* 3: 81-91.
- Watson, A. J. and G. M. Kidder. 1988. Immunofluorescence assessment of the timing of appearance and cellular distribution of Na^+K^+ -ATPase during mouse embryogenesis. *Dev. Biol.* 126: 80-90.
- Watson, A. J., G. M. Kidder and G. A. Schultz. 1992. How to make a blastocyst. *Biochem. Cell Biol.* 70: 849-855.
- Watson, A. J., M. E. Westhusin, P. A. De Sousa, D. H. Betts and L. C. Barcroft. 1999. Gene expression regulating blastocyst formation. *Theriogenology* 51: 117-133.
- Wiebold, J. L. 1988. Embryonic mortality and the uterine environment in first-service lactating dairy cows. *J. Reprod. Fertil.* 84: 393-399.
- Willye, A. H., J. F. R. Kerr and A. R. Currie. 1980. Cell death: the significance of apoptosis. *Int. Rev. Cytol.* 68: 251-306.

- Winkel, G. K., J. E. Ferguson, M. Takeichi and R. Nuccitelli. 1990. Activation of protein kinase C triggers premature compaction in the four-cell stage mouse embryo. *Dev. Biol.* 138: 1-15.
- Wrenzycki, C., D. Herrmann, J. W. Carnwath and H. Neiemann. 1996. Expression of the gap junction gene connexin 43 (Cx43) in preimplantation bovine embryos derived in vitro or in vivo. *J. Reprod. Fertil.* 108: 17-24.
- Wrenzycki, C., D. Herrmann, J. W. Carnwath and H. Niemann. 1999. Alterations in the relative abundance of gene transcripts in preimplantation bovine embryos cultured in medium supplemented with either serum or PVA. *Mol. Reprod. Dev.* 53: 8-18.
- Wu, X. M., M. Sawada and J. C. Carlson. 1992. Simulation of phospholipase A₂ by xanthine oxidase in the rat corpus luteum. *Biol. Reprod.* 47: 1053-1058.
- Zhang, M., S. Thorgeirsson. 1994. Modulation of connexins during differentiation of oval cells into hepatocytes. *Exp. Cell Res.* 213: 37-42.

VITA

Fernando Nestor Scenna was born August 30, 1974 in La Plata, Buenos Aires, Argentina. He attended San Luis High School until graduation in December 1992. Fernando enrolled at the Veterinary Science School of the National University of La Plata in March 1993 and he graduated as D.V.M. in July 1998. Fernando began graduate school in August 2000 at the University of Tennessee, Knoxville under the direction of Dr. F. Neal Schrick. In the Fall 2002, Fernando graduated with a M.S. degree in Animal Science with concentration in Reproductive Physiology.