

## ORIGINAL ARTICLE

# Effects of chronic low- and high-dose ethanol intake on the nitrenergic relaxations of corpus cavernosum and penile nitric oxide synthase in the rabbit

Y Yazir<sup>1</sup>, SS Gocmez<sup>2</sup>, T Utkan<sup>3</sup>, I Komsuoglu-Celikyurt<sup>4</sup>, N Gacar<sup>4</sup> and Y Sarioglu<sup>5</sup>

Epidemiological evidence showed that chronic ethanol consumption is a major risk factor in the development of impotence. The present study investigated the effects of carbachol-, electrical field stimulation (EFS)-, sodium nitroprusside (SNP)- and papaverine-induced relaxant responses in the isolated corpus cavernosum tissues from rabbits submitted to an 12-week course of chronic low (5% v/v) or high ethanol intake (30% v/v). Increased carbachol- and EFS-induced relaxant responses but not SNP and papaverine, were observed in low ethanol-fed rabbits compared with controls. However, impaired carbachol- and EFS-induced relaxant responses were observed in high ethanol-fed rabbits compared with control rabbits. There were no significant differences in SNP- and papaverine-induced relaxant responses between control and high ethanol-fed rabbits. In addition, decreased neuronal nitric oxide synthase (nNOS) and endothelial NOS (eNOS) immunoreactivity in penile tissue were found in high ethanol-fed rabbits, but increased the immunoreactivity in low ethanol-fed group, compared with control group. These results suggest that alterations in nitric oxide (NO) production within the cavernous tissue in the high ethanol-fed rabbits are, at least in part, responsible for the erectile dysfunction.

*International Journal of Impotence Research* (2012) **24**, 185–190; doi:10.1038/ijir.2012.14; published online 10 May 2012

**Keywords:** alcohol consumption; carbachol; corpus cavernosum; impotence; nitric oxide

## INTRODUCTION

Many epidemiological studies have shown that chronic ethanol consumption is associated with increased incidence of impotence.<sup>1</sup> Generally, low levels of alcohol ingestion increase male sexual activity by increasing libido; however, higher alcohol levels impair the penile tumescence and reduce sexual performance.<sup>2</sup> Nitric oxide (NO) is possibly the principal neurotransmitter for cavernous smooth muscle relaxation, the key step in the erectile process.<sup>3</sup> In recent years, impaired NO-mediated relaxant responses of corporal tissue has been recognized as an important cause of impotence in various pathological conditions such as diabetes mellitus,<sup>4</sup> aging,<sup>5</sup> hyperthyroidism,<sup>6</sup> pudental arterial occlusion,<sup>7</sup> hypercholesterolemia,<sup>8</sup> cavernous nerve denervation<sup>9</sup> and smoking.<sup>10</sup> Although chronic alcohol consumption results in impotence, Saito *et al.*<sup>11</sup> demonstrated that *in vivo* chronic ethanol administration (5% v/v) augmented acetylcholine- and electrical field stimulation (EFS)-induced relaxant responses of corporal smooth muscle. Also, many reports in the literature revealed that ethanol increases the synthesis or release of NO in isolated arteries and cultured endothelial cells.<sup>12,13</sup> On the other hand, some investigators noted that ethanol decreased endothelium-dependent relaxations in isolated arteries.<sup>14–16</sup> Recent studies on experimental animals and humans indicated that the effects of ethanol on NO availability are highly dependent on whether it is taken at low or high concentration and acute or chronic intake.<sup>17</sup> Therefore, the aim of the present study was to investigate the effects of chronic

ethanol intake on the nitrenergic-mediated relaxant responses of the rabbit corpus cavernosum and to determine penile endothelial nitric oxide synthase (eNOS) and neuronal NOS (nNOS) expression. To our knowledge, there are no similar studies examining the effect of chronic high dose of ethanol on the nitrenergic relaxation of corpus cavernosum smooth muscle.

## MATERIALS AND METHODS

Experiments were performed on mature male New Zealand White rabbits (2–2.5 kg) obtained from the Experimental Medical Research and Application Center (DETAB, Kocaeli University, Kocaeli, Turkey). The rabbit was chosen as the model based on the close similarities that have been reported in the reactivity *in vitro* of human and rabbit corpus cavernosum.

Ethical approval was granted by the Kocaeli University Animal Research Ethics Committee (Kocaeli, Turkey).

## Treatment schedule

The rabbits were divided into three groups: 9 rabbits in group 1 received tap water ad libitum, and 9 each in groups 2 and 3 were fed 5% ethanol and 30% ethanol, respectively. The model of ethanol feeding was that described previously in which rabbits received 5% ethanol (vol/vol) in the drinking water for the first week, 10% for the next 2 weeks, 20% from weeks 4–6 and 30% from weeks 6–12 in the group 3.<sup>18</sup> In the group 2, rabbits were received 5% ethanol constantly for 12 weeks. All rabbits had constant access to standard laboratory rabbit chow.

<sup>1</sup>Department of Histology and Embryology, Faculty of Medicine, Kocaeli University, Kocaeli, Turkey; <sup>2</sup>Department of Pharmacology, Faculty of Medicine, Namik Kemal University, Tekirdağ, Turkey; <sup>3</sup>Department of Pharmacology and Experimental Medical Research and Application Center, Faculty of Medicine, Kocaeli University, Kocaeli, Turkey; <sup>4</sup>Department of Pharmacology, Faculty of Medicine, Kocaeli University, Kocaeli, Turkey and <sup>5</sup>Department of Pharmacology, Faculty of Medicine, Gazi University, Ankara, Turkey. Correspondence: Professor Dr T Utkan, Department of Pharmacology and Experimental Medical Research and Application Center, Faculty of Medicine, Kocaeli University, 41380, Umuttepe, Kocaeli, Turkey.

E-mail: [tijenutkan@yahoo.com.tr](mailto:tijenutkan@yahoo.com.tr)

Received 14 June 2011; revised 5 October 2011; accepted 11 April 2012; published online 10 May 2012

### Strip preparation and organ bath studies

The rabbits from each group were anesthetized with ketamine (25 mg/kg, i.p.) and xylazine (5 mg/kg, i.p.) and exsanguinated 12 weeks post treatment, as previously described.<sup>19</sup> Briefly, the penis was dissected free and removed at the level of the crural attachments to the puboischial bones. The specimen was immediately placed in Krebs's bicarbonate buffer composed of (mM): NaCl 118, KCl 4.7, CaCl<sub>2</sub> 2.5, NaHCO<sub>3</sub> 25, MgSO<sub>4</sub> 1.2, KH<sub>2</sub>PO<sub>4</sub> 1.2 and glucose 11. The tunica albuginea was cleared of the overlying tissue and opened. The proximal half of the corporal body was dissected free from the tunica and harvested en bloc as previously described.<sup>5,7,9</sup> Each corporal body was cut transversely to obtain two longitudinal strips. Each strip was mounted in a 20-ml organ bath containing Krebs's bicarbonate buffer equilibrated with a gas mixture of 95% O<sub>2</sub> and 5% CO<sub>2</sub> and maintained at 37 °C. Recording of isometric strip tension were made using a transducer (MAY-COM, FDT 10 A, COMMAT ILETISIM, Ankara, Turkey) and recorded online on a computer via a four channel transducer data acquisition system (TDA 94, COMMAT ILETISIM, Turkey) using appropriate software (Polywin 95 1.0 COMMAT ILETISIM, Turkey). The resting load was set at 2 g, a value that has been previously found to be optimal for the measurement of changes in the tension of rabbit corpus cavernosal tissue preparations.<sup>5,7,9</sup> The preparations were allowed to equilibrate in Krebs's bicarbonate buffer for 1 h and during this time Krebs's bicarbonate buffer was replaced every 15 min with fresh solution.

The corpus cavernosum strips in organ chambers were contracted with 10<sup>-6</sup> M phenylephrine and added to carbachol (10<sup>-8</sup>–10<sup>-4</sup> M), sodium nitroprusside (SNP; 10<sup>-8</sup>–10<sup>-4</sup> M) and papaverine (10<sup>-5</sup>–10<sup>-4</sup> M). Transmural EFS was provided by a stimulator (ST95 PT, COMMAT ILETISIM, Turkey) and applied via two platinum wire electrodes set vertically within the organ bath on the opposite sides of the suspended tissue. Before electrical stimulation, the tissue was treated with guanethidine (5 μM; an adrenergic nerve blocker) and atropine (1 μM; a muscarinic receptor blocker) for 30 min. Strips were precontracted with phenylephrine at 10<sup>-5</sup> M, EFS was performed. Square-wave pulses of 10 V with a 0.5-ms duration in 10-s trains with varying frequencies (2–32 Hz) were applied at 5-min intervals. The strips were allowed to return to baseline precontractile tension at each frequency.

### Immunohistochemistry

Cavernous specimens were fixed in 10% neutral-buffered formalin. The specimens were embedded in regular paraffin wax and cut into 3-μm thick sections. Tissue sections were deparaffinized in xylene and rehydrated in ethanol. Antigen retrieval was performed by microwave oven at 600W for 3 × 5 min in the citrate buffer. Endogenous peroxidase was blocked by 3% H<sub>2</sub>O<sub>2</sub> in methanol 15 min and again washed three times in phosphate-buffered saline. Afterwards, sections were incubated in a blocking serum (Histostatin plus kit broad spectrum, Invitrogen, CA, USA) for 10 min at room temperature to block non-specific binding. Subsequently, sections were incubated over night at room temperature with primary goat anti-nNOS polyclonal antibody (abcam-ab1376) at 2 μg/ml and 20 min at room temperature with primary anti-eNOS polyclonal antibody (abcam-ab87750) at 5 μg/ml in a humidified chamber. Negative control incubations were performed by replacing the primary antibody with the appropriate non-immune immunoglobulin G in the same concentrations. Sections were washed three times in phosphate-buffered saline and incubated with the biotinylated secondary antibodies (Histostatin plus kit broad spectrum, Invitrogen, CA, USA) for 20 min at room temperature. After three washes with phosphate-buffered saline, the sections were incubated with peroxidase labeled streptavidin (Histostatin plus kit broad spectrum, Invitrogen, CA, USA) for 10 min. Peroxidase activity was visualized with 3-amino-9-ethylcarbazol chromogen in large-volume 3-amino-9-ethylcarbazol substrate (AEC red kit 00-2007, Invitrogen, CA, USA) for nNOS and with 3, 3'-diaminobenzidine substrate (DAB kit 88-2014, Invitrogen, CA, USA) for eNOS. All incubations were performed in a moist chamber at room temperature using phosphate-buffered saline for washes between incubation steps. The sections were counterstained with Mayer's hematoxylin (Invitrogen, CA, USA) and mounted with Clearmount

(Invitrogen, CA, USA) on glass slides. Slides were examined under light microscope (Olympus BX 50, Tokyo, Japan) and photomicrographs were taken with Leica DM 100 (Leica DFC 290HD, Wetzlar, Hessen, Germany). All samples were treated with exactly the same protocol. Two independent observers graded the staining intensity on a semi-quantitative scale ranging from no expression (–), very weak (+), moderate (++) , strong (+++) to very strong (++++) expression.

### Biochemical parameters

Blood glucose, triglyceride, total cholesterol and high-density lipoproteins (HDL) levels were measured in the rabbit's blood using Abbott Architect c16000 autoanalyzer (Abbott Laboratories, IL, USA). Also testosterone levels were measured in blood using Roche Analytics E170 Immunology Analyzer (Roche, Tokyo, Japan).

### Analysis of data

Experimental values are expressed as the mean ± s.e.m. The relaxant effects of the agonists are expressed as percentage of the precontraction response to phenylephrine. To evaluate the effects of the agonists, pD<sub>2</sub> (that is, the negative logarithm of the concentration for the half-maximal response; EC<sub>50</sub>) and maximum response (E<sub>max</sub>) values were calculated. Agonist pD<sub>2</sub> value was calculated from each agonist dose–response curve by linear portion of the curve and taken as a measure of the sensitivity of the tissues to each agonist.

Statistical comparison between the groups was performed using analysis of variance followed by Tukey's test. The score of immunoreactivity was assessed by the Kruskal–Wallis Test. Probabilities of P < 0.05 were considered significant.

### Drugs

The following drugs were all obtained from Sigma Chemical (St. Louis, MO, USA): carbachol chloride, phenylephrine hydrochloride, SNP, atropine sulfate, guanethidine sulfate and papaverine hydrochloride. All drugs dissolved in distilled water and were freshly prepared on the day of the experiment.

## RESULTS

The contractions elicited by KCl (124 mM) were similar in all the three groups (Table 1).

### Effects of low- and high-dose ethanol consumption on endothelium-dependent relaxation

Carbachol (10<sup>-8</sup>–10<sup>-4</sup> M) produced concentration-dependent relaxation in submaximally precontracted (10<sup>-6</sup> M phenylephrine) corpus cavernosum strips obtained from control, low-dose and high-dose alcohol-fed rabbits. In the low-dose alcohol consumption group, the carbachol concentration–response curve was

**Table 1.** Maximum responses (E<sub>max</sub>) for carbachol, sodium nitroprusside (SNP), papaverine and KCl in corpus cavernosum strips obtained from control and alcohol-fed groups

	Control	5% Alcohol-fed	30% Alcohol-fed
Carbachol (%)	67.9 ± 4	90.4 ± 4.2*	42.8 ± 4.4*
EFS (%)	52.1 ± 8.5	83.7 ± 5.1*	33.6 ± 1.9*
SNP (%)	98.8 ± 0.5	98.7 ± 0.4	97.3 ± 0.7
Papaverine (%)	98.6 ± 1.4	97.5 ± 3.4	98.0 ± 2.7
KCl (mg)	3144.6 ± 466.3	3352.6 ± 402.6	3422.6 ± 266.1

Abbreviations: EFS, electrical field stimulation; KCl, potassium chloride.

\*P < 0.05, statistically different from the response from control rabbits (n = 9 in each group).

E<sub>max</sub> (Percentage of 10<sup>-6</sup> M phenylephrine contractile response) values for carbachol, EFS, SNP, papaverine and E<sub>max</sub> (mg of contractile response) values for KCl. Values are arithmetic means ± s.e.m, n = the number of preparations used.

shifted to the left and  $E_{max}$  values were increased compared with control rabbits ( $P < 0.05$ ; Figure 1, Table 1). There were no significant changes in  $pD_2$  values (Table 2). However, in the high-dose alcohol consumption group, the carbachol concentration–response curve was shifted to the right with significantly lower  $pD_2$  value and  $E_{max}$  values were decreased compared with control rabbits ( $P < 0.05$ ; Figure 1, Tables 1 and 2).

#### Effects of low- and high-dose ethanol consumption on endothelium-independent relaxation


In precontracted strips, SNP ( $10^{-10}$ – $10^{-4}$  M) and papaverine ( $10^{-5}$ – $10^{-4}$  M) produced concentration-dependent relaxation. The relaxation elicited by both SNP and papaverine was similar in low-dose alcohol consumption, high-dose alcohol consumption and control groups, and there were no significant changes in the  $pD_2$  and  $E_{max}$  values (Figure 2, Tables 1 and 2).

#### Effects of low- and high-dose ethanol consumption on neurogenic relaxation

In precontracted strips, EFS (2–32 Hz) evoked frequency-dependent relaxations. In the low-dose alcohol-fed group, EFS responses were higher than those in the control group ( $P < 0.05$ ; Figure 3, Table 1). But in the high-dose alcohol-fed group, EFS responses were inhibited compared with the control group ( $P < 0.05$ ; Figure 3, Table 1).

#### Blood glucose, triglyceride, total cholesterol, HDL and testosterone levels in control and alcohol-fed rabbits

Levels of blood glucose, triglyceride, total cholesterol, HDL and testosterone in control and alcohol-fed rabbits are shown in Table 3. The blood glucose (mg/dl) and HDL levels (mg/dl) of 5% alcohol-fed and 30% alcohol-fed rabbits were similar to the control group. Cholesterol (mg/dl) and triglyceride (mg/dl) levels were significantly higher in 30% alcohol-fed rabbits compared with the control group ( $P < 0.05$ ). Additionally, testosterone level (ng/dl) was significantly lower in 30% alcohol-fed rabbits compared with that of the control group ( $P < 0.05$ ).



**Figure 1.** Carbachol concentration–response in isolated rabbit corpus cavernosum strips precontracted with phenylephrine  $10^{-6}$  M. Each point is expressed as a percentage of the contraction induced by phenylephrine and is given as the mean  $\pm$  s.e.m. Numbers in parentheses indicate the number of preparations used. \* $P < 0.05$ , statistically different from the response from control rabbits.


#### Effects of low- and high-dose ethanol consumption on penile nNOS and eNOS expression

Immunopositive staining was found after nNOS or eNOS immunohistochemistry in penile tissue. Weak immunostaining was observed in penile tissue in high ethanol-fed group (Figures


**Table 2.**  $pD_2$  values for carbachol and sodium nitroprusside (SNP) in corpus cavernosum strips obtained from control and alcohol-fed groups

	Control	5% Alcohol-fed	30% Alcohol-fed
Carbachol	5.92 $\pm$ 0.24	5.79 $\pm$ 0.24	4.82 $\pm$ 0.28*
SNP	5.50 $\pm$ 0.17	5.45 $\pm$ 0.18	5.60 $\pm$ 0.08

\* $P < 0.05$ , statistically different from the response from control rabbits ( $n = 9$  in each group). Values are arithmetic means  $\pm$  s.e.m.



**Figure 2.** Sodium nitroprusside concentration–response in isolated rabbit corpus cavernosum strips precontracted with phenylephrine  $10^{-6}$  M. Each point is expressed as a percentage of the contraction induced by phenylephrine and is given as the mean  $\pm$  s.e.m. Numbers in parentheses indicate the number of preparations used.



**Figure 3.** Relaxation responses evoked by EFS of isolated rabbit corpus cavernosum strips precontracted with phenylephrine  $10^{-6}$  M. Each point is expressed as a percentage of the contraction induced by phenylephrine and is given as the mean  $\pm$  s.e.m. Numbers in parentheses indicate the number of preparations used. \* $P < 0.05$ , statistically different from the response from control rabbits. EFS, electrical field stimulation.

4c and f) compared with control (Figures 4a and d;  $P < 0.05$ ), indicating the NO generation through nNOS or eNOS is impaired in the penile tissue of the high-dose ethanol-fed rabbits. Contrarily, in low ethanol-fed group (Figures 4b and e), increased immunostaining was observed compared with the control group (Figures 4a and d;  $P < 0.05$ ).

## DISCUSSION

The present study addresses the changes in neurogenic and endothelium-dependent relaxation to carbachol in the corporal smooth muscle and its relationship with blood alcohol levels in rabbit-ingested ethanol at different doses for 12 weeks. In most of the human studies, chronic alcohol consumption is a major risk factor in the development of impotence. Also, it is well known that the concentration and duration of ethanol ingestion have an important impact on erectile outcomes in animal models. Experiments on alcohol and erectile response showed that high doses of alcohol attenuate erectile response; however, light-to-moderate alcohol consumption increases male sexual activity.<sup>20–22</sup> The results obtained with the corpus cavernosum were consistent with previous studies showing that light-to-moderate alcohol consumption is inversely associated with cardiovascular diseases.<sup>23–25</sup> It has been proposed that increased NO production is a major physiological system for the protective action of moderate alcohol consumption.<sup>12,13</sup> However, experimental and epidemiological evidence show that chronic high-dose ethanol consumption increases mortality and causes cardiovascular complications, including hypertension.<sup>26</sup> It is hypothesized that

chronic alcohol-induced hypertension is associated with the downregulation of antioxidants and NO production in the aorta of rats.<sup>14–16</sup>

NO plays an essential role in the erection of penis. It is well known that both the autonomic nerves that innervate corpus cavernosum are independent sources of NO.<sup>27,28</sup> In this study, EFS and exogenous application of SNP, papaverine and carbachol produced relaxation responses in both alcohol-treated and control groups. There were no significant differences in response to SNP and papaverine in all the groups. However, the relaxation response to EFS and carbachol were significantly increased in low-dose alcohol-treated rabbits. Therefore, it is speculated that low-dose alcohol consumption increased production of NO. It is also possible that low-dose alcohol consumption may increase relaxation of trabecular smooth muscle or increase the ability to relax via the NO/cGMP pathway; however, this possibility is unlikely as the strips relaxed well to papaverine or SNP. Similarly, we have previously shown increased endothelium-dependent relaxant response to acetylcholine in isolated aortas from 8-week ethanol-treated (7.5%) rats.<sup>29</sup> Our findings are consistent with previous studies demonstrating that chronic moderate alcohol ingestion enhanced NO production by vascular cells *in vitro*.<sup>13</sup> Our evidence indicates that chronic light alcohol ingestion increases trabecular NO production and causes increased trabecular smooth muscle relaxation leading to increase in sexual performance.

Also, in this study, EFS- and carbachol-induced relaxation responses were significantly decreased in the higher-dose alcohol-treated group. The results of the present study further showed that penile nNOS significantly decreased in high ethanol consumption, indicating erectile dysfunction. It is reported that, when rats treated with graded concentrations of alcohol for 12 weeks, higher concentrations of alcohol were associated with hypertension and reduced plasma levels of NO metabolites. Authors recently clarified that daily oral treatment with ethanol for 12 weeks caused significant decreases in aortic NO production, reduced endothelium-dependent vasorelaxation and increased blood pressure that were due to, in part, alcohol-induced reductions in eNOS.<sup>14</sup> This finding is consistent with our results, therefore we suggest that impaired NO production have an important pathophysiological role in the erectile dysfunction after high-dose alcohol consumption in rabbits.

In addition, several mechanisms have been proposed to account for the pathogenesis of impotence associated with chronic alcohol consumption such as impairment of hypothalamic function.<sup>30</sup> Androgens are essential in the maintenance of libido


**Table 3.** Blood glucose, triglyceride, total cholesterol, HDL and testosterone levels in control and alcohol-fed rabbits

	Control	5% Alcohol-fed	30% Alcohol-fed
Glucose (mg/dl)	143.33 ± 1.47	141.67 ± 23.64	149 ± 7.07
Cholesterol (mg/dl)	42 ± 10.14	31.83 ± 4.64	93.3 ± 11.74*
HDL (mg/dl)	11 ± 3.24	17 ± 4.15	19.83 ± 5.05
Triglyceride (mg/dl)	57.83 ± 8.31	74.5 ± 8.50	181.55 ± 21.30*
Testosterone (ng/dl)	4.36 ± 0.56	3.79 ± 0.62	0.95 ± 0.24*

Abbreviation: HDL, high-density lipoprotein.

\* $P < 0.05$ , statistically different from the response from control rabbits ( $n = 9$  in each group).

Values are arithmetic means ± s.e.m.



**Figure 4.** Representative light microscopy of control and chronic ethanol-fed groups in penile tissue. Decreased (c) neuronal nitric oxide synthase (nNOS) and (f) endothelial NOS (eNOS) immunoreactivity at corpus cavernosum in high-dose ethanol-fed rabbits and increased (b) nNOS and (e) eNOS immunoreactivity at penile tissue in low-dose ethanol-fed rabbits compared with (a, d) control groups.

and have an important role in regulating erectile capacity in man. In hypogonadal patients, it is known that exogenous testosterone administration stimulates erectile function.<sup>31</sup> Previous studies have shown that alcohol generally decreases circulating testosterone levels in animals and humans, but many men who are alcohol dependent have normal testosterone and estrogen levels.<sup>32–34</sup> Cicero and Badger<sup>32</sup> reported that low doses of ethanol significantly increased serum testosterone, whereas high doses decreased serum levels of testosterone, in male Sprague–Dawley rats. Moreover, it has been demonstrated that androgens are essential in maintenance of NO-mediated erectile activity in the rat.<sup>35</sup> Animal studies have also revealed that androgens regulate the expression of NOS in the penis.<sup>36,37</sup> These findings are consistent with our results. Although this study was not designed to explore the effects of testosterone on the corpus cavernosum smooth muscle function, their decrease after heavy alcohol consumption may have contributed to the impaired relaxation response to both EFS and carbachol as observed in our study.

Epidemiological evidence suggests that alcohol ingestion modulates the development of cardiovascular disease.<sup>38</sup> Moderate alcohol consumption may prevent atherosclerotic vascular disease, whereas excessive intake may enhance cardiovascular disease.<sup>39</sup> The mechanism which underlie the protective effects of moderate alcohol consumption on cardiovascular disease risk are not fully understood. However, moderate alcohol consumption affects lipoprotein metabolism.<sup>40</sup> Epidemiological<sup>41</sup> and experimental studies<sup>42</sup> have demonstrated that alcohol consumption increases HDL concentration, which is inversely associated with cardiovascular disease risk. In addition, chronic alcohol intake has been shown to increase,<sup>43</sup> decrease<sup>44</sup> or not modify<sup>45</sup> serum cholesterol levels. Furthermore, Azadzi and Saenz de Tejada<sup>8</sup> reported that hypercholesterolemia impairs endothelium-dependent relaxation of rabbit corpus cavernosum smooth muscle. Hence, in our study, changes in blood total cholesterol, HDL and triglyceride levels in chronic high-dose alcohol consumption seem to be important parameters for impaired nitrergic relaxations in corpus cavernosum.

Another popular theory is that acetaldehyde inhibits NO formation, which leads to impaired NO-mediated relaxation in chronic high-dose of alcohol consumption. It is well known that after ingestion, ethanol is metabolized sequentially to acetaldehyde, acetate, CO<sub>2</sub> and water. Previously, Kim *et al*<sup>46</sup> showed that acetaldehyde suppressed neurogenic relaxation induced by transmural EFS in rabbit corpus cavernosum smooth muscle, and noted that increasing acetaldehyde level seen in chronic alcoholism may contribute to male erectile dysfunction mainly by the inhibition of NO formation. Taking these findings together, we speculated that chronic high dose of alcohol consumption impairs the synthesis or availability of NO in corpus cavernosum smooth muscle.

In conclusion, the present study showed that chronic low-dose ethanol consumption leads to increased nitrergic-mediated corpus cavernosum smooth muscle relaxation as well as increased immunostaining in nNOS and eNOS. Our findings suggest that this increment appears to be related to increased NO production and/or release. Also, we demonstrated that nitrergic-mediated relaxations decreased in strips of corpus cavernosum from chronic high-dose ethanol-treated rabbits as well as decreased immunostaining in nNOS and eNOS. This suggests that treatment with high-dose ethanol under the conditions that were used in the study impaired both the neurogenic and endothelial functions. Therefore, we speculated that impaired nitrergic relaxations during high-dose ethanol treatment may contribute to erectile dysfunction through reduced NO production and/or release.

#### ACKNOWLEDGEMENTS

This study was supported by a grant from Kocaeli University Research Fund (Project number: 200319). Research Foundation had no further role in the study design, in the

collection, analysis and interpretation of data, in the writing of the report and in the decision to submit the paper for publication.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### REFERENCES

- Benson GS, Doileau MA. The penis: sexual function and dysfunction. In: Gillenwater JY, Groyhack JT, Howards SS, Duckett JW (eds). *Adult and Pediatric Urology*. Mosby Year Book: Philadelphia, 1991; pp 1599–1642.
- Rubin HB, Henson DE. Effects of alcohol in male sexual responding. *Psychopharmacology* 1976; **47**: 123–134.
- Wespes E. Smooth muscle pathology and erectile dysfunction. *Int J Impot Res* 2002; **14**: S17–S21.
- Azadzi KM, Saenz de Tejada I. Diabetes mellitus impairs neurogenic and endothelium-dependent relaxation of rabbit corpus cavernosum smooth muscle. *J Urol* 1992; **148**: 1587–1591.
- Utkan T, Yildirim S, Sarioglu Y, Yildiz F, Yildirim K. Aging impairs nitric oxide-mediated relaxant responses of rabbit corporal smooth muscle. *Nitric Oxide Biol Chem* 2002; **6**: 342–346.
- Ozdemirci S, Yildiz F, Utkan T, Ulak G, Cetinaslan B, Erden F *et al*. Impaired neurogenic and endothelium-dependent relaxant responses of corpus cavernosum smooth muscle from hyperthyroid rabbits. *Eur J Pharmacol* 2001; **428**: 105–111.
- Utkan T, Sarioglu Y, Utkan NZ, Kurnaz F, Yildirim S. Effects of chronic unilateral arterial occlusion on reactivity of isolated corpus cavernosum strips from rabbits. *Eur J Pharmacol* 1999; **367**: 73–79.
- Azadzi KM, Saenz de Tejada I. Hypercholesterolemia impairs endothelium-dependent relaxation of rabbit corpus cavernosum smooth muscle. *J Urol* 1991; **146**: 238–240.
- Utkan T, Sarioglu Y, Yazir Y. Changes in the neurogenic and endothelium-dependent relaxant responses of rabbit corpus cavernosum smooth muscle after cavernous nerve neurotomy. *Methods Find Exp Clin Pharmacol* 2010; **32**: 151–156.
- Gomez SS, Utkan T, Duman C, Yildiz F, Ulak G, Gacar N *et al*. Secondhand tobacco smoke impairs neurogenic and endothelium-dependent relaxation of rabbit corpus cavernosum smooth muscle: improvement with chronic oral administration of L-arginine. *Int J Impot Res* 2005; **17**: 437–445.
- Saito M, Broderick GA, Levin RM. Effect of chronic ethanol consumption on the pharmacological response of the rabbit corpus cavernosum. *Pharmacology* 1994; **49**: 386–391.
- Abou-Agag LH, Khoo NK, Binsack R, White CR, Darley-Usmar V, Grenett HE *et al*. Evidence of cardiovascular protection by moderate alcohol: role of nitric oxide. *Free Radic Biol Med* 2005; **39**: 540–548.
- Kleinhenz DJ, Sutliff RL, Polikandriotis JA, Walp ER, Dikalov S I, Guidot DM *et al*. Chronic ethanol ingestion increases aortic endothelial nitric oxide synthase expression and nitric oxide production in the rat. *Alcohol Clin Exp Res* 2008; **32**: 1–7.
- Husain K, Vazquez-Ortiz M, Lalla J. Down regulation of aortic nitric oxide and antioxidant systems in chronic alcohol-induced hypertension in rats. *Hum Exp Toxicol* 2007; **26**: 427–434.
- Husain K. vascular endothelial oxidative stress in alcohol-induced hypertension. *Cell Moll Biol* 2007; **53**: 70–77.
- Husain K, Meija J, Lalla J, Kazim S. Dose response of alcohol-induced changes in BP, nitric oxide and antioxidants in rat plasma. *Pharmacol Res* 2005; **51**: 337–343.
- Toda N, Ayajiki K. Vascular actions of nitric oxide as affected by exposure to alcohol. *Alcohol Alcoholism* 2010; **45**: 347–355.
- Chan TCK, Sutter MC. Ethanol consumption and blood pressure. *Life Sci* 1983; **33**: 1965–1973.
- Yildirim MK, Yildirim S, Utkan T, Sarioglu Y, Yalman Y. Effects of castration on adrenergic, cholinergic and nonadrenergic noncholinergic responses of isolated corpus cavernosum from rabbits. *Br J Urol* 1997; **79**: 964–970.
- Rubin HB, Henson DE. Effects of alcohol in male sexual responding. *Psychopharmacology* 1976; **47**: 123–134.
- Wilson GT, Niaraur JL. Alcohol, selective attention and sexual arousal in men. *J Stud Alcohol* 1985; **46**: 107–115.
- Cooper AJ. The effects of intoxication levels of ethanol on nocturnal penile tumescence. *J Sex Marital Ther* 1994; **20**: 14–23.
- Camargo Jr CA, Stampfer MJ, Glynn RJ, Gaziano JM, Manson JE, Goldhaber SZ *et al*. Prospective study of moderate alcohol consumption and risk of peripheral arterial disease in US male physicians. *Circulation* 1997; **95**: 577–580.
- Hill JA. In vino veritas: alcohol and heart disease. *Am J Med Sci* 2005; **329**: 124–135.
- Thun MJ, Peto R, Lopez AD, Monaco JH, Henley SJ, Heat Jr CW *et al*. Alcohol consumption and mortality among middle-aged and elderly US adults. *N Engl J Med* 1997; **337**: 1705–1714.

- 26 Klatsky AL. Alcohol, coronary disease and hypertension. *Annu Rev Med* 1996; **47**: 149–160.
- 27 Bush PA. Nitric oxide is a potent relaxant of human and rabbit corpus cavernosum. *J Urol* 1992; **147**: 1650–1655.
- 28 Azadzi KM. Endothelium derived nitric oxide and cyclooxygenase products modulate corpus cavernosum smooth muscle tone. *J Urol* 1992; **147**: 220–225.
- 29 Utkan T, Yildiz F, Ilbay G, Ozdemirci S, Erden F, Gacar N *et al*. Blood pressure and vascular reactivity to endothelin 1, phenylephrine, serotonin, KCl, and acetylcholine following chronic alcohol consumption *in vitro*. *Fundam Clin Pharmacol* 2001; **15**: 157–165.
- 30 Rivier C. Alcohol rapidly lowers plasma testosterone levels in the rat: evidence that a neural brain-gonadal pathway may be important decreased testicular responsiveness to gonadotropin. *Alcohol Clin Exp Res* 1999; **23**: 38–45.
- 31 Kwan M, Greenleaf WJ, Mann J, Crapo L, Davidson JM. The nature of androgen action on male sexuality: a combined laboratory-self report study on hypogonadal men. *J Clin Endoc Metab* 1983; **57**: 557–562.
- 32 Cicero TJ, Badger TM. Effects of ethanol on the hypothalamic-pituitary-gonadal axis in the male rat. *J Pharmacol Exp Ther* 1997; **201**: 427–433.
- 33 Cicero TJ, Bell RD. Effects of ethanol on acetaldehyde on the biosynthesis of testosterone in the rodent testes. *Biochem Biophys Res Commun* 1980; **94**: 814–819.
- 34 Eriksson CJP, Fukunaga T, Lindman R. Sex hormone response to alcohol. *Nature* 1994; **369**: 711.
- 35 Reilly CM, Zamorano P, Stoper VS, Mills TM. Androgenic regulation of NO availability in rat penile erection. *J Androl* 1997; **18**: 110–115.
- 36 Lugg JA, Rajfer J, Gonzales-Cadavid NE. Dihydrotestosterone is the active androgen in the maintenance of nitric oxide-mediated penile erection in the rat. *Endocrinology* 1995; **136**: 1495–1501.
- 37 Mills TM, Reilly CM, Lewis RW. Androgens and penile erection: a review. *J Androl* 1996; **17**: 633–638.
- 38 Rimm EB. Alcohol consumption and coronary heart disease: good habits may be more important than just good wine. *Am J Epidemiol* 1996; **143**: 1089–1093.
- 39 Camargo CA, Hennekens CH, Gaziano JM, Glynn RJ, Manson JE, Stampfer MJ. Prospective study of moderate alcohol consumption and mortality in US male physicians. *Arch Intern Med* 1997; **157**: 79–85.
- 40 Van der Gaag MS, Van Tol A, Scheek LM, James RW, Urgert R, Schaafsma G *et al*. Daily moderate alcohol consumption increases serum paraoxonase activity: A diet-controlled, randomised intervention study in middle-aged men. *Atherosclerosis* 1999; **147**: 405–410.
- 41 Gaziano JM, Buring JE, Breslow JL, Goldhaber SZ, Rosner B, VanDenBurg M *et al*. Moderate alcohol intake, increased levels of high-density lipoprotein and its subfractions, and decreased risk of myocardial infarction. *N Engl J Med* 1993; **329**: 1829–1834.
- 42 Van der Gaag MS, Ubbink JB, Sillanaukee P, Nikkari S, Hendriks HFJ. Effect of consumption of red wine, spirits, and beer on serum homocysteine. *Lancet* 2000; **255**: 1522.
- 43 Lamisse F, Schellenberg F, Bouyou E, Delarue J, Benard JY, Couet C. Serum lipids and alcohol consumption in alcoholic men: effect of withdrawal. *Alcohol* 1994; **29**: 25–30.
- 44 Taskinen M, Nikkila EA, Valimaki M. Alcohol-induced changes in serum lipoproteins and in their metabolism. *Am Heart J* 1987; **113**: 458–464.
- 45 Marques-Vidal P, Cambou JP, Nicaud V, Luc G, Evans A, Arveiler D. Cardiovascular risk factors and alcohol consumption in France and Northern Ireland. *Atherosclerosis* 1995; **115**: 225–232.
- 46 Kim HJ, Sohng I, Lee G, Kim JJ, Koh SK. Effects of acetaldehyde on responses of rabbit corpus cavernosal smooth muscle. *J Korean Med Sci* 2000; **15**: 295–298.