

p75 neurotrophin receptor mediates apoptosis in transit amplifying cells and its overexpression restores cell death in psoriatic keratinocytes

Running title: p75NTR mediates apoptosis in human keratinocytes

F Truzzi¹, A Marconi¹, P Atzei¹, MC Panza¹, R Lotti¹, K Dallaglio¹, R Tiberio², E Palazzo¹, C Vaschieri¹ and C Pincelli^{*1}.

¹*Institute of Dermatology, School of Biosciences and Biotechnologies, University of Modena and Reggio Emilia, Modena, Italy;* ²*Institute of Dermatology, University of Piemonte Orientale, Novara.*

Correspondence to:

Carlo Pincelli
Institute of Dermatology
School of Biosciences and Biotechnologies,
University of Modena and Reggio Emilia,
Via del Pozzo 71
41100 Modena, Italy.
Tel +390594222931
Fax: +390594224271
carlo.pincelli@unimore.it

Key words: p75NTR; keratinocyte; apoptosis; psoriasis

Abbreviations: p75NTR, p75 neurotrophin receptor; NT, neurotrophin; NGF, nerve growth factor; BDNF, brain-derived neurotrophic factor; NT-3, neurotrophin-3; NT-4, neurotrophin-4; KSC, keratinocyte stem cells; TA, transit amplifying; PM, post mitotic

Abstract

p75 neurotrophin receptor (p75NTR) belongs to the TNF-receptor superfamily and signals apoptosis in many cell settings. In human epidermis, p75NTR is mostly confined to the transit amplifying (TA) subpopulation of basal keratinocytes. Brain-derived neurotrophic factor (BDNF) or neurotrophin-4 (NT-4), that signal through p75NTR, induce keratinocyte apoptosis, while β -amyloid, a ligand for p75NTR, triggers caspase-3 activation to a greater extent in p75NTR transfected cells. Moreover, p75NTR co-immunoprecipitates with NRAGE, induces the phosphorylation of c-Jun N-terminal kinase (JNK) and reduces nuclear factor kappa B (NF- κ B) DNA binding activity. p75NTR also mediates pro-NGF-induced keratinocyte apoptosis through its co-receptor sortilin. Furthermore, BDNF or β -amyloid cause cell death in TA, but not in keratinocyte stem cells (KSC) or in p75NTR silenced TA cells. p75NTR is absent in lesional psoriatic skin and p75NTR levels are significantly lower in psoriatic than in normal TA keratinocytes. The rate of apoptosis in psoriatic TA cells is significantly lower than in normal TA cells. BDNF or β -amyloid fail to induced apoptosis in psoriatic TA cells, and p75NTR retroviral infection restores BDNF- or β -amyloid-induced apoptosis in psoriatic keratinocytes. These results demonstrate that p75NTR plays a proapoptotic role in keratinocytes, and is involved in the maintenance of epidermal homeostasis.

INTRODUCTION

The neurotrophins (NTs) nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophin-3 (NT-3), and neurotrophin-4 (NT-4) are a family of secreted growth factors that exert a wide range of functions in the development and maintenance of the nervous system (1). The activities of NTs are mediated by two structurally unrelated classes of receptors, the common neurotrophin receptor p75 (p75NTR) (2), a member of the tumor necrosis factor (TNF) receptor superfamily (3), and the Trk family receptors tyrosine kinases (4). While p75NTR binds all NTs with equal low affinity, TrkA preferentially interacts with NGF, TrkB with BDNF and NT-4, and TrkC with NT-3 (2). Trk receptor signaling and activities have been well characterized, and it is now widely accepted that the primary role of Trk is the control of neuronal survival and differentiation (5). In contrast, the precise role of p75NTR has been slow to emerge. Recent findings suggest that p75NTR contributes to different signaling pathways. First, p75NTR is a positive modulator of Trk-mediated functions, in that it acts as a co-receptor that refines Trk affinity and specificity for NT (3). On the other hand, p75NTR also controls Trk-independent activities. Numerous studies have demonstrated a pro-apoptotic role of p75NTR in the nervous system. Indeed, the p75NTR cytoplasmic tail contains several potential motifs for interactions with downstream signaling and, similar to other members of the TNF receptor superfamily, it comprises the so-called "death domain" (6). p75NTR can cooperate with many different protein partners and form multimeric receptor complexes to produce a number of

cellular responses, including apoptosis (7). p75NTR signaling involves modulation of nuclear factor kappa B (NF- κ B) and the phosphorylation of c-Jun N-terminal kinase (JNK), as well as increased production of ceramides (8). Moreover, the precursor form of NTs proNGF binds p75NTR, but not TrkA and is a potent inducer of p75NTR-dependent apoptosis. Sortilin, a 95kDa member of the Vps10p domain receptors acts as a co-receptor for p75NTR by binding the pro region of NGF (9).

Human epidermis is a multilayered epithelium where the principal cell type is the keratinocyte. Proliferation takes place in the basal epidermal layer and differentiation begins as keratinocytes migrate through the suprabasal layers. Keratinocyte stem cells (KSC) reside in the basal layer, govern the renewal of epidermis, and generate transit amplifying (TA) cells that terminally differentiate after a number of cell divisions (10). Human basal keratinocytes synthesize and secrete biologically active NGF, NT-3, BDNF and NT-4, and express NT receptors (11,12). K252a, an inhibitor of Trk phosphorylation, induces apoptosis in human keratinocytes, indicating that autocrine NGF protects these cells from programmed cell death through its high affinity receptor (13). On the other hand, the role of p75NTR in human keratinocytes remains to be elucidated. Considering that the balance between Trk and p75NTR seems to be crucial for cell survival or death (14), we wanted to investigate the role of p75NTR in human keratinocytes that express both NT receptors and undergo apoptosis upon various stimuli, including the inhibition of Trk (13).

Hyperproliferation and a distorted differentiation process represent the most prominent features of psoriasis. Psoriasis is an immuno-inflammatory skin disease mediated by the release of Th1 and Th17 T cytokines, which in turn trigger the keratinocyte response (15). However, keratinocytes could carry themselves a defect leading to psoriasis formation (16). In this regard, apoptosis is consistently reduced in psoriasis (17), and psoriatic keratinocytes are abnormally resistant to apoptosis (18). If p75NTR played a proapoptotic role in keratinocyte, we hypothesized that it is also involved in the altered epidermal homeostasis typical of psoriasis.

We present evidence that p75NTR is predominantly expressed in TA cells and induces apoptosis in normal human keratinocytes either alone or in association with sortilin. We also show that, even in the presence of Trk, p75NTR can mediate apoptosis, thus overcoming Trk survival signals. Furthermore, we demonstrate that expression of p75NTR is almost absent in psoriatic lesions. β -amyloid or BDNF induce apoptosis in normal, but not in psoriatic TA cells. Finally, p75NTR infection restores susceptibility to apoptosis in psoriatic keratinocytes.

Results

p75NTR mediates apoptosis in human keratinocytes. While the presence of p75NTR mRNA in human keratinocytes has been previously reported, (11,19) its function has never been elucidated. Given the pro-apoptotic activity

of p75NTR in most cell systems, we investigated whether p75NTR could mediate cell death also in human keratinocytes. To this purpose, and in order to overcome binding to Trk, we needed a specific ligand for p75NTR, which could trigger Trk-independent functions. β -amyloid specifically binds p75NTR with an affinity close to that for NT and mediates apoptosis in neurons and 3T3 cells (20). When β -amyloid was provided to human keratinocytes, survival was significantly affected, as compared to control (Figure 1a). In addition, β -amyloid-induced apoptosis was significantly higher than control, as shown by flow cytometry (subG1 peak, Figure 1b). Caspase-3 is the executioner caspase responsible for the final step of the apoptotic machinery and its activation is a clear marker of cell death. To confirm p75NTR-mediated apoptosis in human keratinocytes, we evaluated the expression of p75NTR and active caspase-3 after addition of β -amyloid. β -amyloid, but not diluent alone induced the activation of caspase-3 in p75NTR positive cells only. In particular, while few p75NTR positive cells are caspase-3 negative, all caspase-3 positive cells express p75NTR (Figure 1c). We have shown previously that human keratinocytes express only the extracellular fragment of TrkB, thus lacking the functional high-affinity signaling domain for BDNF and NT-4 (12). Therefore, we reasoned that BDNF and NT-4 must perform their activities through p75NTR. Indeed, both BDNF and NT-4 significantly induced apoptosis in human keratinocytes, in a time-dependent manner, as shown by TUNEL staining (Figure 1d). Moreover, the addition of anti-p75NTR neutralizing antibody prevented BDNF-induced cell death (Figure 1e). To definitely confirm the pro-

apoptotic role of p75NTR, we present evidence that both BDNF and β -amyloid activated caspase-3 to a greater extent in keratinocytes infected with p75NTR retroviral vector, as compared to mock keratinocytes (Figure 1f, g).

p75NTR signals apoptosis in human keratinocytes Several p75NTR interacting proteins have recently been identified. In particular, NRAGE has been shown to colocalize with p75NTR under physiological conditions and to play a role in p75NTR mediated cell death (21). We demonstrated that, in β -amyloid treated human keratinocytes, p75NTR and NRAGE immunoprecipitates are more abundant than in keratinocytes treated with diluent alone (Figure 2a). Although, the signal transduction involved in p75NTR-mediated apoptosis is still to be completely defined, the critical role of JNK phosphorylation is widely accepted (22). We showed that in human keratinocytes, BDNF induced the phosphorylation of only JNK1 (Figure 2b), that, unlike JNK2, is essential for apoptosis induced by death receptors (23). On the other hand, the role of NF- κ B in p75NTR signal transduction is still controversial. Here, we demonstrate that BDNF reduced the DNA binding activity of NF- κ B. Reduction was more pronounced in p75NTR infected keratinocytes (Figure 2c). This suggests that in human keratinocytes, p75NTR exerts its proapoptotic activity also via the inhibition of NF- κ B.

p75NTR mediates apoptosis in TA cells While p75NTR-induced apoptosis was statistically significant, the low percentage of cells undergoing death

remained to be explained. To better clarify this issue, we investigated the expression of p75NTR in human epidermis. We observed that only a subpopulation of basal keratinocytes expressed p75NTR, whereas no p75NTR expression was detected in suprabasal layers, as ruled out by the simultaneous staining with keratin 10 and transglutaminase I (Figure 3a). To better determine the subpopulation of basal keratinocytes expressing p75NTR, we isolated and characterized KSC and TA cells, as shown previously (24). p75NTR mRNA was significantly more expressed in TA cells than in KSC (Figure 3b). Moreover, p75NTR protein was strongly expressed in TA cells and nearly absent in KSC, while it was not detectable in postmitotic (PM) cells (Figure 3c). If p75NTR is predominantly expressed in TA cells, we predicted that apoptosis would occur only in these cells. Indeed, both BDNF and β -amyloid induced apoptosis in TA cells, whereas they fail to cause cell death in KSC, as shown by the percentage of cells in subG1 peak by flow cytometry (Figure 3d). To confirm that p75NTR is responsible for apoptosis in TA cells, we silenced p75NTR (Figure 3e), and demonstrated that either BDNF or β -amyloid were not able to induce apoptosis in p75NTR siRNA treated TA cells (Figure 3f). It is interesting to note that, upon treatment with β -amyloid, the percentage of apoptosis was strikingly higher in TA cells than in the whole keratinocyte population (Figure 1b), as shown by subG1 peak.

p75NTR mediates pro-NGF-induced keratinocyte apoptosis through its co-receptor sortilin Because sortilin acts as a co-receptor for p75NTR and

mediates apoptosis in other cell settings (9), we first evaluated sortilin expression in human keratinocytes. These cells contained levels of sortilin mRNA which were almost three fold the amount of p75NTR (Figure 4a). Similarly, flow cytometry demonstrated that sortilin protein was expressed in a significantly higher number of cells than p75NTR (Figure 4b), consistent with the wide variety of functions of this protein. As p75NTR was predominantly expressed in TA cells, we examined the pattern of sortilin expression in the same subpopulation of keratinocytes. Consistently with p75NTR, sortilin was expressed mostly in TA cells, while KSC contained a slightly higher amount of sortilin than PM cells, both at the mRNA and at the protein level (Figures 4c and d). It has been proposed that proNGF binds p75NTR, whereas mature NT binds TrkA. Inhibiting sortilin in cells that normally express p75NTR, suppresses proNGF-mediated cell death (9). As sortilin and p75NTR were both predominantly expressed in TA cells, we wanted to evaluate the effect of proNGF in keratinocytes. Increasing doses of proNGF reduced keratinocyte viability (Figure 4e). Moreover, co-addition of proNGF and the Trk inhibitor k252a decreased the rate of cell survival to a greater extent as compared to proNGF or k252a alone (Figure 4f), indicating that simultaneous blockade of Trk signaling and binding of p75NTR is a strong proapoptotic stimulus. This was further confirmed by the TUNEL method, which demonstrated a significant rate of apoptosis induced by proNGF, which was significantly augmented by the addition of k252a (Figure 4g). To definitely confirm the proapoptotic role of p75NTR, we transfected HaCaT keratinocytes with a combination of pcDNA3-

p75NTR and pcDNA3-sortilin (Figure 4h). While transfection alone failed to exert any proapoptotic effect, the addition of proNGF significantly promoted apoptosis to the same level as the positive control (Figure 4i). Taken together, these results indicate that p75NTR together with sortilin mediates the proapoptotic role of proNGF in human keratinocytes, even in the presence of Trk.

p75NTR is reduced in lesional psoriatic epidermis We had previously shown that psoriatic lesions express higher levels of NGF as compared to normal skin (25), which accounts for the increased sprouting of p75NTR and TrkA positive terminal cutaneous nerves (26). In the present study, we focused on the expression of p75NTR and Trk receptors in psoriatic keratinocytes. By immunohistochemistry, p75NTR was detected occasionally in few basal cells from non lesional psoriatic skin, expression being substantially reduced as compared to normal skin. Interestingly, p75NTR protein was completely absent in lesional psoriasis (Figure 5a). To better understand this finding, we evaluated p75NTR expression at the mRNA level. Both Northern Blot (Figure 5b) and Real Time PCR (Figure 5c) revealed that p75NTR mRNA was normally expressed in psoriatic epidermis. Yet, Western blotting on protein lysates from epidermis (Figure 5d) and flow cytometry (Figure 5e) on suspended cells confirmed the lack of p75NTR protein in psoriatic keratinocytes. On the other hand, TrkA protein levels were higher in non lesional and lesional psoriatic epidermis than in normal epidermis. While TrkB was slightly decreased in

psoriatic skin, TrkC remained unchanged. Finally, sortilin was slightly increased in lesional psoriatic epidermis (Figure 5d).

Defective p75NTR renders psoriatic TA cells resistant to apoptosis

Because TA cells express the highest levels of p75NTR, we compared its expression in TA cells in normal and psoriatic keratinocytes by flow cytometry. While no major variations were detected in KSC and PM cells between normal and psoriatic keratinocytes, TA cells from psoriatic keratinocytes presented a four-fold decrease in p75NTR expression as compared to TA from normal cells (Figure 6a). Because TA cells are more prone to cell death than KSC (24) and reduced expression of p75NTR in TA cells seems to account for the low levels of p75NTR in psoriasis, we compared the rate of apoptosis in TA and KSC from normal and psoriatic skin. While percentage of apoptotic cells was similar in KSC from healthy and diseased skin, TA cells from psoriatic skin were statistically less susceptible to apoptosis than the same cells from normal skin, as shown by propidium iodide staining in situ (Figure 6b). As β -amyloid and BDNF induce apoptosis in normal keratinocytes, we analyzed their effect on TA cells from psoriatic and normal keratinocytes. While β -amyloid or BDNF significantly reduced viability and induced apoptosis (subG1 peak) in TA cells from normal epidermis, they failed to exert any effect on TA cells from psoriatic keratinocytes (Figure 6c, d).

Overexpression of p75NTR restores apoptosis in psoriatic keratinocytes

Because psoriatic epidermis is characterized by defective apoptosis (17) and reduced p75NTR, we wanted to investigate the function of this receptor in psoriatic keratinocytes. To this purpose, we infected psoriatic keratinocytes with p75NTR retroviral vector (Figure 7a) and treated cells with either β -amyloid or BDNF. Both stimuli reduced viability in p75NTR infected psoriatic keratinocytes, but not in mock psoriatic cells (Figure 7b). 20% difference in viability roughly corresponds to the percentage of p75NTR infected cells (data not shown). Moreover, while infection with p75NTR increased the rate of apoptosis in untreated cells (cntrl), BDNF induced a significantly higher percentage of cells in subG1 peak in p75NTR infected, as compared to either mock keratinocytes or to infected keratinocytes treated with diluent alone (Figure 7c).

Discussion

NTs and their receptors are expressed in most human skin cells, constituting a network potentially responsible for multiple functions (12, 27). In this context, we detected p75NTR in basal keratinocytes, in agreement with previous reports. While those were mostly incidental observations in studies related to

p75NTR expression in cutaneous nerve fibers or during skin pathologies (28), our work precisely addresses p75NTR localization in human epidermis. We detected p75NTR in basal, but not in suprabasal keratinocytes, as confirmed by the double staining with keratin 10 and transglutaminase I. This is in general agreement with p75NTR expression in other multilayered epithelial cells, including prostate and esophagus. On the other hand, while in other epithelia p75NTR appears to be evenly distributed, the localization pattern in human epidermis seems to be unique, in that only a subpopulation of the germinative basal layer expresses the receptor. We present evidence that p75NTR is almost exclusively expressed in TA cells, with little protein observed in KSC. Although several markers have been proposed, KSC and TA cells are still difficult to be clearly defined. TA cells have been shown to be more susceptible than KSC to apoptosis (29). In fact, TA cells express higher levels of proapoptotic and lower levels of antiapoptotic markers, as opposed to KSC (24). Our finding of p75NTR being predominantly expressed in TA cells seems to be consistent with the proapoptotic role of this receptor in human keratinocytes. Keratinocytes also express Trk receptors (11, 12), and NTs activities strongly depend on the relative expression of the two receptors on the target cell population (30). When both receptors are expressed, NTs bind to Trk-p75NTR high-affinity complex and mediate survival signals in many cell settings (2). In this study, using specific ligands for p75NTR, we were able to induce apoptosis, thus counteracting the survival effects mediated by Trk. Indeed, β -amyloid specifically binds p75NTR and induces apoptosis only in keratinocytes

expressing p75NTR, similarly to β -amyloid peptide-induced cell death in Alzheimer's neuronal cells (31) and in neuroblastoma (32). Because β -amyloid is not a physiologic ligand for p75NTR in human epidermis, we treated keratinocytes with BDNF or NT-4 that bind p75NTR, in absence of TrkB. In fact, BDNF or NT-4 significantly increase the rate of cell death, and induce a high degree of caspase-3 activation in p75NTR overexpressing keratinocytes. The signaling pathways underlying the p75NTR-dependent apoptotic response involve a number of factors. NRAGE, a p75NTR-interacting protein disrupts p75NTR-TrkA complex, and induces cell death through JNK-dependent pathway (21), thus contributing to p75NTR-dependent apoptosis. Consistent with these findings, we provide evidence that upon binding with a specific ligand, p75NTR interacts with NRAGE and phosphorylates JNK in human keratinocytes. While JNK activation was mostly demonstrated by overexpressing p75NTR or after serum withdrawal in the nervous system, the present study indicates that BDNF, released at the skin level, signals apoptosis, by phosphorylating JNK. p75NTR also activates NF- κ B, which is mainly viewed as an anti-apoptotic signalling factor. On the other hand, in neonatal rat oligodendrocytes, activation of NF- κ B was shown to be associated with apoptosis. Our findings indicate that in human keratinocytes, p75NTR signals apoptosis via the reduction of NF- κ B anti-apoptotic function. Taken together, these results suggest that p75NTR has its own signaling pathway also in human keratinocytes.

In the neuronal system, proNGF binds to p75NTR and its co-receptor sortilin, but not to Trk, whereas mature NTs bind to Trk only, leading to the conclusion that the precursors and the mature NTs elicit opposing responses, that is cell death and survival, respectively (9, 33). Here we show that human keratinocytes express sortilin with a pattern similar to that of p75NTR. Indeed, sortilin is predominantly expressed in TA cells, which are more prone to apoptosis (24). This study also shows that proNGF induces keratinocyte apoptosis, while the simultaneous inhibition of Trk phosphorylation by k252a elicits an additive proapoptotic activity, further indicating that the two NT receptors exert opposing effects also in keratinocytes (13). While co-transfection of p75NTR and sortilin per se failed to induce apoptosis, addition of proNGF to transfected cells significantly stimulated caspase activation, consistent with the concept that also in keratinocytes, proNGF acts as a crosslinker, creating a ternary complex with sortilin and p75NTR (9). In substance, we have shown that p75NTR acts as a proapoptotic receptor in human keratinocytes either when it is ligated by proNGF or by mature NTs. This latter mechanism is at variance with the view that, in the presence of Trk, p75NTR is a facilitator of Trk-mediated survival in the nervous system. A possible explanation relies on the following observations: 1. While in the neuronal system, high concentrations of mature NTs required to induce even modest levels of apoptosis are considered non-physiological (34), in human epidermis, the NT-4, BDNF or other yet unidentified ligands could autocrinally and paracrinally be continuously released and in turn bind p75NTR, thus

overcoming Trk functions. 2. Epithelial cells behave differently from neurons, as normal prostate and bladder cells express Trk (35, 36) and still undergo apoptosis via p75NTR binding (37). It has been recently shown that, under certain circumstances and stimuli, a Trk-to-p75NTR molecular switch occurs favoring cell death (38).

p75NTR is nearly absent in psoriatic keratinocytes that are reportedly resistant to apoptosis. On the other hand, restoring p75NTR levels renders psoriatic keratinocytes again susceptible to cell death. This indicates that p75NTR plays a role in epidermal homeostasis and its deficiency accounts for the well-documented resistance of psoriatic keratinocytes to apoptosis. Moreover, p75NTR alteration, coupled with the marked overexpression of NGF and TrkA that mediate survival signals, indicates that the imbalance of NGF receptors underlies the disturbed epidermal homeostasis in psoriasis. Early mathematical models describing cell renewal in psoriasis had demonstrated that the typical psoriatic tissue architecture is based not only on an increased turnover rate in the germinative cell, but also on a defect in cells transiting to the more differentiated state. More recently, Castelijns and co-workers, using kinetics parameters, have proposed that epidermal abnormalities in psoriasis are due to a defect in the transit-amplifying compartment, rather than to a reduction in the cell cycle time. TA cells in psoriasis are thought to be kept in a prolonged proliferative state, which would account for an increased number of cell division, with no reduction of cell cycle time (39). Alternatively, the defect in TA could be due to a lack of differentiation/apoptosis. This has been recently

confirmed by in-silico studies simulating psoriasis by altering TA cells (40). The inherent malfunction in the behavior of this subpopulation is enforced by altered levels of a number of markers in TA from psoriatic lesions as compared to TA from normal skin (41). The present work demonstrates that the most remarkable reduction in p75NTR expression was detected in TA cells. This finding, together with the lesser susceptibility to apoptosis of psoriatic TA cells, strongly indicates that alteration of TA cells in psoriasis is partly based on a defect in p75NTR. While there seems to be a general agreement on the prominent role of TA cells in psoriasis, those works were mostly based on computational and expression data. Here we show that stimuli that induce apoptosis through p75NTR in TA cells from normal epidermis, fail to do so in psoriatic TA, providing the first functional insight into the defect of this keratinocyte subpopulation in the pathogenesis of psoriasis. If p75NTR mRNA is normally expressed in psoriatic keratinocytes, there must be some post-translational alteration leading to the disappearance of the protein, possibly related to a calcium defect in psoriatic keratinocytes (42). In conclusion, p75NTR acts as a proapoptotic receptor in human keratinocytes and plays a crucial role in epidermal homeostasis.

Materials and methods

Cell Culture

Normal human keratinocytes were obtained from foreskin and cultured in serum-free medium KGM (Clonetics, San Diego, CA) as described (12). Freshly

keratinocytes were also divided into three populations and plated onto plastic dishes, coated for 2 hours at 37°C with type IV collagen 100µg/ml (Sigma, St. Louis, MO). They were first allowed to adhere to type IV collagen for 5 min, and the non-adherent cells were then transferred to fresh collagen-coated dishes and allowed to attach overnight. Finally, keratinocytes not yet attached after one night were plated onto type IV collagen to obtain a third population. The three keratinocyte populations were characterized based on β_1 -integrin levels and colony forming efficiency, as previously described (24). This allowed to obtain a population enriched in KSC, a population consisting of TA cells and a third population of PM cells. Keratinocytes were then cultured in serum-free medium (KGM) and used for further experiments. The spontaneously transformed keratinocyte line HaCaT was kindly provided by Dr. N. Fusenig (DKFZ Heidelberg).

Western blotting analysis

Cells or epidermis coming from skin biopsies were washed with phosphate-buffered saline (PBS) and lysed on ice, for Trk, sortilin and JNK detection, in lysis buffer pH 7.4 (50mM Tris-HCl, 1% NP40, 0.25% sodium deoxycholate, 150mM NaCl, 1 mM EGTA, 1mM Na₃VO₄, 1mM NaF), for p75NTR detection in lysis buffer pH 7.5 (150mM NaCl, 15mM MgCl, 1mM EGTA, 50mM Hepes, 10% Glicerolo, 1% Triton) or, for caspase-3, with lysis buffer pH 6.8 (62.5mM Tris-HCl, 2% SDS, 10% glycerol, 50mM DTT, 0,01% bromophenol blue) containing protease inhibitors, as described previously (12). 20µg of total protein were

analyzed under reducing conditions for sortilin and JNK on 7% polyacrylamide gels and blotted onto nitrocellulose membranes. To verify equal loading of total proteins in all lanes, the membranes were stained with Ponceau Red. The blots were blocked for 2 hours in blocking buffer (PBS buffer, pH 7.4, with 0.2% Tween 20 and 5% nonfat milk) and incubated overnight at 4°C with anti-p75NTR mouse monoclonal (1:1000; Upstate, Lake Placid, NY, USA), anti-TrkA rabbit polyclonal (1:1000; Upstate), anti-TrkB rabbit polyclonal (1:750; Upstate), anti-TrkC goat polyclonal (1:750; Upstate), anti-sortilin mouse monoclonal antibody (1:250, BD Biosciences Pharmingen, San Diego, CA, USA), anti-active JNK (1:5000, Promega, Madison, WI, USA), anti-caspase-3 (1:1000, Cell Signaling Technology, Boston, MA, USA) or with anti- β -actin mouse monoclonal antibody (1:5000; Sigma) as a control. Then membranes were washed in PBS/Tween 20, incubated with peroxidase-conjugated goat anti-rabbit antibody, anti-mouse antibody (1:3000; Biorad, Hercules, CA, USA) or donkey anti-goat antibody (1:1000; Santa Cruz Biotechnology Inc, Santa Cruz, CA, USA) for 45 min at room temperature, washed, and developed using the ECL chemiluminescent detection system (Amersham Biosciences UK Limited, Little Chalfont Buckinghamshire, England).

Immunoprecipitation

Human keratinocytes were treated with or without 40 μ M β -amyloid. 24h later cells were harvested with RIPA buffer (10mM Tris, pH 8.0; 150mM NaCl; 1% Nonidet P-40, 0.5% deoxycholate; 0,1% SDS) containing protease inhibitors.

Nuclei were removed by centrifugation (11,000rpm for 3 min). Monoclonal antibody anti-p75NTR (Upstate) or polyclonal serum against NRAGE (kindly provided from professor P.A. Barker) were bound to sepharose beads 1h at 4°C under rotation. Then, pre-cleared lysates were conjugated to sepharose beads-antibody complex or to sepharose beads alone as control, overnight at 4°C under rotation. Immunocomplexes were washed with Port Buffer (10mM Tris-HCl, pH 8.5; 150mM NaCl; 0.1% Renex 30; 0.01% BSA; 2.5% NaN₃) six times and with PT Buffer (10mM Tris-HCl, pH 8.5; 150mM NaCl; 0.5% Tween 20; 2.5% NaN₃) three times. Samples were eluted with 1X Laemmli sample buffer for 5°C at 90°C and Western Blotting for p75NTR or NRAGE was performed as previously described.

Transfection of HaCaT cells

For p75NTR and sortilin transient transfection, cells were seeded in 6-well plates (5×10^5 cells/well) and in 96-well tissue culture plate (4×10^4 cells/well) in antibiotic-free medium. HaCaT cells were double transfected using Lipofectamine Plus reagent (Invitrogen Corporation, Carlsbad, CA, USA) with 2 μ g/well (in 6-well plates) or 0.15 μ g/well (in 96-well plates) of pcDNA3-BimEL or pcDNA3-p75NTR or pcDNA3-sortilin overnight in Opti-MEM medium (Invitrogen Corporation). Then cells were treated with various stimuli for further experiments and p75NTR or sortilin protein level was detected by Western blotting, as described below.

siRNA transfection of TA cells

About 7×10^4 TA cells/well were plated on 6-well plates in penicillin/streptomycin free medium. 24 hours later cells were transfected with 50nM or 100nM p75NTRsiRNA (Dharmacon Inc, Lafayette, CO, USA) or scrambled RNAi as mock control, combined with Lipofectamin 2000 and Opti-MEM (both from Invitrogen Corporation) as datasheet suggests. Cells were transfected twice and used for FACS analysis. p75NTR protein level was detected by Western blotting, as described below.

Infection of normal and psoriatic keratinocytes

10^6 cells from normal or psoriatic biopsies were plated on 75cm^2 flask over a feeder-layer of mitomycin C treated 3T3 fibroblasts (7×10^5 cells for 75cm^2 flask) and mitomycin C treated p75NTR-LNSN packaging cells (3.5×10^5 for 75 flask) or mitomycin C treated LNSN packaging cells (kindly provided from professor F. Mavilio) with medium for keratinocytes (11) without EGF. 48 hours later EGF was added to the cultures. 24 hours after infection cells were plated in KGM for MTT, PI or Western blot analysis.

Real Time PCR

RNA from cells separated into the three populations or from total keratinocytes or from healthy or psoriatic epidermis was extracted using TRI Reagent method performed as described by Sigma. To verify the quality of the RNA, we ran an agarose minigel in RNase-free conditions. Quantitative Real Time PCR

was performed with an ABI 7500 (Applied Biosystems, Foster City, CA, USA) for p75NTR and sortilin. As an internal control, housekeeping gene β -actin mRNA expression was measured in a separated tube. H₂O RNase-free was used as a negative control. We performed three independent experiments, and each one was analysed in a triplicate PCR-reaction. Then, 1 μ g of RNA was subjected to retro-transcription and amplification in a 50 μ l reaction mixture using the One-Step RT-PCR Master Mix Reagents kit (Applied Biosystems). For p75NTR, sortilin and β -actin were used Pre-Developed TaqMan Assay Reagents (MGB™ probes were FAM or VIC dye labelled, Applied Biosystems). Thermal cycling conditions for One-Step RT-PCR were: initial reverse transcription at 48°C for 30 min, then DNA Polymerase activation at 95°C for 10 min, followed by 40 cycles of denaturation at 95°C for 15 s, annealing/extension at 60°C for 1 min. Data from each sample were compared with a calibrator, using the Sequence Detection Software Version 1.2.3 according to the Relative Quantification Study method (Applied Biosystems). Results were obtained as the mean from three independent experiments. Double-sided Student's t-test was performed between samples and calibrator.

Immunohistochemistry

For double staining, cryosections (4 μ m) of normal human skin were acetone-fixed, rehydrated in phosphate-buffered saline (PBS). Quenching of the endogenous peroxidase activity was obtained by treatment with 3% H₂O₂ in PBS. Samples were incubated either with mouse anti-transglutaminase type I

(1:10, Biomedical Technologies Inc, Stoughton, MA, USA) or with mouse anti-cytokeratin 10 antibody (1:50, Dako, Glostrup, Denmark) for 1 hour, then with secondary antibody anti-mouse biotinylated (1:3000, Dako) for 45 min and with a streptavidin-biotin peroxidase complex for 30 min. The reaction products were visualized with amino-methylene-carbazole. Double staining was continued incubating the slides with 1% goat serum. Then the slides were incubated with mouse anti-p75NTR antibody (1:250, Lab Vision Corporation, Neomarkers, Fremont, CA, USA) for 1 hour and, after three washes of 5 min in PBS, slides were incubated with secondary antibody anti-mouse (1:20, Dako) for 45 min at room temperature, followed by incubation with anti-mouse alkaline phosphatase complex for 30 min (1:20, Dako). After three washes of 5 min in PBS, the slides were stained with Fast blue (Sigma). Negative controls were obtained by omitting primary antibody.

For single staining, cryosections (4 μ m) of normal, non lesional and lesional psoriatic human skin were acetone-fixed (10 min), rehydrated in phosphate-buffered saline (PBS). The slides were incubated with mouse anti-p75NTR antibody (1:250, Lab Vision Corporation) for 1 hour and, after three washes of 5 min in PBS, incubated with secondary antibody anti-mouse (1:20, Dako) for 45 min at room temperature, followed by incubation with anti-mouse alkaline phosphatase complex for 30 min (1:20, Dako). After three washes of 5 min in PBS, the slides were stained with New fuchsin (Sigma).

Confocal immunofluorescence

For double staining, keratinocyte cultures were washed in PBS buffer and fixed in situ in 4% paraformaldehyde for 20 min and air dried. After a rehydration in PBS, cells were permeabilized for 10 min with 0.5% Tryton X-100 in 0.1% sodium citrate, treated for 5 min with 50 mM NH₄Cl and incubated with 1% bovine albumin serum for 20 min. Then cells were incubated for 1 hour with the mouse monoclonal anti-caspase-3 active antibody (1:100, R & D Systems, Minneapolis, MN, USA) and for 1 hour with RPE Cy5-conjugated anti-mouse (dilution 1:100, Dako). Then keratinocytes were washed in PBS, labelled with a mouse anti-p75NTR antibody (1:250, Lab Vision Corporation) for 1 hour and with FITC-conjugated anti-mouse antibody (dilution 1:100, Dako) for 45 min. Fluorescent specimens were analyzed by a Confocal Scanning Laser Microscopy (Leica TCS SP2).

MTT assay

Keratinocytes or transfected HaCaT were plated in 96-well tissue culture plate (8000 cells/well). 72 hours after seeding, cells were treated with 40 μ M β -amyloid (25-35) (Bachem, Bubendorf, Switzerland) or 100ng/ml BDNF or 100ng/ml NT-4 or different doses of proNGF (kindly provided from B. Hempstead) for 24, 48 and 72 hours. Proliferative cells were detected by incubating with MTT (Sigma-Aldrich) solution at 37°C for 4 hours. They were solubilized with DMSO and the formazan dye formation was evaluated by scanning multiwell spectrophotometer at 540 nm. The results are expressed as

optical density units (OD) or as viability percentage respect of control. Results are calculated as the mean \pm SD of three different experiments. Student's T-test was used for comparison of the means.

Cell death analysis by Flow cytometry

72 hours after seeding, cells were treated with 40 μ M β -amyloid or 100ng/ml BDNF. 24 or 48 hours later, cells were trypsinized and resuspended (5 \times 10⁵ cells) in 1ml hypotonic fluorochrome solution: 50 μ g/ml propidium iodide containing 0.1% sodium citrate and 0.5% Tryton X-100 (Sigma). After 15 min, cells were analyzed using an Epics XL flow cytometer (Coulter Electronics Inc., Hialeah, FL, USA). Apoptosis was detected by evaluating the reduced fluorescence of the DNA binding dye PI in the apoptotic nuclei.

TUNEL assay

Cells after treatment were fixed in situ in paraformaldehyde (4% in PBS) and air dried. The slides were stained with the "In situ cell death detection kit" (Roche Diagnostics, Basel, Switzerland) as recommended by the manufacturer. Fluorescent specimens were analyzed by Confocal Scanning Laser Microscopy (Leica TCS SP2) in conjunction with a conventional optical microscope (Leica DM IRBE). Results are calculated as the mean \pm SD of three different experiments. Student's T-test was used for comparison of the means.

Detection of apoptotic cells by confocal microscopy

Cells were seeded on glass coverslips coated with poly-L-lysine 100mg/ml for 5 min, fixed in methanol:water (80:20) for 5 min, washed in phosphate-buffered saline (PBS) and subsequently stained with propidium iodide (Sigma, 5 μ g/ml in PBS) for 5 min. Coverslips were examined by laser-scanning confocal microscopy (Leica TCS-SP2) to identify apoptotic nuclei. Nuclei were scored and more than 400 nuclei were counted for each sample. The results are expressed as percentage of the number of apoptotic nuclei over the total number of nuclei examined.

FACS analysis

Keratinocytes were washed in PBS and incubated with mouse biotin-conjugated anti-p75NTR (20 μ l/1 \times 10⁶ cells, BD Biosciences Pharmingen) or rabbit polyclonal anti-sortilin antibody (1:2000, kind gift from Claus M. Petersen, Aarhus University, Denmark) for 20 min at 4°C. For sortilin, after a brief washing in PBS, the cells were incubated with the biotinylated anti-rabbit antibody 1:500 (Dako, Glostrup, Denmark) for 20 min on ice. For both stainings, the incubation with antibodies was followed by incubation with streptavidin/RPE-Cy5 1:50 (Dako) for 20 min on ice. After washing, cells were analyzed using an Epics XL flow cytometer.

Caspase Activity Measurements

The caspase-3-like protease activity in cell lysates was measured using a fluorometric substrate Ac-DEVD-AFC (Alexis Corporation, San Diego, CA, USA), which was dissolved in dimethyl sulfoxide and stored as a 10mM solution at -20°C. HaCaT cells were collected, washed in cold PBS, and lysed at 4°C in buffer A (50mM HEPES, pH 7.5, 150mM NaCl, 20mM EDTA, 1 mM phenylmethylsulfonyl fluoride, 10µg/ml leupeptin, 1µg/ml aprotinin, and 0.3% Nonidet P-40). Homogenates were centrifuged (13,000 rpm, 10min), and the protein concentrations of the supernatants were determined using a Bradford protein assay kit (Bio-Rad). Equal amounts of supernatants (normalized for protein content) were mixed in wells of 96-well plates in a 100µl final volume, in buffer B (50mM HEPES, pH 7.4, 100mM NaCl, 1mM EDTA, 10% sucrose, 0.5% CHAPS, 5mM dithiothreitol, and 100 µM Ac-DEVD-AFC substrate (caspase-3/7). Substrate cleavage was monitored continuously by spectrofluorometry, in kinetic mode, using Fluorolite 1000 (Dynatech Laboratories Inc., Chantilly, VA, USA). Data are reported as relative fluorescence units of product produced/min/10µg of total protein.

Northern blot

Northern Blot was performed using DIG Northern Starter Kit (Roche). Briefly, total cellular epidermal mRNA taken from biopsies of patient with or without psoriasis was isolated with 1 ml of TRI Reagent performed as described by Sigma. p75NTR DNA in pCEP4 vector was linearized at restriction sites

downstream of the cloned insert and purified. PCR of p75NTR vector using T7 RNA polymerase promoter sequences (FW:5'-CTCATCCCTGTCTATTGCTCCA-3';REV:5'-TAATACGACTCACTATAGGAGCTGGAGCATCGGTTGTC-3') was performed using Expand High Fidelity PCR System (Roche). 1 µg of p75NTR DNA template was labeled with T7 RNA polymerase for 1 hour at 42°C, after that DNase I was added for 15 min at 37°C, to remove DNA template, as manufacture's instructions. Yield of DIG-labeled p75NTR RNA was determined comparing it to DIG-labeled actin RNA as control (10ng/µL). 2 µg of total RNA from patients was loaded in MOPS/formaldehyde gel. RNA was blotted over nylon membrane, positively charged, by capillary transfer over night. The RNA was fixed by short-wave UV exposure and RNA hybridization was performed using DIG-labeled p75NTR probe at 68°C over night with gentle agitation. Membrane was incubated with anti-digoxigenin-AP, Fab fragments (1:10000) in blocking solution) 30 min at room temperature. Membrane was washed, equilibrated for 5 min and covered with CDP-Star ready-to-use solution for 5 min at 20°C and signal imaging was developed. To perform actin control, membrane was stripped and hybridized with DIG-labeled actin RNA probe and detection was performed as described before.

Gel mobility shift assay

Nuclear proteins were isolated from the cellular pellets. Samples were suspended with hypotonic buffer (1.5 mM MgCl₂, 0.2 mM PMSF, 1 mM DTT, 10 mM KCl, 10 mM HEPES, pH 7.9) for 15 min. Proteins were centrifuged, then

the pellets were washed with hypotonic buffer and centrifuged. Supernatants were discarded and pellets were suspended in low salt buffer (25% glycerol, 1.5 mM MgCl₂, 0.2 mM EDTA, 0.2 mM PMSF, 1 mM DTT, 20 mM KCl, 20 mM HEPES, pH 7.9). Nuclear proteins were released by adding high salt buffer (25% glycerol, 1.5 mM MgCl₂, 0.2 mM EDTA, 0.2 mM PMSF, 1 mM DTT, 1.2 M KCl, 20 mM HEPES, pH 7.9). Samples were incubated on ice for 30 min, centrifuged and supernatants were collected and stored at -70 °C.

Double-stranded oligonucleotides for the NF-κB-binding site were obtained from Santa Cruz Biotechnology and labeled with [γ -³²P] ATP (Perkin-Elmer, Waltham, MA, USA) and T4 polynucleotide kinase (Promega). The labeled oligonucleotides were separated on a SDS-PAGE gel and eluted out of the gel in distilled H₂O (4 °C, overnight) and the activity of labeled probe counted.

Total of 5 μg of nuclear proteins were used in each reaction. Extracts were incubated with the radiolabeled probe in binding buffer (4 mM Tris-HCl, 12 mM HEPES at pH 7.9, 60 mM KCl, 0.5 mM EDTA, 1 mM dithiothreitol, and 12% glycerol) for 25 min at room temperature. Non-specific binding was blocked by 2 μg of poly(dI-dC):poly(dI-dC) (Sigma). After incubation, samples were fractionated on a 4% polyacrylamide gel in 0.25x Tris-borate-EDTA at 4°C. Following electrophoresis, the gel was dried and autoradiographed.

Acknowledgment

This work was supported by a grant from Ministry of Instruction, University and Research PRIN 2003061751_004.

We wish to thank Chiarella Bozzo for technical advise and scientific support.

Conflict of interest

The authors declare no conflict of interest.

References

1. Dechant G, Molecular interactions between neurotrophin receptors. *Cell Tissue Res.* 2001; 305: 229-238
2. Chao MV and Bothwell M, Neurotrophins: to cleave or not to cleave. *Neuron.* 2002; 33: 9-12
3. Chen Y, Zeng J, Cen L, Chen Y, Wang X, Yao G, et al, Multiple roles of the p75 neurotrophin receptor in the nervous system. *J Intl Med Res.* 2009; 37:281-288
4. Barbacid M, The Trk family of neurotrophin receptors. *J. Neurobiol.* 1994; 25: 1386-1403
5. Lei L and Parada LF, Transcriptional regulation of Trk family neurotrophin receptors. *Cell. Mol. Life Sci.* 2007; 64: 522-532
6. Chapman BS and Kuntz ID, Modeled structure of the 75-kDa neurotrophin receptor. *Protein Sci.* 1995; 4: 1696-1707

7. Gentry JJ, Baker PA and Carter BD, The p75 neurotrophin receptor: multiple interactions and numerous functions. *Prog. Brain Res.* 2004; 146: 25-39
8. Blöchl A and Blöchl R, A cell-biological model of p75NTR signaling. *J. Neurochem.* 2007; 102: 289-305
9. Nykjaer A, Lee R, Teng KK, Jansen P, Madsen P, Nielsen MS et al, Sortilin is essential for proNGF-induced neuronal cell death. *Nature* 2004; 427: 843-848
10. Jones PH, Simons BD, Watt FM, Sic Transit Gloria: Farewell to the Epidermal Transit Amplifying Cell?. *Cell Stem Cell.* 2007; 1(4):371-81. Review.
11. Pincelli C, Sevigiani C, Manfredini R, Grande A, Fantini F, Bracci-Laudiero L, et al, Expression and function of nerve growth factor and nerve growth factor receptor on cultured keratinocytes. *J Invest Dermatol.* 1994; 103: 13-18
12. Marconi A, Terracina M, Fila C, Franchi J, Bonté F, Romagnoli G, et al, Expression and function of neurotrophins and their receptors in cultured human keratinocytes. *J. Invest. Dermatol.* 2003; 121: 1515-21 Erratum in: *J. Invest. Dermatol.* 2004; 123: 803
13. Pincelli C and Yaar M, Nerve growth factor: its significance in cutaneous biology. *J. Invest. Dermatol. Symp. Proc.* 1997; 2: 31-36
14. Yoon SO, Casaccia-Bonnel P, Carter B, Chao MV, Competitive signaling between TrkA and p75 nerve growth factor receptors determines cell survival. *J. Neurosci.* 1998; 18: 3273-3281

15. Nickoloff BJ, Qin JZ, Nestle FO, Immunopathogenesis of psoriasis. *Clin. Rev. Allergy Immunol.* 2007; 33: 45-56
16. Tschachler E, Psoriasis: the epidermal component. *Clin. Dermatol.* 2007; 25: 589-595
17. Heenen M and Simonart T, Apoptosis in psoriatic epidermis. *J. Cutan. Pathol.* 2008; 35: 346
18. Wrone-Smith T, Johnson T, Nelson B, Boise LH, Thompson CB, Núñez G, et al, Discordant expression of Bcl-x and Bcl-2 by keratinocytes in vitro and psoriatic keratinocytes in vivo. *Am. J. Pathol.* 1995; 146: 1079-1088
19. Di Marco E, Mathor M, Bondanza S, Cutuli N, Marchisio PC, Cancedda R, et al, Nerve growth factor binds to normal human keratinocytes through high and low affinity receptors and stimulates their growth by a novel autocrine loop. *J. Biol. Chem.* 1993; 268: 22838-22846
20. Yaar M, Zhai S, Pilch PF, Doyle SM, Eisenhauer PB, Fine RE, et al, Binding of beta-amyloid to the p75 neurotrophin receptor induces apoptosis. A possible mechanism for Alzheimer's disease. *J. Clin. Invest.* 1997; 100: 2333-2340
21. Salehi AH, Xanthoudakis S, Barker PA, NRAGE, a p75 neurotrophin receptor-interacting protein, induces caspase activation and cell death through a JNK-dependent mitochondrial pathway. *J. Biol. Chem.* 2002; 277: 48043-48050
22. Costantini C, Rossi F, Formaggio E, Bernardoni R, Cecconi D, Della-Bianca V, Characterization of the signaling pathway downstream p75

neurotrophin receptor involved in beta-amyloid peptide-dependent cell death.

J. Mol. Neurosci. 2005; 25: 141-156

23. Liu J, Minemoto Y, Lin A, c-Jun N-terminal protein kinase 1 (JNK1), but not JNK2, is essential for tumor necrosis factor alpha-induced c-Jun kinase activation and apoptosis. Mol. Cell Biol. 2004; 24: 10844-10856

24. Tiberio R, Marconi A, Fila C, Fumelli C, Pignatti M, Krajewski S, et al, Keratinocytes enriched for stem cells are protected from anoikis via an integrin signaling pathway in a Bcl-2 dependent manner. FEBS Lett. 2002; 524: 139-144

25. Fantini F, Magnoni C, Bracci-Laudiero L, Pincelli C TE, Nerve growth factor is increased in psoriatic skin. J. Invest. Dermatol. 1995; 105:854-855

26. Raychaudhuri SP, Raychaudhuri SK, Role of NGF and neurogenic inflammation in the pathogenesis of psoriasis. Prog. Brain. Res. 2004; 146:433-437

27. Botchkarev VA, Yaar M, Peters EM, Raychaudhuri SP, Botchkareva NV, Marconi A, et al, Neurotrophins in skin biology and pathology. J. Invest. Dermatol. 2006; 126: 1719-1727

28. Yiangou Y, Facer P, Sinicropi DV, Boucher TJ, Bennett DL, McMahon SB, et al, Molecular forms of NGF in human and rat neuropathic tissues: decreased NGF precursor-like immunoreactivity in human diabetic skin. J. Perpher. Nerv. System. 2002; 7: 190-197

29. Marconi A, Dallaglio K, Lotti R, Vaschieri C, Truzzi F, Fantini F, et al, Survivin identifies keratinocyte stem cells and is downregulated by anti- β 1 integrin during anoikis. *Stem Cells*. 2007; 25: 149-155
30. Lu B, Pang PT, Woo NH, The yin and yang of neurotrophin action. *Nat. Rev. Neurosci*. 2005; 6: 603-614
31. Hashimoto Y, Kaneko Y, Tsukamoto E, Frankowski H, Kouyama K, Kita Y, et al, Molecular characterization of neurohybrid cell death induced by Alzheimer's amyloid-beta peptides via p75NTR/PLAIDD. *J. Neurochem*. 2004; 90: 549-558
32. Perini G, Della-Bianca V, Politi V, Della Valle G, Dal-Pra I, Rossi F, et al, Role of p75 neurotrophin receptor in the neurotoxicity by beta-amyloid peptides and synergistic effect of inflammatory cytokines. *J. Exp. Med*. 2002; 195: 907-918 Erratum in: *J. Exp. Med*. 2002; 195: 1231
33. Teng HK, Teng KK, Lee R, Wright S, Tevar S, Almeida RD, et al, ProBDNF induces neuronal apoptosis via activation of a receptor complex of p75NTR and sortilin. *J. Neurosci*. 2005; 25: 5455-5463
34. Kim JY, Roh JK, Lee SK, Chung CK, Neurotrophin receptor immunoreactivity in severe cerebral cortical dysplasia. *Epilepsia*. 2002; 43 Suppl 5: 220-226
35. Guate JL, Fernández N, Lanzas JM, Escaf S, Vega JA, Expression of p75(LNGFR) and Trk neurotrophin receptors in normal and neoplastic human prostate. *B. J. U. Int*. 1999; 84: 495-502

36. Murray E, Malley SE, Qiao LY, Hu VY, Vizzard MA, Cyclophosphamide induced cystitis alters neurotrophin and receptor tyrosine kinase expression in pelvic ganglia and bladder. *J. Urol.* 2004; 172: 2434-2439
37. Quann EJ, Khwaja F, Zavitz KH, Djakiew D, The aryl propionic acid R-flurbiprofen selectively induces p75NTR-dependent decreased survival of prostate tumor cells. *Cancer Res.* 2007; 67: 3254-3262
38. Costantini C, Weindruch R, Della Valle G, Puglielli L, A TrkA-to-p75NTR molecular switch activates amyloid beta-peptide generation during aging. *Biochem. J.* 2005; 391: 59-67
39. Castelijns FA, Gerritsen MJ, van Erp PE, van de Kerkhof PC, Cell-kinetic evidence for increased recruitment of cycling epidermal cells in psoriasis: the ratio of histone and Ki-67 antigen expression is constant. *Dermatology.* 2000; 201: 105-110
40. Grabe N and Neuber K, Simulating psoriasis by altering transit amplifying cells. *Bioinformatics.* 2007; 23: 1309-131
41. Franssen ME, Zeeuwen PL, Vierwinden G, van de Kerkhof PC, Schalkwijk J, van Erp PE, Phenotypical and functional differences in germinative subpopulations derived from normal and psoriatic epidermis. *J. Invest. Dermatol.* 2005; 124: 373-383
42. McKenzie RC, Oda Y, Szepietowski JC, Behne MJ, Mauro T, Defective cyclic guanosine monophosphate-gated calcium channels and the pathogenesis of psoriasis. *Acta Derm. Venereol.* 2003; 83: 414-418

Legends




Figure 1 p75NTR modulates apoptosis in normal human keratinocytes

in vitro. (a) Confluent keratinocyte cultures, seeded in a 96-well tissue culture plate, were treated with 40 μ M β -amyloid or diluent alone. MTT was performed 24 hrs later. Results are expressed as the mean \pm SD of three different experiments. Student's T-test was used for comparison of the means. (b) Confluent cells were treated as described before, trypsinized at 24 hrs and

stained with PI. Cells in subG1 peak were analyzed by flow cytometry. **(c)** Keratinocytes cultures, seeded on chamber slides, were provided with 40 μ M β -amyloid or diluent for 24 hrs. Cells were fixed, double stained in situ with anti-p75NTR and anti-active caspase-3 antibodies, and analyzed by a Confocal Scanning Laser Microscopy. **(d)** Confluent keratinocyte cultures were treated with BDNF or NT-4 (100ng/ml). At 24, 48 and 72 hrs cells were fixed and stained in situ with TUNEL. Positive cells were counted as described in M&M. **(e)** Confluent keratinocytes were treated individually or in combination with 100ng/ml BDNF and 50 μ g/ml mouse anti-p75NTR neutralizing antibody. After 48 hrs, cells were stained in situ with TUNEL and positive cells were counted as described in M&M. **(f)** Keratinocytes were transiently infected with p75NTR-LNSN packaging cells or LNSN packaging cells as mock control. Both mock and transfected cells were treated with BDNF (100ng/ml) or β -amyloid (40 μ M). 48 hrs after treatment, protein extracts from keratinocytes were analyzed by Western blotting for p75NTR and caspase-3 expression. β -actin was used as internal control. **(h)** Relative intensity of bands on radiograms was quantified by laser scanner densitometry. Values of pro-caspase-3 and caspase-3 fragment are expressed as fold variations compared to mock-diluent keratinocytes.




Figure 2 p75NTR induces apoptosis via JNK and NF-κB (a) Keratinocytes were treated for 24h with β-amyloid (40μM) or diluent. Protein extracts were immunoprecipitated with NRAGE serum and immunoblotted with p75NTR antibody and with NRAGE serum as control. Cell lysates were also immunoprecipitated with p75NTR antibody, immunoblotted with NRAGE serum and with p75NTR antibody as control. (b) Confluent keratinocytes were treated for 5 min with BDNF (100ng/ml) and protein extracts were analyzed by Western blotting using anti-active JNK1 polyclonal antibody. PC12 cells treated with 0.5M sorbitol for 5 minutes were used as positive control. (c) Keratinocytes were infected with p75NTR and treated with with 100ng/ml BDNF or diluent alone for 12h. Nuclear protein extracts were analyzed by Gel mobility shift assay for NF-κB DNA binding activity.




Figure 3 p75NTR-induced apoptosis in TA cells (a) Cryosections of normal skin were acetone-fixed and stained with anti-p75NTR antibody. Fast red was used as chromogen for p75NTR. Cryosections were double stained with anti-p75NTR and anti-keratin 10 (CK10) antibodies or with anti-p75NTR and transglutaminase type I (TgaseI) antibodies. Fast blue was used as chromogen for p75NTR while carbazol for CK10 and TGaseI. (b) Normal human keratinocytes were obtained from neonatal foreskin and separated into three subpopulations on type IV collagen. A Real Time PCR was performed on RNA extracts of the three populations by using primers for p75NTR, as described in M&M. KSC were used as calibrator. Student's T test was performed between

samples and calibrator. **(c)** Protein extracts from the three populations were separated on 7% polyacrylamide gels and transferred onto nitrocellulose membrane. Membranes were immunoblotted with anti-p75NTR antibody. β -actin was used as internal control. **(d)** KSC and TA cells were treated with either BDNF or β -amyloid, stained with P.I., and analyzed by flow cytometry. **(e)** TA cells were transiently transfected with 50nM of p75NTR siRNA, as shown by Western blotting. **(f)** 48 hrs after transfection, both mock and transfected cells were treated with BDNF (100ng/ml) or β -amyloid (40 μ M). SubG1 peak analysis was performed 24 hrs later.




Figure 4 ProNGF mediates apoptosis in human keratinocytes through p75NTR and sortilin. **(a)** Normal human keratinocytes were obtained from neonatal foreskin. A Real Time PCR was performed on RNA extracts by using

primers for p75NTR and sortilin, as described in M&M. p75NTR expression levels were used as calibrator. **(b)** Keratinocyte cultures were trypsinized and stained with mouse monoclonal anti-p75NTR and polyclonal anti-sortilin antibody. Cells were analyzed by flow cytometry. **(c)** Normal human keratinocytes were separated into three subpopulations on type IV collagen. Real Time PCR was performed on RNA extracts of the three populations by using primers for sortilin, as described in M&M. KSC were used as calibrator. **(d)** Protein extracts from the three populations were separated on 7% polyacrylamide gels and transferred onto nitrocellulose membrane. Membranes were immunoblotted with anti-sortilin antibody. **(e)** Keratinocyte cultures, seeded in a 96-well tissue culture plate, were treated with 0.1, 1, 10, 20 ng/ml of proNGF. MTT was performed 72 hrs later. **(f)** Keratinocyte confluent cultures, seeded in a 96-wells tissue culture plate, were treated with 10ng/ml proNGF or 200nM k252a alone, or in combination. MTT was performed 72 hrs later. **(g)** Confluent keratinocytes, seeded in a tissue culture slide-flask, were treated as described above. At 72 hrs cells were fixed and stained in situ with TUNEL. Positive cells were counted as described in M&M. **(h)** HaCaT cells were transfected with 2 μ g of pcDNA3-BimEL (positive control) in combination with 2 μ g of pcDNA3-p75NTR and pcDNA3-sortilin. Transfection was controlled by Western blotting using anti-sortilin and anti-p75NTR antibody. **(i)** After 24 hrs, cells were treated with 10ng/ml proNGF for 24 hrs. Then, HaCaT cells were lysed and caspase activity was assessed using DEVD-AFC substrate. Data represent the mean +/- SE of triplicate determinations.




Figure 5 NT receptor expression in psoriasis. (a) Cryosections of healthy, non lesional and lesional psoriatic skin were acetone-fixed and stained with anti-p75NTR antibody. Fast red was used as chromogen. (b) mRNA extracts from healthy and psoriatic epidermis were analyzed by Northern blot. DIG-labeled probes specific for p75NTR or actin were used. (c) Real Time PCR was performed on healthy and psoriatic epidermal mRNA, as described in M&M. Healthy mRNA was used as calibrator. (d) Protein extracts from healthy, lesional and non lesional psoriatic epidermis were separated on 7% polyacrylamide gels and transferred onto nitrocellulose membrane. Membranes were immunoblotted with anti-p75NTR, anti-sortilin, anti-TrkA, anti-TrkB, anti-

TrkC and β -actin antibodies. **(e)** Keratinocyte extracts from healthy, lesional and non lesional psoriatic skin were stained with mouse monoclonal anti-p75NTR antibody. Cells, obtained from three different patients, were analyzed by flow cytometry.




Figure 6 Reduced levels of p75NTR in TA cells are responsible for resistance to apoptosis in psoriatic keratinocytes **(a)** Normal and psoriatic human keratinocytes were obtained from skin biopsies and separated into three subpopulations on type IV collagen. Keratinocytes were trypsinized and stained with mouse monoclonal anti-p75NTR and analyzed by flow cytometry. **(b)** After 48 hrs, both normal and psoriatic KSC, TA and PM cells were stained in situ with PI and analyzed by Confocal Scanning Laser Microscopy. Data represent the mean \pm SE of triplicate determinations. **(c)** Keratinocyte cultures obtained from normal and psoriatic biopsies were treated with 40 μ M β -amyloid or 100ng/ml BDNF. MTT was performed 48 hrs later. **(d)**

Keratinocytes were treated as in (c) and subG1 peak was evaluated by flow cytometry. Data represent the mean +/- SE from three independent experiments.




Figure 7 p75NTR infection restores apoptosis susceptibility in psoriatic keratinocytes (a) Psoriatic keratinocytes were double infected with mock or p75NTR retroviral vector and infection controlled by Western blotting. (b) 48 hours after infection, cells were treated with 40μM β-amyloid or 100ng/ml BDNF or diluent. MTT was performed 48 hours later. Data represent the mean +/- SE of triplicate determinations. (c) Mock and p75NTR infected keratinocytes were provided 100ng/ml BDNF or diluent and analyzed by flow

cytometry 48 hrs later. Data represent the mean +/- SE of triplicate determinations.