



Journal of Experimental Biology and Agricultural Sciences

<http://www.jebas.org>

ISSN No. 2320 – 8694

COMPARATIVE STUDY OF HAEMATOLOGICAL PARAMETERS IN FISHES FROM THE NATURAL HABITAT, MADURAI REGION

Malathi N*, Rino T, Jaya Supa Sooriya J, Shalini M

Fatima College, Mary Land, Madurai, Tamil Nadu, India

Received – July 18, 2020; Revision – September 17, 2020; Accepted – December 08, 2020

Available Online – March 25, 2021

DOI: [http://dx.doi.org/10.18006/2021.9\(Spl-1-GCSGD_2020\).S176.S181](http://dx.doi.org/10.18006/2021.9(Spl-1-GCSGD_2020).S176.S181)

KEYWORDS

Water quality parameters

Haemoglobin

Labeo rohita

Channa striatus

ABSTRACT

The heamatological profile of two freshwater fishes *Labeo rohita* (Rohu) and *Channa striatus* (Murrel) with a relationship to the sex and the water quality under natural habitat was analyzed. The freshwater fishes were collected from the pond located in Thodaneri, Vadipatti taluk, Madurai district, Tamil Nadu, India. In heamatology studies, no significant variation was observed in Heamoglobin content and found almost the same in both the fish irrespective of the sex and also falls in the range of control value. In differential Leucocyte count (DLC) the following blood components Lymphocyte, Eosinophil, and Monocyte count were significantly higher than the control values and the values are found maximum in Rohu when compared to Murrel. RBC (Erythrocyte) count was almost the same in both the fish and the value falls between the range of control value Packed Cell Volume (PCV) count was almost the same in both the fish and the value falls between the range of control value. The values of Hb and PCV were found higher in male fish as compared to female fish. The leucocyte count is also higher in the males when compared to the females of both Rohu and Murrel. These may be also influenced by physiological activity, feeding habits, infectious agents, water quality parameters, and the environmental stress on the organism.

* Corresponding author

E-mail: malathivishvakshenan@gmail.com ; malvish@rediffmail.com (N. Malathi)

Peer review under responsibility of Journal of Experimental Biology and Agricultural Sciences.

Production and Hosting by Horizon Publisher India [HPI]
(<http://www.horizonpublisherindia.in/>).
All rights reserved.

All the articles published by [Journal of Experimental Biology and Agricultural Sciences](#) are licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](#) Based on a work at www.jebas.org.



1 Introduction

The growing demands of urbanization cause a decrease in the amount of forest and, wetlands. This depleting green cover leads to variation in impervious-pervious which affect the quality of the storm water run-off, leading to poor watershed systems due to loads of pollutants carried by the storm water (Morisawa & La Flure, 1979). In addition to this, the pollutants that are released from the industries are also carried to the riverine system. The overall environment of the aqua system gets affected indefinitely based on the effects of the pollutants, and they directly or indirectly affect the organisms that habitat the wetlands. Fishes are considered valuable resources since they are an important source of food in the world. Fishes as a unit play an important role in maintaining the balance in the ecosystem.

Fishes in the pond are afflicted by the water quality parameters such as pH value, alkalinity, availability of oxygen, level of environmental stress, and hardness which eventually pose a great threat to fish health (Southamani et al., 2015). Rohu (*Labeo rohita*) is the predominant species used in carp polyculture systems. Rohu is a fast-growing species and thrives above 14 °C temperature. Spawning happens in the shallow water and the spawning season of Rohu generally coincides with the south-west monsoon. *Channa striatus* (Ng & Lim, 1990) generally lives in estuaries and can withstand the drought. It can breathe in an anaerobic way and can survive in the dry season (Qin & Fast, 1996; Pradhan et al., 2012). The Murrel involve in reproduction all-round the year (Ali, 1999), and suitable breeding habitat for this fish is the paddy field. Both these fishes are available in almost all seasons and hence chosen for the study.

The heamatology is considered an important indicator to oversee the physiological and pathological changes in the fishes. The total body weight of fish constitutes 1.3–7% of blood which exchanges gas between organism and environment. Studies of haematological parameters are based on health status, age, aquatic biotope, and fish species (Patriche et al., 2011; Kumar et al., 2019). It is greatly influenced by physiological stress, diseases, and the introduction of toxic pollutants in the aqua system. This study aimed to compare the haematological profiles of two freshwater fish in Rohu (*Labeo rohita*) and Murrel (*Channa striatus*) collected from the native environment.

2 Materials and Methods

The haematological profile of the two freshwater fishes *L. rohita* and *C. striatus* in relationship with the sex and the water quality in a natural habitat is analyzed in this work. Ten healthy and disease-free male and female of each *L. rohita* and *C. striatus* species were collected for sampling from the pond located near Thodaneri, Vadipatti Taluk, Madurai, Tamil Nadu, India subjected to haematological studies (Figure 1). Ten male and female fish of *L. rohita* and *C. striatus* served

as a control. The studies include physicochemical parameters of the pond water and heamatology profile of the two selected fishes. The parameters taken for comparison are haematology and sex discrimination factors.




Figure 1 Satellite Map showing the catchment area where fish were collected from Thodaneri pond, Vadipatti Taluk, Madurai district

2.1 Water quality analysis

Pond water from the catchment area was collected in a sterilized PET bottle by immersing it completely in the middle of the pond and screw-capped tightly under the water without lifting the water bottle and it was transported immediately to the laboratory for physicochemical examination of water parameters as described by the standard protocols (APHA, 1989). The sampling was done between 10 am to 2 pm. Water samples in triplicates were collected at different intervals of time for the study. On-site analysis was done for pH and turbidity using a pH meter and turbidity meter. Other physicochemical parameters like dissolved solids, pH, total hardness, and bound minerals were carried out within 24 hrs of sample collection. It was preserved at 4°C. The water quality was analyzed by methods given in APHA (1989) and Trivedy & Goel (1986).

2.2 Collection of Blood

Blood was collected from the punctured caudal vein with the aid of 20 G×1.5 syringes and collected blood samples in falcon tubes containing EDTA from both the sexes of Rohu and Murrel separately. The experiment was repeated twice with ten samples of males and ten samples of the female of each fish. In haematological profile, the differential Leucocyte counts was determined with the help of a Neubauer hemocytometer. Nelson & Morris (1989) method was used to estimate hemoglobin concentrations and hematocrit. Packed cell volume was estimated by adopting the procedure of Wintrobe (1934). Further, erythrocyte indices such as MCV, MCH, and MCHC were calculated as mentioned below:

$$\text{MCH} = \frac{\text{hemoglobin}}{\text{erythrocyte count}} \times 10$$

$$\text{MCV} = \frac{\text{PCV}}{\text{erythrocyte count}} \times 100$$

$$\text{MCHC} = \frac{\text{hemoglobin}}{\text{PCV}} \times 100$$

Sex was also determined for the hematological parameters.

2.3. Statistical analysis

Statistical analysis of hematological parameters in terms of Mean±SD in both sexes of fishes was employing ANOVA.

3 Results

The results are presented as comparative analyses between the species and sexes of Rohu (*Labeo rohita*) and Murrel (*Channa striatus*) the freshwater fish collected from the pond with that of the control.

In the physicochemical examination of water, total dissolved solids were very minimum (211mg/l) when compared to the acceptable limit, the electrical conductivity was 302µS/cm. The pH of the water was 7.10, which is falling under the recommended range.

In a chemical examination of water analysis through total hardness parameter was showing maximum (700 mg/l) the individual chemical components were found minimum when compared to an acceptable limit. Other physio-chemical parameters like calcium, magnesium, iron, ammonia, nitrate, chloride, fluoride, sulphate, and phosphates are below the recommended range (Table 1).

Table 1 Physico-chemical examination of pond water

S.No.	Physical examination	Acceptable limit	Pond Water (Sample)	Control water
1.	Colour	5	Clear & colourless	Clear & transparent
2.	Total Dissolved Solid (mg/l)	500	211	150
Chemical Examination				
3.	pH	6.5–8.5	7.10	6.70
4.	Total hardness (mg/l)	200	700	650
5.	Calcium (mg/l)	75	88	80
6.	Magnesium (mg/l)	30	35	33
7.	Iron (mg/l)	0.3	0	0.1
8.	Ammonia (mg/l)	0.5	0.5	0.1
9.	Nitrate (mg/l)	45	1	0.5
10.	Chloride (mg/l)	250	40	30
11.	Fluoride (mg/l)	1.0	0	0.5
12.	Sulphate (mg/l)	200	15	10
13.	Phosphate (mg/l)	0	0.5	0.1

Haematological profiles were compared between the sex of Rohu and Murrel (Table 2). No significant variation was observed in Hb content and found almost the same in both the fish (11±0.5) and also fell in the range of control value in Rohu female.

Mean corpuscular volume (MCV) were found significantly maximum (188±17) in Rohu male when compared with control range. Mean Corpuscular Hemoglobin (MCH) count was found significantly maximum (47±0.0) in Rohu males when compared to the control range.

In differential leucocytes count the following blood components Lymphocyte (87.0±3), Eosinophil (3.50±0.5) and Monocyte (1.75±1.5) count in Rohu male were significantly higher than the control values. Eosinophil (0.34–0.40) and Monocyte (0.14) and the values were found maximum in Rohu when compared to Murrel. Lymphocyte count was found maximum in both the sex of Rohu and Murrel when compared with control. RBC count was almost same for the both the fish and the value found minimum when compared to the range of control value. PCV count was almost the same in both the fish (40±1) and the value was found between the range of control value.

Table 2 Haematological Profiles of Rohu and Murrel Fish

S. No	Haematology parameters	Rohu		Murrel		Rohu (Control)		Murrel (Control)	
		Male	Female	Male	Female	Male	Female	Male	Female
1.	HB	11±0.5	11.5±1	12±1	11±0.5	11±1	11±0.5	12±1	11±0.5
2.	PCV	40±1	38.5±0.5	41±1	38±1	42±1	40±0.5	43±1	41±2
3.	RBC	2.8±0.1	2.4±0.1	3±0	2.0±2	4±0.1	3.8±0.1	3.9±0	3.7±2
4.	MCV	188±17	122±2	158±12	156±3	93±3	85±2	94±1	81±2
5.	MCH	47±0.0	33±0.2	35±0.3	34±0.2	31±1	29±0.2	33±0.1	30±1
6.	MCHC	30±0.3	32±0.1	31±0.4	30±1.5	37±2	32±0.1	39±1	35±2.5
Differential Leucocyte Count									
8.	Neutrophil	9.5±0.5	6.50±0.5	10±1	9±0.5	16±0.5	11±0.5	15±1	12±0.5
10.	Monocyte	1.75±1.5	1.00±0	1.00±1	1.00±1	0.13±0.05	0.1±0	0.13±0.01	0.11±0.1
11.	Eosinophil	3.50±0.5	2.00±0.5	1.5±0.5	1.75±0.5	0.3±0.1	0.2±0.2	0.4±0.1	0.1±0.2
12.	Lymphocyte	87±2.5	82.5±3	86±2	81±2.5	79±2	78±1	78±1	77±1.5

The given values are Mean± SD

4 Discussion

Haematological results are presented as comparative analyses between Rohu (*L. rohita*) and Murrel (*C. striatus*) freshwater fish. The physico-chemical parameters taken for the study of pond water were above the control range except for iron content. The pH of the water is influenced by the dissolved minerals in the aquatic system, the pH of water influences the solubility of substances thus the presence of these substances influence the aquatic system. Furthermore, pH is additionally emphatically corresponded with electrical conductance and absolute alkalinity (Sahiti et al., 2018; Sarkar et al., 2020). The pH in the present study is alkaline and is also under the optimal range. Huq & Alam (2005) stated that electrical conductivity typically is shown for the dissolved concentration of free radicals in water. The total dissolved solids also influence the electrical conductivity of the water (McNeely et al., 1979).

The amount of chloride was below the optimal level. High Chloride level is an indicator of polluted water (Munawar, 1970). Phosphate was found 0.5 mg/l in the pond water. Phosphates are taken up by plants and are not found in high amounts in water. Phosphates in high quantities indicate pollution in water (WHO, 1993). Presently nitrate level was found to be influenced by the nitrification and denitrification process which is carried out by micro-organisms (Trivedy & Goel, 1986). The sulphates in water determine their use for public and industrial purposes, the sulphate content was 15 mg/l.

According to Svobodova (1994), stream hydrology and capacity are reliant on five significant factors viz., atmosphere, geography, soils, land use, and vegetation. These factors straightforwardly influence the elements of release and silt load, which thus affect the hydrology and morphology of the stream. Additionally, the pesticide and other chemicals leaches out and, in the end, influences the water quality, which thus impacts the miniature atmosphere of the lake and the life form which lives in it.

According to Southamani et al. (2015) the climate of the environment affects the hematology of fish. The specific distinction is seen in the hematological profile of both the species and also concerning the sex. Changes in the environment can induce changes in the hematological profile of fishes as they are very sensitive to their environment (Ahmed et al., 2020).

The values of Hb and PCV have been seen maximum in male fish which is due to the physiologically activeness of male fish. The variation in the erythropoietin is the indicator of higher hematological values in male fish influenced by the water quality of the pond. The leucocyte count is also higher in the males when compared to the females of both Rohu and Murrel. Leukocyte levels in the blood may also fluctuate according to environmental quality, nutritional status, and presence of infectious agents (Sunomonu & Oyelola, 2008).

The authors found that there is an increase in PCV level while decreasing in the erythrocytes due to habitat. Hypoxia results in swelling of RBC (Singh, 1995). Adrenaline stimulates membrane bound Na/K ATPase which is responsible for methemoglobin (Svobodova et al., 2008). This further leads to a decrease in the concentration of hemoglobin and an increase in cell volume. Increased accumulation of CO₂ in the blood causes a decrease in Hb indicating poor oxygen transport. The unexpected occurrence of eosinophils in the blood may be the indicator of homeophysiological demands which is due to the internal and ecophysiological conditions (Qin & Fast, 1996; Tripathi et al., 2003)

Conclusion

The present work involves the comparative study of two freshwater fish Murrel and Rohu from the catchment area (Pond) subjected to haematological profiling of their blood samples. The results of the study showed almost similar data shared for most of the blood parameters among two species irrespective of their sex. On contrary, the values of Hb and PCV have been found maximum in male fish. The leucocyte count is also higher in the males when compared to the females of both Rohu and Murrel. These may be also influenced by the ecological niche and microclimate of the pond, physiological activity, feeding habit, infectious agents, water quality parameters, agricultural leachates, and the environmental stress on the organism. It should be once again evaluated more extensively as they possess many useful genes for fish sustained development.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Ali AB (1999) Aspects of the reproductive biology of female snakehead (*Channa striatus Bloch*) obtained from irrigated rice argro ecosystem, Malaysia. *Hydrobiologia* 411:71-79.
- Ahmed I, Reshi QM, Fazio F (2020) The influence of the endogenous and exogenous factors on hematological parameters in different fish species: a review. *Aquaculture International* 28: 869–899 (2020).
- APHA (1989) Standard Methods for the Examination of Water and Wastewater. 17th edition, American Public Health Association, Washington D.C., 1, Pp.268.
- Huq SMI, Alam MD (2005) A Handbook on Analyses of Soil, Plant, and Water. BACER-DU, University of Dhaka, Dhaka, 1-246.
- Kumar A, Kumar SA, Rakesh B, Singh IJ (2019) Certain hematological and biochemical changes in blood of rohu (*Labeo rohita*) in relation to sex, reproductive status and environmental factors. *Journal of Entomology and Zoology Studies* 7(3): 1484-1490
- McNeely RN, Neimanis VP, Dweyer L (1979) Water Quality Source Book: A Guide to Water Quality Parameters. Inland Waters Directorate, Water Quality Branch Ottawa, Canada, Pp: 88
- Morisawa M, LaFlure E (1979) Hydraulic geometry, stream equilibrium and urbanization. In: Rhodes DD, Williams GP (Eds.) Adjustments of the fluvial systems—Proceedings of the 10th annual Geomorphology Symposium Series, Binghamton, NY.
- Munawar M (1970) A limnological study of fresh water ponds of Hyderabad, India. 1-The biotope. *Hydrobiologia* 35: 127-162.
- Nelson DA, Morris MW (1989) Basic methodology. Hematology and coagulation, part IV. In: Nelson DA, Henry JB (Eds.), Clinical Diagnosis and Management by Laboratory Methods, 17th ed. Saunder Company, Philadelphia, USA, pp. 578– 625 (Chapter 27)
- Ng PKL, Lim KKP (1990) Snakeheads (Pisces: Channidae): natural history, biology and economic importance. Essays in Zoology. Papers Commemorating the 40th Anniversary of the Department of Zoology, National University of Singapore 1990:127-152.
- Patriche T, Patriche N, Bocioc E, Coada MT (2011). Serum biochemical parameters of farmed carp (*Cyprinus carpio*). *Aquaculture, Aquarium, Conservation & Legislation. International Journal of the Bioflux Society* 4(2): 137–140.
- Pradhan CS, Patra K, Sarkar B, Pal A (2012) Seasonal changes in hematological parameter of *Catla catla* (Hamilton 1822). *Comparative Clinical Pathology* 21:1473-1481. 29.
- Qin J, Fast A (1996). Size and feed dependent cannibalism with juvenile snakehead *Channa striatus*. *Aquaculture* 144: 313-320.
- Sahiti H, Bislimi K, Dalo E, Murati K (2018) Effect of water quality in hematological and biochemical parameters in blood of common carp (*Cyprinus carpio*) in two lakes of Kosovo. *International Journal of Natural and Engineering Science* 3(3):323–332
- Sarkar R, Ghosh AR, Mondal NK (2020) Comparative study on physicochemical status and diversity of macrophytes and zooplanktons of two urban ponds of Chandannagar, WB, India. *Applied Water Science* 10: 63.
- Singh M (1995) Haematological responses in a fresh water teleost, *Channa punctatus* to experimental copper and Cr poisoning. *Journal of Environmental Biology* 16:339-341.
- Southamani C, Shanthi G, Deivasigaman M (2015) Heamatological response in three Indian major carps in relation to supplementary

- feeding. *International Journal of Fisheries and Aquatic studies* 3(2): 287–294.
- Sunomonu TO, Oyelola BO (2008) Haematological response of African catfish (*Clarias gariepinus*) and rat to crude oil exposure. *International Journal of Haematology* 4:74-79.
- Svobodova H, Kroupova H, Modra H, Flajshans M, Randak T, Savina LV (2008) Haematological profile of common carp spawners of various breeds. *Journal of applied Ichthyology* 24:55- 59.
- Svobodova ZB (1994) The effects of pollutants on selected haematological and biochemical parameters in fish. In: Müller R Lloyd R (Eds.), *Sub lethal and chronic effects of pollutants on, freshwater*
- fish, Fishing new books, London.
- Tripathi PK, Srivastava VK, Singh A (2003) Toxic Effects of Dimethoate (Organophosphate) on Metabolism and Enzyme System of Freshwater Teleost Fish *Channa punctatus*. *Asian Fisheries Science* 16: 349–359.
- Trivedy RK, Goel PK (1986) Chemical and Biological method for water pollution studies. *Environmental publication* 6: 10-12.
- Winthrobe MM (1933) Variations in the Size and Hemoglobin Content of Erythrocytes in the Blood of Various Vertebrates. *Folia Haematologie*, 51, 32-49.