

Research Article

Bacterial profile of blood stream infections and their antibiograms

Meenakshi Kante*, Muni Lakshmi P, Sreenivasulu Reddy P

Department of Microbiology, Narayana Medical College, Nellore, Andhra Pradesh, India

Received: 20 January 2015

Revised: 23 January 2015

Accepted: 8 February 2015

***Correspondence:**

Dr. Meenakshi Kante,

E-mail: meena.apple86@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Blood culture provides essential information for the evaluation of a variety of diseases like endocarditis, pneumonia, and pyrexia of unknown origin particularly, in patients with suspected sepsis. Septicaemia is one of the leading causes of neonatal mortality along with perinatal hypoxia. Objectives: To study the bacterial flora of blood stream infections in adults & pediatric patients including neonates. To study the risk factors and duration of incubation for obtaining positive cultures. To study their antibiotic resistant pattern and compare results with those of other investigators in this field.

Methods: This prospective study was conducted in the department of microbiology, Narayana medical college and hospital, Nellore over a period of six months, from March 2014 to September 2014. During the six months period, 200 samples received from various departments were processed and relevant findings were noted.

Results: Out of 200 blood samples growth was obtained in 34 samples (17%). Among them Gram positive bacteria were 11 (5.5%). Gram negative bacilli were 23 (11.6%). Among the gram positive bacteria maximum resistance was seen with methicillin and imipenem. No resistance was seen with vancomycin and erythromycin. Most of gram negative bacilli were multidrug resistant. Maximum resistance was seen with piperacillin, amikacin, ceftizoxime and amoxycyclav.

Conclusion: Early diagnosis & appropriate treatment of bacterial infections can make difference between life & death. It would reduce mortality from septicaemia, reduce time & improve patient management.

Keywords: Bacteria, Blood cultures, Sepsis, Antibiotics

INTRODUCTION

Microorganisms present in the circulating blood whether continuously or intermittently are threat to every organ in the body.¹ Approximately 200000 cases of bacteraemia & fungemia occur annually with mortality rates ranging from 20-50%.² The detection of microorganisms in a patient's blood has great diagnostic & prognostic significance.³ Blood culture provides essential information for the evaluation of a variety of diseases like endocarditis, pneumonia, and pyrexia of unknown origin particularly, in patients with suspected sepsis.

Septicaemia is one of the leading causes of neonatal mortality along with perinatal hypoxia.⁴

Many infections in neonatal & pediatric age group can only be established on the basis of etiological agent recovered from blood. Neonatal septicaemia refers to systemic infection in the new born confirmed by a positive blood culture.⁵ It remains a major cause of morbidity & mortality among newborn especially in developing countries, where its incidence is higher than in the developed world.⁶ Apart from the relative immaturity of the immune system of the neonate risk factors predisposing to neonatal septicaemia in

developing countries include prolonged rupture of membranes, prematurity, birth asphyxia, length of time spent in hospital, invasive procedure, delivery outside a hospital, material used in cutting & dressing the cord & maternal infections during pregnancy.⁷ A positive blood culture does not necessarily confirm infection, since contamination of blood can occur. The recovery of organisms traditionally considered as pathogens pose no problems in interpretation.⁸

However, recovery of organisms such as coagulase negative staphylococci (CONS), *Corynebacterium* or *Candida* spp. is often difficult to interpret.⁹ Additional information like the density of bacteraemia, number of positive cultures, duration of incubation of the broth to obtain a positive culture, presence of risk factors or an underlying disease, is required in order to determine whether infection is truly present.¹⁰ Since early 1950's there is striking increase in incidence of bacteraemia caused by members of Enterobacteriaceae & non fermenting gram negative rods.¹¹ *Escherichia coli* which was reported to be common in the past is being replaced by other multidrug resistant bacteria like *Klebsiella*, *enterobacter*, *Salmonella*, *Citrobacter*, *Pseudomonas*, *Acinetobacter* etc.¹²

One key determinant in the ultimate outcome of patients with sepsis is institution of early & appropriate antimicrobial therapy.¹³ It is a common practice to institute early empirical therapy with broad spectrum antibiotics in patients presenting with clinical features suggestive of septicaemia or bacteraemia.¹⁴ Given the severity of septicaemia, such empirical therapy may be justified, but the specific therapy based on the antibiogram of the isolate will definitely improve the therapeutic outcome.¹⁵ Sometimes even after receiving antibiotic susceptibility report, physicians may prefer to maintain the original regimen in setting of clinical improvement.¹⁶ There is conflicting information in how much attention the physician should pay to the cultures & antibiotic susceptibility reports.¹⁷ The infection caused by MDR organism is more likely to prolong the hospital stay, increase the risk of death & require treatment with more expensive antibiotics.¹⁸ Early diagnosis & appropriate treatment of blood stream infections can make the difference between life & death. It would reduce mortality from septicaemia, reduce turnaround time & improve patient management.¹⁹

The present study was undertaken to determine the bacterial flora of the blood stream infections in adults & pediatric patients including neonates & their antibiotic resistant pattern.

METHODS

This prospective study was conducted in the department of microbiology, Narayana medical college and hospital, Nellore over a period of six months, from March 2014 to September 2014. During the six months period, 200

samples received from various departments were processed and relevant findings were noted. Patients presented with prolonged fever or clinical impression of septicaemia/bacteraemia. Patients having prolonged fever in the post-operative period, despite antibiotic coverage were included in the study. Detailed history was taken to identify the possible risk factors. History of antibiotic usage empirically either before or after admission was also obtained. Data was collected on a pre-designed proforma.

Collection and processing of specimen

Blood samples for culture were collected following strict aseptic precautions. If empirical antibiotics were already started, the collection was timed before the next dose of antibiotic was due or about half-an hour before the predicted peak of temperature. A second set was also collected in all patients about an hour later from a different venipuncture site. Three samples were collected in cases of suspected or sonographically diagnosed congenital heart disease. About 1 ml of blood in case of neonates and about 5 ml in case of children was collected in each set. Immediately after collection, the blood was inoculated into BHI broth without switching needles. The bottle containing 10 ml BHI broth was used in case of neonates and 5 ml were used for children and adults to allow 1:10 dilution. The culture bottles were incubated at 37°C aerobically. After overnight incubation, the samples were subcultured onto blood agar, Mac Conkey's agar, and chocolate agar. If no growth observed on plates by the next day, subcultures were again repeated from the broth on day 3, day 4 and finally on day 7. Antibiotic susceptibility tests were performed according to the standard methods.

Procedure

Heat fixed smear is flooded with crystal violet and allowed to act for 30 seconds. The stain is poured off and is washed with iodine solution. Smear is covered with fresh iodine solution & allowed to act for 30 seconds. Iodine solution is drained off and smear is washed freely with water. The smear is flooded with iodine acetone mixture and is allowed to act for 30 seconds. Smear is washed with water & counter stained with carbol fuchsin and is allowed to act for 30 seconds. Smear is again washed with water, dried between folds of filter paper & observed under oil immersion objective.

Culture media

The media used in blood culture bottles are multipurpose & nutritionally enriched. tryptic (or) trypticase soy, supplemented peptone, BHI, Columbia can agar & brucella broths are commonly used. Most commonly used blood culture media contain the anticoagulant sodium polyethanol sulfonate.

- 1) Biphasic media: Brain heart infusion agar and broth: A biphasic (Castenada's) medium is one that combines on agar slope with a broth medium. The blood broth is allowed to run over the slope by tipping the bottle at regular intervals. Microbial activity can be seen by growth on the slope. This avoids the need to subculture from bottle & therefore reduces risk of contaminating the culture.
- 2) Nutrient agar: Sterilized by autoclaving at 15 lbs pressure (121°C) for 15 minutes & poured into sterile petridishes.
- 3) Blood agar: Nutrient agar + 5% sheep blood at 50°C. Final pH - 7.4.
- 4) MacConkey agar: Sterilized by autoclaving at 15 lbs pressure (121°C) for 15 minutes & poured into plates.

General procedure

The inoculated blood culture media should be incubated at 37°C for 7 days. If any growth appears during incubation period, it should be proceed with gram stain. Then based on gram stain smear, it should be subcultured in corresponding media. The further tests such as biochemicals should be performed.

Biochemical reactions

A. Indole test

The test demonstrates the ability of certain bacteria to decompose the amino acid tryptophan to indole, which accumulates in the medium. Medium: The pH was adjusted to 7.4. The medium was dispensed and sterilized by autoclaving at 121°C for 15 minutes.

Kovac's reagent: Aldehyde was dissolved in alcohol layer and slowly the acid was added. The reagent was prepared in small quantities and stored in refrigerator. The medium was inoculated and incubated for 24 hours at 37°C. 0.5 ml of Kovac's reagent was added and shaken gently before use. Interpretation: A red colour in alcohol layer indicates a positive reaction. Yellow colour indicates negative reaction.

B. Methyl red test: (Mackie & MC. Cartney, 14th Edition)

Principle: This test is employed to detect the production of sufficient acid during fermentation of glucose and maintenance of pH below 4.5 The peptone and phosphate were dissolved; the pH was adjusted to 7.6, filtered, dispensed in 5 ml amounts and sterilized at 121°C for 15 minutes. The glucose solution was sterilized by filtration and 0.25 ml was added to each tube.

Methyl red indicator solution: The liquid medium is inoculated lightly & incubated at 37°C for 48 hours.

About 5 drops of methyl red reagent (0.04) was added, mixed & read.

C. Voges-Proskauer (Acetoin production) test: (Mackie & MC. Cartney)

Principle: The test depends on the production of acetoin (acetyl methyl carbinol) from pyruvic acid in media. In presence of alkali and atmospheric oxygen acetoin is oxidized to diacetyl which reacts with alpha-naphthol to give red colour.

Medium: Glucose phosphate peptone water. It is inoculated and incubated at 37°C for 48 hours. 1 ml of 40% KOH & 3 ml of 5 solution of alpha naphthol were added in absolute alcohol. Interpretation: A positive test was indicated by the development of pink colour within 2-5 min, becoming crimson in 30 min.

D. Citrate utilisation test: (Mackie & Mc.Cartney)

Principle: This is a test for ability of an organism to utilize citrate as the sole carbon and energy source for growth and an ammonium salt as the sole source of nitrogen.

Simmon's citrate medium: Organism is inoculated & incubated at 37°C. Interpretation: Blue colour and streak of growth-positive. Original green colour & no growth-negative.

E. Urease test (Mackie & Mc.Cartney, 14th Edition)

Principal: $(\text{NH}_2\text{CO-NH}_2 + \text{H}_2\text{O} \rightarrow 2\text{NH}_3 + \text{CO}_2)$ the glucose & urea solutions were sterilized by filtration. Basal media was prepared & pH is adjusted to 6.8-6.9 and sterilized by autoclaving at 121°C for 15 min & cooled to about 50°C glucose and urea were added and the medium was tubed as deep slopes. The entire slope was inoculated & incubated at 37°C. Interpretation: Urease positive cultures change the colour of indicator to pink.

F. Triple sugar iron agar (Mackie & Mc.Cartney)

Ingredients were heated to dissolve the solids the indicator solution mix was added and tubed. Sterilized at 10 lbs pressure (115°C) for 15 minutes and cooled to form slopes with deep 3cm butts. Heavy inoculum was streaked over the surface of slope and stabbed into the butt. And incubated aerobically at 37°C for 24 h.

Antibiotic sensitivity tests

Carried out by Kirby-Bauer disc diffusion method.

Sensitivity test medium: Muller-Hinton agar medium: The pH was adjusted to 7.4 & sterilized by autoclaving at 121°C for 15 min and poured into plates.

Inoculum: Isolated colonies were inoculated in peptone water & incubated at 35-37°C for 4-6 hours. The density was adjusted to approximately 10 CFU/ML by comparing its turbidity with that of 0.5 McFarland standard opacity tube.

Antibiotic discs: Antibiotic discs are erythromycin, piperacilin, penicillin, cefazolin, vancomycin, imipenem, ciprofloxacin, tobramycin, ceftazidime, amikacin, ceftizoxime. **Method:** A cotton swab was dipped into inoculums. The Muller-Hinton agar plate was inoculated by streaking the swab three times over the entire agar surface. Allowed 3-5 minutes before applying antibiotic discs, using sterile forceps. **Interpretation:** By measuring the diameter of zone of inhibition.

RESULTS

A total of 200 biological specimens were sent for culture to the microbiology lab during the period march 2014 to september 2014. There were 200 blood culture samples, of which 34 (17%) were identified as culture positive samples. Blood culture positive in hospitalised patients and out patients, out of 200 cases studied maximum blood samples 150 (75%) were received from ICU & 24 (12%) from pediatric ward with P.U.O, 22 (11%) from cardiology ward with the history of infective endocarditis & 4 samples from surgery ward with history of wound infection (2%) (Table 1).

Table 1: Studied maximum blood samples numbers of positive, negative and percentages.

	Hospitalised patients	%	Positive culture	%
Pediatrics	24	12%	2	8.33%
Cardiology	22	11%	8	36.3%
Male ICU	84	42%	14	16.6%
Female ICU	66	34%	6	8.82%
Surgery ward	4	2%	4	2%

Out of 200 cases 47 (23.5%) were below the age of 20 years, among them 12 (6%) were neonates, 55 (28%) were between the age of 20-40 years. 48 (24%) were in the age group 41-60 years. No. of cases above 60 years were 50 (25%) (Figure 1).




Figure 1: Age group wise No. of patients in percentages.

Positive cases were predominant in males 21 (61.7%) than in females 13 (38.2%) (Figure 2).




Figure 2: Sex wise distribution.

Out of 200 blood samples, growth was obtained in 34 samples, among them 15 isolates obtained after 48 hours incubation, 6 isolates after 72 hours & 2 isolates after 7 days of incubation (Table 2).

Table 2: Various isolates in relation to the duration of incubation of BHI broth before plating.

Time of inoculation	24 hours	48 hours	72 hours	7 days	Total
Klebsiella	3	1	-	-	4
Acinetobacter	3	1	2	-	6
Pseudomonas	1	8	2	2	13
Staph. aureus	2	5	2	-	9
CONS	2	-	-	-	2
Total	11 (32%)	15	6	2	34

In the present study risk factors were identified in 25 cases out of which 11 (44%) had positive cultures and the following bacteria Pseudomonas, Staphylococcus aureus, Acinetobacter and Klebsiella were isolated. CONS were isolated in 2 cases but repeat isolation confirmed only one. This isolates were from a patient with post-operative wound infection & peritonitis (Table 3).

Table 3: Various risk factors identified & the spectrum of isolates obtained.

Risk factors	No. of patients	No. of isolates	Microorganisms isolated
Endocarditis	6	4	Pseudomonas
Valvular heart disease	4	2	Staph. aureus
Gastroenteritis	6	2	Acinetobacter
UTI	2	1	Klebsiella
Bronchitis pneumonia	4	0	-
Post-operative wound infection	2	1	CONS-1
Peritonitis	1	1	CONS-1
Total	25	11	

Out of 200 blood samples growth was obtained in 34 samples (17%). Among them gram positive bacteria were 11 (5.5%). Gram negative bacilli were 23 (11.6%) The most commonly isolated gram-positive bacteria were Staphylococcus aureus in 9 (4.5%) and coagulase negative Staphylococci in 2 (1%) blood cultures. The most gram-negative bacteria found in positive cultures were Pseudomonas aeruginosa in 13 (6.6%), Acinetobacter in 6 (3%), and Klebsiella pneumoniae in 4 (2%) (Figure 3).




Figure 3: Distribution of bacterial pathogens.

Study area (Gram +ve - Staph. Aureus, CONS) (Gram -ve Pseudomonas, Klebsiella, Acinetobacter)

The antibiotic resistance patterns of gram-positive organisms were obtained from the laboratory reports. Staphylococcus aureus was maximum resistant, seen with Methicillin (100%) followed by imipenem (75%), cefazolin (50%), piperacillin (50%) and penicillin (25%) (Figure 4).




Figure 4: Antibiotic resistant pattern of gram positive bacteria.

Among gram-negative organisms, the resistance patterns were as follows: Pseudomonas was highly resistant to cefepime (90%), ceftizoxime (80%), amikacin (70%), amoxycylav (70%), imipenem (50%), ceftazidime (20%), netillin (20%), ciprofloxacin (20%), piperacillin (10%) and tobramycin (10%). Klebsiella pneumoniae was highly resistant to ceftazidime (100%), and ceftizoxime (100%), followed by piperacillin (90%), amikacin (90%)

and cefepime (90%), ciprofloxacin (40%), tobramycin (20%), amoxycylav (20%) and netillin (20%). Acinetobacter was highly resistant to tobramycin (100%), amikacin (100%), ceftizoxime (100%) and amoxycylav (100%) followed by imipenem (90%), cefepime (71.7%), tobramycin (66.7%), ceftazidime (40%) (Table 4).

Table 4: Antibiotic resistance pattern of gram negative bacteria.

Antibiotics	Pseudomonas N=10	Klebsiella N=5	Acinetobacter N=3
Ciprofloxacin	R-20	R-40	R-0
Tobramycin	R-10	R-20	R-66.7
Piperacillin	R-10	R-90	R-100
Ceftazidime	R-20	R-100	R-40
Imipenem	R-50	R-40	R-90
Amikacin	R-70	R-90	R-100
Ceftizoxime	R-80	R-100	R-100
Cefepime	R-90	R-90	R-71.7
Amoxycylav	R-70	R-20	R-100
Netillin	R-20	R-20	R-0

DISCUSSION

Septicaemia is a clinical syndrome associated with considerable morbidity & mortality. The timely detection of bacteraemia can have profound influence on the final clinical outcome. One blood culture set is rarely sufficient to establish or rule out bacteraemia, and multiple cultures could maximize sensitivity. In our study, the total number of positive cultures was 34 (17%). Whereas Arora,¹ Khanal²⁰ and DS Murthy²¹ et al. have reported 20.02%, 44% and 24.3% respectively. In India, the variation might be due to the fact that most of the patients are given the antibiotics before they come to the tertiary care hospital & other reason is that in most of the cases self-medication is very common.²² In the present study, the highest rate of prevalence 27.5% was found of age group 21-40 years. In age group 61 years & above the prevalence rate was 25% and lowest rate of prevalence 6% was found in age group (0-1 year). Sepsis and septic shock occur at all ages but most often in elderly patients. At present, most sepsis episodes are observed in patients older than 60 years. Advanced age is a risk factor for acquiring nosocomial blood stream infection in the development of severe forms of sepsis. Neonates are vulnerable to infections because of their weak immunological barrier.²³ In our study males were predominantly effected 21 (61.7%) and females 13 (38.2%), it is in accordance with the study of other workers.

Most studies of septic shock report a male preponderance. The percentage of male patients varies from 52-60%. Although any localized infection can disseminate to the blood stream, the most common primary foci are intravascular devices.²⁴ In the respiratory tract, urinary tract & various intra-abdominal sites. In the

present study, risk factors were identified in only cases of which 11 (44%) had positive cultures. The CONS, previously considered as a contaminant, has been recognized increasingly as a cause of bacteraemia. The ascendance of this group of staphylococci has created increased interpretative difficulties for the clinician, since the great majority of CONS isolates continue to represent contamination rather than true bacteraemia. In the present study, CONS was initially isolated in 2 cases but repeat isolation confirmed only one. This isolate was from a patient with postoperative wound infection and peritonitis. The positive predictive value for the isolation of CONS in the present study is 12.5%.²⁴ In their study CONS was initially isolated in 4 cases but repeat isolation confirmed only one (25%). Isolation of *S. aureus* from blood usually signifies infection, but bacteraemia has been observed in only two cases in the present study, which were having underlying cardiac pathology. According to one study, up to 57% of cases where *S. aureus* was repeatedly isolated will have a cardiac pathology and all such patients with *S. aureus* bacteraemia should be thoroughly evaluated for the presence of any cardiac pathology as the cardiac vegetations serve as an important source of persistent *S. aureus* bacteraemia.²⁵ Prior empirical antibiotic therapy before collection of the samples for culture may result in negative blood cultures. In the present study empirical antibiotics were already started by the time of collection of samples for culture in 50 (25%) of the cases of these only 6 (12%) had positive cultures as compared to 26 (17.4%) positive cultures from 150 patients who didn't receive any antibiotics before collection of samples for blood culture. The duration of incubation of the broth to obtain a positive culture was observed to be more among the samples from the patients who were already on antibiotics by the time of collection. This was observed for two isolated of *Acinetobacter* spp. and 4 isolates of *Pseudomonas* so the growth was obtained on third and seventh day respectively. Though this was an interesting finding, the number was too small for a statistical evaluation for its significance. Among the positive cultures, we have observed that (32%) of the cultures were positive by first subculture itself (after 24 hours of incubation of BHI broth) 43.4% and 20% of the culture were positive by second subculture (After 48 hours) & third subculture after 72 hours respectively, while virtually no isolates were obtained later (subcultured on day 7).

In our study the incidence of gram negative bacilli was (11.5%) while 5.5% isolates were gram positive organisms. In our study *Staphylococcus aureus* was isolated in 4.5% of cases & CONS in (1%) of the cases. The CONS (16.5%) and *Staph. aureus* 14% in neonatal septicaemia. 9% of incidence of both CONS and *staph aureus* has been reported by Surinder et al. Amongst gram negative organisms. *Pseudomonas*, one of the important non-fermented was isolated in (6.5%) of cases where as it has been reported to be 7.62% & 5.9 in other studies. It might be because that most of the patients

studied were indoor patients where these non-fermenters are important pathogens, because of invasive procedures used both for diagnostic and treatment purposes.

Acinetobacter was isolated in 3% of the cases. *Acinetobacter* spp. is considered as contaminant but it can cause septicaemia as well. We isolated *Acinetobacter* spp. from a case who presented with gastroenteritis. *Klebsiella pneumoniae* isolation was 2% which is in contrast to other studies who have reported it to be 25.8%.²⁶ Difference in the incidence in different institutes is because of different set up and the type of antibiotics used in those hospitals. Amongst the gram positive organisms maximum resistance was seen with Methicillin (100%) followed by imipenem (75%), cefazoline (50%) and piperacillin (50%). Arora et al. has reported maximum resistance with erythromycin (69.67%). None of our strain showed resistance against vancomycin and teicoplanin and these drugs therefore can be effectively used if methicillin resistance is suspected during treatment.²⁴ Most of the gram negative bacilli were multidrug resistance. Other workers also have reported majority of gram negative isolates in their study as multidrug resistance. Maximum resistance was seen against ceftizoxime (100%), cefepime (90%), amikacin (90%) and piperacillin (10%) was shown by *Pseudomonas*, *Klebsiella* and *Acinetobacter* spp. respectively. Arora et al. have reported Ampicillin resistance (89.96%), cephalexin (68.7%) and piperacillin (51.71%). In this study we found quinolones to be effective in the treatment of multidrug resistant gram negative infections in patients including premature and extremely low birth weight infants. In case of gram negative isolates, which turned out to be the major pathogens ciprofloxacin and tobramycin are good alternatives and they will also provide some economical relief to the patient.

CONCLUSION

Septicaemia is an important cause of morbidity and mortality. The causative agents of septicaemia are acquired in hospital. The prospective study conducted showed both gram positive and gram negative bacteria were responsible for bloodstream infections. Most of the strains were multidrug resistant. None of our strains showed resistance against vancomycin or teicoplanin and these drugs therefore can be effectively used if methicillin resistance is suspected during treatment. In case of gram negative isolates, which turned out to be major pathogens, ciprofloxacin and amikacin are good alternatives and they will also provide some economical relief to the patient. Strict infection control in neonatal units, hand washing combined with judicious policy for antibiotic therapy are the main solution to this problem. Positive blood culture is the gold standard in the diagnosis of neonatal sepsis. A functioning infection surveillance unit is a necessity in the battle against infections in the newborn unit. Provision of running water at all times and aseptic technique in the delivery at

the maternity ward and handling of babies in the neonatal unit will help in reducing acquisition of infection by inborn neonates. Early diagnosis & appropriate treatment of bacterial infections can make difference between life & death. It would reduce mortality from septicaemia, reduce time & improve patient management.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: Not required

REFERENCES

- Arora Usha, Pushpa Devi. Bacterial profile of blood stream infections and antibiotic resistance pattern of isolates. *JK Sci.* 2007;9(4):186-90.
- Bhatta Dharm Raj, Abhishek Gaur, H. S. Supram. Bacteriological profile of blood stream infections among febrile patients attending a tertiary care centre of Western Nepal. *Asian J Med Sci.* 2013;4(3):92-8.
- Weinstein MP, Towns ML, Quartey SM, Mirrett S, Reimer LG, Parmigiani G, et al. The clinical significance of positive blood cultures in the 1990s: a prospective comprehensive evaluation of the microbiology, epidemiology, and outcome of bacteremia and fungemia in adults. *Clin Infect Dis.* 1997;24(4):584-602.
- Beebe, James L, Elmer W. Koneman. Recovery of uncommon bacteria from blood: association with neoplastic disease. *Clin Microbiol Rev.* 1995;8(3):336-56.
- Klein JO, Dashefsky B, Norton CR, Mayer J. Selection of antimicrobial agents for treatment of neonatal sepsis. *Rev Infect Dis.* 1983;5 (Suppl 1):S55-64.
- Lawn, Joy E, Simon Cousens, Jelka Zupan. 4 million neonatal deaths: when? Where? Why? *Lancet.* 2005;365(9462):891-900.
- Elbayoumi Yasser A, Abdelraouf A. Elmanama. Epidemiology of septicemia at neonatal intensive care units in Gaza city hospitals. In: Elbayoumi, Yasser A, Abdelraouf A. Elmanama, eds. *Biological Sciences Master Program. Gaza: Islamic University;* 2011: 1-123.
- Phillips SE, Bradley JS. Bacteremia detected by lysis direct plating in a neonatal intensive care unit. *J Clin Microbiol.* 1990;28(1):1-4.
- Murthy DS, Gyaneshwari M. Blood cultures in pediatric patients: a study of clinical impact. *Indian J Med Microbiol.* 2007;25(3):220.
- Mane AK, Nagdeo NV, Thombare VR. Study of neonatal septicemia in a tertiary care hospital in rural Nagpur. *J Recent Adv Appl Sci.* 2010;25:19-24.
- Leclerc H, Mossel DA, Edberg SC, Struijk CB. Advances in the bacteriology of the coliform group: their suitability as markers of microbial water safety. *Annu Rev Microbiol.* 2001;55(1):201-34.
- Babic Maja, Andrea M. Hujer, Robert A. Bonomo. What's new in antibiotic resistance? Focus on beta-lactamases. *Drug Resistance Updates.* 2006;9(3):142-56.
- Lodise TP Jr, Patel N, Kwa A, Graves J, Furuno JP, Graffunder E, et al. Predictors of 30-day mortality among patients with *Pseudomonas aeruginosa* bloodstream infections: impact of delayed appropriate antibiotic selection. *Antimicrob Agents Chemother.* 2007;51(10):3510-5.
- Sipsas Nikolaos V, Gerald P. Bodey, Dimitrios P. Kontoyiannis. Perspectives for the management of febrile neutropenic patients with cancer in the 21st century. *Cancer.* 2005;103(6):1103-13.
- Owens Jr, Robert C. Antimicrobial stewardship: concepts and strategies in the 21st century. *Diagn Microbiol Infect Dis.* 2008;61(1):110-28.
- Tamma Pranita D, Sara E. Cosgrove. Antimicrobial stewardship. *Infect Disease Clin North Am.* 2011;25(1):245-60.
- Grimwood Keith, Scott C. Bell, Anne B. Chang. Antimicrobial treatment of non-cystic fibrosis bronchiectasis. *Expert Rev Anti-Infect Ther.* 2014;12(10):1277-96.
- French GL. Clinical impact and relevance of antibiotic resistance. *Adv Drug Deliv Rev.* 2005;57(10):1514-27.
- Bhattacharya S. Blood culture in India: a proposal for a national programme for early detection of sepsis. *Indian J Med Microbiol.* 2005;23(4):220.
- Khanal B, Harish BN, Sethuraman KR, Srinivasan S. Infective endocarditis: report of prospective study in an Indian hospital. *Trop Doct.* 2002;32:83-5.
- Murthy DS. Blood cultures in pediatric patients: a study of clinical impact. *Indian J Med Microbiol.* 2007;25(3):220-4.
- Arora U, Jaitwani J. *Acinetobacter* spp.: an emerging pathogen in neonatal septicaemia in Amritsar. *Indian J Med Microbiol.* 2003;21:66-8.
- Roy I, Jain A, Kumar M, Agarwal SK. Bacteriology of neonatal septicaemia in a tertiary care hospital of Northern India. *Indian J Med Microbiol.* 2002;20:156-9.
- Doern GV, Brueggemann AB, Dunne WM, Jenkins SG, Halsted DC, McLaughlin JC. Four-day incubation period for blood culture bottles processed with the Difco ESP blood culture system. *Clin Microbiol.* 1997;35:1290-2.
- Weems J Jr. The many faces of *Staphylococcus aureus* infection. *Postgrad Med.* 2002;3:33-41.
- Kumar Surinder, Rizvi Meher, Vidhani Shalini, Sharma VK. Changing face of septicemia and increasing drug resistance in blood isolates. *Indian J Pathol Microbiol.* 2004;47:441-6.

DOI: 10.5455/2320-6012.ijrms20150332

Cite this article as: Kante M, Muni Lakshmi P, Sreenivasulu Reddy P. Bacterial profile of blood stream infections and their antibiograms. *Int J Res Med Sci* 2015;3:698-704.