

Original Research Article

Management of idiopathic clubfoot with Ponseti technique

Sumeet Singh Charak*, Khalid Muzafar, Omeshwar Singh

Department of Orthopaedic, Government Medical College Jammu, Jammu University, India

Received: 27 June 2017

Accepted: 03 July 2017

***Correspondence:**

Dr. Sumeet Singh Charak,

E-mail: drscharak8@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Idiopathic congenital talipes equinovarus (club foot) is a complex deformity that is difficult to correct. The goal of treatment is to reduce or eliminate its four components so that the patient has a functional foot and leads a normal life.

Methods: Study have treated 20 patients with 32 idiopathic clubfoot deformities using Ponseti method. The severity was assessed by modified pirani scoring.

Results: The mean number of casts that were applied to obtain correction was 7.02 (range four to nine casts). Tenotomy was done in 30 feet. Thirty feet had good results. One patient developed recurrence of the deformity due to non-compliance of the use of orthotics.

Conclusions: The Ponseti method is a safe and effective treatment for congenital idiopathic clubfoot and radically decreases the need for corrective surgery. Non-compliance with orthotics main factor causing failure of the technique.

Keywords: Idiopathic, Ponsetti, Pirani

INTRODUCTION

Idiopathic clubfoot is a complex deformity which is difficult to correct.¹ The deformity has four components: forefoot equinus, hindfootvarus, forefootadductus and midfootcavus. The surgical treatment has multiple complications.² Treatment by Ponseti method involves serial casting, percutaneous tenotomy and application of braces. Due to conservative nature of the treatment method and fewer complications associated with it, the treatment method has become popular.³ The method has been reported to have short-term success rates approaching 90% and the long-term results have been equally impressive.¹

The objective of this study was to present the author's experience with the Ponseti technique in correcting clubfoot and to stress the importance of percutaneous tenotomy in management.

METHODS

This study was done at Government Medical College, Jammu. 32 feet in 20 children were studied. Study design was a prospective case series study. Only idiopathic cases of both gender less than 1 year were included. Syndromic, relapsed, neglected, resistant and recurrent cases were excluded.

After a, thorough clinical examination and confirmation of diagnosis, photographs of the deformity were taken (Figure 1). Severity was assessed by modified Pirani scoring system. Manipulation of foot and long leg plaster of paris application was done (Figure 2). In all patients, the cavus is corrected first by supinating the forefoot and dorsiflexing the first metatarsal. To correct the varus and adduction, the supinated foot was abducted with counter pressure applied with the thumb against the head of the talus. Four to nine long leg casts, changed weekly after

proper manipulation of the foot, were usually sufficient to obtain good correction (Figure 3).



Figure 1: Pre-casting photograph.



Figure 2: Ponseti cast applied.



Figure 3: Ponseti cast applied.

Casting was stopped when midfoot and hindfoot scores were zero with 70 degrees of abduction of the forefoot. With abduction of 70 degrees if dorsiflexion was less than 10 degrees then percutaneous tenotomy of the

Achilles tendon was performed. All cases were done in operation theatre under local anaesthesia. Patients were monitored for 1hour post operatively. A long leg cast was applied in 70 degrees of abduction and maximum available dorsiflexion immediately after tenotomy and maintained for further 3 weeks to allow healing of the tendon. After 3 weeks cast was removed (Figure 4) and Dennis Brown Splint was applied (Figure 5).



Figure 4: Post tenotomy cast removed (desired correction achieved).



Figure 5: Dennis Brown Splint.



Figure 6: 2 years follow up.

Children were reviewed every month and Pirani scores were documented. Dennis brown Splint was worn for 23 hours during first three months after casting and then at night until child is about 4 years old. Each parameter was scored according to the Modified Pirani Score. Six clinical signs are each scored 0 (normal), 0.5 (mildly abnormal) or 1 (severely abnormal). Thus, each foot can receive a Midfoot score between 0-3 and a hindfoot score between 0-3 and a total score between 0-6 (Table 1).

RESULTS

The average age of the patients was 2 months and 3 days (range, 3 days to 9 months of age). Of the 20 infants, 14 had unilateral clubfoot while as 6 had bilateral clubfoot (32 feet). Average Pirani score was 4.92 (range 3.5 to 6). The mean number of cast that were applied to obtain correction was 7.2 (range four to nine casts). More number of casts was required to obtain correction in very severe deformity. 30 of the 32 clubfeet required percutaneous tendo Achilles tenotomy to correct the residual equinus deformity. The average follow-up was for 1.5 years (range 1 to 2 years). Thirty feet were treated successfully using the Ponseti method. These patients obtained complete correction of the deformity with dorsiflexion of $>20^\circ$ and plantar flexion of $>40^\circ$ (Figure 6). In the patient with poor result, correction was obtained after tenotomy but the deformity relapsed. The family of this patient had not complied with the use of orthosis due to inconvenience.

Table 1: Components of pirani severity score.

Parameters	Mild	Moderate	Severe
Mid foot			
Curved lateral border	0	0.5	1
Medial foot crease	0	0.5	1
Talus head coverage	0	0.5	1
Hind foot			
Posterior crease	0	0.5	1
Rigid equines	0	0.5	1
Empty heel	0	0.5	1

DISCUSSION

Ponseti technique of serial manipulation and casting is being employed vastly in many of the CTEV treating centres. Ponseti and Smoley reported that by this method of manipulation surgery was avoided in 89% of cases.⁴ In Ponseti method of management the first element of correction is the cavus deformity by positioning the forefoot in proper alignment with the hindfoot.^{5,6} Cavus, which is due to the pronation of the forefoot in relation to the hindfoot requires only elevating the first ray of the forefoot to achieve a normal longitudinal arch of the foot. The forefoot is supinated not too high or too flat so that the plantar surface of the foot reveals a normal appearing arch. For subsequent correction of adducts and varus, alignment of the forefoot with the hindfoot is necessary

to give an effective abduction movement of the foot. Using the stabilized talar head as fulcrum the foot is abducted. Pronation or eversion of the foot and external rotation of the foot to correct adduction while calcaneus remains in varus are to be avoided. Eversion of the calcaneus to correct heel varus (Kites error) is not possible unless the calcaneus is abducted (i.e., laterally rotated) under the talus. Kite explained in his method of correction to abduct the forefoot against pressure at the calcaneocuboid joint which Ponseti described as Kites error.⁷ It blocked the correction of hindfoot varus and internal rotation. Ponseti technique has been reported with 92 to 98 % successful results for the treatment of idiopathic clubfoot.^{3,8-10} Study have successfully corrected thirty (93.75%) of the 32 clubfoot deformities using Ponseti method. The patient who developed recurrence of the deformity was due to non-compliance with the use of orthotics. This reason has been widely reported to be the main factor causing failure of the technique.^{1,2,8-10} Serious bleeding complications have been reported following percutaneous tendo-achilles tenotomy.¹¹ However, we did not encounter any of the complications and found it very helpful in obtaining full correction. The Ponseti method is a safe and effective treatment for congenital idiopathic clubfoot and radically decreases the need for extensive corrective surgery. Longer follow-up will decide whether we can continue to match Ponseti's results.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

1. Dobbs MB, Rudzki JR, Purcell DB, Walton T, Porter KR, Gurnett CA. Factors predictive of outcome after use of the Ponseti method for the treatment of idiopathic clubfeet. *J Bone Joint Surg.* 2004;86:22-7.
2. Colburn M, Williams M. Evaluation of the treatment of idiopathic clubfoot by using the Ponsetimethod. *J Foot Ankle Surg.* 2003;42(5):259-67.
3. Lehman WB, Mohaideen A, Madan S, Scher DM, Van Bosse HJ, Iannacone M, et al. A method for the early evaluation of the Ponseti (Iowa) technique for the treatment of idiopathic clubfoot. *J Pediatr Orthop.* 2003;12(2):133-40.
4. Ponseti IV, Smoley EN Congenital club foot: the results of treatment. *J Bone Joint Surg Am.* 1963;45:261-75.
5. Ponseti IV. Treatment of congenital club foot. *J Bone Joint Surg Am.* 1992;74:448-53.
6. Clubfoot LS. Ponseti management. 3rd ed. Global Help Publications; 2009.
7. Kite JH. Principles involved in the treatment of congenital clubfoot. *J Bone Joint Surg Am.* 2003;85:1847.

8. Herzenberg JE, Radler C, Bor N. Ponseti versus traditional methods of casting for idiopathic clubfoot. *J Pediatr Orthop.* 2002;22(4):517-21.
9. Goksan SB. Treatment of congenital clubfoot with the Ponseti method. *Acta Orthop Traumatol Turc.* 2002;36(4):281-7.
10. Morcuende JA, Dolan LA, Dietz FR. Radical reduction in the rate of extensive corrective surgery for clubfoot using the Ponseti method. *Pediatr.* 2004;113(2):376-80.
11. Dobbs MB, Gordon JE, Walton T. Bleeding complications following percutaneous tendoachillestenotomy in the treatment of clubfoot deformity. *J Pediatr Orthop.* 2004;24(4):353-7.

Cite this article as: Charak SS, Muzafar K, Singh O. Management of idiopathic clubfoot with Ponseti technique. *Int J Res Med Sci* 2017;5:3355-8.